

ORAL PRESENTATION

Open Access

# A k-population model to calculate the firing rate of neuronal networks with degree correlations


C Schmeltzer<sup>1\*</sup>, A Kihara<sup>2</sup>, I Sokolov<sup>1</sup>, S Rüdiger<sup>1</sup>

From The Twenty Third Annual Computational Neuroscience Meeting: CNS\*2014  
Québec City, Canada. 26-31 July 2014

Revealing the interplay of structure and function of the brain is one of the most intriguing topics in neuroscience.

The theory of complex networks is a promising approach to this aim, where one assumes that high cognitive processes arise as emergent properties of a network, in which many inane neurons are connected by a complex topology [1]. In this regard, we analyze analytically the emerging responses of networks with increasingly complex


connectivity. We present a mathematical theory to calculate the firing rate of a network of leaky integrate-and-fire neurons, taking into account network features such as degree distributions and degree correlations (Figure 1). Heterogeneous connectivity and degree correlations have been shown to heavily influence network function and dynamics [2,3]. Our method is to divide the neuronal network in k-populations according to the number k of



\* Correspondence: schmelt@physik.hu-berlin.de

<sup>1</sup>Institut für Theoretische Physik, Humboldt Universität, Berlin, Germany, 12489, USA

Full list of author information is available at the end of the article



afferent synaptic links that connect to the neuron. Then, the steady state firing rates for these coupled populations can be calculated self-consistently. One of our main findings is that the population heterogeneity yields substantial deviations from mean-field calculations, where one ignores the network properties [4]. Importantly, our analysis shows that networks with assortative degree correlations lead to firing patterns even for sub-threshold inputs, where an uncorrelated network would not fire and thus, to a much larger sensitivity to low stimuli (Figure 2). Using information theory we further find an optimum in assortativity, with larger levels reducing again sensitivity for signal ensembles.

#### Authors' details

<sup>1</sup>Institut für Theoretische Physik, Humboldt Universität, Berlin, Germany, 12489, USA. <sup>2</sup>Universidade Federal do ABC, Santo André, Brazil.

Published: 21 July 2014

#### References

1. Sporns O: *Networks of the Brain*. MIT press; 2011.
2. Newman M: *The structure and function of complex networks*. *SIAM review* 2003, **45**(2):167-256.
3. Schmeltzer C, *et al*: Percolation of spatially constrained Erdős-Rényi networks with degree correlations. *Phys Rev E Stat Nonlin Soft Matter Phys*. 2014, **89**(1):012116.
4. Deco G, *et al*: The dynamic brain: from spiking neurons to neural masses and cortical fields. *PLoS Computational Biology* 2008, **4**(8):e1000092.

doi:10.1186/1471-2202-15-S1-O14

**Cite this article as:** Schmeltzer *et al*: A k-population model to calculate the firing rate of neuronal networks with degree correlations. *BMC Neuroscience* 2014 **15**(Suppl 1):O14.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

