

Research Article

Fucoxanthin Exerts Cytoprotective Effects against Hydrogen Peroxide-induced Oxidative Damage in L02 Cells

Xia Wang ^{1,2}, Yan-jun Cui ³, Jia Qi ³, Min-min Zhu ³, Tian-liang Zhang ⁴,
Min Cheng ⁵, Shun-mei Liu ^{3,4}, and Guang-ce Wang ⁶

¹School of Public Health and Management, Weifang Medical University, Weifang 261053, China

²Collaborative Innovation Center of Prediction and Governance of Major Social Risks in Shandong, Weifang Medical University, Weifang 261053, China

³School of Bioscience and Technology, Weifang Medical University, Weifang 261053, China

⁴Experimental Center for Medical Research, Weifang Medical University, Weifang 261053, China

⁵School of Clinical Medicine, Weifang Medical University, Weifang, 261053, China

⁶Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

Correspondence should be addressed to Shun-mei Liu; liushm@wfmc.edu.cn and Guang-ce Wang; gcwang@qdio.ac.cn

Received 11 September 2018; Revised 20 October 2018; Accepted 28 October 2018; Published 15 November 2018

Academic Editor: Paul W. Doetsch

Copyright © 2018 Xia Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Several previous studies have demonstrated the excellent antioxidant activity of fucoxanthin against oxidative stress which is closely related to the pathogenesis of liver diseases. The present work was to investigate whether fucoxanthin could protect human hepatic L02 cells against hydrogen peroxide- (H_2O_2 -) induced oxidative damage. Its effects on H_2O_2 -induced cell viability, lactate dehydrogenase (LDH) leakage, intracellular reduced glutathione, and reactive oxygen species (ROS) contents, along with mRNA and protein relative levels of the cytoprotective genes including Nrf2, HO-1, and NQO1, were investigated. The results showed that fucoxanthin could upregulate the mRNA and protein levels of the cytoprotective genes and promote the nuclear translocation of Nrf2, which could be inhibited by the PI3K inhibitor of LY294002. Pretreatment of fucoxanthin resulted in decreased LDH leakage and intracellular ROS content but enhanced intracellular reduced glutathione. Interestingly, pretreatment using fucoxanthin protected against the oxidative damage in a nonconcentration-dependent manner, with fucoxanthin of 5 μM demonstrating the optimal effects. The results suggest that fucoxanthin exerts cytoprotective effects against H_2O_2 -induced oxidative damage in L02 cells, which may be through the PI3K-dependent activation of Nrf2 signaling.

1. Introduction

Reactive oxygen species (ROS), including free radical such as hydroxyl radical and non-free radical such as hydrogen peroxide (H_2O_2), are highly reactive byproducts derived from normal cell metabolism, especially from that of mitochondria [1]. Generally, appropriate generation of ROS might be essential for many cellular functions such as phagocytes killing and bacterial ingestion but detrimental to living organisms in the case of overproduction [2]. Oxidative stress represents an imbalance between pro-oxidants and antioxidants. The antioxidative status was found inversely correlated with the occurrence of numerous human diseases [3, 4]. Hepatic cells are rich in mitochondria and prone to generate ROS [5].

Role of oxidative damage in the pathogenesis of various liver injuries has been confirmed. It has been suggested that the overproduction of H_2O_2 contributes to the pathogenesis of many liver diseases such as hepatitis C virus infection, cholestasis and Wilson's disease etc [6]. Enzymatic antioxidants (such as glutathione (GSH) or glutathione-related enzyme system) and non-enzymatic ones are the two major systems to control ROS generation and counteract the oxidative damage [7]. Therefore, it has become an interesting and urgent topic for researchers to find excellent antioxidants for the prevention of liver diseases.

Many synthetic antioxidants, such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA), have been used to slow down the processes of oxidation and

peroxidation in food and pharmaceutical industries. However, their utilizations have been severely restricted due to their potential toxicity to human health because of their carcinogenic effects [8]. And BHT, when applied together with PG, resulted in joint pathology and liver enlargement [9]. Moreover, continuous use of some synthetic phenolic antioxidants exerted teratogenic and carcinogenic effects in laboratory animals and primates [10]. Therefore, natural antioxidants, with minimal side or toxic effects, have been drawing increasing attention from researchers in the past decades in viewing of consumers' preferences and concerns about health.

Carotenoids, natural pigments found in plants, algae, animals, etc., possess the antioxidant ability to protect cells and tissues from detrimental effects caused by ROS [11]. The role of carotenoids to protect against diseases caused by oxidants has been confirmed. Fucoxanthin, mainly existing in brown seaweed, accounts for more than 10% of total carotenoids production in nature [12]. It was speculated that its unique allenic bond and 5, 6-monoepoxide might be crucial for radical scavenging and protecting against cell damages induced by exposure to H_2O_2 or UV-B, with its high antioxidant activity probably attributed to the allenic bond [13].

Nuclear factor-erythroid 2-related factor 2 (Nrf2 or Nfe2l2) is a transcription factor sensitive to cellular stresses such as oxidative stress. Under normal physiological conditions, Nrf2 is mainly in cytoplasm and transcriptionally inactive due to proteasomal degradation mediated by Kelch-like ECH-associated protein 1 (Keap1) which acts as an inhibitor. Under oxidative stress, the sulfhydryl groups of Keap1 can be easily oxidized and its binding affinity to Nrf2 decreases, resulting in the translocation of Nrf2 from the cytoplasm to the nucleus and subsequent binding to antioxidant response elements (ARE), which could enhance the expressions of many stress-induced genes such as HO-1 and NQO1 [14]. It has been reported previously that several dietary compounds could protect cells from oxidative damage through Nrf2-ARE pathway [15].

Several studies have reported the antioxidant effect of fucoxanthin, mainly focusing on the scavenging activity against different free radicals (such as DPPH, ABTS, hydroxyl radical, hydrogen peroxide, and superoxide anion) and the ability to quench singlet oxygen [16–19] using cell-free system or its inhibitory effects on ROS production, DNA damage and cell apoptosis induced by H_2O_2 , UV, and other factors using cell models [20–23]. H_2O_2 , widely used to induce oxidative stress in in vitro models, could easily cross the cellular membranes and generate highly toxic hydroxyl radicals which could react with macromolecules such as DNA and proteins, thus leading to cellular damages [24]. As an essential organ responsible for detoxification and biochemical metabolisms, liver is susceptible to oxidative damage. Moreover, oxidative stress in normal rat hepatic cells in vitro induced by H_2O_2 was similar to that occurred in intact liver [25]. The human L02 cells, commonly accepted in vitro experimental model and used for exploring the pathogenesis of liver diseases [26], were also employed in the present work. Liu et al. suggested the participation of Nrf2-ARE pathway regarding

the antioxidant activity of fucoxanthin in murine hepatic BNL CL2 cells [27]. In consideration of the high incidence of liver diseases in China, we select fucoxanthin due to its high antioxidant activity to investigate its cytoprotective effects against H_2O_2 -induced oxidative damage in hepatic L02 cells and whether it works through the Nrf2-ARE pathway.

2. Materials and Methods

2.1. Cell Culture and Treatments. Normal human hepatic cell line of L02, purchased from the Procell Life Science & Technology Co., Ltd. (Wuhan, Hubei Province, China), was cultured in RPMI 1640 medium (Solarbio, Beijing, China) containing 10% (v/v) fetal bovine serum (FBS), penicillin of 100 U/mL, and streptomycin of 100 μ g/mL in a humidified incubator, with CO_2 as 5% and temperature as 37°C. Cells were transferred to appropriated multiwell plates and used for subsequent experiments when the confluence reached about 70%. Fucoxanthin (Sigma-Aldrich, Steinheim, Germany) was dissolved in DMSO and diluted with culture medium to various concentrations for following tests. Cells from experimental groups were incubated with fucoxanthin for 2 h prior to the oxidative stress induced by H_2O_2 .

2.2. Cell Viability. Cell viability of L02 was evaluated through MTT assay. Briefly, cells were seeded into 96-well plates, with cell concentration as 1×10^5 cells/well, and cultured in serum-free medium for 24 h. Then the cells from experimental groups were pretreated with fucoxanthin at different concentrations (final concentration as 1, 5, 10, or 20 μ M, respectively) or vitamin E (final concentration as 50 μ M) and cultured for 2 h, followed by exposure to H_2O_2 (final concentration as 200 μ M). After incubation for 24 h, MTT (final concentration as 0.5 mg/mL) was added into each well and incubation for another 4 h at 37°C was performed. Subsequently, the plate was centrifuged at a speed of 800 g for 5 min and the supernatant was discarded. Then the formazan crystals formed in each well were dissolved using 100 μ L DMSO and the absorbance was measured using a microplate reader (PowerWave XS, Bio-Tek, Winooski, VT, USA) at a wavelength of 540 nm. The relative cell viability was calculated by comparison with the absorbance of untreated control group.

2.3. Intracellular ROS Measurement. The intracellular ROS level was evaluated by the 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA) method. Cells (1×10^5 cells/mL) were seeded into 96-well plates and cultured for 24 h. Then cells from experimental groups were subjected to various aforementioned concentrations of fucoxanthin or vitamin E (final concentration as 50 μ M) and cultured for 24 h, followed by treatment of 200 μ M H_2O_2 (final concentration) for 30 min at 37°C. After being washed with PBS, cells were treated with 10 μ M DCFH-DA for 20 min at 37°C in a humidified incubator. After the redundant DCFH-DA was removed, cells were washed carefully using PBS three times and the fluorescence intensities of DCF were determined using a fluorescence microplate reader (VICTRO3, PerkinElmer, Waltham, MA, USA), with excitation wavelength at 485 nm and emission

wavelength at 535 nm. And the representative images concerning the levels of intracellular ROS were also captured using an inverted fluorescence microscope (DMI400B; Leica, Wetzlar, Germany) and the fluorescence intensities which reflects the contents of ROS were analyzed using IPP software (Version 6.0, Media Cybernetics, USA) and normalized to that of the control group.

2.4. Lactate Dehydrogenase (LDH) Leakage Assay. The LDH leaked in the culture was evaluated using a LDH activity assay kit (Beyotime Institute of Biotechnology, Shanghai, China) according to the supplier's instructions. Briefly, cells (1×10^5 cells/mL) were seeded into 96-well plates and cultured for 24 h. Then the cells from experimental groups were treated with various aforementioned concentrations of fucoxanthin or vitamin E (final concentration as $50 \mu\text{M}$) and cultured for 2 h, followed by treatment of $200 \mu\text{M}$ H_2O_2 (final concentration) for another 24 h at 37°C in a humidified incubator. The culture was subsequently mixed with the relevant kit reaction solution and incubated at 25°C . Finally, the LDH activity in culture was determined based on the absorbance at 450 nm using a microplate reader (PowerWave XS, Bio-Tek, Winooski, VT, USA). The LDH leakage rate was calculated by comparing with the maximum LDH activity control group.

2.5. Detection of Intracellular Reduced GSH. The experimental grouping and treatment were conducted as described in method 2.4. After incubation, cells were made into cell suspensions by trypsin digestion and washed with PBS three times, followed by homogenate using an ultrasonic cell disruption system (JY92-IIN, Xinzhi, Ningbo, China). Then the intracellular glutathione was measured using a glutathione determination kit (A006-2, Jiancheng, Nanjing, China) in accordance with the supplier's instructions. The intracellular glutathione was evaluated based on the absorbance at 420 nm using a fluorescence microplate reader (SpectraMax M5, MD, CA, USA), with excitation wavelength at 350 nm.

2.6. Real-Time Quantitative PCR. One additional group named L was added for real-time PCR and the following western blot along with the immunofluorescent assay. As to group L, cells were firstly incubated with fucoxanthin of $5 \mu\text{M}$ for 1 h, followed by addition of LY294002 (a specific inhibitor of PI3K, final concentration as $20 \mu\text{M}$) and H_2O_2 of $200 \mu\text{M}$ (final concentration). After being treated as aforementioned in method 2.4 and washed three times using PBS, cells derived from each group were collected and the total RNA was extracted using Trizol reagent (Invitrogen, Carlsbad, CA, USA), followed by the synthesis of cDNA by a commercial first strand cDNA synthesis kit (Takara, Otsu, Japan). The relevant cDNA was kept at -70°C before use. The primers, designed using software of Primer5, were synthesized by Takara. Then the triplicate cDNA samples, included in a reaction system of totally $20 \mu\text{L}$ mainly containing approximately 50 ng cDNA, $10 \mu\text{M}$ primer for each and $10 \mu\text{L}$ premix (Bio-Rad Hercules, CA, USA), were analyzed via a real-time quantitative PCR system (IQ5, Biorad, CA, USA). The quantitative PCR reaction conditions were as following: 95°C

TABLE 1: Primers used for real-time quantitative PCR.

Target	Primer Sequence (5'-3')
Nrf2	F: TCCAGTCAGAAACCAGTGGAT
	R: GAATGTCTGCGCCAAAAGCTG
HO-1	F: CCAGCAACAAAGTGCAAGAT
	R: GTGTAAGGACCCATCGGAGA
NQO1	F: GAAGAGCACTGATCGTACTGGC
	R: GGATACTGAAAGTTCCGCAGGG
β -actin	F: ATTGCCGACAGGATGCAGA
	R: GAGTACTTGCCTCAGGAGGA

for 2 min, followed by 40 cycles of 90°C for 10 s and 60°C for 40 s. The positive and negative controls were utilized to keep the accuracy of the present quantitative PCR. $2^{-\Delta\Delta\text{Ct}}$ method was used to calculate the relative mRNA levels of target genes, with β -actin as the reference gene. And the sequences of primers used for real-time quantitative PCR in the present study were presented in Table 1.

2.7. Western Blot. After being treated as described in method 2.6 and washed using PBS three times, cells derived from each group were collected and lysed using RIPA buffer (Beyotime Institute of Biotechnology, Shanghai, China) containing a protease inhibitor of 1 mM PMSF according to the supplier's instructions. Protein concentrations were determined using BCA method. Equal amounts of proteins were loaded and isolated using a sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and subsequently transferred onto a PVDF membrane. After being blocked with 5% skim milk, membranes were incubated with primary antibody (Nrf2, 1:2000 dilution; HO-1, 1:10000 dilution; NQO1, 1:20000 dilution; Abcam) and goat anti-rabbit secondary antibody were conjugated to horseradish peroxidase (HRP) (1:200 dilution) (Beyotime Institute of Biotechnology, Shanghai, China). β -actin was used as internal reference.

2.8. Immunofluorescent Assay. After being treated as described in method 2.6, cells were washed using 0.5% PBST and fixed with paraformaldehyde of 4% for 10 min, followed by being washed using 0.5% PBST three times and blocked using immunol staining blocking buffer (P0102, Beyotime Institute of Biotechnology, Shanghai, China) for 1.2 h. Then the cells were incubated with Nrf2 rabbit anti-human primary antibody (1:100 dilution) at 4°C overnight, followed by being kept at room temperature for 1 h and washed using 0.5% PBST three times. The cells were subsequently incubated with FITC labeled goat anti-rabbit secondary antibody (1:200 dilution) at 37°C in a humidified incubator for 1.5 h, followed by being washed using 0.5% PBST three times. Finally, cells were stained with Hoechst 33258 of $20 \mu\text{M}$ for 5 min, followed by being washed using 0.5% PBST three times and adding antifade polyvinylpyrrolidone mounting medium (P0123, Beyotime Institute of Biotechnology, Shanghai, China). Then the slices were observed via a fluorescence microscope (Leica DMI400B).




FIGURE 1: Effects of H₂O₂ treatment on the viability of L02 cells. Data were presented as means \pm SD, n=3. ##: $P < 0.01$ (compared with control).

2.9. *Statistical Analyses.* All experiments in the present work were replicated at least three times. All data were presented as means \pm SD and analyzed by one-way ANOVA and *t*-test. All the statistical analyses were performed using SPSS software (version 19.0; IBM, Chicago, IL, USA). And it was considered as statistically significant in the case of *P* value smaller than 0.05.

3. Results

3.1. *Effects of H₂O₂ Treatment on the Viability of L02 Cells.* L02 cells were treated with H₂O₂ at different concentrations (100, 200, 400, 600, 800, 1000, 1200, and 1600 μ M, respectively) and the appropriate concentration used to induce oxidative damage was evaluated by MTT assay. As is shown in Figure 1, the cell viability decreased significantly when treatment concentration of H₂O₂ was equal to or more than 100 μ M (compared with control which is set as 100%, $P < 0.01$). In the presence of 200 μ M H₂O₂, the cell viability demonstrated a moderate decrease. Therefore, treating the L02 cells with H₂O₂ at a final concentration of 200 μ M was selected for subsequent experiments.

3.2. *Effects of Fucoxanthin on the Viability of H₂O₂-Treated L02 Cells.* Effects of fucoxanthin on the viability of H₂O₂-treated L02 cells were also evaluated by MTT. The results were shown in Figure 2 where it could be learned that the viability increased after being treated by 50 μ M vitamin E (VE) or fucoxanthin (1, 5, 10, and 20 μ M, respectively), while there was no significant difference among the groups ($P > 0.05$).

3.3. *Effects of Fucoxanthin on H₂O₂-Induced L02 Cellular LDH Leakage.* The leakage level of maximum LDH leakage group was set as 100% and results were presented in Figure 3. The LDH leakage rate of control group was $16.71 \pm 4.48\%$, with model group as $28.45 \pm 5.13\%$ (compared with control, $P < 0.01$). Compared with model group, VE-treated group exhibited a leakage rate of $18.70 \pm 4.98\%$ ($P < 0.05$), with F5 group demonstrating the lowest LDH leakage ($P < 0.05$).




FIGURE 2: Effects of fucoxanthin on the viability of H₂O₂-treated L02 cells. Con: control; Mod: H₂O₂ model group; F1: 1 μ M fucoxanthin + H₂O₂; F5: 5 μ M fucoxanthin + H₂O₂; F10: 10 μ M fucoxanthin + H₂O₂; F20: 20 μ M fucoxanthin + H₂O₂; VE: 50 μ M vitamin E + H₂O₂. Data were presented as means \pm SD, n=3.




FIGURE 3: Effects of fucoxanthin on H₂O₂-induced L02 cellular LDH leakage. Con: control; Mod: H₂O₂ model group; F1: 1 μ M fucoxanthin + H₂O₂; F5: 5 μ M fucoxanthin + H₂O₂; F10: 10 μ M fucoxanthin + H₂O₂; F20: 20 μ M fucoxanthin + H₂O₂; VE: 50 μ M vitamin E + H₂O₂. Data was shown as percentage of maximum LDH release group and presented as means \pm SD, n=3. ##: $P < 0.01$, compared with control; *: $P < 0.05$, compared with model group.

3.4. *Effects of Fucoxanthin on Intracellular GSH Content in H₂O₂-Treated L02 Cells.* As shown in Figure 4, the GSH content in control group was set as 100% and that of model group dropped to $62.29 \pm 6.92\%$ (compared with control, $P < 0.01$). Treatment of VE (50 μ M) resulted in the GSH content of $106.94 \pm 5.70\%$ (compared with model group, $P < 0.01$). Pretreatment with aforementioned various concentrations of fucoxanthin resulted in the GSH content of $110.69 \pm 4.39\%$, $120.98 \pm 6.72\%$, $103.97 \pm 7.12\%$, and $96.05 \pm 5.59\%$, respectively (compared with model group, $P < 0.01$).

3.5. *Effects of Fucoxanthin on Intracellular ROS Contents in H₂O₂-Treated L02 Cells.* The intracellular ROS contents were presented as a percentage of control. As shown in Figure 5, ROS content of model group increased to $165.38 \pm 16.9\%$, compared with control ($P < 0.01$). Treatment of VE (50 μ M) resulted in ROS content of $103.66 \pm 15.60\%$ (compared with model group, $P < 0.01$). After being pretreated with various aforementioned concentrations of fucoxanthin, the ROS content dropped to $132.60 \pm 16.55\%$, $96.61 \pm 19.72\%$, $105.48 \pm$




FIGURE 4: Effects of fucoxanthin on intracellular GSH content in H_2O_2 -treated L02 cells. Con: control; Mod: H_2O_2 model group; F1: 1 μM fucoxanthin + H_2O_2 ; F5: 5 μM fucoxanthin + H_2O_2 ; F10: 10 μM fucoxanthin + H_2O_2 ; F20: 20 μM fucoxanthin + H_2O_2 ; VE: 50 μM vitamin E + H_2O_2 . Data was shown as a percentage of the maximum LDH release group and presented as means \pm SD, n=3. ##: $P < 0.01$, compared with control; **: $P < 0.01$, compared with model group.




FIGURE 5: Effects of fucoxanthin on intracellular ROS contents in H_2O_2 -treated L02 cells. Con: control; Mod: H_2O_2 model group; F1: 1 μM fucoxanthin + H_2O_2 ; F5: 5 μM fucoxanthin + H_2O_2 ; F10: 10 μM fucoxanthin + H_2O_2 ; F20: 20 μM fucoxanthin + H_2O_2 ; VE: 50 μM vitamin E + H_2O_2 . The data was shown as means \pm SD, n=3. ##: $P < 0.01$, compared with control; *: $P < 0.05$, **: $P < 0.01$, compared with model group.

13.65%, and $110.65 \pm 11.00\%$, respectively (in comparison with model group, $P < 0.01$). The reduced fluorescence intensities, as is shown in Figure 6(h), also suggested the intracellular ROS scavenge capacity of fucoxanthin in H_2O_2 -treated L02 cells.

3.6. Effects of Fucoxanthin on Nuclear Translocation of Nrf2 in H_2O_2 -Treated L02 Cells. Effects of fucoxanthin on nuclear translocation of Nrf2 in H_2O_2 -treated L02 cells were shown in Figure 7. The expression of Nrf2 was shown in green fluorescence and blue fluorescence exhibited the nucleus stained by Hoechst 33258. Cyan represents the occurrence of nuclear transposition. It could be learned that fucoxanthin could enhance the expression of Nrf2 and promote the occurrence of Nrf2 nuclear translocation. LY294002 could inhibit the occurrence of Nrf2 nuclear translocation and reduce the expression of Nrf2.

3.7. Effects of Fucoxanthin on mRNA Relative Levels of Nrf2 Signaling Pathway-Related Proteins. Compared with control group, mRNA relative levels of Nrf2, HO-1 and NQO1 from model group dropped significantly to 0.64 ± 0.06 ($P < 0.01$), 0.85 ± 0.07 ($P < 0.05$), and 0.83 ± 0.06 ($P < 0.01$), respectively (Figures 8–10). After being pretreated with fucoxanthin of 1 or 5 μM , mRNA relative levels of Nrf2, HO-1, and NQO1 were enhanced to 0.81 ± 0.06 and 0.98 ± 0.07 ($P < 0.01$), 1.15 ± 0.11 and 1.61 ± 0.07 ($P < 0.01$), along with 0.95 ± 0.04 ($P < 0.05$), and 1.03 ± 0.08 ($P < 0.01$), respectively, in comparison with model group.

After being treated by LY294002, the mRNA relative levels of Nrf2, HO-1 and NQO1 decreased to 0.65 ± 0.07 ($P < 0.01$), 0.95 ± 0.05 ($P < 0.01$), and 0.91 ± 0.03 ($P < 0.05$), respectively, compared with group F5 (Figures 8–10).

3.8. Effects of Fucoxanthin on Expressions of Nrf2 Signaling Pathway-Related Proteins. As is shown in Figure 11, the expression of Nrf2 in model group dropped significantly when compared with control ($P < 0.05$). After being treated with various aforementioned concentrations of fucoxanthin, the relative protein levels of Nrf2 were enhanced to $1.01 + 0.04$ ($P < 0.05$), $1.14 + 0.05$ ($P < 0.01$), and $1.02 + 0.05$ ($P < 0.05$), respectively, compared with model group. The relative protein level of HO-1, dropping significantly in model group when compared with control ($P < 0.05$), was presented in Figure 12. After being treated with fucoxanthin of 1 or 5 μM , the relative protein level of HO-1 was enhanced to 1.06 ± 0.04 ($P < 0.01$) and 1.23 ± 0.07 ($P < 0.01$), respectively, compared with model group. As to the relative protein level of NQO1, significant difference was only found between group F5 and model group ($P < 0.01$) (Figure 13).

After being treated by LY294002, the relative protein levels of Nrf2, HO-1 and NQO1 decreased to 0.87 ± 0.05 ($P < 0.01$), 0.86 ± 0.07 ($P < 0.01$) and 0.70 ± 0.13 ($P < 0.01$), respectively, compared with group F5.

4. Discussion

Although H_2O_2 physiologically exists in living cells where it can act as a cellular signal transducer below the concentration of 1 μM , at higher concentrations it might result in adverse effects such as growth arrest or cell death caused by the derived oxidative stress [28]. Cell viability can be evaluated by MTT assay and in living cells MTT could be transformed by mitochondria into formazan whose amount is positively related with living cell numbers [29]. Based on cell viability, H_2O_2 at a final concentration of 200 μM was selected for subsequent treatment on L02 cells in the present work. The results showed that fucoxanthin pretreatment for 2 h prior to treatment of H_2O_2 could increase the cell viability compared with model group, suggesting the protecting effects of fucoxanthin against the cellular damages induced by H_2O_2 . Apoptosis of hepatic cells often occur in the case of liver damage [30]; thus the pretreatment of fucoxanthin might suppress the L02 cell death induced by H_2O_2 , though there was no significant difference between the experimental groups and model group (Figure 2) in cell viability.




FIGURE 6: Effects of fucoxanthin on intracellular ROS contents in H_2O_2 -treated L02 cells ($\times 200$ magnification). (a): control; (b): H_2O_2 model group; (c): $1 \mu M$ fucoxanthin + H_2O_2 ; (d): $5 \mu M$ fucoxanthin + H_2O_2 ; (e): $10 \mu M$ fucoxanthin + H_2O_2 ; (f): $20 \mu M$ fucoxanthin + H_2O_2 ; (g): $50 \mu M$ vitamin E + H_2O_2 ; (h): Analysis of intracellular ROS based on fluorescence microscope. ##: $P < 0.01$, compared with control; *: $P < 0.05$, **: $P < 0.01$, compared with model group.




FIGURE 7: Effects of fucoxanthin on nuclear translocation of Nrf2 in H_2O_2 -treated L02 cells ($\times 200$ magnification). Con: control; Mod: H_2O_2 model group; F1: $1 \mu M$ fucoxanthin + H_2O_2 ; F5: $5 \mu M$ fucoxanthin + H_2O_2 ; F10: $10 \mu M$ fucoxanthin + H_2O_2 ; F20: $20 \mu M$ fucoxanthin + H_2O_2 ; L: H_2O_2 + $5 \mu M$ fucoxanthin + LY294002; VE: $50 \mu M$ vitamin E + H_2O_2 ; Nucleus stained with Hoechst appears as blue fluorescence in the first column and stained Nrf2 appears as green fluorescence in the second column, with the merged as the synthesis of them both.




FIGURE 8: Effects of fucoxanthin on mRNA relative level of Nrf2 in H₂O₂-treated L02 cells. Con: control; Mod: H₂O₂ model group; F1: 1 μM fucoxanthin + H₂O₂; F5: 5 μM fucoxanthin+ H₂O₂; F10: 10 μM fucoxanthin +H₂O₂; F20: 20 μM fucoxanthin +H₂O₂; L: 5 μM fucoxanthin + H₂O₂ + LY294002; Data was presented as means ± SD, n=3. ##: P < 0.01, compared with control; **: P < 0.01, compared with model group; &&: P < 0.01, compared with group F5.




FIGURE 9: Effects of fucoxanthin on mRNA relative level of HO-1 in H₂O₂-treated L02 cells. Con: control; Mod: H₂O₂ model group; F1: 1 μM fucoxanthin + H₂O₂; F5: 5 μM fucoxanthin+ H₂O₂; F10: 10 μM fucoxanthin +H₂O₂; F20: 20 μM fucoxanthin +H₂O₂; L: 5 μM fucoxanthin + H₂O₂ + LY294002; Data was presented as means ± SD, n=3. #: P < 0.05, compared with control; **: P < 0.01, compared with model group; &&: P < 0.01, compared with group F5.

LDH, a stable enzyme in the cytoplasm of all cells, could be released promptly into extracellular environment once the cell membrane is damaged [31]. Thus it may act as an important indicator of cell membrane damage extent. Pretreatment of 5 μM fucoxanthin could significantly reduce the H₂O₂-induced intracellular LDH leakage, with effects comparable to VE of 50 μM, suggesting a role of fucoxanthin in protecting the integrity of cell membranes and counteracting the H₂O₂-induced oxidative damage. Similarly, pretreatment of fucoxanthin also led to decline of intracellular ROS. The inhibition of ROS generation by fucoxanthin might be due to its two hydroxyl groups in the ring structure and the number of hydroxyl groups has been reported correlated with inhibited ROS production [32]. GSH, an abundant thiol in liver, could scavenge free radicals or serve as a substrate to glutathione peroxidase (GSHPx) and glutathione s-transferase (GST) to detoxify the H₂O₂-induced damages [33]. The pretreatment of fucoxanthin could significantly reverse the decline of GSH




FIGURE 10: Effects of fucoxanthin on mRNA relative level of NQO1 in H₂O₂-treated L02 cells. Con: control; Mod: H₂O₂ model group; F1: 1 μM fucoxanthin + H₂O₂; F5: 5 μM fucoxanthin+ H₂O₂; F10: 10 μM fucoxanthin +H₂O₂; F20: 20 μM fucoxanthin +H₂O₂; L: 5 μM fucoxanthin + H₂O₂ + LY294002; Data was presented as means ± SD, n=3. ##: P < 0.01, compared with control; *: P < 0.05, compared with model group; **: P < 0.01, compared with model group; &: P < 0.05, compared with group F5.




FIGURE 11: Effects of fucoxanthin on expression of Nrf2 in H₂O₂-treated L02 cells. Con: control; Mod: H₂O₂ model group; F1: 1 μM fucoxanthin + H₂O₂; F5: 5 μM fucoxanthin+ H₂O₂; F10: 10 μM fucoxanthin +H₂O₂; F20: 20 μM fucoxanthin +H₂O₂; L: 5 μM fucoxanthin + H₂O₂ + LY294002; Data was presented as means ± SD, n=3. #: P < 0.05, compared with control; *: P < 0.05, compared with model group; **: P < 0.01, compared with model group; &&: P < 0.01, compared with group F5.

induced by H₂O₂, consistent with what was reported by Zheng J et al. [34]. As to the effects on GSH and intracellular ROS levels, fucoxanthin performed better than VE of 50 μM did. The excellent effects of fucoxanthin aforementioned might be due to its higher antioxidant activity which is based on its allenic bond, epoxide group and hydroxyl group [35].

Lowe et al. found that β-carotene protected HT29 cells from H₂O₂-induced damaging effects only at low concentrations of about 2-3 μM instead of higher concentration of 4-10 μM where the protection ability was rapidly lost [36], similar to what was found in H₂O₂-induced DNA damage in HepG2




FIGURE 12: Effects of fucoxanthin on expression of HO-1 in H_2O_2 -treated L02 cells. Con: control; Mod: H_2O_2 model group; F1: 1 μM fucoxanthin + H_2O_2 ; F5: 5 μM fucoxanthin + H_2O_2 ; F10: 10 μM fucoxanthin + H_2O_2 ; F20: 20 μM fucoxanthin + H_2O_2 ; L: 5 μM fucoxanthin + H_2O_2 + LY294002; Data was presented as means \pm SD, n=3. #: $P < 0.05$, compared with control; **: $P < 0.01$, compared with model group; &&: $P < 0.01$, compared with group F5.




FIGURE 13: Effects of fucoxanthin on expression of NQO1 in H_2O_2 -treated L02 cells. Con: control; Mod: H_2O_2 model group; F1: 1 μM fucoxanthin + H_2O_2 ; F5: 5 μM fucoxanthin + H_2O_2 ; F10: 10 μM fucoxanthin + H_2O_2 ; F20: 20 μM fucoxanthin + H_2O_2 ; L: 5 μM fucoxanthin + H_2O_2 + LY294002; Data was presented as means \pm SD, n=3. **: $P < 0.01$, compared with model group; &&: $P < 0.01$, compared with group F5.

cells by Woods et al. [37]. In the present work, fucoxanthin at concentrations higher than 5 μM also presented lower effects against oxidative damage induced by H_2O_2 in a similar manner. β -Carotene may act as a pro-oxidant at higher concentrations. In vitro, whether β -carotene plays as a pro-oxidant or oxidant depends on oxygen tension and

its concentration [38]. However, β -carotene exhibits pro-oxidant effects at 2.5 μM in LS-174 cells, presumably due to the capability difference of cell to incorporate the carotenoid [39]. Fucoxanthin demonstrated higher DPPH radical scavenging capacity than β -carotene [40] but lower reducing power than ascorbic acid [41]. Superior antioxidant activity of fucoxanthin than β -carotene, along with slightly less potency than ascorbic acid, was also reported and ascorbic acid of 1 μM significantly reduced the H_2O_2 -induced sister chromatid exchanges [35]. Fucoxanthin of 5 μM demonstrated excellent cytoprotective effects against the H_2O_2 -induced oxidative damage in the present work. Hence, based on the aforementioned data, fucoxanthin possesses the prominent antioxidant ability to protect against the cellular damages induced by H_2O_2 and might be a potential therapeutic agent for treating or preventing diseases related to oxidative stress.

Since large quantities of ROS are produced continuously throughout life, antioxidant mechanisms are essential for cells to maintain redox homeostasis [42]. There are many key regulators, such as nuclear factor- (erythroid-derived 2-) like 2 (Nrf2), to protect against oxidative damage. Nrf2 plays a vital role in maintaining cellular redox homeostasis by activating a variety of cytoprotective enzymes including quinone oxidoreductase 1 (NQO1) and heme oxygenase-1 (HO-1) to attenuate liver injury [43]. Under stressed conditions, Nrf2 leaves from its inhibitor of CNC homology- (ECH-) associated protein 1 (Keap1) and is then translocated from the cytoplasm to the nucleus where it could transcriptionally activate its targeted cytoprotective enzymes by binding to the antioxidant response element (ARE) located in their promoter regions [44]. Thus Nrf2 acts as a molecular switch to activate the cytoprotective enzymes defending against oxidative stress. The activation of Nrf2 by some dietary phytochemicals demonstrated chemopreventive effects to suppress oxidative stress [13, 45]. HO-1 has been reported to exert its cytoprotective effects via reducing intracellular pro-oxidant levels while enhancing the levels of carbon monoxide and bilirubin [46]. And the elevated carbon monoxide and bilirubin contribute to antiapoptotic effects and fighting against cellular injury induced by free radical, respectively [47, 48]. NQO1 can be easily induced by various stresses including oxidative stress [49]. The induced high levels of it can be mediated by Nrf2 or Ah receptor [50]. In the present study, pretreatment of 5 μM fucoxanthin prior to exposure to H_2O_2 of 200 μM enhanced the mRNA and protein relative levels of Nrf2, HO-1 and NQO1, consistent with what was reported by Liu et al. [27]. And the nuclear translocation of Nrf2 was also observed (Figure 7). It could be speculated that fucoxanthin might activate the nuclear translocation of Nrf2 and then upregulate the expressions of its targeted genes of HO-1 and NQO1, thereby playing the role of suppressing the oxidative damage through the Nrf2/ARE signaling. Thus it seems that Nrf2 plays a vital role in protecting against the H_2O_2 -induced liver injuries [51].

LY294002 could selectively inhibit the PI3K nexus by competitively and reversibly acting on the ATP-binding site of PI3K [52]. The participations of PI3K pathway in cell survival, proliferation, migration, and metabolism have been reported [53]. Since Nrf2 signaling pathway targeting

the genes of HO-1 and NQO1 may be mediated by PI3K pathway [54], protein kinase C (PKC) pathway [55], c-jun N-terminal kinase (JNK) pathway [56], or ERK pathway [57], etc., LY294002 was used in group L in the present work to investigate whether the Nrf2 signaling pathway involving the antioxidant activity of fucoxanthin was mediated by PI3K pathway. The results of group L exhibited that the mRNA and protein relative levels of Nrf2, HO-1 and NQO1 were significantly downregulated, weakening the antioxidant activity of fucoxanthin. Moreover, the nuclear translocation of Nrf2 was also inhibited by LY294002. Hence, the present work suggests that fucoxanthin may exert its antioxidant effects in L02 cells against H₂O₂-induced oxidative damage through PI3K-dependent activation of Nrf2 signaling, while the exact mechanism of action still needs further study.

It must be noted that pretreatment by 5 μ M fucoxanthin in the present work demonstrated the optimal effects on the parameters such as LDH release, GSH content, and intracellular ROS content, along with the induction of Nrf2, HO-1, and NQO1. Interestingly, it does not work in a concentration-dependent manner, inconsistent with a previous study using a monkey kidney fibroblast line (Vero) [58]. Similarly, some carotenoids, such as β -carotene and lycopene, also presented lower antioxidant or antitumor activities in vitro at concentrations higher than 10 μ M [59, 60]. Cao and Cutler also demonstrated that high concentrations of antioxidants, such as Trolox and uric acid, may not conducive to defense against oxidative stress [61]. It has been reported that many antioxidants at high concentrations might turn out to serve as pro-oxidants and induce large amount of ROS which may lead to oxidative stress and cytotoxicity [62, 63]. There might be three possible explanations for such a bell-shaped effect. Firstly, as a carotenoid, fucoxanthin at higher concentrations (10 μ M and 20 μ M) in the present work might be able to cause the L02 cell membrane more permeabilized in a similar manner to β -carotene [36], consistent with the results of LDH leakage (Figure 3). Secondly, 10 μ M might exceed the capability of L02 cells to incorporate fucoxanthin, just like β -carotene utilized in human colon adenocarcinoma cell lines [39]. Thirdly, fucoxanthin at higher concentrations (more than 10 μ M) might play a role of pro-oxidant instead of antioxidant, and its pro-oxidant activity may arise from the 5,6-monoepoxide which has been shown to undergo ring opening reactions resulting from attacking nucleophiles [64]. However, the mechanism for such a bell-shaped effect needs further investigations to elucidate.

In general, fucoxanthin demonstrated excellent antioxidant activity in defending against the H₂O₂-induced oxidative damage in L02 cells. However, data from the present study should be interpreted with caution since there is a limitation that only the cell experiments were performed. Furthermore, it has been reported that many compounds demonstrated lower antioxidant and disease-prevention activities in vivo than they did in vitro, which is called as “the antioxidant paradox” [63]. Therefore, further intensive research is needed to elucidate the exact antioxidant mechanism of fucoxanthin. However, the prominent effects against H₂O₂-induced oxidative damage demonstrated by

fucoxanthin suggest that it might be a promising agent in protecting against oxidative stress-related diseases in future.

5. Conclusions

In conclusion, the results in the present work suggest that fucoxanthin exerts its cytoprotective effects in L02 cells against H₂O₂-induced oxidative damage, which may be through PI3K-dependent activation of Nrf2 signaling.

Data Availability

The data used to support the findings of this study are included within the article.

Conflicts of Interest

The authors declare no conflicts of interest.

Acknowledgments

This work was supported by National Natural Science Foundation of China (Grant no. 81870237), Shandong Provincial Natural Science Foundation, China (Grant nos. ZR2010DM010, ZR2017LC021), and Higher Educational Science and Technology Program of Shandong Province (Grant no. J10LC20).

References

- [1] R. C. Rabinovitch, B. Samborska, B. Faubert et al., “AMPK Maintains Cellular Metabolic Homeostasis through Regulation of Mitochondrial Reactive Oxygen Species,” *Cell Reports*, vol. 21, no. 1, pp. 1–9, 2017.
- [2] Y. G. Ermakova, D. S. Bilan, M. E. Matlashov et al., “Red fluorescent genetically encoded indicator for intracellular hydrogen peroxide,” *Nature Communications*, vol. 5, no. 1, 2014.
- [3] N. Delanty and M. Dichter, “Antioxidant therapy in neurologic disease,” *American Journal of Ophthalmology*, vol. 131, no. 2, p. 288, 2001.
- [4] J.-Y. Je, P.-J. Park, and S.-K. Kim, “Free radical scavenging properties of hetero-chitoooligosaccharides using an ESR spectroscopy,” *Food and Chemical Toxicology*, vol. 42, no. 3, pp. 381–387, 2004.
- [5] H. Raza, A. John, and F. C. Howarth, “Increased oxidative stress and mitochondrial dysfunction in Zucker diabetic rat liver and brain,” *Cellular Physiology and Biochemistry*, vol. 35, pp. 1241–1251, 2015.
- [6] Y.-S. Yang, T.-H. Ahn, J.-C. Lee et al., “Protective effects of *Pycnogenol* on carbon tetrachloride-induced hepatotoxicity in Sprague—Dawley rats,” *Food and Chemical Toxicology*, vol. 46, no. 1, pp. 380–387, 2008.
- [7] A. Mirshafiey, S. Hosseini, S. Afraei, N. Rastkari, F. T. Zavareh, and G. Azizi, “Anti-aging property of g2013 molecule as a novel immunosuppressive agent on enzymatic and non-enzymatic oxidative stress determinants in rat model,” *Current Drug Discovery Technologies*, vol. 13, no. 1, pp. 25–33, 2016.
- [8] N. Ito, M. Hirose, S. Fukushima, H. Tsuda, T. Shirai, and M. Tatematsu, “Studies on antioxidants: their carcinogenic

- and modifying effects on chemical carcinogenesis," *Food and Chemical Toxicology*, vol. 24, no. 10-11, pp. 1071-1081, 1986.
- [9] J. P. Groten, W. Butler, V. J. Feron, G. Kozianowski, A. G. Renwick, and R. Walker, "An analysis of the possibility for health implications of joint actions and interactions between food additives," *Regulatory Toxicology and Pharmacology*, vol. 31, no. 1, pp. 77-91, 2000.
- [10] D. E. Hathway, "Metabolic Fate in Animals of Hindered Phenolic Antioxidants in Relation to their Safety Evaluation and Antioxidant Function," *Advances in Food and Nutrition Research*, vol. 15, no. C, pp. 1-56, 1966.
- [11] A. Nishino, H. Yasui, and T. Maoka, "Reaction and scavenging mechanism of β -carotene and zeaxanthin with reactive oxygen species," *Journal of Oleo Science*, vol. 66, no. 1, pp. 77-84, 2017.
- [12] T. Tanaka, M. Shnimizu, and H. Moriwaki, "Cancer chemoprevention by carotenoids," *Molecules*, vol. 17, no. 3, pp. 3202-3242, 2012.
- [13] N. M. Sachindra, E. Sato, H. Maeda et al., "Radical scavenging and singlet oxygen quenching activity of marine carotenoid fucoxanthin and its metabolites," *Journal of Agricultural and Food Chemistry*, vol. 55, no. 21, pp. 8516-8522, 2007.
- [14] W. Tang, Y.-F. Jiang, M. Ponnusamy, and M. Diallo, "Role of Nrf2 in chronic liver disease," *World Journal of Gastroenterology*, vol. 20, no. 36, pp. 13079-13087, 2014.
- [15] A. N. Carvalho, O. Firuzi, M. J. Gama, J. van Horssen, and L. Saso, "Oxidative stress and antioxidants in neurological diseases: Is there still hope?" *Current Drug Targets*, vol. 17, no. 16, 2016.
- [16] X. Yan, Y. Chuda, M. Suzuki, and T. Nagata, "Fucoxanthin as the major antioxidant in *Hijikia fusiformis*, a common edible seaweed," *Bioscience, Biotechnology, and Biochemistry*, vol. 63, no. 3, pp. 605-607, 1999.
- [17] S. F. Koduvayur Habeebullah, A. Surendraraj, and C. Jacobsen, "Isolation of Fucoxanthin from Brown Algae and Its Antioxidant Activity: In Vitro and 5% Fish Oil-In-Water Emulsion," *Journal of the American Oil Chemists' Society*, vol. 95, no. 7, pp. 835-843, 2018.
- [18] A. Kawee-ai, A. Kuntiya, and S. M. Kim, "Anticholinesterase and antioxidant activities of fucoxanthin purified from the microalga *Phaeodactylum tricornutum*," *Natural Product Communications (NPC)*, vol. 8, no. 10, pp. 1381-1386, 2013.
- [19] Y. Zhang, H. Fang, Q. Xie et al., "Comparative evaluation of the radical-scavenging activities of fucoxanthin and its stereoisomers," *Molecules*, vol. 19, no. 2, pp. 2100-2113, 2014.
- [20] J. Zheng, M. J. Piao, Y. S. Keum, H. S. Kim, and J. W. Hyun, "Fucoxanthin protects cultured human keratinocytes against oxidative stress by blocking free radicals and inhibiting apoptosis," *Biomolecules & Therapeutics*, vol. 21, no. 4, pp. 270-276, 2013.
- [21] S.-J. Heo and Y.-J. Jeon, "Protective effect of fucoxanthin isolated from *Sargassum siliquastrum* on UV-B induced cell damage," *Journal of Photochemistry and Photobiology B: Biology*, vol. 95, no. 2, pp. 101-107, 2009.
- [22] R. Pangestuti, T.-S. Vo, D.-H. Ngo, and S.-K. Kim, "Fucoxanthin ameliorates inflammation and oxidative responses in microglia," *Journal of Agricultural and Food Chemistry*, vol. 61, no. 16, pp. 3876-3883, 2013.
- [23] C.-L. Liu, A.-L. Liang, and M.-L. Hu, "Protective effects of fucoxanthin against ferric nitrilotriacetate-induced oxidative stress in murine hepatic BNL CL.2 cells," *Toxicology in Vitro*, vol. 25, no. 7, pp. 1314-1319, 2011.
- [24] J. Yin, W. Ren, and X. Wu, "Oxidative stress-mediated signaling pathways: a review," *Journal of Food, Agriculture and Environment (JFAE)*, vol. 11, no. 2, pp. 132-139, 2013.
- [25] S. K. Das, T. Hashimoto, and K. Kanazawa, "Growth inhibition of human hepatic carcinoma HepG2 cells by fucoxanthin is associated with down-regulation of cyclin D," *Biochimica et Biophysica Acta (BBA) - General Subjects*, vol. 1780, no. 4, pp. 743-749, 2008.
- [26] L.-L. Ji, M. Zhang, Y.-C. Sheng, and Z.-T. Wang, "Pyrrolizidine alkaloid clivorine induces apoptosis in human normal liver L-02 cells and reduces the expression of p53 protein," *Toxicology in Vitro*, vol. 19, no. 1, pp. 41-46, 2005.
- [27] C.-L. Liu, Y.-T. Chiu, and M.-L. Hu, "Fucoxanthin enhances HO-1 and NQO1 expression in murine hepatic BNL CL.2 Cells through activation of the Nrf2/ARE system partially by its pro-oxidant activity," *Journal of Agricultural and Food Chemistry*, vol. 59, no. 20, pp. 11344-11351, 2011.
- [28] M. Gülden, A. Jess, J. Kammann, E. Maser, and H. Seibert, "Cytotoxic potency of H₂O₂ in cell cultures: Impact of cell concentration and exposure time," *Free Radical Biology & Medicine*, vol. 49, no. 8, pp. 1298-1305, 2010.
- [29] S. Sabouri-Rad, R. K. Oskuee, A. Mahmoodi, L. Gholami, and B. Malaekheh-Nikouei, "The effect of cell penetrating peptides on transfection activity and cytotoxicity of polyallylamine," *BioImpacts*, vol. 7, no. 3, pp. 139-145, 2017.
- [30] T. Tanoi, T. Tamura, N. Sano et al., "Protecting liver sinusoidal endothelial cells suppresses apoptosis in acute liver damage," *Hepatology Research*, vol. 46, no. 7, pp. 697-706, 2016.
- [31] Z. Liu, Y. Liu, and D. Peng, "Carboxylation of multiwalled carbon nanotube attenuated the cytotoxicity by limiting the oxidative stress initiated cell membrane integrity damage, cell cycle arrestment, and death receptor mediated apoptotic pathway," *Journal of Biomedical Materials Research Part A*, vol. 103, no. 8, pp. 2770-2777, 2015.
- [32] R. Pangestuti and S.-K. Kim, "Neuroprotective effects of marine algae," *Marine Drugs*, vol. 9, no. 5, pp. 803-818, 2011.
- [33] P. M. Riccillo, C. I. Muglia, F. J. De Bruijn, A. J. Roe, I. R. Booth, and O. M. Aguilar, "Glutathione is involved in environmental stress responses in *Rhizobium tropici*, including acid tolerance," *Journal of Bacteriology*, vol. 182, no. 6, pp. 1748-1753, 2000.
- [34] J. Zheng, M. J. Piao, K. C. Kim, C. W. Yao, J. W. Cha, and J. W. Hyun, "Fucoxanthin enhances the level of reduced glutathione via the Nrf2-mediated pathway in human keratinocytes," *Marine Drugs*, vol. 12, no. 7, pp. 4214-4230, 2014.
- [35] R. K. Sangeetha, N. Bhaskar, and V. Baskaran, "Comparative effects of beta-carotene and fucoxanthin on retinol deficiency induced oxidative stress in rats," *Molecular and Cellular Biochemistry*, vol. 331, no. 1-2, pp. 59-67, 2009.
- [36] G. M. Lowe, L. A. Booth, A. J. Young, and R. F. Bilton, "Lycopene and β -carotene protect against oxidative damage in HT29 cells at low concentrations but rapidly lose this capacity at higher doses," *Free Radical Research*, vol. 30, no. 2, pp. 141-151, 1999.
- [37] J. A. Woods, R. F. Bilton, and A. J. Young, " β -Carotene enhances hydrogen peroxide-induced DNA damage in human hepatocellular HepG2 cells," *FEBS Letters*, vol. 449, no. 2-3, pp. 255-258, 1999.
- [38] P. Zhang and S. T. Omaye, " β -Carotene and protein oxidation: Effects of ascorbic acid and α -tocopherol," *Toxicology*, vol. 146, no. 1, pp. 37-47, 2000.
- [39] P. Palozza, S. Serini, N. Maggiano et al., "Induction of cell cycle arrest and apoptosis in human colon adenocarcinoma cell lines

- by β -carotene through down-regulation of cyclin A and Bcl-2 family proteins," *Carcinogenesis*, vol. 23, no. 1, pp. 11–18, 2002.
- [40] T. Nomura, M. Kikuchi, A. Kubodera, and Y. Kawakami, "Proton-donative antioxidant activity of fucoxanthin with 1,1-diphenyl-2-picrylhydrazyl (DPPH)," *International Journal of Biochemistry and Molecular Biology*, vol. 42, no. 2, pp. 361–370, 1997.
- [41] S. Xia, K. Wang, L. Wan, A. Li, Q. Hu, and C. Zhang, "Production, characterization, and antioxidant activity of fucoxanthin from the marine diatom *odontella aurita*," *Marine Drugs*, vol. 11, no. 7, pp. 2667–2681, 2013.
- [42] M. Valko, C. J. Rhodes, J. Moncol, M. Izakovic, and M. Mazur, "Free radicals, metals and antioxidants in oxidative stress-induced cancer," *Chemico-Biological Interactions*, vol. 160, no. 1, pp. 1–40, 2006.
- [43] T. Suzuki, H. Motohashi, and M. Yamamoto, "Toward clinical application of the Keap1-Nrf2 pathway," *Trends in Pharmacological Sciences*, vol. 34, no. 6, pp. 340–346, 2013.
- [44] Z. Luo, W. J. Welch, and C. S. Wilcox, "Abstract 122: activation of nuclear factor erythroid 2-related factor 2 (nrf2) enhances cyclooxygenase 2 expression via promoter antioxidant response element in preglomerular vascular smooth muscle cells (pgvsmcs)," *Acta Radiologica*, vol. 40, no. 5, pp. 545–551, 2015.
- [45] Y.-J. Surh, J. K. Kundu, and H.-K. Na, "Nrf2 as a master redox switch in turning on the cellular signaling involved in the induction of cytoprotective genes by some chemopreventive phytochemicals," *Planta Medica*, vol. 74, no. 13, pp. 1526–1539, 2008.
- [46] R. Gozzelino, V. Jeney, and M. P. Soares, "Mechanisms of cell protection by heme Oxygenase-1," *Annual Review of Pharmacology and Toxicology*, vol. 50, pp. 323–354, 2010.
- [47] J. E. Clark, R. Foresti, C. J. Green, and R. Motterlini, "Dynamics of haem oxygenase-1 expression and bilirubin production in cellular protection against oxidative stress," *Biochemical Journal*, vol. 348, no. 3, pp. 615–619, 2000.
- [48] W. Durante, "Carbon monoxide and bile pigments: Surprising mediators of vascular function," *Vascular Medicine*, vol. 7, no. 3, pp. 195–202, 2002.
- [49] Q.-J. Jin, J.-F. Feng, Y. Sun et al., "Characterization of a heme oxygenase-1 from tobacco that is involved in some abiotic stress responses," *International Journal of Plant Sciences*, vol. 173, no. 2, pp. 113–123, 2012.
- [50] D. Ross and D. Siegel, "Functions of NQO1 in cellular protection and CoQ10 metabolism and its potential role as a redox sensitive molecular switch," *Frontiers in Physiology*, vol. 8, 2017.
- [51] H. Shi, A. Shi, L. Dong et al., "Chlorogenic acid protects against liver fibrosis in vivo and in vitro through inhibition of oxidative stress," *Clinical Nutrition*, vol. 35, no. 6, pp. 1366–1373, 2016.
- [52] H. J. Kim, S. Y. Lee, and S. C. Oh, "The inositide signaling pathway as a target for treating gastric cancer and colorectal cancer," *Frontiers in Physiology*, vol. 7, article 168, 2016.
- [53] K. M. Vasudevan and L. A. Garraway, "AKT signaling in physiology and disease," *Current Topics in Microbiology and Immunology*, vol. 347, no. 1, pp. 105–133, 2010.
- [54] I. Urikura, T. Sugawara, and T. Hirata, "Protective effect of fucoxanthin against UVB-induced skin photoaging in hairless mice," *Bioscience, Biotechnology, and Biochemistry*, vol. 75, no. 4, pp. 757–760, 2011.
- [55] M. Matsui, K. Tanaka, N. Higashiguchi et al., "Protective and therapeutic effects of fucoxanthin against sunburn caused by UV irradiation," *Journal of Pharmacological Sciences*, vol. 132, no. 1, pp. 55–64, 2016.
- [56] H. Maeda, M. Hosokawa, T. Sashima, and K. Miyashita, "Dietary combination of fucoxanthin and fish oil attenuates the weight gain of white adipose tissue and decreases blood glucose in obese/diabetic KK- A^y mice," *Journal of Agricultural and Food Chemistry*, vol. 55, no. 19, pp. 7701–7706, 2007.
- [57] M. Hosokawa, "Fucoxanthin regulates adipocytokine mRNA expression in white adipose tissue of diabetic/obese KK- Ay mice," *Archives of Biochemistry Biophysics*, vol. 504, no. 1, pp. 17–25, 2010.
- [58] S.-J. Heo, S.-C. Ko, S.-M. Kang et al., "Cytoprotective effect of fucoxanthin isolated from brown algae *Sargassum siliquastrum* against H_2O_2 -induced cell damage," *European Food Research and Technology*, vol. 228, no. 1, pp. 145–151, 2008.
- [59] A. J. Young and G. M. Lowe, "Antioxidant and prooxidant properties of carotenoids," *Archives of Biochemistry and Biophysics*, vol. 385, no. 1, pp. 20–27, 2001.
- [60] P. Palozza, "Prooxidant actions of carotenoids in biologic systems," *Nutrition Reviews*, vol. 56, no. 9, pp. 257–265, 1998.
- [61] G. Cao and R. G. Cutler, "High concentrations of antioxidants may not improve defense against oxidative stress," *Archives of Gerontology and Geriatrics*, vol. 17, no. 3, pp. 189–201, 1993.
- [62] M. Valko, M. Izakovic, M. Mazur, C. J. Rhodes, and J. Telser, "Role of oxygen radicals in DNA damage and cancer incidence," *Molecular and Cellular Biochemistry*, vol. 266, no. 1-2, pp. 37–56, 2004.
- [63] B. Halliwell, "The antioxidant paradox: less paradoxical now?" *British Journal of Clinical Pharmacology*, vol. 75, no. 3, pp. 637–644, 2013.
- [64] N. E. Geacintov, H. Hibshoosh, V. Ibanez, M. J. Benjamin, and R. G. Harvey, "Mechanisms of reaction of benzo(a)pyrene-7,8-diol-9,10-epoxide with DNA in aqueous solutions," *Biophysical Chemistry*, vol. 20, no. 1-2, pp. 121–133, 1984.