

Mechanistic Reappraisal of Early Stage Photochemistry in the Light-Driven Enzyme Protochlorophyllide Oxidoreductase

Derren J. Heyes*, Samantha J. O. Hardman, David Mansell, John M. Gardiner, Nigel S. Scrutton*

Manchester Institute of Biotechnology and Photon Science Institute, University of Manchester, Manchester, United Kingdom

Abstract

The light-driven enzyme protochlorophyllide oxidoreductase (POR) catalyzes the reduction of protochlorophyllide (Pchlde) to chlorophyllide (Chlide). This reaction is a key step in the biosynthesis of chlorophyll. Ultrafast photochemical processes within the Pchlde molecule are required for catalysis and previous studies have suggested that a short-lived excited-state species, known as I675*, is the first catalytic intermediate in the reaction and is essential for capturing excitation energy to drive subsequent hydride and proton transfers. The chemical nature of the I675* excited state species and its role in catalysis are not known. Here, we report time-resolved pump-probe spectroscopy measurements to study the involvement of the I675* intermediate in POR photochemistry. We show that I675* is not unique to the POR-catalyzed photoreduction of Pchlde as it is also formed in the absence of the POR enzyme. The I675* species is only produced in samples that contain both Pchlde substrate and Chlide product and its formation is dependent on the pump excitation wavelength. The rate of formation and the quantum yield is maximized in 50:50 mixtures of the two pigments (Pchlde and Chlide) and is caused by direct energy transfer between Pchlde and neighboring Chlide molecules, which is inhibited in the polar solvent methanol. Consequently, we have re-evaluated the mechanism for early stage photochemistry in the light-driven reduction of Pchlde and propose that I675* represents an excited state species formed in Pchlde-Chlide dimers, possibly an excimer. Contrary to previous reports, we conclude that this excited state species has no direct mechanistic relevance to the POR-catalyzed reduction of Pchlde.

Citation: Heyes DJ, Hardman SJO, Mansell D, Gardiner JM, Scrutton NS (2012) Mechanistic Reappraisal of Early Stage Photochemistry in the Light-Driven Enzyme Protochlorophyllide Oxidoreductase. PLoS ONE 7(9): e45642. doi:10.1371/journal.pone.0045642

Editor: Rajagopal Subramanyam, University of Hyderabad, India

Received: July 2, 2012; **Accepted:** August 22, 2012; **Published:** September 26, 2012

Copyright: © 2012 Heyes et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The study was funded by the United Kingdom Biotechnology and Biological Sciences Research Council (BB/F000537/1), the Engineering and Physical Sciences Research Council (EP/J020192/1) and the Royal Society. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: nigel.scrutton@manchester.ac.uk (NSS); derren.heyese@manchester.ac.uk (DJH)

Introduction

The light-driven enzyme protochlorophyllide oxidoreductase (POR) catalyzes the reduction of the C17–C18 double bond of the chlorophyll precursor protochlorophyllide (Pchlde) to form chlorophyllide (Chlide) (Figure 1) [1–4]. Biologically, this reaction is a key regulatory step in the chlorophyll biosynthetic pathway and triggers profound changes in plant development, which result in the modification and reorganization of the plastid membranes [1–3]. The reaction involves the addition of a hydride and proton at the C17 and C18 positions of Pchlde, respectively. Catalysis is initiated by the absorption of light by the Pchlde substrate [3]. For the majority of enzymes, catalysis is generally limited by diffusional processes (e.g. the binding of substrates and coenzymes), conformational changes in the protein, or product release. Mixing strategies are required to derive detailed mechanistic understanding of such reactions and this is often compromised by the temporal resolution of these methods. As POR is light-activated, the reaction can be initiated from a pre-assembled enzyme-substrate complex using a laser pulse. The chemical steps are then monitored on very fast timescales not generally accessible when using rapid mixing strategies with thermally-activated enzymes. Consequently, POR has become an important model system for

studying many aspects of enzyme catalysis [3,4], including the mechanisms and timescales of proton and hydride transfers [4–8], the role of protein dynamics [7,8], the influence of solvent dynamics on reaction chemistry [9,10] and comparative analysis of dynamic properties in PORs from different cyanobacterial strains/plants [11].

The POR-catalyzed reaction involves a highly endergonic light-driven hydride transfer from the *pro-S* face of the nicotinamide ring of NADPH to the C17 position of the Pchlde molecule [6,8], followed by an exergonic thermally-activated proton transfer from a conserved Tyr residue to the C18 position of Pchlde [12] (Figure 1). In POR from *Thermosynechococcus elongatus*, the hydride and proton transfer reactions occur in a sequential mechanism on the microsecond timescale by dynamically coupled nuclear tunneling [8]. Solvent-coupled motions influence the proton transfer reaction but do not affect hydride transfer [9,10]. Coupling of dynamics to the reaction chemistry may have distinct origins in the evolution of POR enzymes [11]. Motions coupled to light-driven hydride transfer are conserved across all POR enzymes and are apparently localized to the enzyme active site. Motions linked to proton transfer vary in different POR enzymes. Extended ‘dynamic networks’ from solvent to active site are thought to optimize proton transfer in cyanobacterial PORs, but




Figure 1. The reaction catalyzed by protochlorophyllide oxidoreductase. The light-driven reduction of the C17–C18 double bond of protochlorophyllide (Pchlde) to form chlorophyllide (Chlide) is catalyzed by protochlorophyllide oxidoreductase (POR) and requires NADPH as a cofactor. The dashed box indicates the double bond that is reduced during the reaction.
doi:10.1371/journal.pone.0045642.g001

are minimized in plant PORs [11]. Following hydride and proton transfer, a series of ordered product release and coenzyme binding steps are required to complete the catalytic cycle. These binding events are linked to major conformational changes in the enzyme [7,13].

Chemical steps in the POR catalytic cycle proceed on a relatively slow microsecond timescale [8], but catalysis is also reliant on picosecond excited-state processes associated with Pchlde [4]. Detailed understanding of the Pchlde photochemistry has proved challenging, but over the last few years a number of time-resolved transient spectroscopy studies have identified different short-lived Pchlde* species [4], both in the isolated pigment [14–19] and in the ternary enzyme-substrate complex [20–22]. Measurements with Pchlde substrate alone have revealed its intrinsic reactivity with multi-exponential excited state dynamics [14–19]. Transient spectral changes following excitation of Pchlde have been interpreted with different models. A number of Pchlde excited state species are likely formed within a few nanoseconds, including an intramolecular charge transfer complex [14–19] and a triplet state [18,19]. The excited state dynamics also depend strongly on solvent polarity [14,15,19].

To investigate the initial ultrafast steps that lead directly to the chemistry of Pchlde photoreduction, additional transient pump-probe experiments have also been performed on the ternary enzyme-substrate (POR-Pchlde-NADPH) complex [20–22]. A putative intermediate with stimulated emission at approximately 675 nm, known as I675*, formed with two time constants of 3 ps and 400 ps [20] and was thought to provide a plausible mechanism for harnessing the light energy to drive the subsequent hydride and proton transfer steps on the microsecond timescale [21,22]. Formation of the I675* intermediate was found to be strongly dependent on the number of pulses applied to the sample. Consequently, the rate and quantum yield of I675* formation increased significantly after the Pchlde substrate had cycled through the excited state at least once [21]. From these observations the reaction was proposed to involve a two photon mechanism. In this mechanism, the initial photon was suggested to convert POR from an inactive to active configuration; the second photon was then required to initiate catalysis i.e. hydride transfer [21]. However, the origin and exact chemical nature of the I675* excited state species remains elusive, although subsequent time-dependent density function theory calculations [23] and solvent isotope effect studies [22] have suggested it may represent

a precursor species in which the Pchlde molecule forms a strongly hydrogen-bonded complex with residues in its direct environment and/or NADPH. Here, we report the use of transient pump-probe absorbance spectroscopy to investigate the formation of the putative I675* intermediate under a range of experimental conditions. In contrast to previous models [21,22], we show that the rate of formation and quantum yield of I675* is independent of POR and has a strong dependence on the wavelength of the pump source used. These new studies lead us to reappraise the mechanism for early stage photochemistry in the POR catalytic cycle. We conclude that the suggested I675* species is in fact caused by excited state energy transfer between neighboring pigment molecules in a Pchlde:Chlide dimer (possibly an excimer), which is not uniquely formed in the enzyme reaction mechanism. We conclude that I675* is not an essential intermediate in POR-catalyzed reduction of Pchlde.

Materials and Methods

Sample Preparation

Recombinant POR from *Thermosynechococcus elongatus* was over-expressed in *Escherichia coli* and purified as described [7]. Pchlde was purified also as described previously [7].

Synthesis of Chlide

Chlide was synthesized by the insertion of magnesium into pheophorbide *a* (Inochem Ltd) using the general protocol described by Lindsey and Woodford [24]. However, under these conditions magnesium insertion was accompanied by the formation of several allomerization by-products, a known occurrence for chlorophyll and its derivatives due to irreversible modification of β -keto ester functionality in the isocyclic ring [25,26]. Consequently, the magnesium insertion reaction was performed within an anaerobic glovebox (Belle Technology Ltd) under a nitrogen atmosphere (<5 ppm oxygen) using anaerobic triethylamine and rigorously degassed dichloromethane (freeze-thaw method). Under these conditions allomerisation products are nearly completely avoided. Pheophorbide *a* (0.05 g, 0.08 mmol) was dissolved in anhydrous dichloromethane (10 mL), followed by addition of triethylamine (0.24 mL, 1.69 mmol) and MgI_2 (0.23 g, 0.84 mmol) and the reaction mixture stirred at room temperature. After 30 minutes the reaction mixture was diluted with dichloromethane and washed with water (3×10 mL). The organic layer was dried over Na_2SO_4 and removed from the anaerobic glovebox for filtration. The solvent was removed under vacuum to give Chlide *a* as the triethylammonium salt in quantitative yield. Mpt $>300^\circ C$. 1H NMR (400 MHz, pyridine- d_5): δ = 9.95 (s, 1H, H-10), 9.81 (s, 1H, H-5), 8.64 (s, 1H, H-20), 8.34 (dd, J 17.8, 11.4 Hz, 1H, H-3a), 6.83 (s, 1H, H-13b), 6.37 (dd, J 17.8, 1.4 Hz, 1H, H-3b'); 6.06 (dd, J 11.4, 1.4 Hz, 1H, H-3b'), 4.60 (br q, J 6.7 Hz, 1H, H-18), 4.47 (br d, J 5.2 Hz, 1H, H-17), 3.81 (s, 3H, OMe), 3.76 (q, J 7.6 Hz, 2H, H-8a), 3.71 (s, 3H, H-12a), 3.37 (s, 3H, H-2a), 3.29 (s, 3H, H-7a), 3.00–2.75 (m, 2H, H-17a), 2.70–2.40 (m, 2H, H-17b), 2.63 (q, J 7.1 Hz, 6H, $(CH_3CH_2)_3N^+H$), 1.66 (t, J 7.8 Hz, 3H, H-8b), 1.63 (d, J 8.0 Hz, 3H, H-18a), 1.03 (t, J 7.2 Hz, 9H, $(CH_3CH_2)_3N^+H$). ^{13}C NMR (100 MHz, pyridine- d_5): δ = 190.3 (C-13a), 175.7 (COO $^-$), 171.5 (COMe), 168.1 (C-19), 161.8 (C-14), 156.4 (C-16), 154.3 (C-1), 152.0 (C-6), 148.2 (C-4), 147.9 (C-11), 146.4 (C-9), 144.6 (C-8), 139.2 (C-3), 136.3 (C-2), 134.6 (C-12), 134.5 (C-7), 131.7 (C-13), 131.0 (C-3a), 119.7 (C-3b), 107.6 (C-10), 106.1 (C-15), 100.1 (C-5), 93.3 (C-20), 66.7 (C-13b), 52.3 (OMe), 51.7 (C-17), 49.6 (C-18), 46.3 ($CH_3CH_2)_3N^+H$), 31.6 (C-17b), 30.8 (C-17a), 23.4 (C-18a), 19.8 (C-8a), 17.8 (C-8b), 12.8 (C-12a), 12.7 (C-2a), 11.2 (C-7a), 11.0 ($CH_3CH_2)_3N^+H$). LRMS (ES $^-$) m/z: 613 [M-

$\text{H}]^-$; HRMS (FTMS NSI^-), $\text{C}_{35}\text{H}_{33}\text{MgN}_4\text{O}_5$ $[\text{M}-\text{H}]^-$; calcd.: 613.2307; obsd.: 613.2293.

Ultrafast Pump-probe Spectroscopy

The laser system used for the transient absorption experiments comprises a Ti:sapphire amplifier (a hybrid Coherent Legend Elite-F-HE) pumped by a Q-switched Nd:YLF laser (Positive Light Evolution-30) and seeded by a Ti:sapphire laser (Spectra-Physics Mai Tai). The amplifier output, which has a wavelength of 800 nm, a 1 kHz repetition rate and ~ 120 fs pulse duration is split, with part of the output used to pump a non-collinear optical parametric amplifier (Light Conversion TOPAS-White), which produces the pump beam with a full-width-half-maximum of ca. 10 nm. The pump wavelengths used are described in the text. Another fraction of the amplifier output is used to generate the white light probe pulse for a Helios (Ultrafast Systems LLC) broadband pump-probe transient absorption spectrometer, with an instrument response function of around 170 fs. Samples of the ternary enzyme-substrate complex contained 200 μM Pchl_{ide}, 500 μM POR and 2 mM NADPH in activity buffer (50 mM Tris pH 7.5, 100 mM NaCl, 1% Triton X-100, 0.1% 2-mercaptoethanol) and were flowed at a rate of approximately 20 ml/min through a 0.2 mm pathlength quartz cell to ensure that a different area of the sample is excited with each pump laser pulse. For enzyme-denatured measurements, samples containing 20 μM Pchl_{ide}, 5 μM POR and 200 μM NADPH in activity buffer were illuminated for different lengths of time, heated to 90°C for 10 mins and centrifuged to remove the precipitated protein. Measurements were performed in a 2 mm pathlength stirred quartz cell. Samples of 'Pchl_{ide} only' and 'Chlide only' contained 20 μM pigment in activity buffer or methanol (as required) in a 2 mm pathlength stirred quartz cell. For all measurements, data points were collected in a random order over a 3 ns time frame.

Global Analysis

Global analysis was performed using the GLOTARAN software package [27]. The number of lifetimes fitted to each data set was determined by increasing the number of components until the residuals were effectively zero. The longest lifetime in each case was set to over 3 μs , essentially infinity on the 3 ns timescale of the experiments. The time-resolved difference spectra were fitted to two different kinetic models. A sequential, unbranched, unidirectional model yielded evolution associated difference spectra (EADS), which represent the spectral evolution of the decay processes (e.g. the second EADS rises with the first lifetime and decays with the second lifetime). A second parallel model of independently decaying components yielded decay associated difference spectra (DADS) which represent the loss or gain of emission or absorption with a certain lifetime.

Fluorescence Measurements

Fluorescence emission end excitation spectra were recorded using a FLS920 fluorescence spectrometer (Edinburgh Instruments Ltd.). Samples contained either 20 μM Pchl_{ide}, 20 μM Chlide or a 50:50 mixture of the 2 pigments in a 0.2 mm pathlength cuvette. Emission spectra were measured using an excitation wavelength of 460 nm and excitation spectra were measured using an emission wavelength of 675 nm. Excitation monochromator slit widths were 1 nm and emission monochromator slit widths were 5 nm.

Results

I675* Formation in POR-Pchl_{ide}-NADPH Complexes

Initial pump-probe measurements were carried out on POR-Pchl_{ide}-NADPH complexes using an excitation wavelength of 475 nm (Figure 2), which yielded similar results to those reported previously [21]. Briefly, time-resolved difference spectra were recorded sequentially so that later scan numbers had been subjected to more laser pulses, resulting in an increased level of the Chlide product in the sample. In the initial scans there is a bleach of the Pchl_{ide} ground state absorption at approximately 640 nm, with only a small amount of the negative signal forming over the first few hundred picoseconds at approximately 675 nm, which represents the I675* intermediate (Figure 2A) [21]. However, in the later scans the negative 675 nm signal forms in higher yields within approximately 500 ps (as shown by the green difference in Figure 2C, Figure S1). In addition, there is also direct excitation of the Chlide product that has been formed during the previous scans, which is shown as a ground state bleach at approximately 675 nm immediately after excitation (Figure 2C).

Subsequently, the formation of the I675* species was investigated by exciting POR-Pchl_{ide}-NADPH complexes with a pump wavelength of 450 nm, where the Pchl_{ide} molecule has




Figure 2. Pump-probe absorption spectroscopy of POR-Pchl_{ide}-NADPH samples after photoexcitation with a laser pulse. The laser pulse was centred at ~ 475 nm. Transient absorption difference spectra at delay times of 2, 17, 492 and 2992 ps after excitation are shown for the average of scans 1–3 (A), scans 8–10 (B) and scans 18–20 (C).

doi:10.1371/journal.pone.0045642.g002

a much higher absorbance (Figure S2). Although photoexcitation of Pchl_{ide} is more efficient at this wavelength the appearance of the negative signal at 675 nm is much less pronounced (Figure S3 and S4), suggesting that I675* formation is not directly related to Pchl_{ide} excitation.

To investigate the role of the enzyme in I675* formation, POR-Pchl_{ide}-NADPH complexes were illuminated for different lengths of time and samples heated to 90°C for 10 mins to denature the POR enzyme. Subsequent pump-probe measurements showed that formation of the I675* intermediate still occurred in these enzyme-denatured samples following excitation at 475 nm (Figure 3). Moreover, the rate of formation and yield of the negative 675 nm signal displays a similar trend to that observed for the native enzyme samples and is dependent on the length of time that samples had been illuminated prior to denaturation (Figure 3). In other words, the level of I675* formation is related to the amount of the Chl_{ide} product in the sample (Figure S5).

Pump-probe Measurements on Pure Pchl_{ide} and Chl_{ide} Samples

The origin of I675* formation was studied in samples containing only the Chl_{ide} product that had been synthesized chemically (see Materials and Methods Section). The time-dependent difference spectra show relatively few spectral features apart from a bleach of the Chl_{ide} ground state absorption at approximately 675 nm at time zero (Figure 4). This slowly returns to the ground state over several ns (time constant of 3.2 ns) and there is no further increase in the amplitude of the negative signal at 675 nm within a few hundred ps. Similar spectral changes were also observed when other pump wavelengths were used (Figure S6), confirming that I675* does not form in ‘Chl_{ide} only’ samples. Moreover, pump-probe experiments on the isolated Pchl_{ide} substrate have also shown that I675* formation does not occur in ‘Pchl_{ide} only’ samples upon photoexcitation at any of the pump wavelengths used [14–19] (Figure S7).

Pump-probe Experiments on Mixtures of Pchl_{ide} and Chl_{ide}

When pump-probe experiments were repeated with mixtures of the isolated Pchl_{ide} and Chl_{ide} pigments (mimicking the enzyme-containing samples, but importantly with enzyme not present) the I675* intermediate was formed in significant quantities with rate constants similar to those observed previously (Figure 5). Moreover, the yield of I675* formation was found to be dependent on the ratio of Pchl_{ide} and Chl_{ide} in the sample with a maximal level of the I675* species found in samples containing a 50:50 mixture of Pchl_{ide} to Chl_{ide} (Figure 5, Table 1). The yield of I675* formation was found to be strongly dependent on the pump wavelength used (Figure S8). The amount of I675* formed was highest when a pump wavelength of 460 nm was used (Figure S8C), with time constants of 10.8 and 258 ps, but this decreases rapidly on the high energy side (very little signal when exciting at 450 nm) and decreases slowly when longer wavelength are used (*i.e.* a relatively large signal remains upon excitation at 475 nm).

In the absence of any additional excited state species the time-resolved difference spectra for the Pchl_{ide} and Chl_{ide} mixtures are expected to be a sum of the ‘Pchl_{ide} only’ and ‘Chl_{ide} only’ data. The time-dependent difference spectra for the Pchl_{ide} and Chl_{ide} mixtures were analyzed by subtracting the sum of the difference spectra of the isolated Pchl_{ide} and Chl_{ide} samples. The residuals ($Z_{residuals}$) from the subtractions were obtained by using the following equation:




Figure 3. Pump-probe absorption spectroscopy of enzyme-denatured POR-Pchl_{ide}-NADPH samples. Spectra are shown after photoexcitation with a laser pulse centred at ~475 nm. The main panels show transient absorption difference spectra at delay times of 2, 16, 516 and 2980 ps after excitation for POR-Pchl_{ide}-NADPH samples that were kept in the dark (A), illuminated for 2 mins (B) and illuminated for 4 mins (C) prior to enzyme denaturation. The insets show the respective kinetic transients at 675 nm (black circles) with a fit of the data to 3 exponentials (solid line). doi:10.1371/journal.pone.0045642.g003

$$Z_{residual} = Z_{mixture} - [(P_{Pchlide} \times Z_{Pchlide}) + (P_{Chlide} \times Z_{Chlide})]$$

where P is the proportion of Pchl_{ide} or Chl_{ide} in the sample mixture (*e. g.* 0.9 and 0.1 for a 90:10% mixture of Pchl_{ide} and Chl_{ide}). The time-resolved difference spectra were also corrected for pump fluence to allow for a more accurate comparison. The residuals calculated after subtraction of the sum of the two component data from the actual time-resolved difference spectra




Figure 4. Pump-probe absorption spectroscopy of 'Chlide only' sample. Spectra are shown after photoexcitation with a laser pulse centred at ~ 475 nm. The main panel shows transient absorption difference spectra at delay times of 2, 16, 516 and 2980 ps after excitation. The inset shows the respective kinetic transients at 675 nm (black circles) with a fit of the data to a single exponential (solid line). doi:10.1371/journal.pone.0045642.g004

are shown for all of the Pchlde and Chlide mixtures at each of the different pump wavelengths (Figure 6 and S9). There is clearly an additional short-lived negative signal at approximately 675 nm in some of the Pchlde and Chlide mixtures that isn't present in the pure pigment samples. Again, the intensity of the negative signal at 675 nm was highest in samples containing a 50:50 mixture of Pchlde to Chlide (indicated by the green trace in Figure 6) and when a pump wavelength of 460 nm was used (Figure 6B).

Global Analysis of Pump-probe Data

Global analysis was used to model the time-resolved difference spectra for the isolated Pchlde and Chlide pigments, and a 50:50 mixture of Pchlde to Chlide after excitation at 450, 460 and 475 nm (Figures 7, S10 and S11). For clarification and to allow a direct comparison between the individual pigments and the Pchlde:Chlide mixture the fitted data for Pchlde only and Chlide only are shown independently and also as a sum of the individual component data. The Pchlde only and Chlide only data could be fitted to 3 exponential functions upon excitation at 450 nm and 460 nm, and 2 exponential functions upon excitation at 475 nm excitation, with the reduced number of components likely to be a result of the low signal intensity at this excitation wavelength. However, an additional component was required to fit the data from the 50:50 mixture of Pchlde:Chlide. A sequential model yields the EADS shown in figure S10, which clearly show the formation and subsequent decay of the negative peak at 675 nm in the Pchlde:Chlide mixture, but not in the individual pigments. However, to gain further insight into the processes that occur upon photoexcitation the data were also fitted to a parallel model, which yielded the DADS shown in figure 7 and S11. Significant differences can be observed between the DADS derived from a sum of the Pchlde only and Chlide only fits, and the DADS fitted to the 50:50 mixture, although the component at approximately 1000 ps is similar at all excitation wavelengths. The spectrum with the infinite lifetime in the 50:50 Pchlde:Chlide mixture shows an increased contribution

Table 1. Amplitudes and time constants for the formation of I675* in samples containing a mixture of Pchlde and Chlide.

Sample	A1 (abs units)	T1 (ps)	A2 (abs units)	T2 (ps)
90% Pchlde, 10% Chlide	0.0016	66.4 \pm 38.1	0.0025	468.1 \pm 138.0
75% Pchlde, 25% Chlide	0.0035	11.5 \pm 3.1	0.0048	279.9 \pm 93.2
50% Pchlde, 50% Chlide	0.0048	10.8 \pm 3.1	0.0045	258.4 \pm 111.2
25% Pchlde, 75% Chlide	0.0049	7.0 \pm 3.2	0.0017	214.5 \pm 252.0
10% Pchlde, 90% Chlide	0.0052	5.0 \pm 3.3	n/a	n/a

doi:10.1371/journal.pone.0045642.t001

from the negative signal at 675 nm (*i.e.* Chlide) relative to the ground state bleach of Pchlde, which is indicative of energy transfer from Pchlde to Chlide (Figure 7 and S11). In addition, the shorter lifetime component of the individual pigments show simple ground state bleaching, whereas the shorter lifetime components of the 50:50 Pchlde:Chlide mixture show a bleach in the region of the Pchlde ground state absorption peak and a positive component in the region of the Chlide ground state absorption peak (Figure 7 and S11). Similar spectral features have also been observed in mixtures of chlorophyll *a* and chlorophyll *b* and are proposed to be characteristic of energy transfer between the two pigments [28–30]. Hence, both the short lifetime and the infinite lifetime DADS indicate that wavelength dependent energy transfer occurs between the Pchlde and Chlide moieties.

To confirm that similar energy transfer processes occur in the ternary enzyme-substrate complex global analysis of the pump-probe data for the POR-Pchlde-NADPH samples revealed that comparable time constants to those of the isolated pigments and pigment mixtures could be obtained. A sequential model yielded the EADS in figure S12, which show the appearance of the I675* species in the later scans of the pump-probe experiment. A parallel model yielded the DADS in figure S13, which shows that the DADS for the initial scans of the pump-probe experiments are very similar to those calculated for the Pchlde only samples at all of the excitation wavelengths. However, as the number of scans increase the spectral features associated with energy transfer becomes more intense and in the final scans (18–20) the DADS are very similar to those calculated for the 50:50 Pchlde:Chlide mixtures.

Effect of Solvent on I675* Formation

Previous studies have shown that the excited state dynamics of isolated Pchlde is strongly dependent on the solvent [14,15,19]. We therefore also investigated the role of solvent on the rate and yield of I675* formation in samples containing a 50:50 mixture of Pchlde and Chlide by repeating transient absorption measurements in the polar solvent methanol (Figure 8). The I675* species was not formed upon photoexcitation at 460 nm in methanol as the amplitude of the negative signal at 675 nm does not increase over the first 500 ps. The residuals calculated after subtraction of the sum of the two component data from the actual time-resolved difference spectra for the Pchlde and Chlide mixture in methanol is negligible over all timescales (Figure 8 inset). Hence, the Pchlde and Chlide mixture does not contain any additional excited state species compared to the 'Pchlde only' and 'Chlide only' samples when methanol is used as solvent.




Figure 5. Pump-probe absorption spectroscopy of samples containing a mixture of Pchlde and Chlide. Spectra are shown after photoexcitation with a laser pulse centred at ~ 460 nm. The main panels show transient absorption difference spectra at delay times of 2, 16, 516 and 2980 ps after excitation for samples containing a mixture of 90% Pchlde and 10% Chlide (A), 50% Pchlde and 50% Chlide (B) and 10% Pchlde and 90% Chlide (C). The insets show the respective kinetic transients at 675 nm (black circles).
doi:10.1371/journal.pone.0045642.g005

Fluorescence Measurements on Pchlde and Chlide Samples

Fluorescence emission spectra were measured for samples containing 'Pchlde only', 'Chlide only' and a 50:50 mixture of Pchlde and Chlide in both aqueous buffer and methanol (Figure 9 and S14). In aqueous buffer there are additional spectral features in the Pchlde and Chlide mixtures, which are not present in the sum of the 'Pchlde only' and 'Chlide only' data (Figure 9A). There




Figure 6. Residuals calculated after subtraction of the sum of the two component data from the time-resolved difference spectra for Pchlde and Chlide mixtures. The residuals calculated at 1 ns after subtraction of the sum of the 'Pchlde only' and 'Chlide only' data from the actual time-resolved difference spectra for all of the Pchlde and Chlide mixtures after photoexcitation at 450 nm (A), 460 nm (B) and 475 nm (C).
doi:10.1371/journal.pone.0045642.g006

is energy transfer from Pchlde to Chlide, as shown by a quenching of the Pchlde fluorescence emission band at approximately 635 nm together with a simultaneous increase in the fluorescence emission from Chlide at approximately 675 nm (Figure 9B). However, in aqueous buffer there are no such differences between the ground state absorption spectra of a 50:50 mixture Pchlde and Chlide and a sum of the 'Pchlde only' and 'Chlide only' spectra (Figure S15), confirming that any changes are due to excited state processes. In addition, fluorescence excitation spectra, recorded using an emission wavelength of 675 nm, show that additional excitation peaks in the Pchlde and Chlide mixture sample are from the Pchlde molecule and provide further evidence of energy transfer between the pigments (Figure S16). This also explains why the I675* species only forms upon photoexcitation at certain wavelengths. Pchlde has a much higher absorbance at 460 nm than Chlide (Figure S2), resulting in a higher level of energy transfer from Pchlde to Chlide. Conversely, at wavelengths below 450 nm there is more direct excitation of Chlide and I675* formation is lower as result of less energy transfer from Pchlde to Chlide. However, when measurements are repeated in methanol the fluorescence emission spectra for the Pchlde and Chlide




Figure 7. Decay associated difference spectra (DADS) resulting from a global analysis of the pump-probe absorption spectroscopy data. Shown are DADS for Pchlde only (A), Chlide only (B), a sum of the 'Pchlde only' and 'Chlide only' data (C) and a mixture of 50% Pchlde and 50% Chlide, after photoexcitation at 460 nm. The data were fitted to a parallel model of independently decaying components as described in the Materials and Methods Section. doi:10.1371/journal.pone.0045642.g007

mixture is simply a sum of the 'Pchlde only' and 'Chlide only' data, suggesting a lack of energy transfer between the two pigments in this solvent (Figure S14).




Figure 8. Pump-probe absorption spectroscopy of samples containing a mixture of 50% Pchlde and 50% Chlide in methanol after photoexcitation with a laser pulse centred at ~460 nm. The main panel show transient absorption difference spectra at delay times of 2, 16, 516 and 2980 ps after excitation. The inset shows the residuals at 675 nm for a mixture of 50% Pchlde and 50% Chlide in methanol and H₂O after subtraction of the sum of the 'Pchlde only' and 'Chlide only' data from the actual time-resolved difference spectra. doi:10.1371/journal.pone.0045642.g008

Discussion

POR is a light-driven enzyme that catalyzes the reduction of the C17–C18 double bond of the chlorophyll precursor molecule, Pchlde [1–4]. Consequently, the catalytic mechanism of POR, which involves sequential hydride and proton transfer steps [8], is completely dependent on the excited-state dynamics of Pchlde, which occur on the picosecond timescale [4]. Although the exact mechanism of photochemistry is not fully understood a short-lived intermediate with a negative signal at approximately 675 nm was previously suggested as the first catalytic intermediate in the reaction cycle, which is formed within a few hundred picoseconds [20–22]. This putative intermediate, referred to as I675*, was proposed to represent an excited state H-bonded intermediate, which harnesses the majority of the excitation energy to drive the subsequent hydride and proton transfer chemistry [21,22]. However, the exact chemical nature of I675* remained unclear and more detailed studies were required to understand the mechanism of photochemistry in the enzyme-substrate complex. We set out to characterize the I675* species in detail using time-resolved pump-probe spectroscopy to study the influence of the POR enzyme, pump wavelength and solvent on its formation. Contrary to previous suggestions [21,22], we find that the I675* species is also formed in the absence of POR. Accordingly, in light of these findings, the mechanism for early stage photochemistry in the light-driven reduction of Pchlde needs to be re-evaluated.

The I675* intermediate has previously only been identified and characterized in transient spectroscopy studies on samples containing POR-Pchlde-NADPH complexes [20–22]. From these studies, formation of I675* was proposed as the first step in the POR-catalytic cycle [21,22]. In turn, formation of I675* was proposed to facilitate subsequent hydride and proton transfer on the microsecond timescale [8]. We have now established that I675* is formed in samples following thermal denaturation of POR and also in Pchlde and Chlide mixtures that contain no POR




Figure 9. Fluorescence emission spectra of samples containing 'Pchlde only', 'Chlide only' and a 50:50 mixture of Pchlde and Chlide. (A) Fluorescence emission spectra of samples containing a mixture of 50% Pchlde and 50% Chlide and the sum of the 'Pchlde only' and 'Chlide only' spectra in H₂O after excitation at 460 nm. (B) Difference spectrum for a mixture of 50% Pchlde and 50% Chlide in H₂O after subtraction of the sum of the 'Pchlde only' and 'Chlide only' data from the actual fluorescence emission spectra. doi:10.1371/journal.pone.0045642.g009

enzyme. Moreover, I675* formation is maximized in 50:50 mixtures of Pchlde and Chlide. It is unlikely therefore that the I675* species is an obligate intermediate in the catalytic cycle of POR. We therefore conclude that formation of I675* is not required for subsequent hydride and proton transfer in the POR catalytic cycle. In POR containing samples, the I675* species has only previously been observed following prior excitation of the enzyme-substrate complex [21]. This has the effect of increasing the level of the Chlide product in the sample prior to further data acquisition. Similarly, in the present work we show that in the absence of POR, the formation of I675* only occurs in samples that contain a mixture of Pchlde and Chlide. This supports our conclusion that the I675* species observed in previous studies [20–22] is not a catalytic intermediate in the POR-catalyzed reaction; it's formation however does require the presence of both the Pchlde substrate and the Chlide product in the sample.

In light of these new findings, it is now possible to propose a new hypothesis regarding the origin and chemical nature of the I675* excited state species. We have shown that I675* only forms when Pchlde and Chlide are both present in the sample and that the level of the I675* species is maximized when a 50:50 ratio of the two pigments is used. This suggests that there are interactions

between Pchlde and Chlide during formation of the I675* species. Moreover, the global analyses and fluorescence measurements reveal that there is direct energy transfer between neighboring Pchlde and Chlide molecules in samples containing a mixture of the two pigments. Previous spectroscopic studies on etiolated beans have shown that Pchlde-Chlide dimers, and possibly larger pigment aggregates, can be formed in etioplast membranes [31–33]. It has also been shown that these Pchlde-Chlide dimers have an absorption peak at 676 nm [31]. Hence, we propose that I675* is likely to represent energy transfer from excited state Pchlde to neighboring Chlide molecules in Pchlde-Chlide dimers. Although the ground state absorption spectra show no evidence of Pchlde-Chlide dimers (see Figure S15), it is possible that I675* is an excimer (*i. e.* the dimer only forms in the excited state), possibly as a result of an increased hydrogen-bonding network. There is precedent for excited state dimerization in similar molecules as excimers have previously been observed in a Chlide analogue as a result of increased H-bonding between neighboring C17 propionic acid sidechains [34].

A dimerization model is also supported by the results on Pchlde and Chlide mixtures in methanol, where no I675* formation was observed. In addition, fluorescence measurements show that there is no energy transfer between Pchlde and Chlide molecules in the excited state. As a polar solvent methanol has been shown to form multiple intermolecular site-specific coordination and H-bonding interactions with the Pchlde and Chlide pigments in the ground state, which are then strengthened in the excited state [23]. This would inhibit any additional interactions between the Pchlde and Chlide molecules and is likely to prevent excited state dimerization in these measurements. Moreover, this may help to explain why the excited state dynamics of Pchlde itself is strongly dependent on the solvent polarity [14,15,19]. In non-polar solvents it would be expected that similar excited state dimerization could occur between neighboring Pchlde molecules, which would then be disrupted in more polar solvents. Hence, this dimerization process may need to be taken into account when interpreting and modeling the excited state dynamics of Pchlde itself [14–19].

Previous studies on the I675* species have led to the suggestion that the POR-catalyzed reaction is a two photon process [21]. The absorption of an initial photon was proposed to induce a more favorable conformation of the active site, which converts the POR enzyme from an inactive to an active configuration. The formation of I675*, which was thought to be necessary for catalysis, only then occurs upon absorption of a second photon [21]. However, this two photon phenomenon for the POR-catalyzed reaction can now be explained using the putative dimerization model for I675* formation. It is likely that a first photon is required to convert Pchlde-Pchlde dimers into Pchlde-Chlide dimers, which can then be converted into Chlide-Chlide dimers upon absorption of the second photon (Figure 10). As we have now shown that the I675* species is likely to result from energy transfer between pigment molecules in Pchlde-Chlide dimers this would explain why it can only be observed in POR-Pchlde-NADPH complexes upon prior excitation with a laser pulse. Indeed, early spectroscopic measurements on etioplast membranes have also indicated that the photoreduction of Pchlde is a two photon process *in vivo* [31,32]. It was shown that enzyme-bound dimers of Pchlde were reduced stepwise to dimers of Chlide in two successive light reactions, with the mixed Pchlde-Chlide dimer formed as a stable intermediate species [31,32].

In conclusion, we have studied the initial excited state processes associated with the light-driven enzyme POR and shown that the previously proposed mechanism of photochemistry [21,22] needs to be re-evaluated. We have shown that an excited state species,




Figure 10. Scheme showing proposed step-wise conversion of Pchlde dimers to Chlide dimers. The I675* species only forms upon excitation of Pchlde-Chlide dimers.
doi:10.1371/journal.pone.0045642.g010

I675*, which was thought to be essential for the subsequent hydride and proton transfer chemistry [21], is not an obligatory catalytic intermediate in Pchlde photoreduction. Instead, we conclude that I675* represents an excited state energy transfer species formed between neighboring pigment molecules in Pchlde-Chlide dimers (possibly excimers) [34]. These findings illustrate the challenges in interpreting complex excited state spectral changes. Our data provide further evidence for the highly reactive nature of these chlorophyll precursor molecules and new mechanistic insight into a natural light-activated enzyme that is crucial to life on Earth.

Supporting Information

Figure S1 The time-dependent absorption changes at 675 nm of POR-Pchlde-NADPH samples after photoexcitation with a laser pulse centred at ~475 nm. The average of scans 1–3, scans 8–10 and scans 18–20 are shown to illustrate the formation of the I675* intermediate within approximately 500 ps in the later scans.
(TIF)

Figure S2 Absorption spectra of Pchlde and Chlide. Both Pchlde and Chlide (5 μ M) were contained in activity buffer (see Experimental Section).
(TIF)

Figure S3 Pump-probe absorption spectroscopy of POR-Pchlde-NADPH samples after photoexcitation with a laser pulse centred at ~450 nm. Transient absorption difference spectra at delay times of 4, 19, 489 and 2989 ps after excitation are shown for the average of scans 1–3 (A), scans 8–10 (B) and scans 18–20 (C).
(TIF)

Figure S4 The time-dependent absorption changes at 675 nm of POR-Pchlde-NADPH samples after photoexcitation with a laser pulse centred at ~450 nm. The average of scans 1–3, scans 8–10 and scans 18–20 are shown to illustrate the lack of I675* formation in approximately 500 ps upon excitation at 450 nm in the later scans.
(TIF)

Figure S5 Absorption spectra and time-dependent absorption changes of enzyme-denatured POR-Pchlde-NADPH samples. (A) Absorption spectra of enzyme-denatured POR-Pchlde-NADPH samples (see Experimental Section) after illumination for varying lengths of time. (B) The time-dependent absorption changes at 675 nm of enzyme-denatured POR-Pchlde-NADPH samples after photoexcitation with a laser pulse centred at ~475 nm. Samples were illuminated for varying lengths of time prior to denaturation. The data are fitted to 3 exponentials (solid lines).
(TIF)

Figure S6 Pump-probe absorption spectroscopy of Chlide only samples after laser pulse photoexcitation. The laser pulse was centred at ~450 nm (A) and 460 nm (B). The main panel shows transient absorption difference spectra at delay times of 2, 16, 516 and 2980 ps after excitation. The insets show the respective kinetic transients at 675 nm (black circles) with a fit of the data to a double exponential function (solid line). Time constants of 136 ps (126 ps upon excitation at 460 nm) and 3.5 ns were calculated.
(TIF)

Figure S7 Pump-probe absorption spectroscopy of Pchlde only samples after laser pulse photoexcitation. The laser pulse was centred at ~450 nm (A), 460 nm (B) and 475 nm (C). Transient absorption difference spectra are shown at delay times of 2, 16, 516 and 2980 ps after excitation.
(TIF)

Figure S8 Pump-probe absorption spectroscopy of samples containing a mixture of 50% Pchlde and 50% Chlide after laser pulse photoexcitation. The laser pulse was centred at ~435 nm (A), 450 nm (B), 460 nm (C), 475 nm (D) and 580 nm (E). The main panels show transient absorption difference spectra at delay times of 2, 16, 516 and 2980 ps after excitation. The insets show the respective kinetic transients at 675 nm (black circles).
(TIF)

Figure S9 The residuals calculated at all time points after subtraction of the sum of the Pchlde only and Chlide only data from the actual time-resolved difference spectra for all of the Pchlde and Chlide mixtures. Residuals calculated after photoexcitation at 450 nm (A), 460 nm (B) and 475 nm (C). The intensity of the blue region indicates the presence of an additional short-lived negative signal at approximately 675 nm.
(TIF)

Figure S10 Evolution associated difference spectra (EADS) for Pchlde, Chlide and mixtures. EADS resulting from a global analysis of the pump-probe absorption spectroscopy data for Pchlde only, Chlide only, a sum of the ‘Pchlde only’ and ‘Chlide only’ data and a mixture of 50% Pchlde and 50% Chlide after photoexcitation at 450 nm (A), 460 nm (B) and 475 nm (C). The data were fitted to a sequential model which represents the spectral evolution of the decay processes as described in the Materials and Methods Section.
(TIF)

Figure S11 Decay associated difference spectra (DADS) for Pchlde, Chlide and mixtures. DADS resulting from a global analysis of the pump-probe absorption spectroscopy data for Pchlde only, Chlide only, a sum of the ‘Pchlde only’ and ‘Chlide only’ data and a mixture of 50% Pchlde and 50% Chlide after photoexcitation at 450 nm (A), 460 nm (B) and 475 nm (C). The data were fitted to a parallel model of independently decaying components as described in the Materials and Methods Section.
(TIF)

Figure S12 Evolution associated difference spectra (EADS) for POR-Pchlde-NADPH samples. EADS resulting from a global analysis of the pump-probe absorption spectroscopy data for POR-Pchlde-NADPH samples after photoexcitation with a laser pulse centred at 450 nm (A), 460 nm (B) and 475 nm (C). The fits for the average of scans 1–3, scans 8–10 and scans 18–20 are shown. The data were fitted to a sequential model which represents the spectral evolution of the decay processes as described in the Materials and Methods Section.
(TIF)

Figure S13 Decay associated difference spectra (DADS) for POR-Pchl_a-NADPH samples. DADS resulting from a global analysis of the pump-probe absorption spectroscopy data for POR-Pchl_a-NADPH samples after photoexcitation with a laser pulse centred at 450 nm (A), 460 nm (B) and 475 nm (C). The fits for the average of scans 1–3, scans 8–10 and scans 18–20 are shown. The data were fitted to a parallel model of independently decaying components as described in the Materials and Methods Section.

(TIF)

Figure S14 Fluorescence emission spectra in methanol. Fluorescence emission spectra of samples containing a mixture of 50% Pchl_a and 50% Chl_a and the sum of the ‘Pchl_a only’ and ‘Chl_a only’ spectra in methanol after excitation at 460 nm.

(TIF)

Figure S15 Absorption spectra in aqueous buffer. Absorption spectra of samples containing a mixture of 50% Pchl_a and 50% Chl_a and the sum of the ‘Pchl_a only’ and ‘Chl_a only’ spectra in aqueous activity buffer.

References

- Lebedev N, Timko MP (1998) Protochlorophyllide photoreduction. *Photosynth Res* 58: 5–23.
- Masuda T, Takamiya K (2004) Novel insights into the enzymology, regulation and physiological functions of light-dependent protochlorophyllide oxidoreductase in angiosperms. *Photosyn Res* 81: 1–29.
- Heyes DJ, Hunter CN (2005) Making light work of enzyme catalysis: protochlorophyllide oxidoreductase. *Trends Biochem Sci* 30: 642–649.
- Scrutton NS, Groot ML, Heyes DJ (2012) Excited state dynamics and catalytic mechanism of the light-driven enzyme protochlorophyllide oxidoreductase. *Phys Chem Chem Phys* 14: 8818–8824.
- Heyes DJ, Ruban AV, Wilks HM, Hunter CN (2002) Enzymology below 200 K: the kinetics and thermodynamics of the photochemistry catalyzed by protochlorophyllide oxidoreductase. *Proc Natl Acad Sci U S A* 99: 11145–11150.
- Heyes DJ, Heathcote P, Rigby SE, Palacios MA, van Grondelle R, et al. (2006) The first catalytic step of the light-driven enzyme protochlorophyllide oxidoreductase proceeds via a charge transfer complex. *J Biol Chem* 281: 26847–26853.
- Heyes DJ, Hunter CN (2004) Identification and characterization of the product release steps within the catalytic cycle of protochlorophyllide oxidoreductase. *Biochemistry* 43: 8265–8271.
- Heyes DJ, Sakuma M, De Visser S, Scrutton NS (2009) Nuclear quantum tunneling in the light-activated enzyme protochlorophyllide oxidoreductase. *J Biol Chem* 284: 3762–3767.
- Durin G, Delaunay A, Darnault C, Heyes DJ, Royant A, et al. (2009) Simultaneous measurements of solvent dynamics and functional kinetics in a light-activated enzyme. *Biophys J* 96: 1902–1910.
- Heyes DJ, Sakuma M, Scrutton NS (2009) Solvent-slaved protein motions accompany proton but not hydride tunneling in light-activated protochlorophyllide oxidoreductase. *Angew Chem Int Ed Engl* 48: 3850–3853.
- Heyes DJ, Levy C, Sakuma M, Robertson DL, Scrutton NS (2011) A twin-track approach has optimized proton and hydride transfer by dynamically coupled tunneling during the evolution of protochlorophyllide oxidoreductase. *J Biol Chem* 286: 11849–11854.
- Menon BRK, Waltho JP, Scrutton NS, Heyes DJ (2009) Cryogenic and laser photoexcitation studies identify multiple roles for active site residues in the light-driven enzyme protochlorophyllide oxidoreductase. *J Biol Chem* 284: 18160–18166.
- Heyes DJ, Sakuma M, Scrutton NS (2007) Laser excitation studies of the product release steps in the catalytic cycle of the light-driven enzyme, protochlorophyllide oxidoreductase. *J Biol Chem* 282: 32015–32020.
- Dietzek B, Maksimenka R, Siebert T, Birckner E, Kiefer W, et al. (2004) Excited-state processes in protochlorophyllide a: a femtosecond time-resolved absorption study. *Chem Phys Lett* 397: 110–115.
- Dietzek B, Kiefer W, Hermann G, Popp J, Schmitt M (2006) Solvent effects on the excited-state processes of protochlorophyllide a: a femtosecond time-resolved absorption study. *J Phys Chem B* 110: 4399–4406.
- Dietzek B, Tschierlei S, Hermann G, Yartsev A, Pascher T, et al. (2006) The excited-state chemistry of protochlorophyllide a: a time-resolved fluorescence study. *ChemPhysChem* 7: 1727–1733.
- Dietzek B, Kiefer W, Yartsev A, Sundstrom V, Schellenberg P, et al. (2009) Protochlorophyllide a: A comprehensive photophysical picture. *ChemPhysChem* 10: 144–150.
- Dietzek B, Tschierlei S, Hanf R, Seidel S, Yartsev A, et al. (2010) Dynamics of charge separation in the excited-state chemistry of protochlorophyllide. *Chem Phys Lett* 492: 157–163.
- Sytina OA, van Stokkum IHM, Heyes DJ, Hunter CN, van Grondelle R, et al. (2010) Protochlorophyllide excited-state dynamics in organic solvents studied by time-resolved visible and mid-infrared spectroscopy. *J Phys Chem B* 114: 4335–4344.
- Heyes DJ, Hunter CN, van Stokkum IHM, van Grondelle R, Groot ML (2003) Ultrafast enzymatic reaction dynamics in protochlorophyllide oxidoreductase. *Nat Struct Biol* 10: 491–492.
- Sytina OA, Heyes DJ, Hunter CN, Alexandre MTA, van Stokkum IHM, et al. (2008) Conformational changes in an ultrafast light-driven enzyme determine catalytic activity. *Nature* 456: 1001–1004.
- Sytina OA, van Stokkum IHM, Heyes DJ, Hunter CN, Groot ML (2012) Spectroscopic characterization of the first ultrafast catalytic intermediate in protochlorophyllide oxidoreductase. *Phys Chem Chem Phys* 14: 616–625.
- Zhao GJ, Han KL (2008) Site-specific solvation of the photoexcited protochlorophyllide a in methanol: formation of the hydrogen-bonded intermediate state induced by hydrogen-bond strengthening. *Biophys J* 94: 38–46.
- Lindsey JS, Woodford JN (1995) A simple method for preparing magnesium porphyrins. *Inorg Chem* 34: 1063–1069.
- Müller AH, Gough SP, Bollivar DW, Meldal M, Willows RD, et al. (2011) Methods for the preparation of chlorophyllide a: An intermediate of the chlorophyll biosynthetic pathway. *Anal Biochem* 419: 271–276.
- Hynninen PH, Hyvärinen K (2002) Tracing the allomerization pathways of chlorophylls by ¹⁸O-labeling and mass spectrometry. *J Org Chem* 67: 4055–4061.
- Snellenburg JJ, Laptinok SP, Seger R, Mullen KM, van Stokkum IHM (2012) Glotaran: A Java-Based Graphical User Interface for the R Package TAMP. *J Statistical Software* 49: 1–22.
- Croce R, Muller MG, Bassi R, Holzwarth AR (2003) Chlorophyll b to chlorophyll a energy transfer kinetics in the CP29 antenna complex: A comparative femtosecond absorption study between native and reconstituted proteins. *Biophys J* 84: 2508–2516.
- Kleima EJ, Gradinaru CC, Calkoen F, van Stokkum IHM, van Grondelle R, et al. (1997) Energy transfer in LHClI monomers at 77K studied by sub-picosecond transient absorption spectroscopy. *Biochemistry* 36: 15262–15268.
- Connelly JP, Muller MG, Bassi R, Croce R, Holzwarth AR (1997) Femtosecond transient absorption study of carotenoid to chlorophyll energy transfer in the light-harvesting complex II of photosystem II. *Biochemistry* 36: 281–287.
- Mathis P, Sauer K (1972) Chlorophyll formation in greening bean leaves during the early stages. *Plant Physiol* 51: 115–119.
- Mathis P, Sauer K (1972) Circular dichroism studies on the structure and photochemistry of protochlorophyllide and chlorophyllide holochrome. *Biochim Biophys Acta* 267: 498–511.
- Vaughan GD, Sauer K (1974) Energy transfer from protochlorophyllide to chlorophyllide during photoconversion of etiolated bean holochrome. *Biochim Biophys Acta* 347: 383–394.
- Wasielewski MR, Studier MH, Katz JJ (1976) Covalently linked chlorophyll a dimer: A biomimetic model of special pair chlorophyll. *Proc Natl Acad Sci U S A* 73: 4282–4286.