

## Editorial

### Aging - How Lifestyle Changes Could Prove to be an Effective Medicine for the Aging Cardiovascular System

Aging is a growing medical concern. Despite more people living longer, the economical impact is likely to be severe, and pressures on healthcare providers are constantly growing to treat aging-related conditions and disorders. It is estimated, that in the US, life expectancy will go from ~76 years of age in 2010 to 80 years in 2030 [1], a pattern that is similar to a host of industrialised countries. Indeed, in the United Kingdom, recent analysis has shown that male life expectancy has increased from 69 years in 1970 to 78 years in 2010, and female life expectancy from 75 years to 82 years in the same time period [2]. As a result, incidence of aging-related diseases such as Cardiovascular Disease (CVD), dementia, and cancer is likely to increase substantially [3]. It is no surprise, therefore, that research into medical advancements and lifestyle behaviours and the impact of these on aging populations has gathered a lot of interest in the academic community, as seen by the Physiological Society announcing a year of 'Understanding Ageing' in 2015. This has led to the growth of research in aging, and also specifically cardiovascular aging.

Lifestyle choices can have significant influence on how our cardiovascular system ages. We see individuals who are fitter and regularly exercise, live longer than unfit/sedentary individuals [4, 5], which could be due to changes in our cardiovascular system. Such favourable changes in our cardiovascular system with simple modifications in our lifestyle thus offer a significant promise in helping to counteract the impact of aging on our bodies. This small collection of reviews, encompasses the effects of diet, exercise and physical activity on the 'aging cardiovascular system', and provides some novel insights into the mechanisms by which these affect our vascular and cardiac tissues.

The first review [6] puts forward the argument that resistance exercise could affect vascular aging via modulation of arterial stiffness and Matrix Metalloproteinases (MMP). Older individuals display elevated levels of arterial stiffness (as measured by pulse wave technology) compared to younger individuals [7], with recent evidence supporting the view that exercise can reduce arterial stiffness in humans [8]. Knox demonstrates some data to suggest that arterial remodelling with exercise may be a result of altered MMP regulation, but more work is needed to observe this link in humans.

The second review [9] discusses the role of aging on the body's endogenous endothelial repair capacity, via the reduction and dysfunction of endothelial progenitor cells, and how exercise and dietary behaviour modulate both the number and function of these cells. The effects vary from improving bone marrow niche environment and pro-apoptotic stimuli to promoting our own stem/progenitor cell repair mechanisms on our ageing vasculature.

The subsequent review by Beaumont *et al.* [10] uses masters athlete's as a model of lifelong exercisers, and the differences in cardiac structure and function between master's athlete's in inactive older people. We observe that masters athletes displays better diastolic function than sedentary age-matched controls, which appears to be maintained into the 8<sup>th</sup> decade of life. Interestingly, the review does not demonstrate much evidence on differences in systolic function, and a much debated measure of systolic function, ejection fraction, does not appear to be different between sedentary controls and masters athletes, being suggestive that lifelong exercise does not improve cardiac health via modulations in ejection fraction.

The final review in our collection, by Peace *et al.* [11], provides an insight into carotid and coronary arterial function, with subsequent focus on the role of advancing age on both, and the potential for exercise to act as a countermeasure for the aging-associated deleterious effects. They also purport that, since coronary artery function is difficult and incredibly invasive to measure and quantify, that we can use carotid artery function as a surrogate, as it appears to show similar structural and functional changes with disease conditions, offering clinicians and researchers a non-invasive global measure of arterial function that could be used to determine risk of myocardial infarction, stroke, and other atherosclerotic-mediated events.

Together, these reviews provide evidence for the importance of lifestyle modifications and behaviour change to treat age-associated cardiovascular problems, which could subsequently have significant effects on healthcare burden and quality of life of patients. However, translation of such work into practice is often poor at best, and thus strategies must be put into place by governing bodies, healthcare providers, and clinicians to encompass such lifestyle strategies to treat disease.

#### DISCLOSURES

The Guest Editor is a contributor to the Special Issue (Ross, 2018).

#### REFERENCES

- [1] Kontis V, Bennett JE, Mathers CD, Li G, Foreman K, Ezzati M. Future life expectancy in 35 industrialised countries: Projections with a Bayesian model ensemble. *The Lancet* 2017; 389(10076): 1323-35.
- [2] Wang H, Dwyer-Lindgren L, Lofgren KT, *et al.* Age-specific and sex-specific mortality in 187 countries, 1970–2010: A systematic analysis for the Global Burden of Disease Study 2010. *The Lancet* 2012; 380(9859): 2071-94.

- [3] Lozano R, Naghavi M, Foreman K, *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010. *The Lancet* 2012; 380(9859): 2095-128.
- [4] Barry VW, Baruth M, Beets MW, Durstine JL, Liu J, Blair SN. Fitness vs. fatness on all-cause mortality: A meta-analysis. *Prog Cardiovasc Dis* 2013; 56(4): 382-90.
- [5] Cabanas-Sánchez V, Guallar-Castillón P, Higuera-Fresnillo S, García-Esquinas E, Rodríguez-Artalejo F, Martínez-Gómez D. Physical activity, sitting time, and mortality from inflammatory diseases in older adults. *Front Physiol* 2018; 9(898).
- [6] Knox A. Arterial aging, metalloproteinase regulation, and the potential of resistance exercise. *Current Cardiology Rev* 2018; 14(4): 227-32.
- [7] Holland AM, Martin JS, Mattson CD, Lohse KR, Finn PR, Stager JM. A cross-sectional study of physical activity and arterial compliance: The effects of age and artery size. *J Am Soc Hypertens*. 2017; 11(2): 92-100.
- [8] Zhang Y, Qi L, Xu L, *et al.* Effects of exercise modalities on central hemodynamics, arterial stiffness and cardiac function in cardiovascular disease: Systematic review and meta-analysis of randomized controlled trials. *PLoS One* 2018; 13(7): e0200829.
- [9] Ross M. Endothelial regenerative capacity and aging: Influence of diet, exercise and obesity. *Curr Cardiol Rev* 2018; 14(4): 233-44.
- [10] Beaumont A, Campbell A, Grace F, Sculthorpe N. Cardiac response to exercise in normal ageing: What can we learn from masters athletes? *Curr Cardiol Rev* 2018; 14(4): 245-53.
- [11] Peace A, Van Mil A, Jones H, Thijssen DHJ. Similarities and differences between carotid artery and coronary artery function. *Curr Cardiol Rev* 2018; 14(4): 254-63.

**Mark. D. Ross**

*(Guest Editors)*

School of Applied Sciences  
Edinburgh Napier University  
Edinburgh  
UK  
E-mail: M.Ross@napier.ac.uk