



The child is encouraged during the art therapy sessions, by exhibiting the works and decorating the work environment.



Through these non-directive methods, the evolution of visual thinking is accelerated. The chromatic diversification, the gestures in painting and the alternation of work techniques such as printing, graphic lines and dripping, are signs of a visual thinking. The child discovers the environment and interacts with it through art.



Observing the potential of the materials around him and also certain physical phenomena, such as a three-dimensional mask that allows the color to outline its volumes by draining it on the shape, the child uses consciously the properties of materials and the movement of the object.

Results: Discovering artistic and decorative skills; Increased self-esteem; Interaction in the artistic environment and even verbal communication in cases of autism.

Conclusions: Through art, the child can get closer to the social life.

Disclosure: No significant relationships.

Keywords: art therapy; social integration; raw art; outstanding skills

EPP0332

Call the Psychiatrist! - Study about Delirium in the context of liaison psychiatry

I. Fonseca Vaz*, S. Mouta, B. Jesus and S. Castro

Unidade Local de Saúde da Guarda, Departamento De Psiquiatria E Saúde Mental, Guarda, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.613

Introduction: Delirium is characterized as a short-term consciousness and cognition disturbance which tends to fluctuate during the course of the day. It is a common and serious problem, mainly in hospitalized older adults, potentially avoidable and often poorly recognized.

Objectives: We propose an analysis on the theme through a work that evaluates the requests for psychiatric consultation made in a district hospital in Portugal during the course of 12 months.

Methods: We identified all patients on the requests for psychiatric consultation and obtained a demographic, clinical and consultation requests by medical specialties data and conducted statistical analysis using Excel.

Results: We identified 106 consultation requests, in which 41 cases were eventually diagnosed as delirium. Most (83%) were hyperactive delirium, 12% were hypoactive delirium and 5% were mixed

delirium. Incidence was higher in males (59%) and in those aged between 66 and 80 years old (56.1%). Most consultation requests were made by Internal Medicine (46.3%), followed by General Surgery (26.8%), Pulmonology (14.6%), Orthopedics (9.8%) and Neurology (2.5%). Finally, we analyzed which symptoms mentioned in the request made physicians consider requesting a psychiatric evaluation. Approximately half of the cases (48.8%) reported psychomotor agitation, followed by temporal/spatial disorientation (41.5%) and aggressive behaviour (17.1%).

Conclusions: We highlight a still notorious lack of proper identification of delirium, resulting in symptoms being incorrectly interpreted as a psychiatric disorder. This may cause a delay in the adequate diagnosis and management of the condition, increasing the morbidity and mortality of patients.

Disclosure: No significant relationships.

Keywords: delirium; symptoms; liaison psychiatry

EPP0333

The relationship between proactive coping and mindfulness: cross-cultural analysis

E. Belinskaya*, M. Djuraeva, O. Tihomandritskaya and E. Dubovskaya

Lomonosov Moscow State University, Faculty Of Psychology, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.614

Introduction: The inconsistency of the available empirical data on personal and situational predictors of effective coping allows us to make an assumption about the presence of mediating variables. Their search can be centered both on the inclusion of parameters of a higher socio-cognitive level in the analysis, and on the procedural characteristics of coping, one of which is proactivity in coping. The construct of mindfulness satisfies both of these requirements.

Objectives: The purpose of this study was to identify cross-cultural differences in the relationship between proactive coping strategies and the level of mindfulness

Methods: Five Facet Mindfulness Questionnaire ("eastern" awareness), Scale of Mindfulness ("western" awareness), Proactive Coping Inventory. The sample was N = 452 (residents of Russia and Uzbekistan, age 18-25)

Results: For Russians and Uzbeks, the indicators of "western" awareness are associated with the attitude to potential difficulties as a source of positive experience and with reflection in case of their occurrence of possible behaviors, cognitive assessment of their own resources and prediction of results, as well as with such a proactivity strategy as the search for instrumental support. Intercultural differences were noted in terms of "eastern" awareness: for Uzbeks, they are associated with a proactive coping strategy, and for Russians – with reflexive coping.

Conclusions: Mindfulness has the potential of meta-cognitive function in a situation of assessing possible life difficulties and choosing proactive coping strategies.

Disclosure: No significant relationships.

Keywords: coping strategies; proactive coping; Mindfulness

EPP0335

Assessment of psychopathological symptoms in patients with primary hypothyroidism

O. Pityk* and N. Karbovskyi

Ivano-Frankivsk National Medical University, Psychiatry, Narcology And Medical Psychology, Ivano-Frankivsk, Ukraine

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.615

Introduction: Thyroid dysfunction such as hypothyroidism, is connected with numerous neurological and psychiatric disorders. However, the importance of assessing the interaction between brain, psyche and thyroid in clinical practice is often underestimated, and this has a direct impact on the planning of therapeutic interventions and treatment efficacy in patients with primary hypothyroidism.

Objectives: We examined 132 patients with primary hypothyroidism.

Methods: Assessment of the presence and severity of psychopathology was performed using the technique SCL-90-R (questionnaire severity of psychopathology).

Results: The results showed the highest scores on the scales of somatization ($3,75 \pm 0,12$), depression ($3,64 \pm 0,13$), interpersonal anxiety ($3,45 \pm 0,19$), phobias ($3,25 \pm 0,31$). High rates of somatization scale showing a violation of bodily dysfunction of various body systems-cardiovascular, gastrointestinal, respiratory and headache, muscular discomfort and other unpleasant sensations in different parts of the body and manifest themselves in a complaint of patients. Scale depression revealed the presence of dysphoria, anhedonia, low affect, loss of vitality and interest in life. Relatively high on a scale of phobias indicate the presence in these patients persistent fear responses to certain situations and objects that are irrational and inadequate and lead to avoiding behavior. General index of severity of symptoms (GSI) and the index of an existing symptomatic distress (PSDI) were significantly higher in the following patients than in hypothyroid patients without mental disorders.

Conclusions: Thus, these results should be taken into account when determining treatment strategy both in psychopharmacotherapy and different methods of psychological correction.

Disclosure: No significant relationships.

Keywords: Hypothyroidism; nonpsychotic mental disorders; psychopathological symptoms

EPP0336

Significant early and long-term improvement of neuropsychiatry symptomatology in HCV-infected patients after viral eradication with DAA

R. Martin-Santos^{1*}, C. Bartrés², L. Nacar³, R. Navinés³, M. Caveró³, S. Lens², S. Rodríguez-Tajes², J.C. Pariente⁴, I. Horrillo⁵, E. Muñoz-Moreno⁴, N. Bargallo⁴, L. Capuron⁶, J. Meana⁵, X. Forn² and Z. Mariño²

¹Hospital Clínic,, 1department Of Psychiatry And Psychology, Barcelona, Spain; ²Hospital Clínic,, UB, IDIBAPS, CIBERSAM, Liver Unit, Barcelona, Spain; ³Hospital Clínic,, UB, IDIBAPS, CIBERSAM, 1department Of Psychiatry And Psychology, Barcelona, Spain;