

RESEARCH

Open Access

Analysis of kinetoplast cytochrome *b* gene of 16 *Leishmania* isolates from different foci of China: different species of *Leishmania* in China and their phylogenetic inference

Bin-Bin Yang^{1,2}, Da-Li Chen², Jian-Ping Chen^{2,3*}, Lin Liao², Xiao-Su Hu² and Jia-Nan Xu²

Abstract

Background: *Leishmania* species belong to the family Trypanosomatidae and cause leishmaniasis, a geographically widespread disease that infects humans and other vertebrates. This disease remains endemic in China. Due to the large geographic area and complex ecological environment, the taxonomic position and phylogenetic relationship of Chinese *Leishmania* isolates remain uncertain. A recent internal transcribed spacer 1 and cytochrome oxidase II phylogeny of Chinese *Leishmania* isolates has challenged some aspects of their traditional taxonomy as well as cladistics hypotheses of their phylogeny. The current study was designed to provide further disease background and sequence analysis.

Methods: We systematically analyzed 50 cytochrome *b* (cyt *b*) gene sequences of 19 isolates (16 from China, 3 from other countries) sequenced after polymerase chain reaction (PCR) using a special primer for cyt *b* as well as 31 sequences downloaded from GenBank. After alignment, the data were analyzed using the maximum parsimony, Bayesian and network methods.

Results: Sequences of six haplotypes representing 10 Chinese isolates formed a monophyletic group and clustered with *Leishmania tarentolae*. The isolates GS1, GS7, XJ771 of this study from China clustered with other isolates of *Leishmania donovani* complex. The isolate JS1 was a sister to *Leishmania tropica*, which represented an *L. tropica* complex instead of clustering with *L. donovani* complex or with the other 10 Chinese isolates. The isolates KXG-2 and GS-GER20 formed a monophyletic group with *Leishmania turanica* from central Asia. In the different phylogenetic trees, all of the Chinese isolates occurred in at least four groups regardless of geographic distribution.

Conclusions: The undescribed *Leishmania* species of China, which are clearly causative agents of canine leishmaniasis and human visceral leishmaniasis and are related to *Sauroleishmania*, may have evolved from a common ancestral parasite that came from the Americas and may have split off earlier than the other old world *Leishmania*. Our results also suggest the following: the isolates GS7, GS1 and XJ771 occur as part of the *L. donovani* complex; the JS1 isolate is *L. tropica*; and the isolate GS-GER20 identified as *Leishmania gerbilli* is close to KXG-2 which is *L. turanica*.

Keywords: *Leishmania*, Phylogeny, cyt *b*, China, *Sauroleishmania*

* Correspondence: jpchen007@163.com

²Department of Parasitology, West China School of Preclinical and Forensic Medicine, Sichuan University, #17, The 3th Section of South Renmin Road, Chengdu, Sichuan 610041, China

³Animal Disease Prevention and Food Safety Key Laboratory of Sichuan Province, #24, The South 1st Section of Yihuan Road, Chengdu, Sichuan 610041, China

Full list of author information is available at the end of the article

Background

The leishmaniasis are a group of vector borne diseases that are caused by flagellate of the genus *Leishmania*, which is transmitted by the bite of the sandfly, and affect as many as 12 million people worldwide with 1.5–2 million new cases each year in 88 countries [1]. The genus *Leishmania* consists of nearly 30 species of morphologically similar kinetoplastid protozoa, and approximately 20 of these species are responsible for a spectrum of human diseases that ranges from mild to fatal infections [2,3].

It is well accepted now that the genus *Leishmania* forms a monophyletic group with three distinct subgenera *Leishmania* (*Leishmania*), *Leishmania* (*Viannia*), and *Leishmania* (*Sauroleishmania*) [4]. Once, the *Leishmania*-like parasites of reptiles were recognized as a separate genus *L. (Sauroleishmania)* [5,6]. But the classification of lizard *Leishmania* as subgenus *L. (Sauroleishmania)* was proposed by Saf'janova [7], Croan *et al.* [8], Orlando *et al.* [9], Zelazny *et al.* [10] and Fraga *et al.* [4] on the basis of biological criteria and analysis of different *Leishmania* gene. According to the absolute chromosomal size difference index (aCSDI) analysis of as many as 31 "conserved" chromosomes, subgenera *L. (Leishmania)* is divided into clusters: (1) Old world representatives of subgenus *Leishmania* (OWL), and (2) New world representatives of subgenus *Leishmania* (NWL) and subgenus *Viannia* (NWV) [6].

The complexity of the taxonomy and phylogenetic relationships of the Chinese *Leishmania* was due to the extensive geographic area and complex ecological environment. Identification of species responsible for different leishmaniasis and clinical manifestation remains uncertain. The strains from cutaneous leishmaniasis (CL) in Xinjiang Uygur Autonomous Region (Xinjiang) especially in Karamay is closely related to *L. tropica* with analysis of SSU rDNA gene [11,12], whereas the pathogen identified as *Leishmania infantum* [13,14] or *Leishmania turanica* [15] from the same geographic region Karamay also could cause CL. However, *L. turanica* is nonpathogenic to humans, according to Strelkova *et al.* [16]. The parasites of some visceral leishmaniasis (VL) cases in Sichuan and Gansu provinces were *L. donovani* or undescribed species *Leishmania* sp. [17-19]. VL and CL have been reported in China to date the species of *Leishmania* comprises much more than that. The isolates in China were more heterogeneous than previously thought, requiring the reassignment of some isolates into different groups as described by Lu *et al.* [20].

Over the past few decades, DNA markers including coding and non-coding genes have become additional information for advancing our understanding of evolutionary and phylogenetic relationships and species differentiation. Data pertaining to the ribosomal RNA (rRNA) gene sequences—in particular, the two non-coding, highly variable internal

transcribed spacer regions (ITS1 and ITS2) are considered to be acceptable molecular criteria for resolving taxonomic questions and determining the phylogenetic affinities among closely related *Leishmania* species [17,21-25]. While coding genes are also widely used for taxonomic studies, such as metabolic enzymes (ICD, ME, MPI, G6PDH, ASAT, GPI, NH1, NH2, PGD and FH) [26,27], heat-shock protein 70 gene (hsp70) [4], cytochrome oxidase II (CO II) [28], the gene encoding the largest subunit of RNA polymerase II (rpoIIIS) [6,8], the glycoprotein 63 gene (*gp63*) [29], cysteine protease B genes (*cpb*) [30] and cytochrome *b* (*cyt b*) [31-35].

In previous ITS1 and CO II study [17,19], we summarized the four endemic *Leishmania* species in China: *L. donovani*, *L. infantum*, *Leishmania gerbilli*, and *L. turanica*. We also noted that there might be an undescribed *Leishmania* species endemic in China and highlighted that the isolate IPHL/CN/77/XJ771 from Bachu County, Xinjiang, is *L. donovani* instead of *L. infantum*. To elucidate the phylogeny, evolution and epidemiology of interesting group of strains in China, further studies of more genes are required.

Cyt b is one of the cytochromes involved in the electron transport process of the mitochondrial respiratory chain is considered one of the most useful genes for phylogenetic work [34]. Marco *et al.* proved that the *cyt b* gene sequencing can precisely identify the *Leishmania* spp. for all of the local stocks that are well characterized by multi-locus enzyme electrophoresis (MLEE), the current gold standard [32]. Phylogeny and sequence variation of the genus *Leishmania* has also been discussed successfully with *cyt b* sequencing [34,35]. In this paper, the *cyt b* gene of *Leishmania* from China was sequenced and analyzed using bioinformatics methods. Moreover, the phylogenetic relationships were reconstructed using *cyt b* sequences obtained by this study and download from the GenBank database. We then discuss in detail the implications of relationships between strains in China and other locations.

Methods

Leishmania isolates

A population of cloned promastigotes (including 16 Chinese *Leishmania* isolates and three isolates from other countries) was stored in liquid nitrogen, and kept at the Department of Parasitology, Western China School of Preclinical and Forensic Medicine, Sichuan University. All of the *Leishmania* isolates used in this study are listed in Table 1. The promastigotes were cultivated in medium 199 supplemented with 15% heat-inactivated fetal bovine serum (HIFBS) at 28°C. Approximately 1–5 × 10⁹ promastigotes were collected at room temperature by centrifugation at 3300 × g for 10 min and washed with phosphate-buffered saline.

Table 1 List of *Leishmania* strains, origin, and database accession numbers, including sequences of *Leishmania* retrieved from GenBank

| Sequence length (bp) | GenBank accession numbers | MLEE-based species assignment | WHO code | Origin | Reference |
|----------------------|---------------------------|-------------------------------|----------------------|-----------------|------------------------------|
| 1079 | HQ908255 ^a | <i>Leishmania</i> sp. | MHOM/CN/84/SD1 | Shandong, China | This study |
| 1078 | HQ908260 ^a | <i>Leishmania</i> sp. | MHOM/CN/90/SC10H2 | Sichuan, China | This study |
| 1079 | HQ908263 ^a | <i>Leishmania</i> sp. | MHOM/CN/89/GS6 | Gansu, China | This study |
| 1079 | HQ908264 ^a | <i>Leishmania</i> sp. | MHOM/CN/86/SC6 | Sichuan, China | This study |
| 1080 | HQ908271 ^a | <i>Leishmania</i> sp. | MHOM/CN/84/GS3 | Gansu, China | This study |
| 1080 | HQ908273 | <i>Leishmania</i> sp. | MHOM/CN/83/GS2 | Gansu, China | This study |
| 1079 | HQ908266 | <i>Leishmania</i> sp. | MHOM/CN/80/XJ801 | Xinjiang, China | This study |
| 1079 | HQ908272 | <i>Leishmania</i> sp. | MHOM/CN/90/SC11 | Sichuan, China | This study |
| 979 | HQ908268 | <i>Leishmania</i> sp. | MCAN/CN/86/SC9 | Sichuan, China | This study |
| 1079 | HQ908269 | <i>Leishmania</i> sp. | MHOM/CN/89/GS5 | Gansu, China | This study |
| 1080 | HQ908259 | <i>L. gerbilli</i> | MGER/CN/60/GS-GER20 | Gansu, China | This study |
| 1080 | HQ908256 | <i>L. turanica</i> | MRHO/CN/88/KXG-2 | Karamay, China | This study |
| 1079 | HQ908262 | <i>L. donovani</i> complex | MCAN/CN/60/GS1 | Gansu, China | This study |
| 1079 | HQ908261 | <i>L. donovani</i> complex | MHOM/CN/93/GS7 | Gansu, China | This study |
| 1079 | HQ908267 | <i>L. donovani</i> complex | IPHL/CN/77/XJ771 | Xinjiang, China | This study |
| 1060 | HQ908265 | <i>L. tropica</i> | MHOM/CN/84/JS1 | China | This study |
| 1080 | HQ908270 ^b | <i>L. tropica</i> | MHOM/SU/74/K27 | Soviet Union | This study |
| 1080 | HQ908257 ^b | <i>L. tropica</i> | - | - | This study |
| 1080 | AB095960 ^b | <i>L. tropica</i> | MHOM/SU/58/Strain OD | Soviet Union | Luyo-Acero et al., 2004 [35] |
| 1079 | AB095965 ^c | <i>L. garnhami</i> | MHOM/VE/76/JAP78 | Venezuela | Luyo-Acero et al., 2004 [35] |
| 1078 | HQ908258 ^c | <i>L. mexicana</i> | - | - | This study |
| 1079 | AB095957 ^d | <i>L. donovani</i> | MHOM/SD/62/2S-25M-C2 | Sudan | Luyo-Acero et al., 2004 [35] |
| 1079 | AB095958 ^d | <i>L. infantum</i> | MHOM/TN/80/IPT1 | Tunisia | Luyo-Acero et al., 2004 [35] |
| 1080 | AB095959 ^e | <i>L. chagasi</i> | MHOM/BR/74/PP75 | Brazil | Luyo-Acero et al., 2004 [35] |
| 872 | EF579896 ^e | <i>L. donovani</i> | MHOM/IN/80/DD8 | India | Foulet et al., 2007 [33] |
| 1079 | AB434677 | <i>L. archibaldi</i> | MHOM/ET/72/GEBRE1 | Ethiopia | Asato et al., 2009 [34] |
| 872 | EF579897 ^f | <i>L. chagasi</i> | MHOM/BR/74/PP75a | Brazil | Foulet et al., 2007 [33] |
| 872 | EF579913 ^f | <i>L. infantum</i> | MCAN/GR/94/CRE69 | Greece | Foulet et al., 2007 [33] |
| 1080 | AB095962 | <i>L. aethiopica</i> | MHOM/ET/72/L100 | Ethiopia | Luyo-Acero et al., 2004 [35] |
| 872 | EF579908 | <i>L. aethiopica</i> | MHOM/ET/72/L100 | Ethiopia | Foulet et al., 2007 [33] |
| 1080 | AB095970 | <i>L. major</i> | MHOM/EC/88/PT-115 | Ecuador | Luyo-Acero et al., 2004 [35] |
| 1080 | AB434675 | <i>L. turanica</i> | MRHO/SU/80/CLONE3720 | Soviet Union | Asato et al., 2009 [34] |
| 1080 | AB434674 | <i>L. arabica</i> | MPSA/SA/83/JISH220 | Saudi Arabia | Asato et al., 2009 [34] |
| 1104 | M97357 | <i>L. mexicana</i> | - | - | Lee et al., 1992 [36] |
| 1089 | M92829 | <i>L. mexicana</i> | - | - | Lee et al., 1992 [36] |
| 1078 | AB095964 | <i>L. amazonensis</i> | MHOM/BR/73/M2269 | Brazil | Luyo-Acero et al., 2004 [35] |
| 1079 | AB095963 ^g | <i>L. mexicana</i> | MHYC/BZ/62/M379 | Belize | Luyo-Acero et al., 2004 [35] |
| 872 | EF579909 ^g | <i>L. amazonensis</i> | LMAMPRO/BR/72/M1841 | Brazil | Foulet et al., 2007 [33] |
| 872 | EF579902 ^g | <i>L. amazonensis</i> | MHOM/BR/73/M2269 | Brazil | Foulet et al., 2007 [33] |
| 1078 | AB434678 | <i>L. aristidesi</i> | MORY/PA/69/GML | Panama | Asato et al., 2009 [34] |
| 1078 | M10126 | <i>L. tarentolae</i> | - | - | de la Cruz et al., 1984 [37] |
| 1078 | AB095966 | <i>L. braziliensis</i> | MHOM/BR/75/M2904 | Brazil | Luyo-Acero et al., 2004 [35] |
| 1078 | AB434682 | <i>L. braziliensis</i> | MHOM/BR/75/M2903 | Brazil | Asato et al., 2009 [34] |

Table 1 List of *Leishmania* strains, origin, and database accession numbers, including sequences of *Leishmania* retrieved from GenBank (Continued)

| | | | | | |
|------|-----------------------|---------------------------|-------------------|---------------|--------------------------------------|
| 1078 | AB095967 | <i>L. braziliensis</i> | MHOM/EC/88/INH-03 | Ecuador | Luyo-Acero <i>et al.</i> , 2004 [35] |
| 872 | EF579905 ^h | <i>L. guyanensis</i> | MHOM/GF/79/LEM85 | French Guiana | Foulet <i>et al.</i> , 2007 [33] |
| 872 | EF579912 ^h | <i>L. guyanensis</i> | MHOM/BR/75/M4147 | Brazil | Foulet <i>et al.</i> , 2007 [33] |
| 1078 | AB095968 | <i>L. panamensis</i> | MHOM/BR/71/LS94 | Brazil | Luyo-Acero <i>et al.</i> , 2004 [35] |
| 1078 | AB434680 | <i>L. shawi</i> | MHOM/BR/79/M15065 | Brazil | Asato <i>et al.</i> , 2009 [34] |
| 1080 | AB434686 | <i>L. equatorensis</i> | MCOH/EC/82/LSP-1 | Ecuador | Asato <i>et al.</i> , 2009 [34] |
| 1078 | M94286 | <i>Trypanosoma brucei</i> | - | - | Feagin <i>et al.</i> , 1987 [38] |

a, b, c, d, e, f, g, and h shared the same haplotype.

DNA extraction and polymerase chain reaction (PCR)

Total genomic DNA of the parasite was extracted by proteinase K digestion and phenol/chloroform/isoamyl alcohol extraction procedures followed by ethanol precipitation to purify the extracted DNA as described by Sambrook and Russell [39]. PCR was performed to generate a fragment spanning *cyt b* kinetoplast DNA (kDNA) between the forward primer COIIF (5'-TAAT ACGACTCACTATAGTTTATATTG ACATTTTGTWG ATT-3') and the reverse primer MURF4R (5'-GGGTTT TCCCAG TCACGACGAATCTCTCTCTCCCTT -3') [35]. The PCR protocols for amplification were: 94°C for 3 min followed by 35 cycles of 94°C for 30s, 58°C for 30s, and 72°C for 1.5 min, followed by a final elongation step at 72°C for 10 min. The amplified products were purified on a 2.0% agarose gel stained with ethidium bromide, using a commercial DNA purification kit according to the manufacturer's protocol. The purified PCR product was then sequenced. The DNA sequences of each individual and each species were deposited in the GenBank database under accession numbers (HQ908255-HQ908273).

Sequence alignment and analyses

A set of *cyt b* sequences of *Leishmania* were retrieved from GenBank, included 29 sequences of genus *Leishmania* (AB095957-AB095960, AB095962-AB095968, AB095970, EF579896-EF579897, EF579902, EF579905, EF579908, EF579909, EF579912, EF579913, AB434674, AB434675, AB434677, AB434678, AB434680, AB434682, M97357, M92829, M10126), *Leishmania equatorensis* (AB434686) and *Trypanosoma brucei* (M94286) (Table 1). The sequences were first aligned using Clustal X 1.83 [40] with a gap-opening penalty of 5 and gap-extension penalty of 1 following the small gap costs recommendation of Hickson *et al.* [41]. The aligned matrix from this procedure was verified to have the same length, and minor adjustments were then made manually using SeaView v.4.2.5 [42]. The data matrices are available from the corresponding author. The haplotype analyses were performed to 37 sequences using DAMBE software for 50 sequences [43]. Compositional heterogeneity was

evaluated using chi-square (χ^2) tests implemented in PAUP* 4.0b10 [44]. Distances from the predicted amino acid sequences were determined with the p-distance models which were computed by MEGA v. 4.1 [45].

Phylogenetic analyses

Phylogenetic hypotheses of *Leishmania* were generated with *cyt b* kDNA segments using two types of commonly applied phylogenetic techniques: heuristic searches using maximum parsimony (MP) analyses performed with the program PAUP* program and Bayesian inference (BI) using the MrBayes v.3.2 program [46]. In both MP and BI analyses, gaps were treated as missing data. For heuristic searches under parsimony, invariant characters were removed from the dataset. Each search involved 10 random additional replicates, one tree held at each step, with tree bisection and reconnectin branch swapping, steepest descent on, and a maximum of 10,000 saved trees. Non-parametric bootstrapping was used to generate phylogeny confidence values [47], with 1,000 pseudoreplicates using a heuristic tree search for each pseudoreplicate. *Trypanosoma brucei* (M94286) was used to root the trees.

Prior to Bayesian analyses, the best-fit model of evolution, TIM3 + G, was selected using jModeltest v. 0.1.1 [48] under the Bayesian information criterion [49], following recent recommendations [50]. We estimated the posterior probability distributions by allowing four incrementally heated Markov chains (default heating values) to proceed to four million generations, and with samples were taken every 200 generations. Analyses were repeated beginning with different starting trees to ensure that the analyses were not restricted from the global optimum [51]. Convergence was first tested by examining the average deviation of the split frequencies of the two runs, in order to determine whether the two runs had converged. MCMC convergence was also explored by examining the potential scale reduction factor (PSRF) convergence diagnostics for all model parameters (provided by the *sump* and *sumt* commands). The first one million generations before this chain became stationary were discarded, and the remaining samples from the independent runs were

pooled to obtain the final approximation of the posterior tree distribution.

Sequence alignments were additionally inferred from uncorrected p-distances through NJ networks (Neighbor Net) obtained by SplitsTree 4 [52,53]. This software can detect the alternative evolutionary paths supported by the sequence alignments, and as such, they do not enforce the single bifurcating dendrogram. To yield a single phylogeny hypothesis, the posterior distribution was summarized as a 50% majority rule consensus.

Results

Base composition and nucleotide substitution patterns

The size of the newly determined *cyt b* fragments is shown in the Table 1. Of the 911 aligned characters, 324 were variable, including 225 that were parsimony-informative. Percentage base compositions were as follows: A, 27.8; C, 7.2; G, 15.6; and T, 50.4. The average maximum likelihood estimated Ti/Tv ratio was 1.2.

A base stationarity test showed insignificant differences among the taxa in base composition bias in the data ($\chi^2 = 85.386150$, $df = 108$, $p = 0.94687017$). The p-distances among the 10 isolates (10 isolates, SD1, SC10H2, GS6, SC6, GS3, GS2, XJ801, SC11, SC9, GS5; the isolates SD1, SC10H1, GS6, SC6 and GS3 share the same sequence) in China were ranged from 0.000 to 0.023 (mean = 0.010), which are smaller than the distances between these isolate and any other known species. These 10 isolates were then classified into the *Leishmania* sp. group. The divergence between *Leishmania* sp. and other *Leishmania* species ranged from 0.051 (*Leishmania* sp. versus *L.tarentolae*) to 0.131 (*Leishmania* sp. vs. *L. turanica* and *L. gerbilli*), with an average of 0.096, a value that is larger than that within *Leishmania* sp. group. The *L. donovani* complex group contains seven haplotypes: GS1, GS7, XJ771, PP75a (the

same sequence with CRE69), 2S-25M-C2 (the same sequence shared with IPT1), PP75 (the same sequence with DD8). The distance within the strains of *L. donovani* complex ranged from 0.001 (the strain GS7 of China vs. *L. chagasi* PP75a and *L. infantum* CRE69) to 0.013 (*L. archibaldi* GEBRE1 vs. *L. chagasi* PP75 and *L. donovani* DD8), which are smaller than the distances between these strains and other known species. The average distance in this group as a whole is 0.006. The p distances among all species except *Leishmania* sp. and *T. brucei* were from 0.002 (between *L. turanica* and *L. gerbilli*) to 0.136 (between *Leishmania arabica* and *L. equatoresnsis*). Most pairwise comparisons mentioned above had divergence values < 0.136, with an average of 0.106 (Table 2).


Phylogenetic relationships

The heuristic search of the *cyt b* matrix resulted in 10,000 equally parsimonious trees of 15519 steps with high CI (0.6112) and RI (0.8427) values. In the strict consensus phylogram (Figure 1), three clades (BP = 88%) were formed. *Leishmania* sp., *L. tarentolae* and *L. braziliensis* complex form a clade (BP = 68%); six haplotypes in China formed a strongly cluster *Leishmania* sp. (BP = 88%), and clustered with *L. tarentolae* (BP = 99%); and next joined by the *L. braziliensis* complex (BP = 99%) containing three haplotypes of *L. braziliensis*, *L. guyanensis*, *L. panamensis* and *L. shawi*. The OWL clade (BP = 86%) clustered with the following OWL species: *L. donovani*, *L. major*, *L. tropica*, *L. arabica*, *L. turanica* and *L. gerbilli*. *Leishmania chagasi* (with the synonymous *L. infantum*), *L. donovani*, *Leishmania archibaldi* (with the synonymous *L. donovani*) [54], *L. infantum*, and the isolates GS7, GS1 and XJ771 of this study from China formed a monophyletic clade *L. donovani* complex. The strain JS1 from Jiangsu province of China clustered with *L. tropica* from the Soviet Union (BP = 100%)

Table 2 Pairwise genetic distances for *cytb* segments among *Leishmania* species

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
|----------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|----|
| 1 <i>Leishmania</i> sp. | - | | | | | | | | | | |
| 2 <i>L. tarentolae</i> | 0.051 | - | | | | | | | | | |
| 3 <i>L. braziliensis</i> complex | 0.095 | 0.095 | - | | | | | | | | |
| 4 <i>L. mexicana</i> complex | 0.104 | 0.106 | 0.107 | - | | | | | | | |
| 5 <i>L. donovani</i> complex | 0.110 | 0.113 | 0.102 | 0.095 | - | | | | | | |
| 6 <i>L. turanica</i> | 0.131 | 0.131 | 0.122 | 0.107 | 0.092 | - | | | | | |
| 7 <i>L. gerbilli</i> | 0.131 | 0.131 | 0.123 | 0.107 | 0.093 | 0.002 | - | | | | |
| 8 <i>L. arabica</i> | 0.130 | 0.130 | 0.124 | 0.106 | 0.101 | 0.085 | 0.085 | - | | | |
| 9 <i>L. tropica</i> complex | 0.116 | 0.119 | 0.113 | 0.095 | 0.081 | 0.088 | 0.088 | 0.095 | - | | |
| 10 <i>L. equatoresnsis</i> | 0.126 | 0.128 | 0.119 | 0.112 | 0.120 | 0.126 | 0.127 | 0.136 | 0.125 | - | |
| 11 <i>Trypanosoma brucei</i> | 0.177 | 0.176 | 0.150 | 0.158 | 0.158 | 0.183 | 0.183 | 0.185 | 0.176 | 0.155 | - |

Pairwise genetic distances between different groups are detected by MEGA4.




and then clustered with *L. aethiopica* (BP = 100%), which formed the *L. tropica* complex except *L. major* (BP = 89%). *L. turanica* from the Soviet Union clustered with strains GS-GER20 and KXG-2 (BP = 63%) in China from this study clustered together (BP = 100%), which next joined by *L. arabica* (BP = 94%) and *L. major* (BP = 54%). The clade NWL (BP = 100%) is clustered with *L. mexicana*, *L. garnhami*, *L. amazonensis*, and *Leishmania aristides*. Information about the strains is shown in Table 1.

For the BI analyses, the likelihood value of the 50% majority consensus tree (Figure 2) was $\ln L = -4132.1156$, while the average PSRF was 1.001. The topology of the BI tree is a little different from that of MP tree. Two robust clades were formed. In one robust clade (PP = 0.95), *Leishmania* sp. (PP = 0.99) was a sister of *L. tarentolae* (PP = 1.00) and forms one clade with the species of *L. (Leishmania)* (LL), the other clade was consis of the species of *L. (Viannia)* (LV) (PP = 1.00)

(see Figure 3). The group of *Leishmania* sp. and *L. tarentolae* was basal to subgenus *Leishmania* and *L. equatorensis*. *L. mexicana*, *L. garnhami*, *L. amazonensis*, and *L. aristidesi* formed the *L. mexicana* complex (PP = 1.00). The *L. mexicana* and *L. equatorensis* are fundamental to all remaining subgenus *Leishmania* species. Within the other members of subgenus *Leishmania*, the *L. donovani* complex is sister to the clade (PP = 1.00) that clustered with *L. tropica* (PP = 1.00) plus *L. aethiopica* (PP = 1.00), and the clade (PP = 0.97) that consists of *L. turanica*, *L. gerbilli* and *L. arabica*, next joined by *L. major* (PP = 1.00).

In addition to the common phylogenetic relationships among the different species shown by the MP tree and BI tree, the network (Figure 3) calculated by SplitsTree 4 also indicated a clear evolutionary path with a high value. *Leishmania* sp. and *L. tarentolae* share most of their evolutionary paths.



Discussion

As a part of worldwide *Leishmania* population, the phylogenetics of Chinese isolates with analysis of the *cyt b* genetic sequences of 16 *Leishmania* isolates was discussed in this paper which demonstrated similarities and differences compared with previous data [17,19] and keep the genus evolutionary unity and integrity over large geographic ranges and time periods.

Leishmania sp. of China

Most interestingly 10 Chinese strains, representing 6 closely related haplotypes, could not be assigned to any of the so far described species of *Leishmania*, a finding that is congruent with our earlier ITS1 and COII studies [17,19]. These *Leishmania sp.* isolates were most closely related to the lizard-infecting *L. tarentolae* (Figures 1, 2, 3).

It was reported that one of these isolates SC6 was collected from patients with VL in Nanping County of Sichuan Province, was infected successfully 8 dogs (8/12) and its amastigotes were detected in their bone marrow smears [55]. Another isolate SC10H2 was proved that it clustered with the pathogen of canine leishmaniasis in Beichuan County, Sichuan Province, China based on the 17S RNA gene [18]. The non-pathogenic to humans *L. tarentolae* has been classified as subgenus *L. (Sauroleishmania)* on the basis of biological

criteria and different genes [4,8,10,34]. In such cases, we can conclude that the undescribed *Leishmania* species which is clearly a causative agent of canine leishmaniasis and human VL do exist in China are related to the *Sauroleishmania*. However, the more lizard parasites are required to confirm whether *Leishmania sp.* is assigned to the *Sauroleishmania*.

The pairwise genetic distance analysis (Table 2) and phylogenetic network (Figure 3) suggest that the *cyt b* sequences of the Chinese/*tarentolae* group (*Leishmania sp.* and *L. tarentolae*) are closer to the *Viannia* clade than the older world *Leishmania*. This finding is in contrast to that of our ITS1 study [17] and other studies: as an OWL species branching from within New World taxa, *L. tarentolae* (*Sauroleishmania*) are closer to the *Leishmania* subgenus than to the *Viannia* subgenus based on different DNA marks (polA and RNA polymerase II, 7SL RNA, hsp 70) [4,8,10]. It is well known that different genes can have different evolutionary histories and be influenced by selection and horizontal gene transfer, and the phylogenies are also prone to sampling bias; therefore, more genes of diverse geographic original strains would be needed to elucidate the phylogeny, evolution, and epidemiology of the Chinese/*tarentolae* group.

The isolates of *Leishmania sp.* were collected from different foci (plain, desert and hill), and the longest




Figure 3 The phylogenetic network of the *Leishmania* *cyt b* sequences shown in Table 1 and Figures 1, 2 was built with 1000 bootstrap replicates. It was algorithm, excluding all conserved site. Distances were calculated using the Kimura 2-parameter distance. *Trypanosoma brucei* (M94286) is the outgroup. Each A-F panel is drawn to the scale indicated and expressed as dissimilarity per nucleotide counted over variable sites (Figures 1–2) in *cyt b* alignment. The dots indicate the sequence position in the network. **A:** Complete network with representation of the five groups shown in detail in the remaining panels; the whole network excluding the *L. equatorensis* and *T. brucei* is divided into four segments: OWL, NWL, the *Leishmania* sp. and *L. tarentolae* group, and NWV. **B:** The *L. donovani* complex includes three Chinese isolates. **C:** Includes the species *L. tropica*, *L. aethiopica*, *L. major*, *L. turanica*, *L. gerbilli*, and *L. arabica*. **D:** The New World *Leishmania* subgenus. **E:** The *Leishmania* sp. of this study. **F:** The *L. (Viannia)* subgenus. Strain information is shown in Table 1.

distance between isolates is more than 2000 miles (from Shandong to the Xinjiang) (Figure 4). Meanwhile, different species were found in the same area. The isolate XJ801 of *Leishmania* sp. is from Kashi city of Xinjiang. The isolate 801 identified as *L. donovani* based on ITS1 sequences by Wang et al. [16] and Yang et al. [17] is also

from Kashi city. Another two isolates MHOM/CN/76/BT013 and MHOM/CN/81/812 which is differs from *L. donovani* (PHON/CN/77/771), *L. turanica* (MRHO/CN/88/KXG-2) and *L. gerbilli* (MRHO/CN/62/1) based on polymorphisms in both kinetoplast (kDNA) and nuclear (nDNA) DNAs that also collected from the same area

Kashi [20]. As such, the *Leishmania* isolates in China were more heterogeneous, further epidemiologic survey and more strains are required in Kashi.


L. donovani complex of China

Analysis in the current study revealed that the *cyt b* sequences of GS1, GS7 and XJ771 clustered with other species of *L. donovani* complex (PP = 1.00). On the basis of MLEE of the representative isolates from the plain, hill, and desert regions of China, Xu *et al.* were first to identify the causative agents responsible of VL as *Leishmania donovani sensu lato* and *L. infantum* [56]. The results based on sequences of *cyt b*, ITS1 [17] and COII [19] sequences clearly proved the existence of *L. donovani* in China. However, *L. donovani* or *L. infantum* standard isolates cannot be distinguished from *L. donovani* complex isolate using the *cyt b* gene in the BI and MP trees. These findings aren't consistent the ITS1 study showing three isolates clustered with *L. donovani* and a clear classification within subspecies between

Leishmania donovani donovani and *Leishmania donovani infantum*. Therefore, the inter-specific variation of the ribosomal RNA gene ITS1 was inferred to be more suitable than mtDNA segment *cyt b* for studying the phylogenetic relationships among subspecies. Of course, we can't exclude the possibility that the different inter-specific variation between ITS1 and *cyt b* are calculated by choosing the different samples or numbers of the isolates or strains.

L. turanica of China

Our *cyt b* data demonstrate that the isolates KXG-2 and GS-GER20 clustered with *L. turanica* (AB434675) from central Asia, findings that are congruent with those of our earlier studies [17,19] and then clustered with *L. arabica* from western Asia, a finding that agrees with that of Asato *et al.* [34] (Figures 1, 2, 3). The definitive hosts of *L. gerbilli*, *L. turanica*, *L. arabica* are rodents of the Old World [57]. Using MLEE methods, the isolate KXG-2 was identified as *L. turanica* [16], and the isolate GS-GER20 was



identified as *L. gerbilli* [58]. In the 1990s, *L. turanica* and *L. gerbilli* were identified in rodents or sandflies in Karamay, Xinjiang and *L. turanica* was proved to be pathogenic in both monkeys and humans in the laboratory, *Phlebotomus mongolensis* and *Phlebotomus andrejevi* were its major vectors [16]. We considered the isolates KXG-2 and GS-GER20 to be *L. turanica* and *L. gerbilli*, respectively via the *cyt b* gene sequences.

L. tropica of China

The species of the *L. tropica* complex cause the urban form of Old World CL. In Iran, Iraq, and India, it is transmitted by *Phlebotomus papatasi*. This species is rarely reported in China. The fact that the isolate JS1 was collected from Jiangsu Province clustered with *L. tropica*, which agrees with the results of our earlier study based on the COII gene [19]. Lu *et al.* used random amplified polymorphic DNA data to suggest a close relationship between the isolate JS1 and *L. tropica* (K27) [59]. Thus, we infer that the isolate JS1 may be *L. tropica*. However to further confirm this inference, more data such as host specificity, life cycle, and biochemical analysis will be needed.

Evolution inference and epidemiology of China

In our analysis, *Leishmania* *cyt b* sequences are consistent with the genus *Leishmania* that contains three subgenera: *Leishmania*, *Sauroleishmania* and *Viannia* [4]. Based on the suggestion that mammalian *Leishmania* did not evolve from those of lizards but vice versa [60,61], Lukeš *et al.* [54] proposed that the ancestor of the new world *Leishmania* evolved in South America and then migrated via the Bering land bridge to Asia via multiple independent genetic loci. The *Leishmania* lineage would have been dispersed throughout central and/or Southeast Asia, where a major diversification gave rise to *L. aethiopica*, *L. major*, *L. gerbilli*, *L. turanica*, *L. tropica*, and the *L. donovani* complex. The isolates from China were absent in this analysis. However Fraga *et al.* thought this theory puts *L. tarentolae* (*Sauroleishmania*) in an illogical position. Our data suggest that *Leishmania* sp. of the pathogen of VL and CanL clustering with *L. tarentolae* (*Sauroleishmania*) was in the same “illogical position”. The maximum parsimony consensus tree (Figure 1) and splittree (Figure 3) supports the idea of a common origin with the *Viannias* subgenus, whereas the Bayesian tree (Figure 2) show the Chinese/tarentolae group clustered together with species of *Leishmania* subgenus. This ambiguous position of *L. tarentolae* had been discussed by Luyo-Acero *et al.* based on the same DNA marker *cyt b* [35] that *L. tarentolae* clustered with *Viannia* in the NJ tree consisting with the minicircle phylogenetic analysis [62], and clustered with *Leishmania* in the MP tree supported by ATPase 6 gene [63]. The position of

L. equatorensis as falling outside the *Leishmania* clade in the parsimony tree is supported by the phylogeny suggested by Cupolillo *et al.* [64]. However the Chinese/tarentolae group which was not described by Lukeš *et al.* [54], may have evolved from a common ancestral parasite that came from the Americas and may split off earlier than the other OWL.

Leishmaniasis remains endemic in China, especially in the west and northwest frontier regions. The epidemic foci of VL in China were classified into three types according to different geographical origin, infective agent, and clinical evidences, i.e., plain foci, hill foci, and desert foci [20]. Human VL and CL occur in China, most being VL along with rare CL cases [56,65-67]. VL was one of the most important parasitic diseases occurring in over 17 Chinese provinces in 1951 [68]. Since the condition has come under control, currently, VL is mainly prevalent in six provinces in northwest China [69] (Figure 4). This study proved that the evolution hypothesis of Tian and Chen related to the Chinese *Leishmania* isolates from different epidemic foci was limited and lacked integrity [70]. In fact, the Chinese *Leishmania* species occurs as the multiple species *L. donovani* (*L. donovani donovani*, *L. donovani infantum*), *L. turanica*, *L. tropic*, *Leishmania* sp. and so forth, and some of these such as *L. donovani* and *L. turanica* were shared with neighbouring countries including India, Russia, and Uzbekistan.

Conclusions

The current study investigated the Chinese *Leishmania* parasites using *cyt b* sequence data. Undescribed *Leishmania* species which are clearly causative agents of CanL and human VL do exist in China and are related to the *Sauroleishmania* subgenus, may have evolved from a common ancestral parasite that came from the Americas and split off earlier than the other OWL. Our *cyt b* results also suggest the following: the isolates GS7, GS1 and XJ771 occur as part of the *L. donovani* complex; the isolate JS1 is *L. tropica*; and the isolate KXG-2 is close to the isolate GS-GER20, which is *L. turanica* and *L. gerbilli* respectively. The results of the current study indicate that the isolates from China may have had a more complex evolutionary history. In the future, we will build upon the currently described data set to gain more insight into the fascinating spectrum of Chinese *Leishmania*.

Competing interests

The authors have no competing interests to declare.

Authors' contributions

JPC and BBY conceived, designed and coordinated the field study, while DLC, LL, XSH and JNX participated in the study design and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by the National Natural Science Foundations of China (NO: 30771883, 30800094 and 81171607) and the Chinese Important Scientific Research Project on Infectious Diseases (NO 2008-ZX10004-011).

Author details

¹Department of Medical Laboratory, Weifang Medical University, #7166, The West Baotong street, Weifang, Shandong 261053, China. ²Department of Parasitology, West China School of Preclinical and Forensic Medicine, Sichuan University, #17, The 3th Section of South Renmin Road, Chengdu, Sichuan 610041, China. ³Animal Disease Prevention and Food Safety Key Laboratory of Sichuan Province, #24, The South 1st Section of Yihuan Road, Chengdu, Sichuan 610041, China.

Received: 5 April 2012 Accepted: 20 January 2013

Published: 5 February 2013

References

- World Health Organization: **Control of the leishmaniase**. In *Proceedings of a meeting of the WHO Expert Committee, Geneva, 22–26 March 2010*. Geneva: World Health Organization; 2010:22–26.
- Shaw JJ: **Taxonomy of the genus *Leishmania*: Present and future trends and their implications**. *Mem Inst Oswaldo Cruz* 1994, **89**:471–478.
- World Health Organization: **Control of the leishmaniases**. *Proceedings of a meeting of the WHO Expert Committee, Geneva, 1–15 March 1990*. Geneva: World Health Organization; 1990.
- Fraga J, Montalvo AM, de Doncker S, Dujardin J-C, der Auwera GV: **Phylogeny of *Leishmania* species based on the heat-shock protein 70 gene**. *Infect Genet Evol* 2010, **10**:238–245.
- Lainson R, Shaw JJ: **Evolution, classification and geographical distribution of *Leishmania***. In *The Leishmaniasis in Biology and Medicine*. Edited by Killick W, Kendrick R.: London Academic Press; 1987:1–120.
- Dujardin JC, Henriksson J, Victoir K, Brisse S, Gamboa D, Arevalo J, Le Ray D: **Genomic Rearrangements in *Trypanosomatids*: an Alternative to the 'One Gene' Evolutionary Hypotheses?** *Mem Inst Oswaldo Cruz* 2000, **95**:527–534.
- Safjanova VM: **The problem of taxonomy with *Leishmania***. *Ser Protozool Sov Acad Sci Leningr* 1982, **7**:5–109.
- Croan DG, Morrison DA, Ellis JT: **Evolution of the genus *Leishmania* revealed by comparison of DNA and RNA polymerase gene sequences**. *Mol Biochem Parasitol* 1997, **89**:149–159.
- Orlando TC, Rubio MA, Sturm NR, Campbell DA, Floeter-Winter LM: **Intergenic and external transcribed spacers of ribosomal RNA genes in lizard-infecting *Leishmania*: molecular structure and phylogenetic relationship to mammal-infecting *Leishmania* in the subgenus *Leishmania*(*Leishmania*)**. *Mem Inst Oswaldo Cruz* 2002, **97**:695–701.
- Zelazny AM, Fedorko DP, Li L, Neva FA, Fisher SH: **Evaluation of 7SL RNA gene sequences for the identification of *Leishmania* spp.** *Am J Trop Med Hyg* 2005, **72**:415–420.
- Chen JP, Hu XS, Jing BQ, Yang WT: **Studies of SSUrDNA sequence homologies of *Leishmania* species causing cutaneous leishmaniasis in Xinjiang, China by PCR and hybridization**. *Acta Parasitol Med Entomol Sin* 1996, **3**:89–92.
- Zhang T, Hu XS, Jing BQ, Zhen XL, Ma Y, Li F: **PCR Amplification and Cloning of SSU rDNA Variable Region of Cutaneous Leishmaniasis Pathogen from Xinjiang**. *China J Fujian Med Univ* 1998, **32**:1–5.
- Ren HY, Guan LR, Yang YQ, Zuo XP, Li F, Xu YX, Abulikemujiang, Deng LQ: **Clinicopathologic analysis on ninety cases of cutaneous leishmaniasis in Karamay region of Xinjiang**. *Chin J Parasitol Parasit Dis* 1996, **14**:226–230.
- Wang JY, Qu JQ, Wang RQ, Guan LR, Ren HY, Chang KP: **Analysis on homology in several isolates of *Leishmania* from Karamay Xinjiang**. *Chin J Parasitol Parasit Dis* 1996, **14**:266–269.
- Guan LR, Yang YQ, Xu YX, Wu JT: **Leishmaniasis in Karamay XI. The development of cutaneous Leishmaniasis in monkey and man experimentally infected with *Leishmania* from Karamay big gerbil**. *Bull Dis Control Prev* 1992, **10**:263–266.
- Strelkova MV, Shurkhal AV, Kellina OI, Eliseev LN, Evans DA, Peters W, Chapman CJ, Le Blancq SM, van Eys GJ: **A new species of *Leishmania* isolated from the great gerbil *Rhombomys opimus***. *Parasitology* 1990, **101**:327–335.
- Yang BB, Guo XG, Hu XS, Zhang JG, Liao L, Chen DL, Chen JP: **Species discrimination and phylogenetic inference of 17 Chinese *Leishmania* isolates based on internal transcribed spacer 1 (ITS1) sequences**. *Parasitol Res* 2010, **107**:1049–1065.
- Sun K, Guan W, Zhang JG, Wang YJ, Tian Y, Liao L, Yang BB, Chen DL, Chen JP: **Prevalence of canine leishmaniasis in Beichuan County, Sichuan, China and phylogenetic evidence for an undescribed *Leishmania* sp. in China based on 7SL RNA**. *Parasit Vectors* 2012, **5**:75.
- Cao DP, Guo XG, Chen DL, Chen JP: **Species delimitation and phylogenetic relationships of Chinese *Leishmania* isolates reexamined using kinetoplast cytochrome oxidase II gene sequences**. *Parasitol Res* 2011, **109**:163–173.
- Lu HG, Zhong L, Guan LR, Qu JQ, Hu XS, Chai JC, Xu ZB, Wang CT: **Separation of Chinese *Leishmania* isolates into five genotypes by kinetoplast and chromosomal DNA heterogeneity**. *Am J Trop Med Hyg* 1994, **50**:763–770.
- Dávila AMR, Memen H: **Internaltranscribedspacer (ITS) sequences used to explore phylogenetic relationships within *Leishmania***. *Ann Trop Med Parasitol* 2000, **94**:651–654.
- Schönian G, Akuffo H, Lewin S, Maasho K, Nylén S, Pratlong F, Eisenberger CL, Schnur LF, Presber W: **Genetic variability within the species *Leishmania aethiops* does not correlate with clinical variations of cutaneous leishmaniasis**. *Mol Biochem Parasitol* 2000, **106**:239–248.
- El Tai NO, El Fari M, Mauricio I, Miles MA, Oskam L, El Safi SH, Presber WH, Schönian G: ***Leishmania donovani*: intraspecific polymorphisms of Sudanese isolates revealed by PCRbased analyses and DNA sequencing**. *Exp Parasitol* 2001, **97**:35–44.
- Kuhls K, Mauricio IL, Pratlong F, Presber W, Schönian G: **Analysis of ribosomal DNA internal transcribed spacer sequences of the *Leishmania donovani* complex**. *Microbes Infect* 2005, **7**:1224–1234.
- Parvizi P, Moradi G, Akbari G, Farahmand M, Ready PD, Piazak N, Assmar M, Amirkhani A: **PCR detection and sequencing of parasite ITSrDNA gene from reservoirs host of zoonotic cutaneous leishmaniasis in central Iran**. *Parasitol Res* 2008, **103**:1273–1278.
- Mauricio IL, Yeo M, Baghaei M, Doto D, Pratlong F, Zemanova E, Dedet JP, Lukes J, Miles MA: **Towards multilocus sequence typing of the *Leishmania donovani* complex: Resolving genotypes and haplotypes for five polymorphic metabolic enzymes (ASAT, GPI, NH1, NH2, PGD)**. *Int J Parasitol* 2006, **36**:757–769.
- Zemanová E, Jirků M, Mauricio IL, Horák A, Miles MA, Lukeš J: **The *Leishmania donovani* complex: Genotypes of five metabolic enzymes (ICD, ME, MPI, G6PDH, and FH), new targets for multilocus sequence typing**. *Int J Parasitol* 2007, **37**:149–160.
- Ibrahim ME, Barker DC: **The origin and evolution of the *Leishmania donovani* complex as inferred from a mitochondrial cytochrome oxidase II gene sequence**. *Infect Genet Evol* 2001, **1**:61–68.
- Mauricio IL, Gaunt MW, Stothard JR, Miles MA: **Glycoprotein 63 (gp63) genes show gene conversion and reveal the evolution of Old World *Leishmania***. *Int J Parasitol* 2007, **37**:565–576.
- Hidde M, Bras-Goncalves R, Bañuls AL: **Specific cpb copies within the *Leishmania donovani* complex: evolutionary interpretations and potential clinical implications in humans**. *Parasitology* 2007, **134**:379–389.
- Marco JD, Uezato H, Mimori T, Barroso PA, Korenaga M, Nonaka S, Baosombrio MA, Taranto NJ, Hashiguchi Y: **Are cytochrome b gene sequencing and polymoraw chain reaction as reliable as multilocus enzyme electrophoresis for identifying *Leishmania* sp from Argentina?** *Am J Trop Med Hyg* 2006, **75**:256–260.
- Marco JD, Bhutto AM, Soomro FR, Baloch JH, Barroso PA, Kato H, Uezato H, Katakura K, Korenaga M, Nonaka S, Hashiguchi Y: **Multilocus enzyme electrophoresis and cytochrome b gene sequencingbased identification of *Leishmania* isolates from different foci of cutaneous leishmaniasis in Pakistan**. *Am J Trop Med Hyg* 2006, **75**:261–266.
- Foulet F, Botterel F, Buffet P, Morizot G, Rivollet D, Deniau M, Pratlong F, Costa JM, Bretagne S: **Detection and identification of *Leishmania* species from clinical specimens by using a realtime PCR assay and sequencing of the cytochrome B gene**. *J Clin Microbiol* 2007, **45**:2110–2115.
- Asato Y, Oshiro M, Myint CK, Yamamoto Y, Kato H, Marco JD, Mimori T, Gomez EA, Hashiguchi Y, Uezato H: **Phylogenetic analysis of the genus *Leishmania* by cytochrome b gene sequencing**. *Exp Parasitol* 2009, **121**:352–361.
- LuyoAcero GE, Uezato H, Oshiro M, Takei K, Kariya K, Katakura K, GomezLandires E, Hashiguchi Y, Nonaka S: **Sequence variation of the cytochrome b gene of various human infecting members of the genus *Leishmania* and their phylogeny**. *Parasitology* 2004, **128**:483–491.

36. Lee ST, Tarn C, Wang CY: Characterization of sequence changes in kinetoplast DNA maxicircles of drugresistant *Leishmania*. *Mol Biochem Parasitol* 1992, **56**:197–207.
37. de la Cruz VF, Neckelmann N, Simpson L: Sequences of six genes and several open reading frames in the kinetoplast maxicircle DNA of *Leishmania tarentolae*. *J Biol Chem* 1984, **259**:15136–15147.
38. Feagin JE, Jasmer DP, Stuart K: Developmentally regulated addition of nucleotides within apocytocrome *b* transcripts in *Trypanosoma brucei*. *Cell* 1987, **49**:337–345.
39. Sambrook J: *Russell D W: Molecular cloning: a laboratory manual*. New York: Cold Spring Harbor Laboratory Press; 2001.
40. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG: The Clustal X Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 1997, **25**:4876–4882.
41. Hickson RE, Simon C, Perrey SW: The performance of several multiple sequence alignment programs in relation to secondary structure features for an rRNA sequence. *Mol Biol Evol* 2000, **17**:530–539.
42. Gouy M, Guindon S, Gascuel O: SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 2010, **27**:221–224.
43. Xia X, Xie Z: DAMBE: data analysis in molecular biology and evolution. *J Hered* 2001, **92**:371–373.
44. Swofford DL: *PAUP* Phylogenetic analysis using parsimony (* and other methods)*. version 4 Sinauer, MA: Sunderland; 2002.
45. Tamura K, Dudley J, Nei M, Kumar S: MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 2007, **24**:1596–1599.
46. Ronquist F, Huelsenbeck JP: MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 2003, **19**:1572–1574.
47. Felsensten JP: Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 1985, **39**:783–791.
48. Posada D: j ModelTest: phylogenetic model averaging. *Mol Biol E* 2008, **25**:1253–1256.
49. Schwarz G: Estimating the dimension of a model. *Ann Stat* 1978, **6**:461–464.
50. Posada D, Buckley TR: Model selection and model averaging in phylogenetics: advantages of akaike information criterion and Bayesian approach. *Syst Biol* 2004, **53**:793–808.
51. Huelsenbeck JP, Larget B, Miller RE, Ronquist F: Potential applications and pitfalls of Bayesian inference of phylogeny. *Syst Biol* 2002, **51**:673–688.
52. Huson DH: SplitTree: analyzing and visualizing evolutionary data. *Bioinformatics* 1998, **14**:68–73.
53. Huson DH, Bryan D: Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol* 2006, **23**:254–267.
54. Lukeš J, Mauricio IL, Schönián G, Dujardin JC, Soteriadou K, Dedet JP, Kuhls K, Tintaya KWQ, Jirků M, Chocholová E, Haralambous C, Pralong F, Oborník M, Horák A, Ayala FJ, Miles MA: Evolutionary and geographical history of the *Leishmania donovani* complex with a revision of current taxonomy. *Proc Natl Acad Sci USA* 2007, **104**:9375–9380.
55. Hu XS, Yang WT, Lu HG, Yan HP, Chen JP, Ma Y, Jing BQ, Zhang T: Sequencing a specific kinetoplast DNA fragment of *Leishmania donovani* for polymerase chain reaction amplification in diagnosis of Leishmaniasis in bone marrow and blood samples. *J Parasito* 2000, **84**:822–826.
56. Xu ZB, Le Blancq S, Evans DA, Peters W: The characterization by isoenzyme electrophoresis of *Leishmania* isolated in the People's Republic of China. *Trans R Soc Trop Med Hyg* 1984, **78**:689–693.
57. Ashford RW: The leishmaniasis as emerging and reemerging zoonoses. *Int J Parasitol* 2000, **30**:1269–1281.
58. Guan LR, Yang YQ, Ren HY, Shen WX: Discovery and study of *Leishmania turanica* for the first time in China. *Bull World Health Organ* 1995, **69**:595–601.
59. Lu DM, Hu XS, Qiao ZD: Analysis of *Leishmania* species and strains from China by RAPD technique. *Chin J Parasitol Parasit Dis* 2001, **19**:290–293.
60. Shaw JJ: Ecological and evolutionary pressures on leishmanial parasites. *Braz J Gen* 1997, **20**:123–128.
61. Noyes H: Implications of a Neotropical origin of the genus *Leishmania*. *Memórias do Instituto Oswaldo Cruz* 1998, **93**:657–661.
62. Yurchenko V, Kolesnikov A, Lukes J: Phylogenetic analysis of Trypanosomatina (Protozoa: Kinetoplastida) based on minicircle conserved regions. *Folia Parasitol* 2000, **47**:1–5.
63. Brewster S, Barker DC: The ATPase subunit 6 gene sequence predicts that RNA editing is conserved between lizard-and human-infecting *Leishmania*. *Gene* 1999, **235**:77–84.
64. Cupolillo E, Medina-Acosta E, Noyes H, Momen H, Grimaldi G Jr: A revised classification for *Leishmania* and *Endotrypanum*. *Parasitol Today* 2000, **16**:142–144.
65. Guan LR, Yang YQ, Ren HY, Chai JJ: Eight cutaneous Leishmaniasis cases detected in Karamay, Xinjiang Uygur Autonomous Region, China. *Southeast Asian J Trop Med Public Health* 1992, **23**:803–804.
66. Zheng CJ, Wang LY, Xu X, Zhu XH, Wu WP: Visceral Leishmaniasis in China during 2004–2007. *Chin J Parasitol Parasit Dis* 2009, **27**:344–346.
67. Chen J, Xu MJ, Zhou DH, Song HQ, Wang CR, Zhu XQ: Canine and feline parasitic zoonoses in China. *Parasit Vectors* 2012, **5**:152.
68. Wang JY, Cui G, Chen HT, Zhou XN, Gao CH, Yang YT: Current epidemiological profile and features of visceral leishmaniasis in People's Republic of China. *Parasit Vectors* 2012, **5**:31.
69. Zhou XN, Lv S, Yang GJ, Kristensen TK, Bergquist NR, Utzinger J, Malone JB: Spatial epidemiology in zoonotic parasitic diseases: insights gained at the 1st International Symposium on Geospatial Health in Lijiang, China, 2007. *Parasit Vectors* 2009, **2**:10.
70. Tian Y, Chen JP: Cloning and sequence analysis of ITS gene of *Leishmania donovani* isolates from different epidemic foci in China. *J Biomed* 2005, **22**:540–544.

doi:10.1186/1756-3305-6-32

Cite this article as: Yang et al.: Analysis of kinetoplast cytochrome *b* gene of 16 *Leishmania* isolates from different foci of China: different species of *Leishmania* in China and their phylogenetic inference. *Parasites & Vectors* 2013 **6**:32.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

