



POSTER PRESENTATION

Open Access

Disproportionately enlarged subarachnoid space hydrocephalus presence in patients with idiopathic normal pressure hydrocephalus

Tomas Radovnický^{1*}, Daniel Adamek², Milous Derner², Martin Sames¹

From Hydrocephalus 2015
Banff, Canada. 18-21 September 2015

Introduction

Idiopathic normal pressure hydrocephalus (iNPH) is a treatable disease of an elderly people. It's known for 50 years, but there are still some controversies, especially in the diagnostic algorithm. Disproportionately enlarged subarachnoid space hydrocephalus (DESH) seems to be important radiological feature of iNPH, but it's not present in all patients. The aim of our study was to analyse appearance of DESH in our set of patients.

Methods

We retrospectively analysed 1, 5 T MRI in 27 iNPH patients before surgery and MRI in 24 healthy controls. Evaluation was performed by neurosurgeon and radiologist independently and blindly. We assessed tight high convexity and medial subarachnoid space, dilatation of Sylvian fissure and focal dilatation of sulci. iNPH patients were identified by clinical examination, dilatation of ventricles on MRI defined by Evans' ratio > 0, 30 and positive tap test and/or lumbar infusion test. Patient outcome was measured by iNPH grading scale one year after surgery.

Results

In the group of 27 iNPH patients, we have found DESH presence in 15 cases (55, 6%), all 15 patients were shunt responders. In the group of 12 iNPH patients without DESH 4 of them did not respond to surgery (33, 3%).

Among the 24 healthy controls, we haven't found any DESH appearance.

Conclusion

We confirmed, that DESH appearance on MRI is supportive for the iNPH diagnosis, but it should not be used as a single predictor. Patients without DESH are at a higher risk to be shunt non-responders. Indication for shunt surgery should be still based on correlation between the radiological and clinical evaluation and supplementary CSF dynamics tests.

Supported by IGA NT14448-3/2013.

Authors' details

¹Department of Neurosurgery, Masaryk Hospital, Usti nad Labem, Czech Republic. ²Department of Radiology, Masaryk Hospital, Usti nad Labem, Czech Republic.

Published: 18 September 2015

doi:10.1186/2045-8118-12-S1-P43

Cite this article as: Radovnický et al.: Disproportionately enlarged subarachnoid space hydrocephalus presence in patients with idiopathic normal pressure hydrocephalus. *Fluids and Barriers of the CNS* 2015 12 (Suppl 1):P43.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: radovnický@gmail.com

¹Department of Neurosurgery, Masaryk Hospital, Usti nad Labem, Czech Republic

Full list of author information is available at the end of the article