

2,4-Dichlorobenzaldehyde 2,4-dinitrophenylhydrazone

Feng-yu Bao

Department of Applied Chemistry, College of Sciences, Henan Agricultural University, Zhengzhou 450002, People's Republic of China
Correspondence e-mail: bfyu2008@yahoo.com.cn


Received 4 June 2008; accepted 2 July 2008

Key indicators: single-crystal X-ray study; $T = 291\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$;
 R factor = 0.062; wR factor = 0.188; data-to-parameter ratio = 12.2.

The asymmetric unit of the title compound, $\text{C}_{13}\text{H}_8\text{Cl}_2\text{N}_4\text{O}_4$, contains two independent but similar and almost planar molecules. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond is observed in each molecule.

Related literature

For background, see: Okabe *et al.* (1993); Ohba (1996).



Experimental

Crystal data

$\text{C}_{13}\text{H}_8\text{Cl}_2\text{N}_4\text{O}_4$
 $M_r = 355.14$

Monoclinic, $P2_1/c$
 $a = 13.3814(7)\text{ \AA}$

$b = 28.9980(13)\text{ \AA}$
 $c = 7.3996(3)\text{ \AA}$
 $\beta = 92.422(4)^\circ$
 $V = 2868.7(2)\text{ \AA}^3$
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 0.48\text{ mm}^{-1}$
 $T = 291(2)\text{ K}$
 $0.16 \times 0.11 \times 0.09\text{ mm}$

Data collection

Bruker SMART APEX CCD
diffractometer
Absorption correction: multi-scan
(SADABS; Bruker, 1998)
 $R_{\text{int}} = 0.092$, $T_{\text{min}} = 0.927$, $T_{\text{max}} = 0.958$

21827 measured reflections
5060 independent reflections
2794 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.092$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$
 $wR(F^2) = 0.187$
 $S = 1.02$
5060 reflections

415 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.70\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.36\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N6—H6A \cdots O3	0.86	2.05	2.645 (5)	126
N8—H8A \cdots O8	0.86	2.03	2.627 (5)	126

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2743).

References

- Bruker (1998). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Ohba, S. (1996). *Acta Cryst. C*52, 2118–2119.
Okabe, N., Nakamura, T. & Fukuda, H. (1993). *Acta Cryst. C*49, 1678–1680.
Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o1433 [doi:10.1107/S1600536808020266]

2,4-Dichlorobenzaldehyde 2,4-dinitrophenylhydrazone

F. Bao

Comment

Several phenylhydrazone derivatives have been shown to be potentially DNA-damaging and are mutagenic agents (Okabe *et al.*, 1993). As part of our work in this area, we have synthesized the title compound, (I), and we report its crystal structure here.

The two molecules in the asymmetric unit of (I) are almost planar, the dihedral angles between the dichlorobenzene ring and the dinitrobenzene ring are 4.4 (2) $^{\circ}$ and 3.7 (2) $^{\circ}$, in the C1 and C14 molecules, respectively. Otherwise, bond lengths and angles agree with those of other dinitrophenylhydrazone derivatives (Ohba, 1996). Intramolecular N—H···O hydrogen bonds (Fig. 1, Table 1) help to establish these molecular conformations.

Experimental

2,4-Dinitrophenylhydrazine (1 mmol, 0.198 g) was dissolved in anhydrous methanol, H₂SO₄ (98% 0.5 ml) was added to this, the mixture was stirred for several minutes at 351 K, then 2,4-dichlorobenzaldehyde (1 mmol, 0.175 g) in methanol (8 ml) was added dropwise and the mixture was stirred at refluxing temperature for 2 h. The product was isolated and recrystallized from DMF, yielding brown blocks of (I) after 6 d.

Refinement

All H atoms were placed in calculated positions (C—H = 0.93 Å, N—H = 0.86 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$.

Figures



Fig. 1. The molecular structure of (I) with displacement ellipsoids drawn at the 30% probability level (arbitrary spheres for the H atoms). The hydrogen bonds are indicated by dashed lines.

2,4-Dichlorobenzaldehyde 2,4-dinitrophenylhydrazone

Crystal data

C₁₃H₈Cl₂N₄O₄

$F_{000} = 1440$

$M_r = 355.14$

$D_x = 1.645 \text{ Mg m}^{-3}$

Monoclinic, $P2_1/c$

Mo $K\alpha$ radiation

Hall symbol: -P 2ybc

$\lambda = 0.71073 \text{ \AA}$

$a = 13.3814 (7) \text{ \AA}$

Cell parameters from 721 reflections

$\theta = 2.5\text{--}20.6^\circ$

supplementary materials

$b = 28.9980 (13) \text{ \AA}$	$\mu = 0.48 \text{ mm}^{-1}$
$c = 7.3996 (3) \text{ \AA}$	$T = 291 (2) \text{ K}$
$\beta = 92.422 (4)^\circ$	Block, brown
$V = 2868.7 (2) \text{ \AA}^3$	$0.16 \times 0.11 \times 0.09 \text{ mm}$
$Z = 8$	

Data collection

Bruker SMART APEX CCD diffractometer	5060 independent reflections
Radiation source: sealed tube	2794 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.093$
$T = 291(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
ω scans	$\theta_{\text{min}} = 1.4^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 1998)	$h = -15 \rightarrow 15$
$T_{\text{min}} = 0.927, T_{\text{max}} = 0.958$	$k = -34 \rightarrow 34$
21827 measured reflections	$l = -8 \rightarrow 8$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.061$	H-atom parameters constrained
$wR(F^2) = 0.187$	$w = 1/[\sigma^2(F_o^2) + (0.0939P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\text{max}} = 0.018$
5060 reflections	$\Delta\rho_{\text{max}} = 0.70 \text{ e \AA}^{-3}$
415 parameters	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
-----	-----	-----	----------------------------------

C1	0.5169 (3)	0.79565 (15)	0.8938 (6)	0.0525 (11)
C2	0.4504 (4)	0.76034 (15)	0.8543 (6)	0.0568 (12)
H2A	0.4681	0.7299	0.8798	0.068*
C3	0.3594 (4)	0.77030 (16)	0.7782 (6)	0.0567 (12)
C4	0.3307 (4)	0.81536 (17)	0.7419 (7)	0.0649 (13)
H4A	0.2670	0.8217	0.6926	0.078*
C5	0.3956 (4)	0.85013 (17)	0.7786 (6)	0.0630 (13)
H5A	0.3755	0.8802	0.7532	0.076*
C6	0.4926 (4)	0.84239 (14)	0.8539 (6)	0.0508 (11)
C7	0.5828 (4)	0.95479 (14)	0.8532 (6)	0.0522 (11)
H7A	0.6458	0.9503	0.9085	0.063*
C8	0.5499 (3)	1.00103 (14)	0.7955 (5)	0.0483 (11)
C9	0.4563 (3)	1.00735 (15)	0.7104 (6)	0.0526 (11)
H9A	0.4151	0.9818	0.6913	0.063*
C10	0.4222 (3)	1.05004 (15)	0.6533 (6)	0.0535 (11)
H10A	0.3595	1.0534	0.5959	0.064*
C11	0.4836 (4)	1.08737 (14)	0.6836 (6)	0.0495 (11)
C12	0.5769 (3)	1.08302 (15)	0.7673 (6)	0.0519 (11)
H12A	0.6181	1.1086	0.7848	0.062*
C13	0.6078 (3)	1.03998 (14)	0.8247 (6)	0.0496 (11)
C14	0.0562 (3)	0.40012 (14)	0.2229 (5)	0.0453 (10)
C15	-0.0068 (3)	0.43582 (15)	0.1708 (6)	0.0501 (11)
H15A	0.0142	0.4662	0.1854	0.060*
C16	-0.1000 (3)	0.42650 (15)	0.0978 (6)	0.0509 (11)
C17	-0.1325 (4)	0.38143 (16)	0.0780 (6)	0.0572 (12)
H17A	-0.1972	0.3754	0.0330	0.069*
C18	-0.0705 (4)	0.34611 (16)	0.1240 (6)	0.0590 (12)
H18A	-0.0930	0.3160	0.1062	0.071*
C19	0.0269 (3)	0.35334 (14)	0.1980 (5)	0.0479 (11)
C20	0.1045 (4)	0.23951 (15)	0.2590 (6)	0.0535 (12)
H20A	0.1681	0.2440	0.3117	0.064*
C21	0.0647 (3)	0.19295 (14)	0.2269 (5)	0.0472 (11)
C22	0.1187 (3)	0.15381 (15)	0.2772 (5)	0.0506 (11)
C23	0.0818 (4)	0.10936 (14)	0.2470 (6)	0.0517 (11)
H23A	0.1193	0.0836	0.2818	0.062*
C24	-0.0113 (4)	0.10474 (15)	0.1644 (6)	0.0513 (11)
C25	-0.0668 (4)	0.14258 (16)	0.1099 (6)	0.0588 (12)
H25A	-0.1296	0.1388	0.0528	0.071*
C26	-0.0283 (4)	0.18613 (15)	0.1409 (6)	0.0552 (12)
H26A	-0.0657	0.2116	0.1031	0.066*
N1	0.1514 (3)	0.41294 (14)	0.3070 (5)	0.0577 (10)
N2	-0.1667 (3)	0.46460 (16)	0.0431 (5)	0.0612 (11)
N3	0.2910 (3)	0.73221 (16)	0.7304 (6)	0.0708 (12)
N4	0.6139 (3)	0.78294 (14)	0.9750 (6)	0.0659 (11)
N5	0.5238 (3)	0.92107 (12)	0.8261 (5)	0.0546 (10)
N6	0.5553 (3)	0.87810 (12)	0.8820 (5)	0.0541 (10)
H6A	0.6134	0.8742	0.9336	0.065*
N7	0.0487 (3)	0.27372 (12)	0.2123 (5)	0.0529 (10)
N8	0.0870 (3)	0.31688 (11)	0.2431 (5)	0.0532 (9)

supplementary materials

H8A	0.1465	0.3208	0.2890	0.064*
O1	0.2127 (3)	0.74172 (14)	0.6467 (6)	0.0950 (13)
O2	0.3142 (3)	0.69353 (13)	0.7743 (6)	0.0985 (13)
O3	0.6762 (3)	0.81245 (11)	1.0089 (5)	0.0726 (10)
O4	0.6286 (3)	0.74302 (14)	1.0074 (8)	0.135 (2)
O5	-0.1387 (3)	0.50380 (12)	0.0700 (5)	0.0796 (11)
O6	-0.2487 (3)	0.45419 (12)	-0.0264 (5)	0.0833 (11)
O7	0.1687 (3)	0.45318 (12)	0.3374 (6)	0.0913 (13)
O8	0.2111 (3)	0.38253 (11)	0.3504 (5)	0.0731 (10)
Cl1	0.44331 (10)	1.14207 (4)	0.61744 (18)	0.0689 (4)
Cl2	-0.05792 (11)	0.04957 (4)	0.12777 (17)	0.0700 (4)
Cl3	0.23487 (9)	0.15728 (4)	0.38772 (18)	0.0705 (4)
Cl4	0.72292 (10)	1.03677 (4)	0.94132 (19)	0.0724 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.054 (3)	0.047 (3)	0.057 (3)	0.004 (2)	0.000 (2)	-0.002 (2)
C2	0.061 (3)	0.041 (3)	0.069 (3)	0.001 (2)	0.002 (3)	0.002 (2)
C3	0.055 (3)	0.052 (3)	0.064 (3)	-0.006 (2)	0.004 (2)	0.004 (2)
C4	0.054 (3)	0.060 (3)	0.080 (3)	0.007 (3)	-0.005 (3)	0.012 (3)
C5	0.066 (3)	0.049 (3)	0.073 (3)	0.002 (3)	-0.005 (3)	0.008 (2)
C6	0.061 (3)	0.038 (3)	0.053 (3)	0.000 (2)	0.005 (2)	-0.0013 (19)
C7	0.058 (3)	0.045 (3)	0.053 (3)	0.001 (2)	0.004 (2)	0.002 (2)
C8	0.050 (3)	0.047 (3)	0.048 (2)	0.001 (2)	0.003 (2)	0.0021 (19)
C9	0.055 (3)	0.048 (3)	0.055 (3)	-0.011 (2)	-0.001 (2)	0.000 (2)
C10	0.050 (3)	0.055 (3)	0.055 (3)	0.001 (2)	-0.002 (2)	0.008 (2)
C11	0.060 (3)	0.039 (2)	0.050 (3)	0.004 (2)	0.003 (2)	0.0070 (19)
C12	0.054 (3)	0.045 (3)	0.057 (3)	-0.002 (2)	0.003 (2)	0.006 (2)
C13	0.048 (3)	0.048 (3)	0.052 (3)	-0.002 (2)	-0.005 (2)	0.003 (2)
C14	0.051 (3)	0.044 (2)	0.041 (2)	-0.003 (2)	0.003 (2)	-0.0004 (18)
C15	0.055 (3)	0.044 (3)	0.051 (3)	-0.007 (2)	0.002 (2)	-0.003 (2)
C16	0.052 (3)	0.048 (3)	0.053 (3)	0.000 (2)	0.004 (2)	0.003 (2)
C17	0.051 (3)	0.058 (3)	0.062 (3)	-0.008 (2)	-0.007 (2)	-0.001 (2)
C18	0.063 (3)	0.048 (3)	0.066 (3)	-0.010 (2)	-0.002 (3)	-0.002 (2)
C19	0.053 (3)	0.046 (3)	0.045 (2)	-0.006 (2)	0.003 (2)	0.0022 (19)
C20	0.062 (3)	0.047 (3)	0.052 (3)	-0.004 (2)	0.002 (2)	0.000 (2)
C21	0.056 (3)	0.044 (3)	0.042 (2)	0.004 (2)	0.003 (2)	0.0053 (18)
C22	0.054 (3)	0.055 (3)	0.043 (2)	0.001 (2)	0.002 (2)	-0.0008 (19)
C23	0.059 (3)	0.042 (3)	0.054 (3)	0.006 (2)	-0.002 (2)	0.006 (2)
C24	0.058 (3)	0.051 (3)	0.044 (2)	-0.006 (2)	-0.001 (2)	0.001 (2)
C25	0.058 (3)	0.064 (3)	0.053 (3)	-0.001 (3)	-0.008 (2)	0.003 (2)
C26	0.066 (3)	0.045 (3)	0.054 (3)	-0.004 (2)	-0.003 (2)	0.003 (2)
N1	0.058 (3)	0.049 (3)	0.065 (2)	-0.001 (2)	-0.007 (2)	-0.0022 (19)
N2	0.057 (3)	0.063 (3)	0.062 (2)	0.008 (2)	-0.008 (2)	0.007 (2)
N3	0.059 (3)	0.064 (3)	0.089 (3)	-0.011 (2)	0.000 (3)	0.002 (2)
N4	0.069 (3)	0.039 (2)	0.088 (3)	0.008 (2)	-0.014 (2)	0.006 (2)
N5	0.068 (3)	0.035 (2)	0.060 (2)	0.0043 (19)	0.0047 (19)	0.0060 (17)

N6	0.064 (3)	0.043 (2)	0.056 (2)	0.0031 (19)	0.0001 (19)	0.0027 (17)
N7	0.068 (3)	0.038 (2)	0.053 (2)	-0.0054 (19)	0.0012 (19)	-0.0002 (16)
N8	0.058 (2)	0.039 (2)	0.063 (2)	-0.0008 (18)	-0.0017 (19)	0.0015 (17)
O1	0.072 (3)	0.090 (3)	0.121 (3)	-0.011 (2)	-0.027 (3)	0.002 (2)
O2	0.085 (3)	0.053 (2)	0.157 (4)	-0.008 (2)	-0.013 (3)	0.006 (2)
O3	0.069 (2)	0.052 (2)	0.096 (3)	-0.0040 (18)	-0.019 (2)	-0.0004 (17)
O4	0.094 (3)	0.048 (3)	0.256 (6)	0.001 (2)	-0.075 (4)	0.023 (3)
O5	0.087 (3)	0.046 (2)	0.105 (3)	0.0062 (19)	-0.012 (2)	0.0089 (19)
O6	0.066 (3)	0.084 (3)	0.098 (3)	0.005 (2)	-0.016 (2)	0.007 (2)
O7	0.081 (3)	0.047 (2)	0.142 (4)	-0.0054 (19)	-0.036 (3)	-0.012 (2)
O8	0.070 (2)	0.053 (2)	0.095 (3)	0.0059 (18)	-0.019 (2)	-0.0025 (18)
Cl1	0.0721 (9)	0.0489 (7)	0.0843 (9)	0.0067 (6)	-0.0118 (7)	0.0137 (6)
Cl2	0.0824 (10)	0.0512 (7)	0.0750 (8)	-0.0133 (6)	-0.0150 (7)	-0.0007 (6)
Cl3	0.0564 (8)	0.0704 (9)	0.0833 (9)	-0.0032 (6)	-0.0139 (7)	-0.0004 (6)
Cl4	0.0567 (8)	0.0676 (8)	0.0911 (9)	-0.0022 (6)	-0.0189 (7)	0.0157 (7)

Geometric parameters (Å, °)

C1—C2	1.379 (6)	C16—N2	1.467 (6)
C1—C6	1.422 (6)	C17—C18	1.352 (6)
C1—N4	1.455 (6)	C17—H17A	0.9300
C2—C3	1.352 (6)	C18—C19	1.408 (6)
C2—H2A	0.9300	C18—H18A	0.9300
C3—C4	1.385 (6)	C19—N8	1.361 (5)
C3—N3	1.468 (6)	C20—N7	1.280 (5)
C4—C5	1.351 (7)	C20—C21	1.467 (6)
C4—H4A	0.9300	C20—H20A	0.9300
C5—C6	1.409 (7)	C21—C26	1.388 (6)
C5—H5A	0.9300	C21—C22	1.388 (6)
C6—N6	1.343 (5)	C22—C23	1.395 (6)
C7—N5	1.267 (5)	C22—Cl3	1.728 (5)
C7—C8	1.469 (6)	C23—C24	1.371 (6)
C7—H7A	0.9300	C23—H23A	0.9300
C8—C13	1.381 (6)	C24—C25	1.376 (6)
C8—C9	1.390 (6)	C24—Cl2	1.734 (4)
C9—C10	1.379 (6)	C25—C26	1.380 (6)
C9—H9A	0.9300	C25—H25A	0.9300
C10—C11	1.372 (6)	C26—H26A	0.9300
C10—H10A	0.9300	N1—O7	1.209 (4)
C11—C12	1.375 (6)	N1—O8	1.224 (5)
C11—Cl1	1.739 (4)	N2—O5	1.211 (5)
C12—C13	1.376 (6)	N2—O6	1.229 (5)
C12—H12A	0.9300	N3—O2	1.204 (5)
C13—Cl4	1.736 (5)	N3—O1	1.226 (5)
C14—C15	1.379 (6)	N4—O4	1.197 (5)
C14—C19	1.422 (6)	N4—O3	1.213 (5)
C14—N1	1.443 (6)	N5—N6	1.373 (5)
C15—C16	1.365 (6)	N6—H6A	0.8600
C15—H15A	0.9300	N7—N8	1.368 (5)

supplementary materials

C16—C17	1.384 (6)	N8—H8A	0.8600
C2—C1—C6	121.6 (4)	C18—C17—H17A	119.9
C2—C1—N4	117.1 (4)	C16—C17—H17A	119.9
C6—C1—N4	121.3 (4)	C17—C18—C19	122.2 (4)
C3—C2—C1	119.4 (4)	C17—C18—H18A	118.9
C3—C2—H2A	120.3	C19—C18—H18A	118.9
C1—C2—H2A	120.3	N8—C19—C18	120.5 (4)
C2—C3—C4	121.3 (4)	N8—C19—C14	123.5 (4)
C2—C3—N3	118.7 (4)	C18—C19—C14	116.0 (4)
C4—C3—N3	119.9 (5)	N7—C20—C21	117.8 (4)
C5—C4—C3	119.6 (5)	N7—C20—H20A	121.1
C5—C4—H4A	120.2	C21—C20—H20A	121.1
C3—C4—H4A	120.2	C26—C21—C22	116.9 (4)
C4—C5—C6	122.3 (4)	C26—C21—C20	121.2 (4)
C4—C5—H5A	118.8	C22—C21—C20	121.9 (4)
C6—C5—H5A	118.8	C21—C22—C23	122.4 (4)
N6—C6—C5	119.8 (4)	C21—C22—Cl3	121.8 (3)
N6—C6—C1	124.6 (4)	C23—C22—Cl3	115.8 (3)
C5—C6—C1	115.6 (4)	C24—C23—C22	118.1 (4)
N5—C7—C8	118.7 (4)	C24—C23—H23A	120.9
N5—C7—H7A	120.6	C22—C23—H23A	120.9
C8—C7—H7A	120.6	C23—C24—C25	121.5 (4)
C13—C8—C9	116.8 (4)	C23—C24—Cl2	118.3 (3)
C13—C8—C7	122.8 (4)	C25—C24—Cl2	120.2 (4)
C9—C8—C7	120.4 (4)	C24—C25—C26	119.2 (4)
C10—C9—C8	122.5 (4)	C24—C25—H25A	120.4
C10—C9—H9A	118.8	C26—C25—H25A	120.4
C8—C9—H9A	118.8	C25—C26—C21	121.9 (4)
C9—C10—C11	118.0 (4)	C25—C26—H26A	119.0
C9—C10—H10A	121.0	C21—C26—H26A	119.0
C11—C10—H10A	121.0	O7—N1—O8	121.9 (4)
C12—C11—C10	121.8 (4)	O7—N1—C14	119.2 (4)
C12—C11—Cl1	118.4 (3)	O8—N1—C14	118.8 (4)
C10—C11—Cl1	119.8 (4)	O5—N2—O6	124.3 (4)
C11—C12—C13	118.5 (4)	O5—N2—C16	118.7 (4)
C11—C12—H12A	120.8	O6—N2—C16	116.9 (4)
C13—C12—H12A	120.8	O2—N3—O1	123.4 (5)
C12—C13—C8	122.3 (4)	O2—N3—C3	119.0 (5)
C12—C13—Cl4	116.8 (3)	O1—N3—C3	117.6 (4)
C8—C13—Cl4	120.9 (3)	O4—N4—O3	122.3 (4)
C15—C14—C19	121.2 (4)	O4—N4—C1	117.7 (4)
C15—C14—N1	116.4 (4)	O3—N4—C1	120.0 (4)
C19—C14—N1	122.4 (4)	C7—N5—N6	118.1 (4)
C16—C15—C14	119.9 (4)	C6—N6—N5	118.1 (4)
C16—C15—H15A	120.0	C6—N6—H6A	120.9
C14—C15—H15A	120.0	N5—N6—H6A	120.9
C15—C16—C17	120.5 (4)	C20—N7—N8	117.0 (4)
C15—C16—N2	119.7 (4)	C19—N8—N7	117.2 (4)
C17—C16—N2	119.8 (4)	C19—N8—H8A	121.4

supplementary materials

C18—C17—C16	120.1 (4)	N7—N8—H8A	121.4
-------------	-----------	-----------	-------

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N6—H6A···O3	0.86	2.05	2.645 (5)	126
N8—H8A···O8	0.86	2.03	2.627 (5)	126

supplementary materials

Fig. 1

