



# Genetics, genomics and clinical features of adenomatous polyposis

Jihoon E. Joo<sup>1,2</sup> · Julen Viana-Errasti<sup>3,4,5,6</sup> · Daniel D. Buchanan<sup>1,2,7</sup> · Laura Valle<sup>3,4,5</sup>

Received: 9 February 2025 / Accepted: 16 March 2025 / Published online: 16 April 2025  
© The Author(s) 2025

## Abstract

Adenomatous polyposis syndromes are hereditary conditions characterised by the development of multiple adenomas in the gastrointestinal tract, particularly in the colon and rectum, significantly increasing the risk of colorectal cancer and, in some cases, extra-colonic malignancies. These syndromes are caused by germline pathogenic variants (PVs) in genes involved in Wnt signalling and DNA repair. The main autosomal dominant adenomatous polyposis syndromes include familial adenomatous polyposis (FAP) and polymerase proofreading-associated polyposis (PPAP), caused by germline PVs in *APC* and the *POLE* and *POLD1* genes, respectively. Autosomal recessive syndromes include those caused by biallelic PVs in the DNA mismatch repair genes *MLH1*, *MSH2*, *MSH6*, *PMS2*, *MSH3* and probably *MLH3*, and in the base excision repair genes *MUTYH*, *NTHL1* and *MBD4*. This review provides an in-depth discussion of the genetic and molecular mechanisms underlying hereditary adenomatous polyposis syndromes, their clinical presentations, tumour mutational signatures, and emerging approaches for the treatment of the associated cancers. Considerations for genetic testing are described, including post-zygotic mosaicism, non-coding PVs, the interpretation of variants of unknown significance and cancer risks associated with monoallelic variants in the recessive genes. Despite advances in genetic testing and the recent identification of new adenomatous polyposis genes, many cases of multiple adenomas remain genetically unexplained. Non-genetic factors, including environmental risk factors, prior oncologic treatments, and bacterial genotoxins colonising the intestine - particularly colibactin-producing *Escherichia coli* - have emerged as alternative pathogenic mechanisms.

**Keywords** Hereditary colorectal cancer · Gastrointestinal polyposis · Familial adenomatous polyposis · APC · MUTYH · NTHL1 · MBD4 · POLE · POLD1 · MMR

---

Jihoon E. Joo, Julen Viana-Errasti equally contributed to this work (first authors).

---

Daniel D. Buchanan, Laura Valle equally contributed to this work (senior and corresponding authors).

---

✉ Daniel D. Buchanan  
daniel.buchanan@unimelb.edu.au

✉ Laura Valle  
lvalle@idibell.cat

<sup>1</sup> Colorectal Oncogenomics Group, Department of Clinical Pathology, The University of Melbourne, Parkville, VIC, Australia

<sup>2</sup> Collaborative Centre for Genomic Cancer Medicine, Victorian Comprehensive Cancer Centre, Parkville, VIC, Australia


<sup>3</sup> Hereditary Cancer Program, Catalan Institute of Oncology, IDIBELL, Hospitalet de Llobregat, Av. Gran Via 199- 203, Hospitalet de Llobregat 08908, Spain

<sup>4</sup> Program in Molecular Mechanisms and Experimental Therapy in Oncology (Oncobell), IDIBELL, Hospitalet de Llobregat, Barcelona, Spain

<sup>5</sup> Centro de Investigación Biomédica en Red de Cáncer (CIBERONC), Madrid, Spain

<sup>6</sup> Doctoral Program in Biomedicine, University of Barcelona, Hospitalet de Llobregat, Barcelona, Spain

<sup>7</sup> Genomic Medicine and Family Cancer Clinic, Royal Melbourne Hospital, Parkville, VIC, Australia



**Fig. 1** Adenomatous polyposis syndromes, mode of inheritance, causal genes, affected molecular pathways and associated COSMIC tumour mutational signatures, and possible causes of adenomatous polyposis in patients without germline PVs in known polyposis genes. *Abbreviations:* CMMRD, constitutional mismatch repair deficiency; CRC,

colorectal cancer; FAP, familial adenomatous polyposis; LS, Lynch syndrome; MANS, MBD4-associated neoplasia syndrome; MAP, MUTYH-associated polyposis; MMR, mismatch repair; NTS, NTHL1 tumour syndrome; PPAP, polymerase proofreading-associated polyposis; ED, exonuclease domain

## Adenomatous polyposis syndromes: causal genes and clinical characteristics

Adenomatous polyposis syndromes are characterised by the development of multiple adenomatous polyps in the colon and rectum, significantly increasing the risk of colorectal cancer (CRC), including the potential to develop multiple CRCs either as synchronous or metachronous events. Depending on the underlying aetiology, patients may have an increased risk of developing extra-colonic cancers.

Adenomatous polyposes are distinguished from other polyposis syndromes including serrated polyposis syndrome, the hamartomatous polyposis syndromes Peutz-Jeghers, Juvenile Polyposis and PTEN-hamartoma-tumour, and mixed polyposis, by the morphology of the predominant polyp type and the underlying causes [1]. Adenomatous polyposis syndromes are primarily caused by the (dominant or recessive) inheritance of constitutional pathogenic variants (PVs) in several genes involved in Wnt signalling and DNA repair mechanisms (Fig. 1).

### Autosomal dominant syndromes

Familial adenomatous polyposis (FAP), the best characterised and most common polyposis syndrome, has a birth incidence of around one in 8500, manifests equally in both sexes, and accounts for 0.5% or less of CRC cases. FAP is caused by monoallelic constitutional PVs in *APC*, a negative regulator of Wnt signalling [2–5]. In FAP, somatic inactivation of the wild-type allele (“somatic second hit”) leads to complete inactivation of *APC*, causing constitutional activation of Wnt signalling due to the lack of degradation of the transcriptional coactivator  $\beta$ -catenin, which translocates to the nucleus where it binds TCF/LEF family members and

activates transcription. This eventually results in the activation of oncogenes and other cancer-related genes, triggering tumourigenesis [6]. The importance of the *APC* gene in colorectal tumourigenesis is highlighted by the fact that it is one of the most commonly somatically mutated genes in CRC [7]. Up to 30% of affected individuals present ostensibly *de novo APC* PVs and thus have no family history of the disease in previous generations. When these PVs occur post-zygotically in embryonic tissues, they result in *APC* mosaicism [8, 9]. The extent of mosaicism in colonic tissues seems to determine the severity of the polyposis phenotype in these cases [9, 10].

Clinically, the classic form of FAP is characterised by the presence of hundreds to thousands of colorectal adenomas, with a disease onset in the late childhood or adolescence, with 100% risk of developing CRC if untreated. This condition is also associated with extracolonic features, some of which are highly relevant, specifically upper gastrointestinal (GI) tumours and desmoid disease, which are the main causes of FAP-related mortality [11]. In addition to CRC, but to a much lesser extent, increased cancer risk has been observed for duodenal, gastric, pancreatic, small intestine, and pancreatic cancers, hepatoblastoma, medulloblastoma, and cribriform-morular variant of papillary thyroid cancer [12]. Also, non-malignant extracolonic manifestations may occur and help with a diagnosis. These include congenital hypertrophy of the retinal pigment epithelium (CHRPE), dermoid cysts, osteomas, dental abnormalities, benign cutaneous lesions, and adrenal masses [12].

FAP is a spectrum disease, with high variability in the clinical phenotype, including attenuated forms of the disease characterised by lower polyp burden (20–100 adenomas) and later age of onset [1, 13]. A recent analysis of the data from the Danish Polyposis Registry, comprising 311 patients with classical FAP and 134 patients with attenuated

FAP diagnosed in Denmark since 1974, indicated that patients with both classic and attenuated forms of FAP were at higher overall cancer risk, including CRC risk, compared with population individuals, although these risks were not different between the two groups of FAP patients. While extracolonic cancers and mortality were higher in patients with classic FAP compared with population individuals, no statistically significant increase was observed for attenuated FAP compared with population individuals [14]. The location of PVs within the *APC* gene sequence may determine the phenotypic severity of FAP, although variability between patients exists. Exon 15 of the *APC* gene contains a mutation cluster region (MCR) linked to severe phenotypes. In contrast, PVs in the 5' and 3' regions are associated with less severe phenotypes.

An extremely rare gastric phenotype, termed gastric adenocarcinoma and proximal polyposis syndrome (GAPPS), is caused by germline single nucleotide PVs affecting the YY1 binding site of exon 1B (promoter) of *APC* [15]. A systematic review and meta-analysis of 113 patients with a PV in the promoter 1B region showed that GAPPS is a gastric polyposis syndrome with a substantial risk of developing gastric cancer from an early age, with a remarkable variability in clinical expression within and among families. The presence of corpus fundic polyposis in addition to a family history of gastric cancer, regardless of the age at diagnosis, should be considered suggestive of GAPPS. These patients have no other manifestations resembling FAP, including intestinal, i.e., colorectal adenomatous polyposis, or extraintestinal manifestations [16].

Polymerase proofreading-associated polyposis (PPAP) is caused by constitutional PVs in *POLE* or *POLD1*. Cancer-associated PVs occur within the exonuclease domain of polymerases  $\epsilon$  and  $\delta$ , affecting their proofreading activity, and thus resulting in a particular type of DNA repair deficiency [17].

PPAP is clinically characterised by adenomatous polyposis (<100 adenomas), and increased risk of colorectal, endometrial, ovarian, breast, brain, and upper GI cancers. The presence of café-au-lait macules (CALMs) is often observed in PPAP patients. Typically, benign and malignant neoplasms are diagnosed in the adulthood (35–60 years of age) [18]. Several PPAP cases with particularly aggressive phenotypes have been reported in the literature, with both polyposis and cancer diagnosed early in life [19–24]. These cases are characterised by early-onset diagnoses (medulloblastomas diagnosed in early childhood, and polyposis and CRC in the adolescence or young adulthood), and the presence of CALMs and non-malignant tumours (pilomatricomas, etc.), mimicking the Constitutional Mismatch Repair Deficiency (CMMRD) phenotype. Patients with such phenotypes have been associated with specific variants, including

*POLE* p.E277G, p.S297F, p.V411L, p.P436R, p.M444K, p.A456P, and p.S461T, likely to induce more severe effects on the proofreading activity of the polymerase, and with the co-occurrence of constitutional PVs in *POLE* or *POLD1* exonuclease domain and in a DNA mismatch repair (MMR) gene [19–25].

Constitutional PVs in *AXIN2*, another Wnt signalling negative regulator, have been associated with oligodontia, ectodermal dysplasia (sparse hair and eyebrows) and increased risk of adenomatous polyposis [26]. Although only a few cases have been reported, the phenotypic variability is shown to be broad between heterozygotes. Some individuals exhibit one or more clinical manifestations, while those with polyps show variation in polyp number and histologic types, with most being adenomas, although serrated polyps have also been observed [27].

### Autosomal recessive syndromes

MUTYH-associated polyposis (MAP) is a recessive adenomatous polyposis syndrome caused by constitutional PVs in *MUTYH*; a gene that encodes MYH glycosylase, which is part of the DNA base excision repair (BER) system [28, 29]. After FAP, MAP is the second most common adenomatous polyposis syndrome, estimated to occur in 1:20,000 to 1:60,000 individuals due to the presence of relatively prevalent founder mutations in different populations [30].

Clinical features of MAP are mostly restricted to the GI tract. As in other polyposis syndromes, people with MAP have a broad phenotypic variability: from mild to profuse colorectal polyposis, and in some cases presenting with CRC without a polyposis phenotype. MAP patients also have increased risk of duodenal cancer, and to a lesser extent, of non-melanoma skin cancer, ovarian, endometrial or bladder cancer. Although rare, extracolonic non-malignant features including CHRPE, thyroid nodules, benign adrenal lesions, or jawbone cysts, have been reported in MAP patients [1, 12].

Biallelic PVs in *NTHL1* and *MBD4*, also BER glycosylases, cause two different ultra-rare recessive adenomatous polyposis syndromes. *NTHL1* tumour syndrome (NTS) is characterised by high risk of gastrointestinal tumours, including CRC and polyposis, endometrial cancer, and breast cancer, and increased risk of other cancer types and non-neoplastic manifestations. *NTHL1* biallelic carriers have very high risk of developing multiple primary tumours [31–33]. *MBD4*-associated neoplasia syndrome (MANS) causes increased risk of acute myeloid leukaemia (AML)-preceded by myelodysplastic syndrome (MDS)-, adenomatous polyposis and CRC, and to a lesser extent, of uveal melanoma and schwannomas [34, 35].

Biallelic PVs in DNA mismatch repair (MMR) genes, including four MMR genes associated with Lynch syndrome (*MLH1*, *MSH2*, *MSH6* and *PMS2*) and other recessive MMR genes such as *MSH3* or *MLH3* cause adenomatous polyposis, usually as an attenuated polyposis phenotype. Constitutional mismatch repair deficiency (CMMRD) is a severe childhood cancer predisposition syndrome caused by biallelic PVs in the Lynch syndrome-related MMR genes. Most MMR variants identified in CMMRD patients are in the least penetrant Lynch syndrome genes, i.e., *PMS2* and *MSH6*. CMMRD typically presents with haematological, brain, and gastrointestinal cancers in childhood or adolescence, with a median age of onset <10 years. CMMRD patients are highly likely to develop multiple malignancies and at increased risk of embryonal tumours, germ cell tumours, sarcomas, ganglioneuroma, melanoma, urinary tract, prostate, breast, endometrial and ovarian cancers. CMMRD is also associated with distinctive non-neoplastic features, most commonly CALMs, other skin pigmentation alterations, and multiple developmental venous anomalies [36, 37]. Mild CMMRD phenotypes resembling Lynch syndrome in tumour spectrum and age at cancer diagnosis have been reported in patients carrying biallelic hypomorphic MMR gene variants [38].

Constitutional biallelic PVs in *MSH3* have been reported in a few adult-onset adenomatous polyposis cases [39–43]. While available information points towards mostly gastrointestinal tract phenotypes, with CRC and adenomas as the main clinical features, data is still scarce to determine if there is an increased risk of tumours in other organs. Preliminary data suggests that a similar phenotype may be linked to biallelic PVs in *MLH3* although only a small number of cases have been reported to date [44, 45].

## Considerations for genetic testing

### Somatic or post-zygotic mosaicism

As mentioned before, up to 30% of FAP patients carry *de novo* PVs in *APC*, and a relevant part of these (estimated between 10% and 25%) may be the result of post-zygotic mosaicism in *APC* [8]. As a general rule, these patients lack a family history of the disease. However, in exceptional cases, somatic mosaicism may be observed in two affected members (siblings) of the same family [46]. Mosaicism occurs when the variant arises *de novo* in embryonic tissue. The presence of the disease-causing variant in different organs or tissue types depends on the timing of the mutation during the embryonic stages. The detection of mosaicism may require lowering the variant allele frequency threshold ( $\leq 5\%$ ) when analysing germline sequencing data; i.e.,

when multi-gene panel, exome or genome sequencing data obtained from blood-derived DNA is analysed. If no variants are uncovered or are difficult to discern from low level artefacts, testing other tissues, such as normal colorectal mucosa or multiple gastrointestinal polyps is highly recommended since the mosaic mutation might not be present in haematopoietic cells, and sometimes even be restricted to the colorectal epithelium [47, 48]. If the only available non-haematopoietic tissue source consists of gastrointestinal neoplasms (adenomas or cancer), at least two independent lesions should be analysed to confirm the presence of a common *APC* variant in all of them. As most adenomatous polyps acquire somatic *APC* mutations, testing multiple neoplasms is important to differentiate somatic *APC* mutations from a likely mosaic *APC* variant. Furthermore, testing for the *APC* mosaic variant using sensitive loci-specific detection techniques (e.g. digital PCR) in different sources/lineages of DNA will further enable confirmation and differentiation of localised versus soma-wide *APC* mosaicism, which has important implications for subsequent cancer risk and risk-appropriate clinical management.

Although likely to be less common than observed with *APC*, somatic mosaicism in other adenomatous polyposis genes is also expected. Somatic mosaicism has been reported in MMR genes (Lynch syndrome) [49], and other non-adenomatous polyposis genes such as *PTEN* or *STK11* [8, 50].

### Pathogenic variants in non-coding regions and structural/copy number variants affecting *APC*

Another potential source of missed diagnoses in adenomatous polyposis individuals is the presence of deep-intronic variants that have aberrant effects on splicing or variants that affect regulatory regions of the gene, particularly in *APC*. Although rare, intronic PVs in other (non-adenomatous) polyposis syndrome genes have also been reported [51–53].

Several recent studies have shown that deep-intronic variants in *APC* that alter splicing may be a common cause of missed diagnoses of FAP [54–57]. The use of whole-genome sequencing or gene panels that capture the intronic regions of the genes, followed by the evaluation of potential effects on splicing using *in silico* predictors, helps identify deep intronic variants with potential spliceogenic effects. Subsequent RNA studies are needed to clarify the actual deleterious effect, and if so, assess a potential hypomorphic nature. DNA-RNA paired testing is a straightforward approach to identify deep intronic PVs, not only in *APC* but in all risk genes included in the panels [54].

Complex and large structural variants involving *APC*, which can originate from both homologous and non-homologous recombination events mediated by Alu elements,

have also been identified in FAP patients [58–63]. Multi-modal approaches, including long-read genome and RNA sequencing, optimal mapping, and chromosomal microarray, may be required when FAP is genetically unexplained, as exemplified by a case of constitutional chromothripsis involving the *APC* locus [62], or a complex rearrangement between *APC* and *TP63* [60], both of which were undetected by standard short-read multi-gene panel sequencing.

Current clinical tests primarily involve multi-gene panel testing that utilises short-read sequencing technology due to its high accuracy, high throughput capability, and cost-effectiveness. However, these approaches have limited sensitivity for detecting structural variants, particularly in low-complexity regions (e.g., repetitive sequences) [64]. As a result, a significant proportion of FAP diagnoses linked to copy number alterations and complex rearrangements may be missed by current clinical testing. A small study reported that large duplications or deletions in the *APC* gene were identified in approximately 24% of patients undergoing multi-ligation probe amplification (MLPA) testing [65], although lower detection rates have been observed in other studies [66–68]. Clinical testing using whole-genome sequencing and long-read genome and/or RNA sequencing technologies (e.g., Oxford Nanopore Technologies [69], single-molecule real-time sequencing [70]) could improve the detection of this type of variants.

The effect of certain non-coding variants on regulatory regions of the gene has been less studied, probably due to the difficulty in demonstrating their actual effect on the allelic expression. The easiest to identify are the ones located in promoter regions. Variants in the promoter 1B of *APC* have been identified in individuals with FAP, and more specifically with a rare gastric phenotype (GAPPS) [15, 71]. The recent identification and characterisation of a likely pathogenic variant in the 5' UTR of the *APC* gene promoter region in a multi-generational family with FAP supports the inclusion of this region in multi-gene panel testing for adenomatous polyposis cases [72]. Further investigation is warranted to assess the relevance of variants in regulatory regions of polyposis genes in adenomatous polyposis patients without a known genetic cause.

In conclusion, the expansion of current clinical testing methodologies and analytical approaches beyond short-read sequencing will enable the detection of splicing-altering and non-coding regulatory variants as well as large structural and/or complex rearrangements, thereby increasing the yield of *APC*-related diagnoses.

## Specific considerations for *POLE* and *POLD1* genetic testing

Unlike any other hereditary cancer genes, loss-of-function variants in *POLE* and *POLD1* are not associated with cancer predisposition or PPAP. Only variants that affect the exonuclease activity of the polymerases, not affecting their DNA replicative ability, are the ones that should be clinically actioned. So far, these correspond to specific missense variants within the exonuclease domain [18]. Theoretically, in-frame insertions-deletions (indels) or variants that cause an in-frame splicing defect within the exonuclease domain might also be considered potentially pathogenic for PPAP.

Constitutional loss-of-function variants and variants located outside the exonuclease domain may predispose to very rare and severe autosomal recessive or dominant congenital disorders [73–78]. When identified in cancer patients, in absence of severe congenital problems, these variants should not be taken into consideration for genetic counselling or clinical purposes.

## Variant classification

An additional challenge for the accurate identification of hereditary adenomatous polyposis syndromes is the classification of germline variants of uncertain significance (VUS). The increasing application of germline multi-gene panel testing together with an increasing number of genes assessed by these panels and less phenotype-driven decision making on who gets testing has contributed to an increasing number of VUS. Uncertainty regarding the pathogenicity of a VUS impacts clinical management and the decision to test relatives for the variant.

The InSiGHT/ClinGen Hereditary Colorectal Cancer and Polyposis Variant Curation Expert Panel (<https://clinicalgenome.org/affiliation/50099/>; accessed 31/1/2025) was established by the International Society for Gastrointestinal Hereditary Tumours (InSiGHT) and the Clinical Genome Resource (ClinGen) with the task of developing gene-specific recommendations for variant interpretation. The panel has recently published gene-specific recommendations, based on the guidelines of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology (ACMG/AMP) [79], for the classification of variants in *APC* [80], representing gene- and disease-informed specifications. The criteria were applied to 10,228 unique *APC* variants with 41% and 61% of the VUS from ClinVar and LOVD databases, respectively, reclassified into clinically actionable classes of (likely) benign and (likely) pathogenic [81]. Current gene- and disease-informed specifications are under development for the *MUTYH*, *POLE* and *POLD1* genes, and other adenomatous and hamartomatous



polyposis genes will follow to optimize clinical management and opportunities for cancer prevention. Independently from the ClinGen panel, recommendations for the classification of *POLE* and *POLD1* variants in the context of cancer predisposition have also been published [18].

### Identification of monoallelic pathogenic variants in recessive adenomatous polyposis genes

Detecting heterozygous PVs in the *MUTYH*, *NTHL1*, *MSH3*, and *MBD4* genes poses a clinical challenge for counseling monoallelic carriers about their cancer risk and for establishing rational surveillance protocols.

The identification of heterozygous PVs in *MUTYH* through multi-gene panel testing is extremely common due to the high prevalence of heterozygotes in the general population (1–2%) [82]. Previous studies have reported a small but significantly increased risk of CRC in monoallelic *MUTYH* PV carriers [83–85]. However, a recent association study involving 58,998 CRC patients and 71,171 controls showed no association with CRC risk for the two most common PVs in *MUTYH* in European populations: p.Gly396Asp (rs36053993) and p.Tyr179Cys (rs34612342) [86]. Even if a low/moderate increased risk of CRC is considered, this would occur at population screening ages [87], and therefore, should not be over-interpreted. A large pan-cancer study of 10,389 patients involving 33 different types of cancer and 117,000 healthy controls revealed associated extracolonic risks for adrenocortical carcinoma, oesophageal carcinoma, sarcoma, prostate adenocarcinoma and kidney renal clear cell carcinoma, but not for colon or rectal cancers. The extra-colonic tumours from monoallelic carriers showed somatic loss of the wildtype *MUTYH* allele and presence of the characteristic *MUTYH*-associated mutational signatures [88].

The mentioned association study in 58,998 CRC patients and 71,171 controls found no association with CRC for the most commonly reported PVs in *NTHL1*-associated polyposis: c.268C>T, p.Gln90Ter (rs150766139) and c.859C>T, p.Gln287Ter (rs146347092) [86]. Likewise, analysis of 5,942 individuals with unexplained polyposis or familial CRC found no enrichment of monoallelic *NTHL1* PVs when compared to the general population [89]. No evidence of a somatic second hit affecting the wildtype allele of *NTHL1* or evidence of the SBS30 mutational signature associated with *NTHL1* deficiency, was observed in 11 CRCs and two adenomas from monoallelic *NTHL1* carriers [89]. Several studies have suggested an increased risk of breast cancer in *NTHL1* heterozygotes [90, 91]. In a pan-cancer analysis of 11,081 patients, 39 were heterozygous for *NTHL1* PVs, which was similar to the frequency observed in gnomAD non-Finnish European controls. Moreover, only two cancers

had the SBS30 mutational signature: a serous ovarian carcinoma with LOH of the *NTHL1* wildtype allele, and a prostate cancer without evidence of LOH [92]. In this study, no *NTHL1* heterozygous carriers that underwent colonoscopy were identified with polyposis ( $\geq 10$  polyps) [92].

Heterozygous germline *MBD4* PVs are associated with genetic predisposition to uveal melanoma. In these cases, the associated tumours show loss of the wildtype *MBD4* allele and, like in neoplasms from individuals with MANS, showed elevated tumour mutation burden enriched in CpG>TpG mutations (signature SBS1/SBS96) [35, 93–96]. The limited data suggests that constitutional monoallelic inactivation of *MBD4* does not increase the risk of CRC and/or polyposis [34, 35]. However, there has been a report of a monoallelic carrier of an *MBD4* pathogenic variant that developed ~30 adenomatous polyps and a CRC at 42 years of age with loss of *MBD4* protein expression, LOH of the wildtype *MBD4* allele and an enrichment of C>T transitions within CpG sites within the CRC tissue, consistent with biallelic inactivation of *MBD4* via a somatic second hit [97].

Available evidence for *MSH3*, although limited, suggests no increased risk of cancer in *MSH3* heterozygotes. In the mentioned pan-cancer study that included 11,081 patients, 12 carried monoallelic PVs in *MSH3*, a similar frequency observed in gnomAD non-Finnish European controls. A breast cancer and a prostate cancer developed by *MSH3* heterozygotes had *MSH3* LOH causing the loss of the wildtype allele. In those tumours that could be tested, no evidence of EMAS or loss of the *MSH3* protein was detected [92], as was observed in a patient with a monoallelic deletion of multiple *MSH3* exons [41].

In summary, the cumulative evidence, including a comprehensive genome-wide association study (GWAS) of imputed rare variants in hereditary adenomatous polyposis syndrome genes [86], suggests that monoallelic PVs in the recessively inherited genes *MUTYH*, *NTHL1*, *MSH3*, and *MBD4* do not increase the risk of CRC or polyposis. However, in monoallelic carriers, a somatic second hit can lead to biallelic inactivation, however, this occurs in a tissue- or organ-dependent manner, which may explain some of the extra-colonic cancer risks observed in heterozygotes.

### Tumour molecular features

Morphologically, the adenomatous polyps from people with constitutional PVs in *APC*, *AXIN2*, *POLE*, *POLD1*, MMR genes, *MUTYH*, *NTHL1*, *MBD4*, *MSH3* and *MLH3* are indistinguishable among the different syndromes and morphologically indistinguishable from sporadic adenomatous polyps. However, neoplasms that arise in each of these

adenomatous polyposis syndromes can present with unique molecular characteristics.

### Somatic second hits

For the autosomal dominant genes, somatic second hits via somatic mutations or loss of heterozygosity inactivating the wildtype allele, have been demonstrated for *APC*- and *AXIN2*-related adenomas and CRCs [98, 99]. In the case of *POLE*, tumour and normal tissue data, as well as evidence in yeast, indicate that polymerase  $\epsilon$  proofreading deficiency is haploinsufficient, not requiring the inactivation of the second allele to promote DNA damage (hypermutation) and adenoma/cancer initiation [17, 100]. Emerging evidence suggests that *POLD1* exonuclease domain mutations are haplosufficient, requiring a second hit (LOH of the wildtype allele) or MMR deficiency to cause hypermutability [100–102], although further research is needed.

### Gene-specific tumour mutational signatures associated with DNA repair deficiencies

Unique molecular characteristics have been described in neoplasms from defective DNA repair-related polyposis involving the *MUTYH*, *NTHL1*, *MBD4*, *MSH3*, *POLE* and *POLD1* genes and the MMR genes in CMMRD (Fig. 1). Except for CMMRD-related neoplasms, tumours (adenomas and cancer) developed in the context of these particular syndromes are predominantly MMR-proficient, although a subset of tumours that develop in patients with biallelic PVs in *MUTYH* or PVs in *POLE* or *POLD1* may show MMR deficiency as a result of biallelic somatic MMR gene mutations [103–105]. In general, defective DNA repair syndromes tend to result in higher tumour mutation burden (TMB) when compared to DNA repair-proficient neoplasms.

### MMR deficiencies

CMMRD tumours are characterised for a high TMB due to the presence of MMR deficiency. MMR deficiency may be identified using either microsatellite instability (MSI) testing (by PCR or next generation sequencing (NGS) analysis) or immunohistochemical (IHC) staining for the corresponding MMR proteins MLH1, MSH2, MSH6 and PMS2. In general, MMR-deficient tumours show specific COSMIC mutational signatures that represent the highly specific DNA repair deficiency, including signatures SBS6, 15, 21, 26, and 44, and ID2 and ID7. In CMMRD, detection of MMR deficiency in non-neoplastic tissues may be used as diagnostic test and to help interpret uncertain results from genetic testing [106].

MSH3 is also part of the DNA MMR machinery, detecting and repairing replication errors in long microsatellite repeats ( $\geq 2$  nucleotides). While *MLH1*, *MSH2*, *MSH6* or *PMS2* deficiencies affect predominantly mononucleotide and dinucleotide DNA repeat regions, biallelic *MSH3* deficiency affects tetranucleotide repeats, referred to as EMAST (Elevated Microsatellite Alterations at Selected Tetranucleotide repeats). EMAST is characteristic of tumours that develop in the context of MSH3-associated polyposis [39]. These tumours show a similar TMB and single base substitution mutation spectra to sporadic adenomas, but the proportion of small insertion/deletion (indels) is significantly higher [107]. The limited data indicates that *MLH3*-deficient tumours do not show MSI [44, 45], and it is still unknown if they have a unique mutational signature as observed in neoplasms associated with other DNA repair defects.

### BER deficiencies

Despite not reaching the high mutational burdens of MMR- or polymerase proofreading-defective tumours, BER-deficient tumours show elevated mutational burdens and glycosylase-specific mutational signatures. *MUTYH*-deficient adenomas and cancers developed in the context of MAP are characterised by COSMIC mutational signatures SBS18 and SBS36, related to an abundance C>A nucleotide transversions [103, 108–111]. *NTHL1*-deficient tumours harbour the highly specific signature SBS30, associated with an enrichment of C>T nucleotide transitions [32, 103, 112]. The high discriminatory accuracy of these mutational signatures have shown to be particularly useful for reclassifying VUS in these genes and has better characterised the tumour spectrum of the associated syndromes.

*MBD4* preferentially binds to 5-methylcytosine CpG:TpG mismatches which are the primary product of deamination at methyl CpG sites [113]. Biallelic loss of *MBD4* causes accumulation of G:T mismatches at CpG dinucleotides giving rise to excessively high levels of COSMIC mutational signature SBS1, or the highly similar SBS96 [35, 114], typically observed as having > 60% of single nucleotide variants being mCpG>TpG [34].

### Polymerase proofreading deficiency

The presence of PVs in *POLE* and *POLD1* affecting the exonuclease activity of polymerases  $\epsilon$  and  $\delta$ , respectively, causes uncorrected errors during DNA replication resulting in the highest tumour mutational burden, often exceeding 100 mut/Mb, and referred to as ultra-hypermutator phenotype. The pattern of somatic mutations is dominated by C>A transversions that are recognised as the COSMIC mutational signature SBS10, which has been further

differentiated as signatures associated with *POLE* deficiency (SBS10a, SBS10b and SBS28) or *POLD1* deficiency (SBS10c and SBS10d). When proofreading deficiency coexists with MMR deficiency, the tumour mutational spectra shift to SBS14 in the case of *POLE* mutations and SBS20 for *POLD1* [17, 18].

## Precision oncology in the context of adenomatous polyposis syndromes

Immunotherapy, i.e., immune checkpoint inhibitors, has proven highly effective for the treatment of cancers with high tumour mutational burdens due to higher neoantigen loads [115]. This is particularly relevant in MMR-deficient and polymerase proofreading-deficient cancers, regardless of tumour type [116–118]. Good responses have also been observed in *MUTYH*- and *MBD4*-deficient cancers [119, 120]. Despite the lack of published evidence, similar responses may be expected for *NTHL1*-deficient cancers.

The deregulation of Wnt/ $\beta$ -catenin signalling is related to the initiation and progression of various types of cancers, including the majority of sporadic CRCs. Despite the pivotal role of  $\beta$ -catenin in cancer development and progression, the Wnt/ $\beta$ -catenin pathway is highly conserved and is involved in numerous physiological processes, making it difficult to selectively target this pathway without causing significant off-target effects and toxicity. Additionally,  $\beta$ -catenin lacks an easily druggable binding site, further complicating the development of inhibitors against this protein. Nevertheless, inhibitors, antagonists and agonists have been designed to target different components of the Wnt signalling pathway [121]. Although the focus on Wnt research has advanced significantly, no Wnt-targeted treatments have yet been approved. There are, however, several preclinical investigations and clinical trials underway with molecules targeting Wnt signalling that include anti-FZD antibodies, FZD domain-containing recombinant proteins, inhibitors of  $\beta$ -catenin/CBP interactions, and Porcupine inhibitors [122]. Activation of the Wnt pathway in tumours can lead to an immunosuppressive tumour microenvironment, limiting the effectiveness of immune checkpoint inhibitors [123], however, this opens a therapeutic opportunity by combining Wnt pathway inhibitors with immunotherapy to enhance anti-tumour immune responses in these patients.

Readthrough of premature stop codons in *APC* appears as a promising therapeutic strategy for tumours with *APC* truncating mutations in preclinical studies. In particular, macrolide ZKN-0013 can suppress premature termination of protein translation induced by nonsense mutations in the *APC* gene, resulting in the restoration of active APC protein. In vivo efficacy of ZKN-0013 was demonstrated in

the *Apc*<sup>min</sup> mouse model, where it significantly reduced the number of intestinal polyps [124].

## Adenomatous polyposis with unknown genetic cause

The presence of multiple adenomas in the absence of constitutional PVs in known polyposis or CRC genes is relatively common in older screening age populations [125], likely reflecting an environmental component. While older age and male sex are non-modifiable adenoma risk factors, obesity, smoking and unhealthy (western) diet have been identified as the most significant modifiable risk factors for adenoma development [126–128]. Previous treatment with abdominopelvic radiotherapy or chemotherapy with alkylating agents for childhood or young adulthood cancer has also been associated with increased risk of polyposis of varying polyp histology [129].

Recent data suggests that the presence of intestinal Colibactin-producing *pks+Escherichia coli* might explain a relevant proportion of genetically unexplained adenomatous polyposes [130, 131]. *Pks+E. coli* corresponds to specific strains of the bacteria that harbour the polyketide synthases complex, known as the *pks* island [132, 133]. *Pks+E. coli* produces colibactin, a genotoxin that induces specific patterns of DNA damage, that correspond to the colibactin-associated COSMIC mutational signatures SBS88 and ID18 [134, 135]. The colibactin-associated mutation signatures may be detected in ~12% of sporadic CRCs [135] and in normal colonic mucosa from CRC patients [136]. The *APC* gene is one of the key targets of colibactin-related DNA damage [134]. Specifically, the *APC* c.835-8A>G splice variant occurs in the specific nucleotide sequence context targeted by colibactin-induced DNA damage and is significantly associated with the intra-tumoural presence of *pks+E. coli* in sporadic CRCs [135, 137]. The colibactin-associated *APC* mutations have been identified in a relevant number of adenomas and tumours from patients with unexplained adenomatous polyposis, demonstrating a bacterial aetiology. In a recent study, approximately 30% of unexplained adenomatous polyposis had at least one tumour harbouring the *APC* c.835-8A>G mutation, suggesting that *pks+E. coli* might be a key contributor underlying the aetiology of idiopathic adenomatous polyposes [130, 131].

The importance of adverse lifestyle and environmental factors on the risk of CRC has been extensively documented [138], but for adenomatous polyposis, the role of these factors has not been studied. However, it should be considered given their potential to influence the gut microbiome. The gut microbiome in FAP patients differs from that of the healthy population. It has been observed that



FAP patients have bacterial biofilms enriched with *E. coli* and *Bacteroides fragilis* [139], followed by the demonstration that co-colonising *Apc*<sup>min/+</sup> mice with *pks*+*E. coli* and enterotoxigenic *Bacteroides fragilis* accelerated tumourigenesis [140]. Similarly, transplanting *Apc*<sup>min/+</sup> mice with gut microbiota from CRC-affected patients activated the Wnt pathway and increased tumourigenic rates [140]. Other gut bacteria, including *Campylobacter jejuni* and *Fusobacterium nucleatum*, also showed similar effects [141, 142]. These studies not only highlight the importance of the gut microbiome in mitigating CRC risk among FAP patients but also warrant further studies investigating the interplay between the gut microbiome and environmental exposures in the pathogenesis of genetically unexplained adenomatous polyposis.

Recent efforts to identify new adenomatous polyposis genes have led to the identification of biallelic *MCM9* variants. MCM9, with MCM8, form a helicase hexameric complex involved in DNA replication (initiation), meiosis, repair of double-strand breaks via homologous recombination and DNA mismatch repair. Recessive inheritance of PVs has been linked to adenomatous polyposis, gastric cancer, and early-onset CRC, in addition to its prior documented association with infertility in both males and females related to hypogonadism and early development of germ cell tumours [143].

*BMPR2*, in addition to predisposing to pulmonary arterial hypertension, has been postulated as a gene involved in autosomal dominant polyposis, based on the recent identification of several polyposis patients with genetic variants in this gene [144]. Likewise, *FOCAD*, and Wnt signalling genes, including Wnt negative regulators *DKK4*, *HECW1*, *ITPR3*, and *WNT9B*, have been proposed as candidate causal genes for polyposis predisposition [145–147]. The evidence gathered to date for these genes is still insufficient to include them in clinical genetic diagnostics.

## Conclusions

Adenomatous polyposis occurs in the context of hereditary genetic syndromes caused by PVs in genes involved in the Wnt signalling pathway (*APC*, *AXIN2*), or DNA repair mechanisms (*POLE*, *POLD1*, *MUTYH*, *NTHL1*, *MBD4* and several MMR genes). Germline PVs in *APC* underlie the majority of adenomatous polyposis cases, followed by biallelic *MUTYH* PVs, and to a much lesser extent, PVs in the exonuclease domain of *POLE* or *POLD1*. Each adenomatous polyposis syndrome has its own distinct phenotypic features including variations in onset age, risks for different extracolonic cancers, clinical manifestations, and tumour molecular characteristics. In this regard, DNA repair-deficient tumours

exhibit gene-specific mutational signatures, which may aid in variant interpretation and classification.

Genetic testing for adenomatous polyposis can present challenges, such as the identification of somatic mosaicism, complex structural variants, and deep-intronic PVs, particularly in the *APC* gene. Another challenge is interpreting monoallelic PVs in recessively inherited genes. Over the past few decades, rapidly advancing genomic technologies have enabled the identification of bona-fide and candidate risk genes, and improved our understanding of genetic and non-genetic aetiologies. Despite the discovery of new polyposis predisposition genes, up to one-third of adenomatous polyposis patients do not receive a definitive genetic diagnosis, most of which are presumed to have a non-genetic cause. Through better molecular/mutational characterisation of neoplasms using new sequencing technologies and a deeper understanding of the gut microbiome, we anticipate continued discovery of novel molecular mechanisms associated with the development of adenomatous polyposis, paving the way for personalised medicine in these patients.

**Author contributions** J.E.J., J.V-E, D.D.B. and L.V. wrote the main manuscript text and contributed to the preparation of Fig. 1. All authors reviewed the manuscript.

**Funding** Open Access funding enabled and organized by CAUL and its Member Institutions

Laura Valle's research activity is funded by the Spanish Ministry of Science and Innovation (Agencia Estatal de Investigación), co-funded by FEDER funds -a way to build Europe- [PID2020-112595RB-I00, doi: AEI/<https://doi.org/10.13039/501100011033>, and “contrato predoctoral para la formación de doctores” (JV-E)]; Instituto de Salud Carlos III [CIBERONC CB16/12/00234, and PMP22/00064, co-funded by NextGenerationEU (MRR)/PRTR]; and Government of Catalonia [AGAUR 2021SGR01112, and CERCA Program for institutional support (IDIBELL)]. Daniel D. Buchanan is supported by a National Health and Medical Research Council of Australia (NHMRC) Investigator grant GNT1194896, a University of Melbourne Dame Kate Campbell Fellowship and by funding awarded to the The Colon Cancer Family Registry (CCFR, [www.coloncf.org](http://www.coloncf.org)) from the National Cancer Institute (NCI), National Institutes of Health (NIH) [award U01 CA167551].

**Data availability** No datasets were generated or analysed during the current study.

## Declarations

**Competing interests** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended

use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Valle L, Monahan KJ (2024) Genetic predisposition to Gastrointestinal polyposis: syndromes, tumour features, genetic testing, and clinical management. *Lancet Gastroenterol Hepatol* 9(1):68–82. [https://doi.org/10.1016/S2468-1253\(23\)00240-6](https://doi.org/10.1016/S2468-1253(23)00240-6)
- Kinzler KW, Nilbert MC, Su LK et al (1991) Identification of FAP locus genes from chromosome 5q21. *Science* 253(5020):661–665. <https://doi.org/10.1126/science.1651562>
- Nishisho I, Nakamura Y, Miyoshi Y et al (1991) Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. *Science* 253(5020):665–669. <https://doi.org/10.1126/science.1651563>
- Groden J, Thliveris A, Samowitz W et al (1991) Identification and characterization of the Familial adenomatous polyposis coli gene. *Cell* 66(3):589–600. [https://doi.org/10.1016/0092-8674\(81\)90021-0](https://doi.org/10.1016/0092-8674(81)90021-0)
- Joslyn G, Carlson M, Thliveris A et al (1991) Identification of deletion mutations and three new genes at the Familial polyposis locus. *Cell* 66(3):601–613. [https://doi.org/10.1016/0092-8674\(81\)90022-2](https://doi.org/10.1016/0092-8674(81)90022-2)
- Clevers H (2006) Wnt/beta-catenin signaling in development and disease. *Cell* 127(3):469–480. <https://doi.org/10.1016/j.cell.2006.10.018>
- (2012) Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 487(7407): 330–337
- Jansen AML, Goel A (2020) Mosaicism in patients with colorectal cancer or polyposis syndromes: A systematic review. *Clin Gastroenterol Hepatol* 18(9):1949–1960. <https://doi.org/10.1016/j.cgh.2020.02.049>
- Spier I, Drichel D, Kerick M et al (2016) Low-level APC mutational mosaicism is the underlying cause in a substantial fraction of unexplained colorectal adenomatous polyposis cases. *J Med Genet* 53(3):172–179. <https://doi.org/10.1136/jmedgenet-2015-103468>
- Aretz S, Stienen D, Friedrichs N et al (2007) Somatic APC mosaicism: a frequent cause of Familial adenomatous polyposis (FAP). *Hum Mutat* 28(10):985–992. <https://doi.org/10.1002/humu.20549>
- Yen T, Stanich PP, Axell L, Patel SG (1998) [Updated 2022]] APC-Associated Polyposis Conditions. In Adam MP ED, Mirzaa GM, (ed) *GeneReviews*® [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2023
- Zaffaroni G, Mannucci A, Koskenvuo L et al (2024) Updated European guidelines for clinical management of Familial adenomatous polyposis (FAP), MUTYH-associated polyposis (MAP), gastric adenocarcinoma, proximal polyposis of the stomach (GAPPS) and other rare adenomatous polyposis syndromes: a joint EHTG-ESCP revision. *Br J Surg* 111(5). <https://doi.org/10.1093/bjs/znae070>
- Knudsen AL, Bülow S, Tomlinson I et al (2010) Attenuated familial adenomatous polyposis: results from an international collaborative study. *Colorectal Dis* 12(10 Online): e243–9 <https://doi.org/10.1111/j.1463-1318.2010.02218.x>
- Beck SH, Karstensen JG, Bülow S et al (2024) Cancer risks in attenuated and classical Familial adenomatous polyposis: A nationwide cohort with matched, nonexposed individuals. *Am J Gastroenterol*. <https://doi.org/10.14309/ajg.0000000000003167>
- Li J, Woods SL, Healey S et al (2016) Point mutations in exon 1B of APC reveal gastric adenocarcinoma and proximal polyposis of the stomach as a Familial adenomatous polyposis variant. *Am J Hum Genet* 98(5):830–842. <https://doi.org/10.1016/j.ajhg.2016.03.001>
- Skat-Rørdam PA, Kaya Y, Qvist N et al (2024) Gastrointestinal manifestations in patients with gastric adenocarcinoma and proximal polyposis of the stomach (GAPPS): a systematic review with analysis of individual patient data. *Hered Cancer Clin Pract* 22(1):12. <https://doi.org/10.1186/s13053-024-00284-6>
- Palles C, Cazier JB, Howarth KM et al (2013) Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet* 45(2):136–144. <https://doi.org/10.1038/ng.2503>
- Mur P, Viana-Errasti J, García-Mulero S et al (2023) Recommendations for the classification of germline variants in the exonuclease domain of POLE and POLD1. *Genome Med* 15(1):85. <https://doi.org/10.1186/s13073-023-01234-y>
- Wimmer K, Beilken A, Nustede R et al (2017) A novel germline POLE mutation causes an early onset cancer prone syndrome mimicking constitutional mismatch repair deficiency. *Fam Cancer* 16(1):67–71
- Rosner G, Gluck N, Carmi S et al (2018) POLD1 and POLE gene mutations in Jewish cohorts of Early-Onset colorectal cancer and of multiple colorectal adenomas. *Dis Colon Rectum* 61(9):1073–1079. <https://doi.org/10.1097/DCR.0000000000001150>
- Lindsay H, Scollon S, Reuther J et al (2019) Germline POLE mutation in a child with hypermutated Medulloblastoma and features of constitutional mismatch repair deficiency. *Cold Spring Harb Mol Case Stud* 5(5). <https://doi.org/10.1101/mcs.a004499>
- Galati MA, Hodel KP, Gams MS et al (2020) Cancers from novel Pole-mutant mouse models provide insights into polymerase-mediated hypermutagenesis and immune checkpoint Blockade. *Cancer Res* 80(24):5606–5618. <https://doi.org/10.1158/0008-5472.CAN-20-0624>
- Sehested A, Meade J, Scheie D et al (2022) Constitutional POLE variants causing a phenotype reminiscent of constitutional mismatch repair deficiency. *Hum Mutat* 43(1):85–96. <https://doi.org/10.1002/humu.24299>
- Schamschula E, Kinzel M, Wernstedt A et al (2022) Teenage-Onset colorectal cancers in a digenic cancer predisposition syndrome provide clues for the interaction between mismatch repair and polymerase  $\Delta$  proofreading deficiency in tumorigenesis. *Biomolecules* 12(10). <https://doi.org/10.3390/biom12101350>
- Michaeli O, Ladany H, Erez A et al (2022) Di-genic inheritance of germline POLE and PMS2 pathogenic variants causes a unique condition associated with pediatric cancer predisposition. *Clin Genet* 101(4):442–447. <https://doi.org/10.1111/cge.14106>
- Lammi L, Arte S, Somer M et al (2004) Mutations in AXIN2 cause Familial tooth agenesis and predispose to colorectal cancer. *Am J Hum Genet* 74(5):1043–1050. <https://doi.org/10.1086/386293>
- Chan JM, Clendenning M, Joseland S et al (2021) Rare germline variants in the AXIN2 gene in families with colonic polyposis and colorectal cancer. *Fam Cancer*. <https://doi.org/10.1007/s10689-021-00283-9>
- Al-Tassan N, Chmiel NH, Maynard J et al (2002) Inherited variants of MYH associated with somatic G:C→T:A mutations in colorectal tumors. *Nat Genet* 30(2):227–232. <https://doi.org/10.1038/ng828>
- Cleary SP, Cotterchio M, Jenkins MA et al (2009) Germline muty human homologue mutations and colorectal cancer: a multisite case-control study. *Gastroenterology* 136(4):1251–1260. <https://doi.org/10.1053/j.gastro.2008.12.050>
- Nielsen M, Infante E, Brand R (1993) MUTYH polyposis. In: Adam MP, Mirzaa GM, Pagon RA et al (eds) *GeneReviews*®.

- Seattle (WA), University of Washington, Seattle. Copyright © 1993–2022, University of Washington, Seattle. GeneReviews is a registered trademark of the University of Washington, Seattle. All rights reserved
31. Belhadj S, Mur P, Navarro M et al (2017) Delineating the phenotypic spectrum of the NTHL1-Associated polyposis. *Clin Gastroenterol Hepatol* 15(3):461–462. <https://doi.org/10.1016/j.cgh.2016.09.153>
  32. Grolleman JE, de Voer RM, Elsayed FA et al (2019) Mutational Signature Analysis Reveals NTHL1 Deficiency to Cause a Multi-tumor Phenotype. *Cancer Cell* 35(2): 256–66.e5 <https://doi.org/10.1016/j.ccell.2018.12.011>
  33. Weren RD, Ligtenberg MJ, Kets CM et al (2015) A germline homozygous mutation in the base-excision repair gene NTHL1 causes adenomatous polyposis and colorectal cancer. *Nat Genet* 47(6):668–671 <https://doi.org/10.1038/ng.3287>
  34. Palles C, West HD, Chew E et al (2022) Germline MBD4 deficiency causes a multi-tumor predisposition syndrome. *Am J Hum Genet* 109(5):953–960. <https://doi.org/10.1016/j.ajhg.2022.03.018>
  35. Terradas M, Gonzalez-Abuin N, Garcia-Mulero S et al (2023) MBD4-associated neoplasia syndrome: screening of cases with suggestive phenotypes. *Eur J Hum Genet* 31(10):1185–1189. <https://doi.org/10.1038/s41431-023-01418-5>
  36. Wimmer K, Kratz CP, Vasen HF et al (2014) Diagnostic criteria for constitutional mismatch repair deficiency syndrome: suggestions of the European consortium ‘care for CMMRD’ (C4CMMRD). *J Med Genet* 51(6):355–365. <https://doi.org/10.1136/jmedgenet-2014-102284>
  37. Colas C, Guerrini-Rousseau L, Suerink M et al (2024) ERN GEN-TURIS guidelines on constitutional mismatch repair deficiency diagnosis, genetic counselling, surveillance, quality of life, and clinical management. *Eur J Hum Genet* 32(12):1526–1541. <https://doi.org/10.1038/s41431-024-01708-6>
  38. Gallon R, Brekelmans C, Martin M et al (2024) Constitutional mismatch repair deficiency mimicking Lynch syndrome is associated with hypomorphic mismatch repair gene variants. *NPJ Precis Oncol* 8(1):119. <https://doi.org/10.1038/s41698-024-00603-z>
  39. Adam R, Spier I, Zhao B et al (2016) Exome sequencing identifies biallelic MSH3 germline mutations as a recessive subtype of colorectal adenomatous polyposis. *Am J Hum Genet* 99(2):337–351. <https://doi.org/10.1016/j.ajhg.2016.06.015>
  40. Gavric A, Krajc M, Strnisa L, Gavric AU, Plut S (2024) MSH3-related adenomatous polyposis in a patient with the negative family history of colorectal polyps. *Gastroenterol Hepatol* 47(4):397–400. <https://doi.org/10.1016/j.gastrohep.2023.08.006>
  41. Villy MC, Masliah-Planchon J, Schnitzler A et al (2023) MSH3: a confirmed predisposing gene for adenomatous polyposis. *J Med Genet* 60(12):1198–1205. <https://doi.org/10.1136/jmg-2023-109341>
  42. Koi M, Leach BH, McGee S, Tseng-Rogenski SS, Burke CA, Carethers JM (2024) Compound heterozygous MSH3 germline variants and associated tumor somatic DNA mismatch repair dysfunction. *NPJ Precis Oncol* 8(1):12. <https://doi.org/10.1038/s41698-024-00511-2>
  43. Aelvoet AS, Hoekman DR, Redeker BJW et al (2023) A large family with MSH3-related polyposis. *Fam Cancer* 22(1):49–54. <https://doi.org/10.1007/s10689-022-00297-x>
  44. Olkinuora A, Nieminen TT, Martensson E et al (2019) Biallelic germline nonsense variant of MLH3 underlies polyposis predisposition. *Genet Med* 21(8):1868–1873. <https://doi.org/10.1038/s41436-018-0405-x>
  45. Johannessen KM, Karstensen JG, Rasmussen AO et al (2025) A novel case of biallelic MLH3 variants in a patient with rectal cancer and polyps. *Clin Genet*. <https://doi.org/10.1111/cge.14689>
  46. Terlouw D, Hes FJ, Suerink M et al (2023) APC mosaicism, not always isolated: two first-degree relatives with apparently distinct. *Gut* 72(11):2186–2187. <https://doi.org/10.1136/gutjnl-2022-328540>
  47. Jansen AM, Crobach S, Geurts-Giele WR et al (2017) Distinct patterns of somatic mosaicism in the APC gene in neoplasms from patients with unexplained adenomatous polyposis. *Gastroenterology* 152(3):546–9e3. <https://doi.org/10.1053/j.gastro.2016.10.040>
  48. Rofes P, González S, Navarro M et al (2021) Paired Somatic-Germline testing of 15 polyposis and colorectal Cancer-Predisposing genes highlights the role of APC mosaicism in de Novo Familial adenomatous polyposis. *J Mol Diagn* 23(11):1452–1459. <https://doi.org/10.1016/j.jmoldx.2021.07.024>
  49. Walker R, Joo JE, Mahmood K et al (2025) DNA mismatch repair (MMR) gene mosaicism is rare in people with MMR-deficient cancers. *Gastroenterology*. <https://doi.org/10.1053/j.gastro.2024.12.027>
  50. de Moraes FCA, Moretti NR, Sano VKT, Ngan CWT, Burbano RMR (2024) Genomic mosaicism in colorectal cancer and polyposis syndromes: a systematic review and meta-analysis. *Int J Colorectal Dis* 39(1):201. <https://doi.org/10.1007/s00384-024-04776-8>
  51. Terkelsen T, Larsen OH, Vang S, Jensen UB, Wikman F (2020) Deleterious mis-splicing of STK11 caused by a novel single-nucleotide substitution in the 3' polypyrimidine tract of intron five. *Mol Genet Genomic Med* 8(9):e1381. <https://doi.org/10.1002/mgg3.1381>
  52. Jelsig AM, Rønlund K, Gede LB et al (2023) Identification of a novel pathogenic deep intronic variant in PTEN resulting in pseudoexon inclusion in a patient with juvenile polyps. *J Hum Genet* 68(10):721–724. <https://doi.org/10.1038/s10038-023-01174-w>
  53. Forte G, Buonadonna AL, Fasano C et al (2024) Clinical and molecular characterization of SMAD4 splicing variants in patients with juvenile polyposis syndrome. *Int J Mol Sci* 25(14). <https://doi.org/10.3390/ijms25147939>
  54. Young CC, Horton C, Grzybowski J et al (2024) Solving missing heritability in patients with Familial adenomatous polyposis with DNA-RNA paired testing. *JCO Precis Oncol* 8:e2300404. <https://doi.org/10.1200/PO.23.00404>
  55. Bozsik A, Butz H, Grolmusz VK, Polgár C, Patócs A, Papp J (2023) Genome sequencing-based discovery of a novel deep intronic APC pathogenic variant causing exonization. *Eur J Hum Genet* 31(7):841–845. <https://doi.org/10.1038/s41431-023-01322-y>
  56. Nieminen TT, Pavicic W, Porkka N et al (2016) Pseudoexons provide a mechanism for allele-specific expression of APC in Familial adenomatous polyposis. *Oncotarget* 7(43):70685–70698. <https://doi.org/10.18632/oncotarget.12206>
  57. Spier I, Horpaopan S, Vogt S et al (2012) Deep intronic APC mutations explain a substantial proportion of patients with Familial or early-onset adenomatous polyposis. *Hum Mutat* 33(7):1045–1050. <https://doi.org/10.1002/humu.22082>
  58. Garza-Rodríguez ML, Treviño V, Pérez-Maya AA et al (2021) Identification of a novel pathogenic rearrangement variant of the APC gene associated with a variable spectrum of Familial cancer. *Diagnostics (Basel)* 11(3). <https://doi.org/10.3390/diagnostics11030411>
  59. Nakamura W, Hirata M, Oda S et al (2024) Assessing the efficacy of target adaptive sampling long-read sequencing through hereditary cancer patient genomes. *NPJ Genom Med* 9(1):11. <https://doi.org/10.1038/s41525-024-00394-z>
  60. Oda S, Ushiyama M, Nakamura W et al (2023) A complex rearrangement between APC and TP63 associated with Familial adenomatous polyposis identified by multimodal genomic analysis: a



- case report. *Front Oncol* 13:1205847. <https://doi.org/10.3389/fonc.2023.1205847>
61. Tanabe H, Koshizuka Y, Tanaka K et al (2024) High-resolution genetic analysis of whole APC gene deletions: a report of two cases and patient characteristics. *Hum Genome Var* 11(1):46. <http://doi.org/10.1038/s41439-024-00301-z>
  62. Scharf F, Leal Silva RM, Morak M et al (2022) Constitutional chromothripsis of the APC locus as a cause of genetic predisposition to colon cancer. *J Med Genet* 59(10):976–983. <https://doi.org/10.1136/jmedgenet-2021-108147>
  63. Su LK, Steinbach G, Sawyer JC, Hindi M, Ward PA, Lynch PM (2000) Genomic rearrangements of the APC tumor-suppressor gene in Familial adenomatous polyposis. *Hum Genet* 106(1):101–107. <https://doi.org/10.1007/s004399900195>
  64. Sedlazeck FJ, Lee H, Darby CA, Schatz MC (2018) Piercing the dark matter: bioinformatics of long-range sequencing and mapping. *Nat Rev Genet* 19(6):329–346. <https://doi.org/10.1038/s41576-018-0003-4>
  65. Lee JK, Kwon WK, Hong SN et al (2022) Necessity of multiplex ligation probe amplification in genetic tests: germline variant analysis of the APC gene in Familial adenomatous polyposis patients. *Cancer Genet* 262–263:95–101. <https://doi.org/10.1016/j.cancergen.2022.02.002>
  66. Friedl W, Aretz S (2005) Familial adenomatous polyposis: experience from a study of 1164 unrelated German polyposis patients. *Hered Cancer Clin Pract* 3(3):95–114. <https://doi.org/10.1186/1897-4287-3-3-95>
  67. Aretz S, Stienen D, Uhlhaas S et al (2005) Large submicroscopic genomic APC deletions are a common cause of typical Familial adenomatous polyposis. *J Med Genet* 42(2):185–192. <https://doi.org/10.1136/jmg.2004.022822>
  68. Sieber OM, Lamlum H, Crabtree MD et al (2002) Whole-gene APC deletions cause classical Familial adenomatous polyposis, but not attenuated polyposis or multiple colorectal adenomas. *Proc Natl Acad Sci U S A* 99(5):2954–2958. <https://doi.org/10.1073/pnas.042699199>
  69. Jain M, Koren S, Miga KH et al (2018) Nanopore sequencing and assembly of a human genome with ultra-long reads. *Nat Biotechnol* 36(4):338–345. <https://doi.org/10.1038/nbt.4060>
  70. Roberts RJ, Carneiro MO, Schatz MC (2013) The advantages of SMRT sequencing. *Genome Biol* 14(7):405. <https://doi.org/10.1186/gb-2013-14-6-405>
  71. Dixon K, Senz J, Kaurah P, Huntsman DG, Schrader KA (2021) Rare APC promoter 1B variants in gastric cancer kindreds unselected for fundic gland polyposis. *Gut* 70(7):1415–1416. <http://doi.org/10.1136/gutjnl-2020-321990>
  72. Young B, Neklasen DW, Clark K et al (2025) Functional characterization of a genetic variant in the 5' UTR of APC 1B promoter in a Familial adenomatous polyposis family. *Am J Med Genet A*:e63992. <https://doi.org/10.1002/ajmg.a.63992>
  73. Eason C, Aleisa A, Jones JR, Prijoles EJ, Wine Lee L (2020) Filling in the gaps on FLS syndrome: A case report and literature review. *Pediatr Dermatol* 37(5):915–917. <https://doi.org/10.1111/pde.14274>
  74. Thiffault I, Saunders C, Jenkins J et al (2015) A patient with polymerase E1 deficiency (POLE1): clinical features and overlap with DNA breakage/instability syndromes. *BMC Med Genet* 16:31. <http://doi.org/10.1186/s12881-015-0177-y>
  75. Logan CV, Murray JE, Parry DA et al (2018) DNA polymerase epsilon deficiency causes image syndrome with variable immunodeficiency. *Am J Hum Genet* 103(6):1038–1044. <https://doi.org/10.1016/j.ajhg.2018.10.024>
  76. Nakano T, Sasahara Y, Kikuchi A et al (2022) Novel POLE mutations identified in patients with IMAGE-I syndrome cause aberrant subcellular localisation and protein degradation in the nucleus. *J Med Genet* 59(11):1116–1122. <https://doi.org/10.1136/jmedgenet-2021-108300>
  77. Shastry S, Simha V, Godbole K et al (2010) A novel syndrome of mandibular hypoplasia, deafness, and progeroid features associated with lipodystrophy, undescended testes, and male hypogonadism. *J Clin Endocrinol Metab* 95(10):E192–E197. <https://doi.org/10.1210/jc.20100419>
  78. Murdocca M, Spitalieri P, D'Apice MR, Novelli G, Sangiuolo F (2023) From cue to meaning: the involvement of POLD1 gene in DNA replication, repair and aging. *Mech Ageing Dev* 211:111790. <https://doi.org/10.1016/j.mad.2023.111790>
  79. Richards S, Aziz N, Bale S et al (2015) Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American college of medical genetics and genomics and the association for molecular pathology. *Genet Med* 17(5):405–424. <https://doi.org/10.1038/gim.2015.30>
  80. Spier I, Yin X, Richardson M et al (2024) Gene-specific ACMG/AMP classification criteria for germline APC variants: recommendations from the ClinGen insight hereditary colorectal cancer/polypsis variant curation expert panel. *Genet Med* 26(2):100992. <https://doi.org/10.1016/j.gim.2023.100992>
  81. Yin X, Richardson M, Laner A et al (2024) Large-scale application of ClinGen-InSiGHT APC-specific ACMG/AMP variant classification criteria leads to substantial reduction in VUS. *Am J Hum Genet* 111(11):2427–2443. <https://doi.org/10.1016/j.ajhg.2024.09.002>
  82. Win AK, Jenkins MA, Dowty JG et al (2017) Prevalence and penetrance of major genes and polygenes for colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 26(3):404–412. <https://doi.org/10.1158/1055-9965>
  83. Win AK, Cleary SP, Dowty JG et al (2011) Cancer risks for monoallelic MUTYH mutation carriers with a family history of colorectal cancer. *Int J Cancer* 129(9):2256–2262. <https://doi.org/10.1002/ijc.25870>
  84. Ma X, Zhang B, Zheng W (2014) Genetic variants associated with colorectal cancer risk: comprehensive research synopsis, meta-analysis, and epidemiological evidence. *Gut* 63(2):326–336. <http://doi.org/10.1136/gutjnl-2012-304121>
  85. Patel R, McGinty P, Cuthill V, Hawkins M, Clark SK, Latchford A (2021) Risk of colorectal adenomas and cancer in monoallelic carriers of MUTYH pathogenic variants: a single-centre experience. *Int J Colorectal Dis* 36(10):2199–2204. <https://doi.org/10.1007/s00384-021-03983-x>
  86. Mahmood K, Thomas M, Qu C et al (2023) Elucidating the Risk of Colorectal Cancer for Variants in Hereditary Colorectal Cancer Genes. *Gastroenterology* 165(4): 1070-6 e3 <https://doi.org/10.1053/j.gastro.2023.06.032>
  87. Breen KE, Katona BW, Catchings A et al (2022) An updated counseling framework for moderate-penetrance colorectal cancer susceptibility genes. *Genet Med* 24(12):2587–2590. <https://doi.org/10.1016/j.gim.2022.08.027>
  88. Barreiro RAS, Sabbaga J, Rossi BM et al (2022) Monoallelic deleterious MUTYH germline variants as a driver for tumorigenesis. *J Pathol* 256(2):214–222. <https://doi.org/10.1002/path.5829>
  89. Elsayed FA, Grolleman JE, Ragunathan A et al (2020) Monoallelic NTHL1 Loss-of-Function variants and risk of polyposis and colorectal cancer. *Gastroenterology* 159(6):2241–3e6. <https://doi.org/10.1053/j.gastro.2020.08.042>
  90. Nurmi AK, Peltari LM, Kiiski JI et al (2023) NTHL1 is a recessive cancer susceptibility gene. *Sci Rep* 13(1):21127. <https://doi.org/10.1038/s41598-023-47441-w>
  91. Li N, Zethoven M, McInerney S et al (2021) Evaluation of the association of heterozygous germline variants in NTHL1 with breast cancer predisposition: an international multi-center study of 47,180 subjects. *NPJ Breast Cancer* 7(1):52. <https://doi.org/10.1038/s41523-021-00255-3>



92. Salo-Mullen EE, Maio A, Mukherjee S et al (2021) Prevalence and characterization of biallelic and monoallelic NTHL1 and MSH3 variant carriers from a Pan-Cancer patient population. *JCO Precis Oncol* 5. <https://doi.org/10.1200/PO.20.00443>
93. Derrien AC, Rodrigues M, Eeckhoutte A et al (2021) Germline MBD4 mutations and predisposition to uveal melanoma. *J Natl Cancer Inst* 113(1):80–87. <https://doi.org/10.1093/jnci/djaa047>
94. Rodrigues M, Mobuchon L, Houy A et al (2018) Outlier response to anti-PD1 in uveal melanoma reveals germline MBD4 mutations in hypermutated tumors. *Nat Commun* 9(1):1866. <https://doi.org/10.1038/s41467-018-04322-5>
95. Johansson PA, Stark A, Palmer JM et al (2019) Prolonged stable disease in a uveal melanoma patient with germline MBD4 nonsense mutation treated with pembrolizumab and ipilimumab. *Immunogenetics* 71(5–6):433–436. <https://doi.org/10.1007/s00251-019-01108-x>
96. Villy MC, Le Ven A, Le Mentec M et al (2024) Familial uveal melanoma and other tumors in 25 families with monoallelic germline MBD4 variants. *J Natl Cancer Inst* 116(4):580–587. <https://doi.org/10.1093/jnci/djad248>
97. Tanakaya K, Kumamoto K, Tada Y et al (2019) A germline MBD4 mutation was identified in a patient with colorectal oligopolyposis and earlyonset cancer: A case report. *Oncol Rep* 42(3):1133–1140. <https://doi.org/10.3892/or.2019.7239>
98. Chan JM, Clendenning M, Joseland S et al (2022) Rare germline variants in the AXIN2 gene in families with colonic polyposis and colorectal cancer. *Fam Cancer* 21(4):399–413. <https://doi.org/10.1007/s10689-021-00283-9>
99. Moller P, Ahadova A, Kloor M et al (2025) Colorectal carcinogenesis in the Lynch syndromes and Familial adenomatous polyposis: trigger events and downstream consequences. *Hered Cancer Clin Pract* 23(1):3. <https://doi.org/10.1186/s13053-025-00305-y>
100. Robinson PS, Coorens THH, Palles C et al (2021) Increased somatic mutation burdens in normal human cells due to defective DNA polymerases. *Nat Genet* 53(10):1434–1442. <https://doi.org/10.1038/s41588-021-00930-y>
101. Andrianova MA, Seplyarskiy VB, Terradas M et al (2024) Discovery of recessive effect of human polymerase  $\Delta$  proofreading deficiency through mutational analysis of POLD1-mutated normal and cancer cells. *Eur J Hum Genet* 32(7):837–845. <https://doi.org/10.1038/s41431-024-01598-8>
102. Haradhvala NJ, Polak P, Stojanov P et al (2016) Mutational strand asymmetries in cancer genomes reveal mechanisms of DNA damage and repair. *Cell* 164(3):538–549. <https://doi.org/10.1016/j.cell.2015.12.050>
103. Walker R, Joo JE, Mahmood K et al (2025) Adenomas from individuals with pathogenic biallelic variants in the MUTYH and NTHL1 genes demonstrate base excision repair tumour mutational signature profiles similar to colorectal cancers, expanding potential diagnostic and variant classification applications. *Transl Oncol* 52:102266. <https://doi.org/10.1016/j.tranon.2024.102266>
104. Elsayed FA, Kets CM, Ruano D et al (2015) Germline variants in POLE are associated with early onset mismatch repair deficient colorectal cancer. *Eur J Hum Genet* 23(8):1080–1084. <https://doi.org/10.1038/ejhg.2014.242>
105. Morak M, Heidenreich B, Keller G et al (2014) Biallelic MUTYH mutations can mimic Lynch syndrome. *Eur J Hum Genet* 22(11):1334–1337. <https://doi.org/10.1038/ejhg.2014.15>
106. Chung J, Negm L, Bianchi V et al (2023) Genomic microsatellite signatures identify germline mismatch repair deficiency and risk of cancer onset. *J Clin Oncol* 41(4):766–777. <https://doi.org/10.1200/JCO.21.02873>
107. Perne C, Peters S, Cartolano M et al (2021) Variant profiling of colorectal adenomas from three patients of two families with MSH3-related adenomatous polyposis. *PLoS ONE* 16(11):e0259185. <https://doi.org/10.1371/journal.pone.0259185>
108. Georgeson P, Pope BJ, Rosty C et al (2021) Evaluating the utility of tumour mutational signatures for identifying hereditary colorectal cancer and polyposis syndrome carriers. *Gut* 70(11):2138–2149. <https://doi.org/10.1136/gutjnl-2019-320462>
109. Viel A, Bruselles A, Meccia E et al (2017) A specific mutational signature associated with DNA 8-Oxoguanine persistence in MUTYH-defective colorectal cancer. *EBioMedicine* 20:39–49. <https://doi.org/10.1016/j.ebiom.2017.04.022>
110. Georgeson P, Harrison TA, Pope BJ et al (2022) Identifying colorectal cancer caused by biallelic MUTYH pathogenic variants using tumor mutational signatures. *Nat Commun* 13(1):3254. <https://doi.org/10.1038/s41467-022-30916-1>
111. Pilati C, Shinde J, Alexandrov LB et al (2017) Mutational signature analysis identifies MUTYH deficiency in colorectal cancers and adrenocortical carcinomas. *J Pathol* 242(1):10–15. <https://doi.org/10.1002/path.4880>
112. Drost J, van Boxtel R, Blokzijl F et al (2017) Use of CRISPR-modified human stem cell organoids to study the origin of mutational signatures in cancer. *Science* 358(6360):234–238. <https://doi.org/10.1126/science.aao3130>
113. Millar CB, Guy J, Sansom OJ et al (2002) Enhanced CpG mutability and tumorigenesis in MBD4-deficient mice. *Science* 297(5580):403–405. <https://doi.org/10.1126/science.1073354>
114. Degasperis A, Zou X, Amarante TD et al (2022) Substitution mutational signatures in whole-genome-sequenced cancers in the UK population. *Science* 376(6591). <https://doi.org/10.1126/science.abl9283>
115. Roudko V, Bozkus CC, Orfanelli T et al (2020) Shared Immunogenic Poly-Epitope frameshift mutations in microsatellite unstable tumors. *Cell* 183(6):1634–49e17. <https://doi.org/10.1016/j.cell.2020.11.004>
116. Sha D, Jin Z, Budczies J, Kluck K, Stenzinger A, Sinicrope FA (2020) Tumor mutational burden as a predictive biomarker in solid tumors. *Cancer Discov* 10(12):1808–1825. <https://doi.org/10.1158/2159-8290.CD-20-0522>
117. Cecchini M, Sundar R (2024) Proofreading the way: immune checkpoint inhibitors in polymerase  $\epsilon$ /polymerase  $\delta$  (POLE/POLD1)-altered colorectal cancer. *Ann Oncol* 35(7):582–584. <https://doi.org/10.1016/j.annonc.2024.04.006>
118. Tommasi O, Marchetti M, Tripepi M et al (2025) PD-1 and PD-L1 expression in endometrial cancer: A systematic review of the literature. *J Clin Med* 14(2). <https://doi.org/10.3390/jcm14020401>
119. Volkov NM, Yanus GA, Ivantsov AO et al (2020) Efficacy of immune checkpoint blockade in MUTYH-associated hereditary colorectal cancer. *Invest New Drugs* 38(3):894–898. <https://doi.org/10.1007/s10637-019-00842-z>
120. Saint-Ghislain M, Derrien AC, Geoffrois L et al (2022) MBD4 deficiency is predictive of response to immune checkpoint inhibitors in metastatic uveal melanoma patients. *Eur J Cancer* 173:105–112. <https://doi.org/10.1016/j.ejca.2022.06.033>
121. Zhang Y, Wang X (2020) Targeting the Wnt/ $\beta$ -catenin signaling pathway in cancer. *J Hematol Oncol* 13(1):165. <https://doi.org/10.1186/s13045-020-00990-3>
122. Iluta S, Nistor M, Buruiana S, Dima D (2024) Wnt signaling pathway in tumor biology. *Genes (Basel)* 15(12). <https://doi.org/10.3390/genes15121597>
123. Luke JJ, Bao R, Sweis RF, Spranger S, Gajewski TF (2019) WNT/ $\beta$ -catenin pathway activation correlates with immune exclusion across human cancers. *Clin Cancer Res* 25(10):3074–3083. <https://doi.org/10.1158/1078-0432.CCR-18-1942>
124. Graf MR, Apte S, Terzo E et al (2023) Novel read through agent: ZKN-0013 demonstrates efficacy in APC<sup>min</sup> model of Familial

- adenomatous polyposis. *J Mol Med (Berl)*. <https://doi.org/10.1007/s00109-023-02291-x>
125. Alexander JL, Johnston BJ, Smith TJ et al (2021) Low referral rates for genetic assessment of patients with multiple adenomas in united Kingdom bowel cancer screening programs. *Dis Colon Rectum* 64(9):1058–1063. <https://doi.org/10.1097/DCR.00000000000001972>
  126. Choe AR, Song EM, Seo H et al (2024) Different modifiable risk factors for the development of non-advanced adenoma, advanced adenomatous lesion, and sessile serrated lesions, on screening colonoscopy. *Sci Rep* 14(1):16865. <https://doi.org/10.1038/s41598-024-67822-z>
  127. Xu J, Chi P, Qin K et al (2023) Association between lifestyle and dietary preference factors and conventional adenomas and serrated polyps. *Front Nutr* 10:1269629. <https://doi.org/10.3389/fnu.2023.1269629>
  128. Kim J, Nath K, Schmidlin K et al (2023) Hierarchical contribution of individual lifestyle factors and their interactions on adenomatous and serrated polyp risk. *J Gastroenterol* 58(9):856–867. <https://doi.org/10.1007/s00535-023-02004-8>
  129. Biller LH, Ukaegbu C, Dhingra TG et al (2020) A Multi-Institutional cohort of Therapy-Associated polyposis in childhood and young adulthood cancer survivors. *Cancer Prev Res (Phila)* 13(3):291–298. <https://doi.org/10.1158/1940-6207.CAPR-19-0416>
  130. Terlouw D, Suerink M, Boot A, van Wezel T, Nielsen M, Morreau H (2020) Recurrent APC splice variant c.835-8A>G in patients with unexplained colorectal polyposis fulfilling the colibactin mutational signature. *Gastroenterology* 159(4):1612–4e5. <https://doi.org/10.1053/j.gastro.2020.06.055>
  131. Terlouw D, Boot A, Ducarmon QR et al (2024) Enrichment of colibactin-associated mutational signatures in unexplained colorectal polyposis patients. *BMC Cancer* 24(1):104. <https://doi.org/10.1186/s12885-024-11849-y>
  132. Nougayrede JP, Homburg S, Taieb F et al (2006) *Escherichia coli* induces DNA double-strand breaks in eukaryotic cells. *Science* 313(5788):848–851. <https://doi.org/10.1126/science.1127059>
  133. Buc E, Dubois D, Sauvanet P et al (2013) High prevalence of mucosa-associated *E. coli* producing cyclomodulin and genotoxin in colon cancer. *PLoS ONE* 8(2):e56964. <https://doi.org/10.1371/journal.pone.0056964>
  134. Pleguezuelos-Manzano C, Puschhof J, Rosendahl Huber A et al (2020) Mutational signature in colorectal cancer caused by genotoxic pks(+) *E. coli*. *Nature* 580(7802):269–273. <https://doi.org/10.1038/s41586-020-2080-8>
  135. Rosendahl Huber A, Pleguezuelos-Manzano C, Puschhof J et al (2024) Improved detection of colibactin-induced mutations by genotoxic *E. coli* in organoids and colorectal cancer. *Cancer cell* 42(3): 487–96 e6 <https://doi.org/10.1016/j.ccell.2024.02.009>
  136. Chen B, Ramazzotti D, Heide T et al (2023) Contribution of pks(+) *E. coli* mutations to colorectal carcinogenesis. *Nat Commun* 14(1):7827. <https://doi.org/10.1038/s41467-023-43329-5>
  137. Joo JE, Chu YL, Georgeson P et al (2024) Intratumoral presence of the genotoxic gut bacteria pks(+) *E. coli*, enterotoxigenic *Bacteroides fragilis*, and *Fusobacterium nucleatum* and their association with clinicopathological and molecular features of colorectal cancer. *Br J Cancer*. <https://doi.org/10.1038/s41416-023-02554-x>
  138. Gunter MJ, Alhomoud S, Arnold M et al (2019) Meeting report from the joint IARC-NCI international cancer seminar series: a focus on colorectal cancer. *Ann Oncol* 30(4):510–519. <https://doi.org/10.1093/annonc/mdz044>
  139. Dejea CM, Fathi P, Craig JM et al (2018) Patients with Familial adenomatous polyposis harbor colonic biofilms containing tumorigenic bacteria. *Science* 359(6375):592–597. <https://doi.org/10.1126/science.aah3648>
  140. Li L, Li X, Zhong W et al (2019) Gut microbiota from colorectal cancer patients enhances the progression of intestinal adenoma in apc. *EBioMedicine* 48:301–315. <https://doi.org/10.1016/j.ebiom.2019.09.021>
  141. He Z, Gharaibeh RZ, Newsome RC et al (2019) *Campylobacter jejuni* promotes colorectal tumorigenesis through the action of cytolethal distending toxin. *Gut* 68(2):289–300. <https://doi.org/10.1136/gutjnl-2018-317200>
  142. Kostic AD, Chun E, Robertson L et al (2013) *Fusobacterium nucleatum* potentiates intestinal tumorigenesis and modulates the tumor-immune microenvironment. *Cell Host Microbe* 14(2):207–215. <https://doi.org/10.1016/j.chom.2013.07.007>
  143. Helderman NC, Yang T, Palles C et al (2024) Clinical syndromes linked to biallelic germline variants in MCM8 and MCM9. *MedRxiv*: 2024.10.30.24315828. <https://doi.org/10.1101/2024.10.30.24315828>
  144. Bonjoch L, Fernandez-Rozadilla C, Alvarez-Barona M et al (2023) BMPR2 as a novel predisposition gene for hereditary colorectal polyposis. *Gastroenterology*. <https://doi.org/10.1053/j.gastro.2023.03.006>
  145. Terradas M, Munoz-Torres PM, Belhadj S et al (2019) Contribution to colonic polyposis of recently proposed predisposing genes and assessment of the prevalence of NTHL1- and MSH3-associated polyposes. *Hum Mutat* 40(11):1910–1923. <https://doi.org/10.1002/humu.23853>
  146. Weren RD, Venkatachalam R, Cazier JB et al (2015) Germline deletions in the tumour suppressor gene FOCAD are associated with polyposis and colorectal cancer development. *J Pathol* 236(2):155–164. <https://doi.org/10.1002/path.4520>
  147. Quintana I, Terradas M, Mur P et al (2023) Wnt genes in colonic polyposis predisposition. *Genes Dis* 10(3):753–757. <https://doi.org/10.1016/j.gendis.2022.12.002>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.