



# Management of inoperable endometrial cancer

Supakorn Pitakkarnkul, MD<sup>1</sup>, Saranya Chanpanitkitchot, MD<sup>2</sup>, Siriwan Tangjitgamol, MD<sup>3,4</sup>

<sup>1</sup>Division of Gynecologic Oncology, National Cancer Institute, <sup>2</sup>Department of Obstetrics and Gynecology, Rajavithi Hospital, Rangsit University College of Medicine, <sup>3</sup>Women's Care Center, MedPark Hospital, <sup>4</sup>Department of Obstetrics and Gynecology, Faculty of Medicine Vajira Hospital, Navamindradhiraj University, Bangkok, Thailand

Some endometrial cancer (EMC) patients are not good candidates for primary surgery. The three major types of treatment for inoperable EMC are radiation therapy, chemotherapy, or their combination as neoadjuvant treatment before surgery. Radiation therapy alone (of different modes) has been used as the sole definitive therapeutic modality, particularly for early-stage disease that is limited to the uterine body and cervix with or without parametrial invasion. The most common treatment modality is neoadjuvant treatment before surgery. Postoperative adjuvant treatment is also occasionally used, depending mainly on the sites of the disease and the results of surgery. Data on neoadjuvant hormonal or radiation therapy are limited, with studies focusing on laboratory outcomes or having only a small number of patients. Most neoadjuvant treatments before surgery involved chemotherapy and fewer combined chemoradiotherapy. Surgery was generally performed, particularly in patients who had shown responses or at least stable disease to neoadjuvant treatment. Perioperative outcomes after neoadjuvant treatment were superior to those after primary surgery, whereas survival data were still inconsistent. Features that had or tended to have a favorable prognosis were younger age, early-stage disease, response to neoadjuvant treatment, low preoperative cancer antigen-125 level, and optimal surgery. Among different modalities of neoadjuvant treatment, which has become a frequent mode of treatment, neoadjuvant chemotherapy was more common than radiation therapy alone or chemoradiation.

**Keywords:** Endometrial cancer; Radiotherapy; Chemotherapy; Quality of life

## Introduction

Surgical treatment with total hysterectomy and bilateral salpingo-oophorectomy is the standard primary treatment for endometrial cancer (EMC). Unless the patients show some risk features necessitating adjuvant treatment, surgery may be the only treatment they will receive.

The prognosis of patients with EMC is generally good because most patients have early-stage disease at presentation. However, approximately 10-15% of EMC patients present with advanced-stage cancer [1]. This latter group of patients is unlikely to undergo safe or successful surgical removal; therefore, they are not good candidates for primary surgery. Moreover, the general condition of some patients at the time of diagnosis may not be optimal for performing major surgery. The combined incidence of these two conditions is also not an uncommon phenomenon. Therefore, surgery is not an appropriate primary treatment option for such patients. In these two scenarios, neoadjuvant therapy before definitive surgical treatment is likely to be a better option. Another

relevant scenario involves the presence of severe comorbid illnesses that are frequently found in elderly and frail EMC patients. This latter group of patients may not be able to undergo definite surgical treatment at any time, especially when their life expectancy may be already shortened by

Received: 2021.07.05. Revised: 2021.12.18. Accepted: 2022.03.07.  
Corresponding author: Siriwan Tangjitgamol, MD  
Department of Obstetrics and Gynecology, Faculty of Medicine Vajira Hospital, 3333 Rama IV Rd, Khlong Toei, Bangkok 10110, Thailand  
E-mail: siriwanonco@yahoo.com  
<https://orcid.org/0000-0001-5367-8884>

Siriwan Tangjitgamol has been an Editorial Board of Obstetrics & Gynecology Science; however, he was not involved in the peer reviewer selection, evaluation, or decision process of this article. Otherwise, no other potential conflicts of interest relevant to this article were reported.

Articles published in Obstet Gynecol Sci are open-access, distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2022 Korean Society of Obstetrics and Gynecology

medical illnesses themselves.

For cases showing advanced disease and an equivocal possibility of optimal cytoreduction on physical examination, several imaging studies, such as computerized tomography (CT) scans, magnetic resonance imaging (MRI), and positron emission tomography scans are currently available to help surgeons evaluate the extent of disease prior to surgery [2-5]. This approach is especially helpful for high-grade or serous carcinomas, which frequently show a higher probability of extrauterine or distant metastasis than other types of EMC [2,6].

Most of the existing information regarding the treatment of inoperable EMC is derived from retrospective studies or case series. Without evidence-based data from randomized studies, combined data from the available reports may help gynecologic oncologists to select an alternative treatment option for EMC patients who are not candidates for primary surgery. This study collected and summarized data from a literature review of relevant studies based on the treatment intention (definitive or neoadjuvant). Treatment administered with a palliative aim, either chemotherapy, hormonal therapy, or best supportive care, is beyond the scope of this review.

## Methods

We searched the MEDLINE database for studies and reports published from January 1997 to December 2020. We searched for all studies that reported treatments for EMC patients who did not receive surgery as the primary treatment due to advanced disease or medical illnesses. Only studies with curative intent (definitive treatment) were included. Studies that provided palliative treatment were excluded.

## Results

A review of the literature revealed that most of the studies on this topic were retrospective studies without randomized trials. The three major types of treatment used for inoperable EMC were radiation therapy, chemotherapy, combinations of radiation therapy and chemotherapy, and interval surgery. The findings for these types of treatment are described below.

### 1. Radiation as a definitive therapy

Radiation has long been used as a definitive therapy, particularly for EMC patients with comorbid illnesses that preclude surgery and those with disease that is limited to the pelvis. The different modes of radiation therapy that have been used are external pelvic radiation therapy (EPRT), brachytherapy (intracavitary radiation therapy [ICRT]), or their combination.

One study suggested aggressive (multimodal) radiation therapy even for early-stage EMC [7]. This study described the findings for 101 EMC patients aged 39-94 years who could not undergo surgery due to medical problems [7]. All patients had clinical stage I disease, with the majority (82%) showing cancer invading less than half of the myometrium, while the remaining patients showed no myometrial invasion. The modes of radiation used were EPRT plus split-field and ICRT (61%), ICRT alone (26%), EPRT plus ICRT (10%), and EPRT alone (3%). The 5-year disease-free survival (DFS) and overall survival (OS) rates were >80%. The survival rates of patients who were older than 75 years were lower than those of younger patients: DFS, 55% vs. 84%; OS, 78% vs. 84%.

Other studies have reported inconsistent data regarding the benefits of adding ICRT to EPRT [8,9]. One study used the National Cancer Institute's Surveillance, Epidemiology, and End Results public database to analyze the prognostic factors and outcomes in 997 stage I-II EMC patients who did not undergo surgical treatment [8]. Among these patients, 39% received radiation therapy as EPRT alone (21%) or ICRT with or without EPRT (18%). The type of radiation therapy was not associated with OS or EMC-specific survival. On the other hand, EPRT alone was sufficient, since the additional use of ICRT did not improve survival in comparison with EPRT alone [8]. However, in another small study involving 11 EMC patients with stage I-III disease who could not undergo ICRT after EPRT and image-guided stereotactic body radiation [9], nearly half (45%) of the patients experienced locoregional disease progression at a median follow-up of 10 months. Progression-free survival (PFS) at 18 months was only 41%. Stage IB disease was much worse than stage IA disease, with a PFS of 33% in comparison with 100%.

The type of ICRT used appeared to have no major influence on the outcome. One study reported successful treatment of 45 clinical low-risk inoperable EMC patients (stage I, small grade 1-2 tumors <2 cm, with less than 50% myo-

metrial invasion) with the use of only high-dose rate image-guided tandem and cylinder ICRT as a definitive treatment [10]. Vaginal bleeding was controlled in 97% of patients. Clinical complete radiologic response from MRI and 2-year locoregional control were achieved in 90% of the patients, with cancer-specific survival and OS rates of 86% and 97%, respectively [10]. Another study also reported a high success rate of treatment with low-dose rate ICRT using Heyman's capsules in 44 cases of medically inoperable clinical stage I EMC [11], with 5-year disease-specific survival and OS rates of 88% and 61%, respectively. The authors concluded that low dose rate ICRT yielded outcomes similar to those of surgery and high dose rate ICRT.

Data from the studies that used radiation therapy as a definitive treatment for EMC are shown in Table 1 [7,9-29]. Aside from the disease stage, the detailed characteristics of EMC may facilitate tailoring of the type of radiation therapy. The American Brachytherapy Society has supported the use of MRI and 3D planning for definitive radiation treatment of medically inoperable EMC [30]. The gross tumor volume, depth of myometrial invasion, clinical target volume, and organs at risk can be assessed with this approach. For clinical stage I tumors without deep myometrial invasion and lymph node involvement, ICRT alone can be used. On the other hand, additional EPRT may be considered when MRI is not available to assess these features. In brief, the society summarized that ICRT is sufficient for stage I disease with tumor grade 1-2, and no deep myometrial invasion on MRI. External pelvic radiation along with ICRT is recommended in stage I disease with deep myometrial invasion or no MRI assessment (or only CT scan), and in stage II-III EMC that is still limited within the pelvis [30].

The American Brachytherapy Society also reported survival outcomes according to the type of radiation therapy from studies over different periods of time [30]. In summary, the 5-year cancer-specific survival of patients with stage I EMC treated with only ICRT (either low- or high-dose) or with additional EPRT ranged from 65-85% and up to 95%, respectively. The survival rates with only ICRT or with additional EPRT were, respectively, 40-69% and 88% for stage II disease and 14% and 57% for stage III disease. The differences in the ranges of survival rates obtained with different modes of radiation therapy across various studies may be partly due to unavailable or inconsistent use of MRI to assess tumor features (especially in clinical stage I EMC) and thereby tailor

the type of radiation in each study, especially in studies conducted in the remote past.

## 2. Neoadjuvant treatment prior to surgery

In the modern era, with rapid advancements in medical technology and medications (chemotherapy), most gynecologic oncologists may opt to use neoadjuvant treatment prior to definitive surgery. Neoadjuvant treatment is usually aimed at reducing the bulk or extent of the tumor, making surgery possible. On the other hand, EMC patients with some medical illnesses may need medical care to improve their condition before major surgery.

For physicians (mostly gynecologic oncologists), the selection of an appropriate treatment, which is usually a choice between definitive treatment or only palliative management, for patients with inoperable EMC is rather difficult. In such cases, the potential survival benefit has to be weighed against potential adverse reactions and treatment futility. Such evaluations should take several factors into consideration, including the patient's general health status, extent of disease, chance of undergoing surgery, surgical outcomes, and clinicopathological features of EMC itself.

Histologic type and grade are important prognostic factors for EMC [31-33]. Tumor histopathology and grade have been shown to be associated with the response to chemotherapy in the neoadjuvant setting [34,35]. One study found that grade 2 endometrioid tumors had a significantly better response to chemotherapy than grade 3 endometrioid tumors (72% vs. 43%, respectively) [35]. Surprisingly, the response rate for grade 1 tumors was only 46%. These findings require confirmation in additional studies because only 16% of the cases were in the primary setting (adjuvant or neoadjuvant therapy) and 84% involved chemotherapy in the recurrent setting, when tumor grade and biological behavior may be different from those in primary tumors. Another study found associations among pathological features, chemotherapy response score, and survival [34].

Aside from low-grade and less-aggressive histopathology, the ability to undergo surgical treatment appears to be an important favorable prognostic factor for EMC patients who receive neoadjuvant treatment of any type. One study reviewed the data of 59 patients with unresectable stage II to IVA EMC who underwent curative-intent treatment with neoadjuvant radiation (more than half also received concurrent chemotherapy) (47%), neoadjuvant chemotherapy (NACT)

**Table 1.** Studies which reported radiation as a definitive therapy for endometrial cancer patients

Study	Year of study	Stage	Number of patients	Age (yr)	Radiation type	5 years PC (%)	5 years DFS/DSS (%)	5 years OS (%)	Late complication (%)
LDR brachytherapy with or without EBPR									
Kupelian et al. (1993) [12]	1960-1986	I-IV	152	69	LDR±EBPRT	86	81	55	5
		I	120						
		II	17						
		III-IV	15						
Fishman et al. (1996) [13]	1975-1992	I	39	72	LDR	NR	80	30	0
		II	15	70	LDR±EBPRT		85	24	
Chao et al. (1996) [7]	1965-1990	I	101	71	LDR±EBPRT	90	84	66	5
Churn et al. (1999) [14]	1989-1996	I-II	25	66	LDR/HDR±EBPRT	100	68	48	3
		III-IV	12			50	33	33	
Shenfield et al. (2009) [11]	1986-2006	I	44	NR	LDR	88.6	88	61	2
Podzielinski et al. (2012) [15]	1997-2009	I-II	74	65	LDR/HDR±EBPRT, EBPR alone	82	82	NR	8
		I	59						
		II	15						
HDR brachytherapy with or without EBPR									
Nguyen et al. (1995) [16]	1984-1992	I-II	27	74	HDR±EBPRT	85	76 (8 years)	NR	11
		I	20				95 (8 years)		
		II	7				21 (8 years)		
Knocke et al. (1997) [17]	1981-1992	I-III	280	73	HDR	75.4	777	53	5
		IA	116			86	85	64	
		IB	119			68.8	73	47	
		II	37			60.5	69	40	
		III	8			NR	NR	NR	
Kucera et al. (1998) [18]	1981-1992	I	228	NR	HDR	83	85	60	5
		IA	113				89		
		IB	115				80		
Nguyen et al. (1998) [19]	1989-1997	I	36	65	HDR	88 (3 years)	85 (3 years)	65 (3 years)	15 (3 years)
Niazi et al. (2005) [20]	1984-2003	I-II	38	74	HDR±EBPRT	NR	78 (15 years)	NR	8
		I	29				90 (15 years)		
		II	9				42 (15 years)		

**Table 1.** Continued

Study	Year of study	Stage	Number of patients	Age (yr)	Radiation type	5 years PC (%)	5 years DFS/DSS (%)	5 years OS (%)	Late complication (%)
Weitmann et al. (2005) [21]	1997-2001	I-II	13	79	HDR±EBPRT	100	100	39	0
		I	12						
		II	1						
Coon et al. (2008) [22]	1997-2007	I-III	49	65	HDR±EBPRT	94	87	42	8
		I	42						
		II	5						
Inciura et al. (2010) [23]	1995-1998	I-III	29	75	HDR±EBPRT	83	74	48	14
		I	14						
		II	9						
Wegner et al. (2010) [24]	1997-2008	I-III	26	83	HDR±EBPRT	100 (1 year)	93 (1 year)	89 (1 year)	8
		I	19	92 (2 years)		73 (2 years)	28 (2 years)		
		II	5						
Gill et al. (2015) [25]	2007-2013	I	38	69	HDR±EBPRT	90.6 (2 years)	NR	94 (2 years)	0
		II	6						
		III	1						
Acharya et al. (2016) [26]	2003-2015	I-III	43	62	HDR±EBPRT	91.7 (2 years)	NR	65 (2 years)	5
		I	36						
		II	6						
Draghini et al. (2017) [27]	2005-2016	I, III	17	79	HDR±EBPRT	86 (3 years)	93 (1 year)	NR	12
		I	15	69 (6 years)		85 (2 years)			
		III	2	85 (6 years)					
Gebhardt et al. (2017) [10]	2007-2016	Tis	2	63	HDR	90 (2 years)	86 (2 years)	97 (2 years)	0
		I	43						
Gannavarapu et al. (2020) [28]	2012-2019	I-III	29 <sup>a)</sup>	59	HDR±EBPRT	7 (1 year, LR)	100 (2 years, LR & HR)	92 (1 year, LR)	7
		I	17	25 (1 year, HR)		84 (1 year, HR)			
		II	3	7 (2 years, LR)		77 (2 years, LR)			
		III	9	44 (2 years, HR)		73 (2 years, HR)			

**Table 1.** Continued

Study	Year of study	Stage	Number of patients	Age (yr)	Radiation type	5 years PC (%)	5 years DFS/DSS (%)	5 years OS (%)	Late complication (%)
EBPRT with SBRT									
Kemmerer et al. (2013) [9]	2006-2011	I-III	11	78	EBPRT±SBRT boost	55	41 (18 months)	57 (18 months)	0
		II	1						
		III	1						
Carbon ion radiotherapy									
Irie et al. (2018) [29]	1998-2014	I-III	14	70	C-ion RT	86	73	68	0 (≥grade 3)
		I	1						
		II	9						
		III	4						

PC, pelvic control; DFS, disease-free survival; DSS, disease-specific survival; OS, overall survival; LDR, low dose rate; EBPRT, external beam pelvic radiation therapy; HDR, high dose rate; NR, not reported; LR, low risk (stage 1-2 and grade 1/2); HR, high risk (stage 3 and/or grade 3); SBRT, stereotactic body radiation therapy; RT, radiation therapy.

<sup>a</sup>Among 29 patients in the study of Gannavarapu et al. [28], 25 had definite radiation therapy whereas four had radiation as neoadjuvant treatment before surgery.

(29%), or definitive radiation (24%) [36]. The overall surgery rate was 66%:82% in the NACT group and 79% in the neoadjuvant radiation group. Overall, 19% of the patients experienced at least grade 2 gastrointestinal, genitourinary, vaginal, or musculoskeletal toxicities. The survival benefit of surgery was also demonstrated. The 3-year survival rates of patients who could undergo surgery (66%) were significantly higher than those who were not able to undergo surgery: 84% vs. 41% for OS and 56% vs. 11% for DFS, respectively.

A positive effect of surgical treatment on survival was also found in studies that focused only on chemotherapy as a neoadjuvant treatment. One study found that patients who could undergo interval debulking surgery (IDS) had superior OS than other patients: 15-16 months compared to 5-6 months [37]. Some features that were reported to be associated with a higher chance of surgical treatment were younger age, earlier stage, or endometrioid histopathology [36].

The surgical outcome is even more important than the ability to undergo surgery. Although the benefit of optimal cytoreduction in EMC is not as well recognized as in ovarian cancer, few studies have identified optimal cytoreduction as a favorable prognostic factor for EMC patients. Rajkumar et al. [38] described optimal cytoreduction ( $\leq 1$  cm) as an independent good prognostic factor for survival in 45 patients with stage IIIC/IV EC patients who received neoadjuvant therapy

(38% received chemotherapy). Patients who had suboptimal cytoreduction showed a 3.55-fold increased risk of death, with significantly shorter survival than those who received optimal surgery: 12 months vs. 16 months for PFS and 18 months vs. 29 months for OS. Another study in 30 serous stage IV (transperitoneal spread) patients, which was the only prospective study to date, reported successful surgery (80% optimal cytoreduction and 92% without residual tumor) in 28 patients who showed some response after NACT (complete, partial response, or stable disease). The authors also reported a PFS of 13 months and OS of 23 months [39].

Predictive factors for successful surgery are useful considering their important influence on surgical outcomes. Some clinical features associated with a higher chance of surgical treatment, such as young age, lower stage, endometrioid histopathology, or response to neoadjuvant therapy, have also been reported to be associated with better surgical outcomes [36,37]. A reduction in preoperative cancer antigen (CA)-125 levels is another factor. One study in 115 patients with advanced-stage Müllerian carcinoma who had NACT demonstrated that a CA-125 reduction of at least 90% was a favorable prognostic feature associated with complete IDS, fewer bowel resections, and absence of viable tumor or presence of only microscopic disease [40].

These previously described predictive factors for good sur-



gical outcomes (younger age, endometrioid histology) as well as low-grade tumors were also good prognostic features for survival [7,36]. Other favorable factors were preoperative CA 125 level <20 mIU/mL (HR 0.37 for survival) [40], absence of tumor infiltration, presence of tumor necrosis [39], and high chemotherapy response score [34]. On the other hand, unfavorable prognostic factors that have been reported to date include poor performance status and the presence of bowel disease [38].

Although the Cancer Genome Atlas Research Network has categorized EMC on the basis of distinct molecular and genetic features to guide appropriate therapy for individual patients [41], this classification was not widely practiced, especially in the remote past and in settings with limited resources. Hence, data on individualized therapy, especially in neoadjuvant settings for inoperable EMC, are not currently available.

A physician may consider all currently available prognostic factors, counsel the patient, and make appropriate decisions regarding the type of neoadjuvant treatment. Data on hormonal therapy, radiation, chemotherapy, or their combinations as neoadjuvant treatment prior to surgery are provided below.

## 1) Neoadjuvant hormonal therapy

Since EMC is a hormone-related cancer, many studies have reported the use of hormonal therapy for advanced or recurrent EMC [42], medical unfit early-stage EMC [43] or in adjuvant settings [44]. However, only a few studies have assessed the role of hormonal agents in neoadjuvant treatment [45,46]. One study compared the effects of letrozole, anastrozole, and exemestane in 38 patients [45]. The findings showed an improvement in clinical signs (decreased endometrial thickness) and symptoms (relief of abdominal pain and uterine discharge), most frequently with anastrozole. Laboratory investigations revealed reductions in intratumoral aromatase and blood estradiol levels more frequently after anastrozole and letrozole administration, whereas tumor progesterone receptor levels were markedly lower after exemestane administration [45]. A subsequent report from the same group of authors described superior effects of 2-4 weeks of aromatase inhibitor treatment in comparison with 2-9 weeks of metformin treatment in terms of endometrial thickness, grade change, and proliferative index among 38 EMC patients (it was unclear whether the patients were the same as

those in the authors' preceding report) [47]. Another small trial randomized 24 EMC patients with any histopathology to receive anastrozole (n=16) or placebo (n=8) for at least 14 days prior to surgery [46]. Expression of estrogen receptor alpha, androgen receptor, and Ki-67 (but not Bcl2 and progesterone receptor) was significantly lower after treatment with anastrozole.

Advances in knowledge from clinical studies on neoadjuvant hormonal therapy for EMC are unlikely, probably because the cytostatic effect of hormonal agents to slow tumor progression is not the aim in a neoadjuvant setting. In these settings, the induction of a cytotoxic effect to reduce the tumor load through the more remarkable effects of chemotherapy or radiotherapy in reducing tumor bulk is probably preferred.

## 2) Neoadjuvant radiation therapy

Few studies have reported the use of radiation as a neoadjuvant treatment in locally advanced EMC, wherein the disease has spread beyond the uterine body to the cervix or parametrium. In these studies, the mode of radiation was either EPRT combined with ICRT or ICRT alone.

Some studies have used preoperative ICRT alone for EMC that was limited to the uterus or cervix (stage I or II) [48-50]. One small study included 12 patients who received high-dose ICRT for clinical stage I-II [50]. In that study, surgery revealed a complete pathological response in two patients and only microscopic residual disease in the other five patients, and the 2-year disease-free and cause-specific survival rates were 88% and 100%, respectively. Two other non-English reports also described success after preoperative ICRT in EMC [48,49]. One study from Poland showed that 51% of the 35 patients who underwent surgery had no residual cancer after ICRT [48]. Another study from France described the findings for 780 clinical stage I-II EMC patients who received preoperative low-dose ICRT and underwent long-term follow-up of over 10 years [49]. Postoperatively, pelvic radiation was used if necessary. The 5-year local control rate in that study was 93%, with an 86% DFS and 84% OS.

Other studies have used combinations of EPRT and ICRT for the treatment of EMC with cervical or parametrial extension. One study by Vargo et al. [51] reported the use of neoadjuvant EPRT (predominantly intensity-modulated radiation therapy) and image-based high-dose rate ICRT with or without chemotherapy, followed by surgery in 36 patients

with EMC with or without parametrium involvement. Extrafascial hysterectomy performed at a median of 6 weeks after radiotherapy revealed a clinical complete response in 91% of cervical lesions, whereas pathological complete response was achieved in 24% of patients. The 3-year DFS and OS rates were 73% and 100%, respectively. Another study by the same group of authors focused on the safety of extrafascial hysterectomy either by laparotomy or laparoscopy after neoadjuvant external beam radiotherapy followed by image-based high-dose rate ICRT in 29 EMC patients [52]. Except for vaginal complications, which occurred frequently (33%) and were particularly prominent in patients who underwent laparoscopic surgery, the incidence of other perioperative complications was low in both laparotomy and laparoscopic approaches (3-6%). The authors also noted no significant correlation between radiotherapy dose and postoperative complication rates.

### 3) Neoadjuvant chemotherapy

The role of NACT followed by delayed primary surgery, which is more commonly called IDS, in ovarian cancer has been well-recognized and commonly practiced [53,54]. This is because ovarian cancer commonly presents with advanced-stage disease, for which primary surgery is not suitable. Unlike ovarian cancer, most patients with EMC usually present with early-stage disease, leading to limited evidence-based data regarding NACT in EMC.

Nevertheless, several case reports [55,56], retrospective studies [57-60], and one prospective study [61] describing the use of NACT before IDS in advanced-stage EMC have been identified. In the mid-2000s, Despierre et al. [55] and Takami et al. [56] reported the findings for three EMC patients with advanced-stage clear cell or serous carcinoma who showed complete or remarkable response (disappearance of lung metastasis) after three cycles of NACT. The patient underwent successful surgery without residual disease.

Generally, IDS is performed in patients who show some responses (preferably complete or partial response rather than stable disease) after a few cycles of NACT. Response to chemotherapy is a key factor determining whether patients can undergo IDS. Aside from the efficacy of NACT, another important factor influencing the response is probably the aggressiveness of the EMC itself (as reflected by the histopathological findings). This was demonstrated in several studies [37,59]. One small study by Khouri et al. [37] included 39

EMC patients showing unresectable tumors with aggressive histology of clear cells, carcinosarcoma, or neuroendocrine carcinomas aside from endometrioid carcinoma (only 28%). After NACT (85% of the patients received paclitaxel and carboplatin), 59% of the patients could not undergo IDS because of disease progression (70%) or persistent unresectable disease (17%). Another recent study collected data for 102 advanced-stage EMCs of all histologic types [59]. Outcomes after NACT (89% of the patients received paclitaxel and carboplatin) differed by histopathological subgroup. The response rate to NACT is 64% in endometrioid carcinoma and 80% in serous carcinoma [59].

The surgical or perioperative outcomes of IDS appear to be better than those of primary surgery. Two retrospective studies compared the treatment outcomes of stage IV serous EMC patients who underwent NACT followed by IDS or primary surgery: 15 patients in each group using a propensity-matched algorithm in one study [60] and 10 patients (NACT) or 34 patients (primary surgery) in another study using a simple cohort design [58,60]. Better operative and postoperative outcomes were demonstrated in patients who underwent NACT and IDS than in those who underwent primary surgery followed by chemotherapy, in terms of shorter operative time, lower transfusion rate, and a shorter hospital stay [58,60]. Although not statistically significant, the rate of no gross residual disease was 2-fold higher in the NACT/IDS (70%) group than in the primary (32%) group (same figures in both studies) [58,60].

Studies that compared the survival of EMC patients who had undergone NACT/IDS after primary surgery/chemotherapy yielded inconsistent findings. Among these, two retrospective studies that reported improved operative and postoperative outcomes as described earlier demonstrated comparable survival between the patients who underwent NACT or primary surgery: PFS, 10-12 months vs. 12-15 months; OC, 17 months vs. 18-21 months [58,60]. However, other studies found lower survival rates in patients who underwent NACT/ICT than in those who had primary surgery/chemotherapy [57,61]. One large multicenter study from Japan reviewed the data of 426 patients with clinical or surgical stage IVB EMC [57]. The median OS was the longest after primary surgery (279 patients; 21 months), followed by primary chemotherapy (NACT) (125 patients; 12 months), and the lowest with only palliative care (125 patients; 1 month). The survival difference between patients who underwent



primary surgery or primary chemotherapy in this study must be interpreted with caution because patients in the primary surgery group had more favorable prognostic features (better performance status and fewer extra-abdominal metastases). Among a subset of patients who responded to NACT and were able to undergo surgery, the OS was similar to that in patients who underwent primary surgery.

Another study retrieved data from the National Cancer Database to assess the influence of primary treatment on the survival of over 48,000 patients with stage III/IV EMC between 2004 and 2015 [61]. The majority of the patients underwent primary surgery followed by adjuvant chemotherapy (83%), whereas the others received only chemotherapy (12%) or NACT/IDS (5%). The OS was the highest among patients who underwent surgery and adjuvant chemotherapy, followed by those who received NACT before surgery, and was the worst among those who received chemotherapy alone: 61 months vs. 25 months vs. 11 months, respectively. In comparison with chemotherapy alone, the hazard ratios were 0.32 for primary surgery and adjuvant chemotherapy and 0.44 for NACT followed by IDS. Subgroup analyses by stage (stage III or stage IV) or histology (type I or type II EMC) showed a significant OS benefit in patients who underwent primary surgery in comparison with the other two groups, regardless of stage or histology. Nevertheless, NACT showed a significantly superior survival benefit over chemotherapy alone, except for cases with type I stage III cancer, which showed no survival difference. One observation from this study was that although the study included a large number of patients, the percentage of patients who had NACT/IDS was small (5%). Furthermore, data on the prognostic features of patients in each group were not available for analysis. Hence, a definite conclusion is unlikely.

Although there are no solid evidence-based data from randomized controlled trials for the use of NACT in EMC, findings from several reports were consistent, showing more favorable surgical outcomes. Although survival data of NACT/IDS compared to primary surgery/chemotherapy are still inconsistent, NACT appears to be a common treatment option for inoperable EMC in recent years.

We were aware of the activity of immunotherapeutic agents (anti-programmed death-1 monoclonal antibody) either alone [62] or in combination with tyrosine kinase inhibitors in advanced EMC that had failed prior to systemic therapy [63]. The demonstrated efficacy of these agents in phase I/II studies

led to phase III trials that had recently reported superior survival outcomes over standard chemotherapy with paclitaxel and carboplatin [64] or are still ongoing [65]. Whether these agents will be used in the future as neoadjuvant treatment in advanced and inoperable EMC is yet to be seen.

#### 4) Neoadjuvant chemoradiation therapy

The use of combined chemoradiation, especially in a concurrent pattern, is well recognized as a standard treatment for locally advanced-stage cervical cancer. However, this is not the situation in EMC, since the activity of concurrent chemoradiation in cases of EMC has been tested as an adjuvant therapy after surgery in only a few randomized trials [66-68]. In fact, there are even fewer data regarding the use of concurrent chemoradiation for EMC in a neoadjuvant setting.

To date, only three reports on this topic have been published. One case report described a patient with stage IVB EMC who presented with a large uterine tumor with widespread retroperitoneal and inguinal nodal metastases that became resectable after chemotherapy (carboplatin and 5-fluorouracil) followed by radiation therapy [69]. The patient achieved a 4-year recurrence-free outcome. Combinations of chemotherapy and radiation can be tailored according to the characteristic features of the cancer. This was supported by another case report in which multimodality treatment was performed in an EMC patient with grade 3 cancer extending to the cervix, parametrium, pelvic side wall, and pelvic and para-aortic nodal metastases [70]. The patient received a course of neoadjuvant chemoradiotherapy with pelvic and para-aortic field intensity-modulated radiation therapy, in combination with low-dose weekly cisplatin. This was followed by intrauterine tandem and interstitial needles based on the initial extent of the disease, with an additional high-dose rate ICRT prior to surgery. Complete surgical staging 1 month after the radiation revealed a complete pathological response.

The third report was a retrospective study of 34 patients with type II EMC (serous, carcinosarcoma, and clear cell carcinoma) of stage II-III [71] who received external pelvic radiation plus ICRT concurrent with platinum chemotherapy (mainly cisplatin) as neoadjuvant treatment. Over 90% of cancer cases were downstaged. The surgical approach was laparoscopic (41%), laparotomic (38%), or robotic (21%). Free surgical margins were achieved in 94% of patients. Pathologic complete response was demonstrated in 15% of

the patients, while 35% showed only microscopic residual disease. While 6% of the patients required blood transfusion, no intraoperative complications were encountered. Postoperative complications were found in 39% of the patients: readmission or ileus (12% each), wound infection (9%), and pulmonary emboli or sepsis (3% each). The 2-year local, regional, and distant control rates were approximately 80%, with 53% DFS and 64% OS rates. The results of this study should be interpreted with caution because they were not simply due to the effect of concurrent chemoradiation treatment. Neoadjuvant or adjuvant treatment was also administered to 12% and 79% of the patients, respectively. Notably, 12% of patients received NACT before chemoradiation and surgery followed by adjuvant chemotherapy.

Although not directly relevant to the topic of this review, one more study merits a mention in the present investigation. This study used a national registry of over 20,000 stage III EMC patients diagnosed between 2004 and 2013 to assess the patterns-of-care and OS benefits of chemoradiation versus chemotherapy or radiation alone as adjuvant therapy [72]. The use of chemoradiation increased significantly during the study period. A significantly decreased risk of death (hazard ratio, 0.62) was associated with the use of treatment combinations over a single mode of treatment. However, the trend for combined chemoradiation as neoadjuvant therapy for primary inoperable EMC remains unclear.

One major caveat is the tolerability of EMC patients who are old and frequently have comorbid illnesses. Some patients may not be able to endure the adverse reactions of combined treatments such as neoadjuvant therapy. Furthermore, some patients may require adjuvant therapy postoperatively. Thus, neoadjuvant concurrent chemoradiation should not be performed until more studies can demonstrate its clear benefit.

## Discussion

Radiation therapy alone can be administered as definitive treatment for early-stage primary inoperable EMC. MRI should be used to tailor the mode of radiation therapy (according to the extent of tumor) when it is administered as the sole treatment. Neoadjuvant treatment with radiation, chemotherapy, or a combination of these prior to surgery can be used for all cases of early- or advanced-stage disease.

The most common neoadjuvant treatment is chemotherapy, whereas less data is available regarding radiation or chemoradiation. Neoadjuvant treatment yielded better perioperative outcomes, but the survival rates associated with neoadjuvant treatment were inconsistent in comparison with those obtained with primary surgery.

## Conflict of interest

No potential conflict of interest relevant to this article was reported.

## Ethical approval

This study does not require approval of the Institutional Review Board because no patient data is contained in this article. The study was performed in accordance with the principles of the Declaration of Helsinki.

## Patient consent

Written informed consent and the use of images from patients are not required for the publication.

## Funding information

None.

## Author contributions

SP: Data curation, writing original draft, review, and editing; SC: Data curation, writing original draft; ST: Conceptualization, data curation, supervision, writing original draft, review, and editing. All authors have read and approved the final manuscript.

## Data availability

The authors declare that data supporting the findings of this

study are available in the article.

## References

1. Creasman WT, Odicino F, Maisonneuve P, Quinn MA, Beller U, Benedet JL, et al. Carcinoma of the corpus uteri. FIGO 26th annual report on the results of treatment in gynecological cancer. *Int J Gynaecol Obstet* 2006;95 Suppl 1:S105-43.
2. Bogani G, Gostout BS, Dowdy SC, Multinu F, Casarin J, Cliby WA, et al. Clinical utility of preoperative computed tomography in patients with endometrial cancer. *Int J Gynecol Cancer* 2017;27:1685-93.
3. Dai S, Nahas S, Murphy JK, Lawrence J, May T, Feigenberg T. Impact and cost of preoperative computed tomography imaging on the management of patients diagnosed with high-grade endometrial cancer. *Int J Gynaecol Obstet* 2019;145:219-24.
4. Chao A, Chang TC, Ng KK, Hsueh S, Huang HJ, Chou HH, et al. 18F-FDG PET in the management of endometrial cancer. *Eur J Nucl Med Mol Imaging* 2006;33:36-44.
5. Teng F, Zhang YF, Wang YM, Yu J, Lang X, Tian WY, et al. Contrast-enhanced MRI in preoperative assessment of myometrial and cervical invasion, and lymph node metastasis: diagnostic value and error analysis in endometrial carcinoma. *Acta Obstet Gynecol Scand* 2015;94:266-73.
6. Bristow RE, Asrari F, Trimble EL, Montz FJ. Extended surgical staging for uterine papillary serous carcinoma: survival outcome of locoregional (stage I-III) disease. *Gynecol Oncol* 2001;81:279-86.
7. Chao CK, Grigsby PW, Perez CA, Mutch DG, Herzog T, Camel HM. Medically inoperable stage I endometrial carcinoma: a few dilemmas in radiotherapeutic management. *Int J Radiat Oncol Biol Phys* 1996;34:27-31.
8. Yoo S, Hegarty SE, Mishra MV, Patel N, Cantrell LA, Showalter TN. Definitive radiation therapy for stage I-II endometrial cancer: an observational study of nonoperative management. *Am J Clin Oncol* 2017;40:582-9.
9. Kemmerer E, Hernandez E, Ferriss JS, Valakh V, Miyamoto C, Li S, et al. Use of image-guided stereotactic body radiation therapy in lieu of intracavitary brachytherapy for the treatment of inoperable endometrial neoplasia. *Int J Radiat Oncol Biol Phys* 2013;85:129-35.
10. Gebhardt B, Gill B, Glaser S, Kim H, Houser C, Kelley J, et al. Image-guided tandem and cylinder brachytherapy as monotherapy for definitive treatment of inoperable endometrial carcinoma. *Gynecol Oncol* 2017;147:302-8.
11. Shenfield CB, Pearcey RG, Ghosh S, Dundas GS. The management of inoperable stage I endometrial cancer using intracavitary brachytherapy alone: a 20-year institutional review. *Brachytherapy* 2009;8:278-83.
12. Kupelian PA, Eifel PJ, Tornos C, Burke TW, Delclos L, Oswald MJ. Treatment of endometrial carcinoma with radiation therapy alone. *Int J Radiat Oncol Biol Phys* 1993;27:817-24.
13. Fishman DA, Roberts KB, Chambers JT, Kohorn EI, Schwartz PE, Chambers SK. Radiation therapy as exclusive treatment for medically inoperable patients with stage I and II endometrioid carcinoma with endometrium. *Gynecol Oncol* 1996;61:189-96.
14. Churn M, Jones B. Primary radiotherapy for carcinoma of the endometrium using external beam radiotherapy and single line source brachytherapy. *Clin Oncol (R Coll Radiol)* 1999;11:255-62.
15. Podzielinski I, Randall ME, Breheny PJ, Escobar PF, Cohn DE, Quick AM, et al. Primary radiation therapy for medically inoperable patients with clinical stage I and II endometrial carcinoma. *Gynecol Oncol* 2012;124:36-41.
16. Nguyen C, Souhami L, Roman TN, Clark BG. High-dose-rate brachytherapy as the primary treatment of medically inoperable stage I-II endometrial carcinoma. *Gynecol Oncol* 1995;59:370-5.
17. Knocke TH, Kucera H, Weidinger B, Höller W, Pötter R. Primary treatment of endometrial carcinoma with high-dose-rate brachytherapy: results of 12 years of experience with 280 patients. *Int J Radiat Oncol Biol Phys* 1997;37:359-65.
18. Kucera H, Knocke TH, Kucera E, Pötter R. Treatment of endometrial carcinoma with high-dose-rate brachytherapy alone in medically inoperable stage I patients. *Acta Obstet Gynecol Scand* 1998;77:1008-12.
19. Nguyen TV, Petereit DG. High-dose-rate brachytherapy for medically inoperable stage I endometrial cancer. *Gynecol Oncol* 1998;71:196-203.
20. Niazi TM, Souhami L, Portelance L, Bahoric B, Gilbert L, Stanimir G. Long-term results of high-dose-rate brachytherapy in the primary treatment of medically inoperable

- stage I-II endometrial carcinoma. *Int J Radiat Oncol Biol Phys* 2005;63:1108-13.
21. Weitmann HD, Pötter R, Waldhäusl C, Nechvile E, Kirisits C, Knocke TH. Pilot study in the treatment of endometrial carcinoma with 3D image-based high-dose-rate brachytherapy using modified Heyman packing: clinical experience and dose-volume histogram analysis. *Int J Radiat Oncol Biol Phys* 2005;62:468-78.
  22. Coon D, Beriwal S, Heron DE, Kelley JL, Edwards RP, Sukumvanich P, et al. High-dose-rate Rotte "Y" applicator brachytherapy for definitive treatment of medically inoperable endometrial cancer: 10-year results. *Int J Radiat Oncol Biol Phys* 2008;71:779-83.
  23. Inciura A, Atkocius V, Juozaityte E, Vaitkiene D. Long-term results of high-dose-rate brachytherapy and external-beam radiotherapy in the primary treatment of endometrial cancer. *J Radiat Res* 2010;51:675-81.
  24. Wegner RE, Beriwal S, Heron DE, Richard SD, Kelly JL, Edwards RP, et al. Definitive radiation therapy for endometrial cancer in medically inoperable elderly patients. *Brachytherapy* 2010;9:260-5.
  25. Gill BS, Chapman BV, Hansen KJ, Sukumvanich P, Beriwal S. Primary radiotherapy for nonsurgically managed stage I endometrial cancer: utilization and impact of brachytherapy. *Brachytherapy* 2015;14:373-9.
  26. Acharya S, Esthappan J, Badiyan S, DeWees TA, Tandrup K, Schwarz JK, et al. Medically inoperable endometrial cancer in patients with a high body mass index (BMI): patterns of failure after 3-D image-based high dose rate (HDR) brachytherapy. *Radiother Oncol* 2016;118:167-72.
  27. Draghini L, Maranzano E, Casale M, Trippa F, Anselmo P, Arcidiacono F, et al. Definitive three-dimensional high-dose-rate brachytherapy for inoperable endometrial cancer. *J Contemp Brachytherapy* 2017;9:118-23.
  28. Gannavarapu BS, Hrycushko B, Jia X, Albuquerque K. Upfront radiotherapy with brachytherapy for medically inoperable and unresectable patients with high-risk endometrial cancer. *Brachytherapy* 2020;19:139-45.
  29. Irie D, Okonogi N, Wakatsuki M, Kato S, Ohno T, Karasawa K, et al. Carbon-ion radiotherapy for inoperable endometrial carcinoma. *J Radiat Res* 2018;59:309-15.
  30. Schwarz JK, Beriwal S, Esthappan J, Erickson B, Feltmate C, Fyles A, et al. Consensus statement for brachytherapy for the treatment of medically inoperable endometrial cancer. *Brachytherapy* 2015;14:587-99.
  31. Creasman WT, Morrow CP, Bundy BN, Homesley HD, Graham JE, Heller PB. Surgical pathologic spread patterns of endometrial cancer. A gynecologic oncology group study. *Cancer* 1987;60(8 Suppl):2035-41.
  32. Creasman WT, Kohler MF, Odicino F, Maisonneuve P, Boyle P. Prognosis of papillary serous, clear cell, and grade 3 stage I carcinoma of the endometrium. *Gynecol Oncol* 2004;95:593-6.
  33. Brinton LA, Felix AS, McMeekin DS, Creasman WT, Sherman ME, Mutch D, et al. Etiologic heterogeneity in endometrial cancer: evidence from a gynecologic oncology group trial. *Gynecol Oncol* 2013;129:277-84.
  34. Williams AT, Ganesan R. Role of the pathologist in assessing response to treatment of ovarian and endometrial cancers. *Histopathology* 2020;76:93-101.
  35. Davidson BA, Foote J, Clark LH, Broadwater G, Ehrisman J, Gehrig P, et al. Tumor grade and chemotherapy response in endometrioid endometrial cancer. *Gynecol Oncol Rep* 2016;17:3-6.
  36. Conway JL, Lukovic J, Ferguson SE, Zhang J, Xu W, Dhani N, et al. Clinical outcomes of surgically unresectable endometrial cancers. *Am J Clin Oncol* 2019;42:777-82.
  37. Khouri OR, Frey MK, Musa F, Muggia F, Lee J, Boyd L, et al. Neoadjuvant chemotherapy in patients with advanced endometrial cancer. *Cancer Chemother Pharmacol* 2019;84:281-5.
  38. Rajkumar S, Nath R, Lane G, Mehra G, Begum S, Sayasneh A. Advanced stage (IIIC/IV) endometrial cancer: role of cytoreduction and determinants of survival. *Eur J Obstet Gynecol Reprod Biol* 2019;234:26-31.
  39. Vandenput I, Van Calster B, Capoen A, Leunen K, Berteloot P, Neven P, et al. Neoadjuvant chemotherapy followed by interval debulking surgery in patients with serous endometrial cancer with transperitoneal spread (stage IV): a new preferred treatment? *Br J Cancer* 2009;101:244-9.
  40. Mahdi H, Maurer KA, Nutter B, Rose PG. The impact of percent reduction in CA-125 levels on prediction of the extent of interval cytoreduction and outcome in patients with advanced-stage cancer of Müllerian origin treated with neoadjuvant chemotherapy. *Int J Gynecol Cancer* 2015;25:823-9.
  41. Cancer Genome Atlas Research Network, Kandoth C,

- Schultz N, Cherniack AD, Akbani R, Liu Y, et al. Integrated genomic characterization of endometrial carcinoma. *Nature* 2013;497:67-73.
42. Thigpen JT, Brady MF, Alvarez RD, Adelson MD, Homesley HD, Manetta A, et al. Oral medroxyprogesterone acetate in the treatment of advanced or recurrent endometrial carcinoma: a dose-response study by the Gynecologic Oncology Group. *J Clin Oncol* 1999;17:1736-44.
43. Nikolopoulos M, Godfrey MAL, Wuntakal R. Medically unfit women with early-stage endometrial cancer treated with the levonorgestrel intrauterine system. *Obstet Gynecol Sci* 2020;63:337-45.
44. Martin-Hirsch PP, Bryant A, Keep SL, Kitchener HC, Lilford R. Adjuvant progestagens for endometrial cancer. *Cochrane Database Syst Rev* 2011;(6):CD001040.
45. Bershtein LM, Danilova MA, Kovalevskii Alu, Gershfel'd ED, Poroshina TE, Tsyrlina TE, et al. Criteria for evaluating the effectiveness of aromatase inhibitors in the neoadjuvant treatment of patients with endometrial carcinoma. *Vopr Onkol* 2009;55:314-8.
46. Thangavelu A, Hewitt MJ, Quinton ND, Duffy SR. Neoadjuvant treatment of endometrial cancer using anastrozole: a randomised pilot study. *Gynecol Oncol* 2013;131:613-8.
47. Bershtein LM, Maksimov Sla, Danilova MA, Gershfel'd ED, Boiarkina MP, Khadzhimba AS, et al. Effectiveness of aromatase inhibitors in comparison with metformin for neoadjuvant treatment in patients with endometrial cancer. *Vopr Onkol* 2011;57:737-41.
48. Sobotkowski J, Zielinska M, Grzelak M, Pietraszek A. Preoperative high-dose brachytherapy in cases of endometrial carcinoma--preliminary assessment of outcomes and safety. *Med Wieku Rozwoj* 2004;8(2 Pt 1):309-16.
49. Hoffstetter S, Brunaud C, Marchal C, Luporsi E, Guillemain F, Leroux A, et al. Preoperative brachytherapy for clinical stage I and II endometrial carcinoma: results from a series of 780 patients with a 10-year follow-up. *Cancer Radiother* 2004;8:178-87.
50. Shukla G, Beriwal S, Krivak TC, Kelley JL, Sukumvanich P, Edwards RP, et al. Preoperative high dose rate brachytherapy for clinical stage II endometrial carcinoma. *J Contemp Brachytherapy* 2011;3:70-3.
51. Vargo JA, Boisen MM, Comerci JT, Kim H, Houser CJ, Sukumvanich P, et al. Neoadjuvant radiotherapy with or without chemotherapy followed by extrafascial hysterectomy for locally advanced endometrial cancer clinically extending to the cervix or parametria. *Gynecol Oncol* 2014;135:190-5.
52. Boisen MM, Vargo JA, Beriwal S, Sukumvanich P, Olawaiye AB, Kelley JL, et al. Surgical outcomes of patients undergoing extrafascial hysterectomy after neoadjuvant radiotherapy with or without chemotherapy for locally advanced endometrial cancer clinically extending to the cervix or parametria. *Int J Gynecol Cancer* 2017;27:1149-54.
53. Rose PG, Nerenstone S, Brady MF, Clarke-Pearson D, Olt G, Rubin SC, et al. Secondary surgical cytoreduction for advanced ovarian carcinoma. *N Engl J Med* 2004;351:2489-97.
54. Vergote I, Tropé CG, Amant F, Kristensen GB, Ehlen T, Johnson N, et al. Neoadjuvant chemotherapy or primary surgery in stage IIIc or IV ovarian cancer. *N Engl J Med* 2010;363:943-53.
55. Despierre E, Moerman P, Vergote I, Amant F. Is there a role for neoadjuvant chemotherapy in the treatment of stage IV serous endometrial carcinoma? *Int J Gynecol Cancer* 2006;16 Suppl 1:273-7.
56. Takami M, Ohta Y, Nakayama Y, Fukai H, Matsumoto H, Takimoto T, et al. A case of advanced clear cell carcinoma of the endometrium that responded remarkably to neoadjuvant chemotherapy of combination carboplatin plus weekly paclitaxel. *Gan To Kagaku Ryoho* 2007;34:457-60.
57. Eto T, Saito T, Shimokawa M, Hatae M, Takeshima N, Kobayashi H, et al. Status of treatment for the overall population of patients with stage IVb endometrial cancer, and evaluation of the role of preoperative chemotherapy: a retrospective multi-institutional study of 426 patients in Japan. *Gynecol Oncol* 2013;131:574-80.
58. Wilkinson-Ryan I, Frolova AI, Liu J, Massad LS, Thaker PH, Powell MA, et al. Neoadjuvant chemotherapy versus primary cytoreductive surgery for stage IV uterine serous carcinoma. *Int J Gynecol Cancer* 2015;25:63-8.
59. de Lange NM, Ezendam NPM, Kwon JS, Vandenput I, Mirchandani D, Amant F, et al. Neoadjuvant chemotherapy followed by surgery for advanced-stage endometrial cancer. *Curr Oncol* 2019;26:e226-32.
60. Bogani G, Ditto A, Maggiore ULR, Scaffa C, Mosca L, Chiappa V, et al. Neoadjuvant chemotherapy followed by interval debulking surgery for unresectable stage IVb



- serous endometrial cancer. *Tumori* 2019;105:92-7.
61. Chambers LM, Jia X, Rose PG, AlHilli M. Impact of treatment modality on overall survival in women with advanced endometrial cancer: a national cancer database analysis. *Gynecol Oncol* 2021;160:405-12.
62. Marabelle A, Le DT, Ascierto PA, Di Giacomo AM, De Jesus-Acosta A, Delord JP, et al. Efficacy of pembrolizumab in patients with noncolorectal high microsatellite instability/mismatch repair-deficient cancer: results from the phase II KEYNOTE-158 study. *J Clin Oncol* 2020;38:1-10.
63. Makker V, Taylor MH, Aghajanian C, Oaknin A, Mier J, Cohn AL, et al. Lenvatinib plus pembrolizumab in patients with advanced endometrial cancer. *J Clin Oncol* 2020;38:2981-92.
64. Makker V, Colombo N, Casado Herráez A, Santin A, Colomba A, Miller D, et al. A multicenter, open-label, randomized, phase 3 study to compare the efficacy and safety of lenvatinib in combination with pembrolizumab vs treatment of physician's choice in patients with advanced endometrial cancer. *Gynecol Oncol* 2021;162 Suppl 1:S4.
65. Marth C, Tarnawski R, Tyulyandina A, Pignata S, Gilbert L, Kaen D, et al. Phase 3, randomized, open-label study of pembrolizumab plus lenvatinib versus chemotherapy for first-line treatment of advanced or recurrent endometrial cancer: ENGOT-en9/LEAP-001. *Int J Gynecol Cancer* 2022;32:93-100.
66. Randall ME, Filiaci V, McMeekin DS, von Gruenigen V, Huang H, Yashar CM, et al. Phase III trial: adjuvant pelvic radiation therapy versus vaginal brachytherapy plus paclitaxel/carboplatin in high-intermediate and high-risk early stage endometrial cancer. *J Clin Oncol* 2019;37:1810-8.
67. Matei D, Filiaci V, Randall ME, Mutch D, Steinhoff MM, DiSilvestro PA, et al. Adjuvant chemotherapy plus radiation for locally advanced endometrial cancer. *N Engl J Med* 2019;380:2317-26.
68. de Boer SM, Powell ME, Mileskin L, Katsaros D, Besette P, Haie-Meder C, et al. Adjuvant chemoradiotherapy versus radiotherapy alone in women with high-risk endometrial cancer (PORTEC-3): patterns of recurrence and post-hoc survival analysis of a randomised phase 3 trial. *Lancet Oncol* 2019;20:1273-85.
69. Kaneyasu Y, Okawa T, Yajima M, Saito R, Nakabayashi M, Seshimo A, et al. Stage IVB uterine endometrial cancer successfully salvaged by chemoradiotherapy and surgery. *Int J Clin Oncol* 2003;8:60-4.
70. Horne Z, Vargo JA, Comerchi JT, Beriwal S. Complete pathologic response following neoadjuvant chemoradiotherapy and high-dose-rate brachytherapy for locally advanced endometrial carcinoma. *Cureus* 2015;7:e407.
71. Iheagwara UK, Vargo JA, Chen KS, Burton DR, Taylor SE, Berger JL, et al. Neoadjuvant chemoradiation therapy followed by extrafascial hysterectomy in locally advanced type II endometrial cancer clinically extending to cervix. *Pract Radiat Oncol* 2019;9:248-56.
72. Boothe D, Orton A, Odei B, Stoddard G, Suneja G, Poppe MM, et al. Chemoradiation versus chemotherapy or radiation alone in stage III endometrial cancer: patterns of care and impact on overall survival. *Gynecol Oncol* 2016;141:421-7.