

United States Epilepsy Center Characteristics

A Data Analysis From the National Association of Epilepsy Centers

Adam P. Ostendorf, MD,* Stephanie M. Ahrens, DO,* Fred Alexander Lado, MD, PhD, Susan T. Arnold, MD, Shasha Bai, PhD, Meriem K. Bensalem Owen, MD, Kevin E. Chapman, MD, Dave F. Clarke, MD, Mariah Eisner, MS, Nathan B. Fountain, MD, Johanna M. Gray, MPA, Jennifer L. Hopp, MD, Ellen Riker, MHA, Stephan U. Schuele, MD, MPH, Barbara V. Small, BA, and Susan T. Herman, MD

Neurology® 2022;98:e449-e458. doi:10.1212/WNL.0000000000013130

Correspondence

Dr. Ostendorf
adam.ostendorf@
nationwidechildrens.org

Abstract

Background and Objectives

Patients with drug-resistant epilepsy (DRE) may benefit from specialized testing and treatments to better control seizures and improve quality of life. Most evaluations and procedures for DRE in the United States are performed at epilepsy centers accredited by the National Association of Epilepsy Centers (NAEC). On an annual basis, the NAEC collects data from accredited epilepsy centers on hospital-based epilepsy monitoring unit (EMU) size and admissions, diagnostic testing, surgeries, and other services. This article highlights trends in epilepsy center services from 2012 through 2019.

Methods

We analyzed data reported in 2012, 2016, and 2019 from all level 3 and level 4 NAEC accredited epilepsy centers. Data were described using frequency for categorical variables and median for continuous variables and were analyzed by center level and center population category. EMU beds, EMU admissions, epileptologists, and aggregate procedure volumes were also described using rates per population per year.

Results

During the period studied, the number of NAEC accredited centers increased from 161 to 256, with the largest increases in adult- and pediatric-only centers. Growth in EMU admissions (41%), EMU beds (26%), and epileptologists (109%) per population occurred. Access to specialized testing and services broadly expanded. The largest growth in procedure volumes occurred in laser interstitial thermal therapy (LiTT) (61%), responsive neurostimulation (RNS) implantations (114%), and intracranial monitoring without resection (152%) over the study period. Corpus callosotomies and vagus nerve stimulator (VNS) implantations decreased (−12.8% and −2.4%, respectively), while growth in temporal lobectomies (5.9%), extratemporal resections (11.9%), and hemispherectomies/otomies (13.1%) lagged center growth (59%), leading to a decrease in median volumes of these procedures per center.

Discussion

During the study period, the availability of specialty epilepsy care in the United States improved as the NAEC implemented its accreditation program. Surgical case complexity increased while aggregate surgical volume remained stable or declined across most procedure types, with a corresponding decline in cases per center. This article describes recent data trends and current state of resources and practice across NAEC member centers and identifies several future directions for driving systematic improvements in epilepsy care.

*Adam P. Ostendorf and Stephanie M. Ahrens are co-first authors.


From the Department of Pediatrics (A.P.O., S.M.A.), Nationwide Children's Hospital and Ohio State University, Columbus; Department of Neurology (F.A.L.), Hofstra Northwell Comprehensive Epilepsy Center, Great Neck, NY; Department of Pediatrics (S.T.A.), Children's Medical Center, Dallas, TX; Pediatric Biostatistics Core (S.B.), Emory University School of Medicine, Atlanta, GA; Department of Neurology (M.K.B.O.), University of Kentucky, Lexington; Department of Child Health (K.E.C.), University of Arizona College of Medicine and Barrow's Neurological Institute at Phoenix Children's Hospital; Division of Pediatric Neurology (D.F.C.), Department of Neurology, Dell Medical School, University of Texas at Austin; Biostatistics Resource at Nationwide Children's Hospital (M.E.), Columbus, OH; Department of Neurology (N.B.F.), Comprehensive Epilepsy Program, University of Virginia, Charlottesville; National Association of Epilepsy Centers (J.M.G., E.R., B.V.S.), Washington, DC; University of Maryland Medical Center (J.L.H.), Baltimore, MD; Ken and Ruth Davee Department of Neurology (S.U.S.), Northwestern University Feinberg School of Medicine, Chicago, IL; and Department of Neurology (S.T.H.), Barrow Neurological Institute, Phoenix, AZ.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by the National Association of Epilepsy Centers.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

RELATED ARTICLE

 **Editorial**
Epilepsy Centers in the
United States: The Times
They Are A-Changing
Page 175

MORE ONLINE

 **Infographic**
links.lww.com/WNL/B761

 **CME Course**
NPub.org/cmelist

Glossary

AAN = American Academy of Neurology; ASM = antiseizure medication; CLTE = certified long-term EEG technologist; DRE = drug-resistant epilepsy; EMU = epilepsy monitoring unit; IQR = interquartile range; LiTT = laser interstitial thermal therapy; MEG = magnetoencephalography; NAEC = National Association of Epilepsy Centers; REEGT = registered EEG technologist; RNS = responsive neurostimulation; VNS = vagus nerve stimulator.

Epilepsy affects 1%–2% of the global population, with an estimated prevalence of 3.4 million persons in the United States.¹ About 30% continue to have refractory seizures despite treatment with antiseizure medications (ASMs),² leading to increased morbidity and mortality, decreased quality of life, and increased health care utilization.^{3,4} Persons with drug-resistant epilepsy (DRE) may benefit from surgical intervention, dietary therapy, or access to investigational trials.^{5,6} Therefore, the American Academy of Neurology (AAN) and other professional organizations recommend persons with DRE discuss or undergo evaluation at a comprehensive epilepsy center for consideration of epilepsy surgery evaluation and other specialized services.^{7–9} Epilepsy center practices are steered by national and international practice guidelines and quality measures establishing standards in care delivery and patient safety.^{8,10,11}

Most evaluations and procedures in the United States for DRE are performed at National Association of Epilepsy Centers (NAEC) member institutions. The NAEC is a nonprofit association with a membership of more than 260 epilepsy centers. NAEC requires completion of an annual accreditation process by every member center to evaluate certain criteria of specialized epilepsy centers as outlined by NAEC.⁸ Centers are recognized as level 3 or level 4 centers based on center resources, with level 4 centers serving as regional or national referral sites with comprehensive diagnostic and surgical treatment capabilities.⁸ In general, level 3 epilepsy centers are facilities with EEG, neuroimaging, interdisciplinary epilepsy care, vagus nerve stimulator (VNS) insertion, and epilepsy surgeries not requiring invasive monitoring. Level 4 centers are distinguished by expertise with specialized neuroimaging, intracranial EEG, and more complex surgical techniques. Formal accreditation was implemented in 2015 based on expanded criteria.⁸ Member centers self-report data regarding services provided, procedure volumes, and staff numbers. Sample diagnostic and surgical reports are uploaded and reviewed. Credentials of listed personnel are verified by NAEC staff to meet required levels of certification.

Kaiboriboon and colleagues¹² reviewed NAEC annual report data between 2003 and 2012 and reported growth in number of epilepsy centers, total surgeries, and total VNS insertions across NAEC centers. Since then, NAEC implemented its accreditation program and the number of centers accredited by NAEC greatly expanded. This article describes recent data trends and current state of resources and practice across NAEC member centers and identifies several future directions for driving systematic improvements in epilepsy care.

Methods

We analyzed data obtained from annual reports submitted for 2012, 2016, and 2019 from all level 3 and level 4 NAEC epilepsy centers (eAppendices 1–3, links.lww.com/WNL/B678). These years were selected to assess trends beginning from the last NAEC data publication to the most recently available data.¹² All reported data were collected prior to the COVID-19 pandemic.

Statistical Methods


Data were analyzed using frequency (%) for categorical variables and median (interquartile range [IQR]) for continuous variables. Select variables per capita were described using rates based on US total population data collected from the US Census Bureau (data.census.gov/cedsci). Dependent variables were analyzed by center level (level 3 vs level 4) and by center population (adult only, adult/pediatric combined, or pediatric only). Procedure data in tables are presented 2 ways. First, the frequency (percentage) of centers that performed at least one procedure was displayed. Then, the median (IQR) of procedures performed among those centers that performed at least one of the procedures (excluding zeros) was calculated.

Minor changes in the way data were collected occurred during the study time frame. For example, in 2012, centers with >9 epileptologists were all marked as having ≥ 10 (censored). Thus, both the median for uncensored values and the frequency/percent of censored epileptologist values were reported. This also caused the rate of epileptologists per 1,000,000 population to appear artificially low that year.

If a variable was not collected in one of the years, for example, the 2012 annual report did not include data on center access to magnetoencephalography (MEG), neuropsychology testing, hemispherotomies, laser interstitial thermal therapy (LiTT), or responsive neurostimulation (RNS) implantation (approved by the Food and Drug Administration in 2013), these observations were not considered missing. However, rarely, data points were missing from individual center annual reports and the percent missing data per field was calculated. The intracranial monitoring total was the sum of reported temporal lobectomies with intracranial electrodes, extratemporal resections with intracranial electrodes, and intracranial electrodes with no resection. Missing values were ignored in these sums if one or more component procedure volumes was reported.

Due to the low number of years being compared, we applied nonparametric statistical tests to test for any increasing or

Figure 1 Number of National Association of Epilepsy Centers Member Centers Over Time



(A) By Level. (B) Demographic.

decreasing trend across 3 years. For variables that are expressed as medians or rates per populace, the Jonckheere-Terpstra test¹³ was used. For variables that are expressed as binomial proportions, the Cochran-Armitage test was used.^{14,15} For variables collected only in 2016 and 2019, comparisons between years were conducted using the 2-sided Mood¹⁶ median test for medians and the χ^2 test for proportions. Mood test is a nonparametric alternative to the *t* test for small sample size and non-normally distributed data. If only one procedure was performed in a combination of year and accreditation level/patient demographic, a statistical test could not be computed, and no *p* values were calculated. Due to the descriptive nature of the study, no multiple testing correction was applied. Two-sided

p values less than 0.05 were considered statistically significant. All analyses were conducted using R 4.0 (R Core Team).

Standard Protocol Approvals, Registrations, and Patient Consents

The ethical standards committee at Nationwide Children's Hospital determined this study exempt from institutional review board approval.

Data Availability

Qualified researchers may request data using the NAEC policy governing the release of member center data (naec-epilepsy.org/wp-content/uploads/NAECBoardPoliciesforDataAccess.pdf).

Table 1 Staffing and Length of Stay by Center Level Over Time

Characteristic	Level 3			<i>p</i> Value	Level 4			<i>p</i> Value
	2012 (n = 25) ^b	2016 (n = 40) ^b	2019 (n = 58) ^b		2012 (n = 136) ^b	2016 (n = 184) ^b	2019 (n = 198) ^b	
EEG staff FT	4.0 (2.0, 6.0)	4.0 (2.8, 6.0)	6.0 (4.0, 8.8)	<0.01 ^c	8.0 (5.0, 11.5)	9.0 (6.0, 13.0)	11.0 (7.0, 16.0)	<0.01 ^c
EEG staff REEGT	2.0 (1.0, 3.0)	3.0 (2.0, 4.2)	3.5 (2.0, 5.0)	0.04 ^c	4.0 (2.0, 7.0)	5.0 (3.0, 8.0)	6.5 (4.0, 10.0)	<0.01 ^c
EEG staff CLTM	0.0 (0.0, 0.0)	0.0 (0.0, 1.0)	0.0 (0.0, 1.0)	0.06	0.0 (0.0, 0.0)	0.0 (0.0, 1.0)	1.0 (0.0, 2.0)	<0.01 ^c
Number of epileptologists ^a	2.0 (1.0, 3.0)	3.0 (2.0, 3.0)	3.0 (2.0, 3.8)	0.06	5.0 (3.0, 7.0)	5.0 (3.0, 7.0)	6.0 (4.0, 9.0)	<0.01 ^c
Epileptologists ≥10, n (%) ^a	0.0 (0.0)	—	—	—	10.0 (7.4)	—	—	—
EMU beds, all	4.0 (3.0, 5.0)	4.0 (3.0, 6.0)	4.0 (4.0, 6.0)	0.05	8.0 (5.8, 11.0)	7.0 (5.0, 10.0)	8.0 (6.0, 11.0)	0.48
EMU admissions, all	129.0 (79.0, 207.2)	154.0 (78.5, 228.0)	127.0 (70.5, 194.0)	0.89	340.0 (200.0, 549.0)	310.0 (183.0, 492.8)	330.5 (208.2, 545.5)	0.83
Average LOS	3.0 (2.8, 4.2)	3.1 (2.3, 4.0)	3.3 (2.6, 4.0)	0.80	4.0 (3.0, 4.6)	3.5 (3.0, 4.2)	3.5 (2.9, 4.0)	<0.01 ^c

Abbreviations: CLTM = certified long-term monitoring; EMU = epilepsy monitoring unit; FT = full time; LOS = length of stay; REEGT = registered EEG technologist.

^a Due to the design limitation, the 2012 survey had a predefined choice, but did not record the number of epileptologists when there are more than 10.

^b Statistics presented: median (interquartile range).

^c Significant.

Table 2 Staffing and Length of Stay by Center Population Over Time

Characteristic	Adult			Adult/pediatric			Pediatric					
	2012 (n = 52) ^b	2016 (n = 88) ^b	2019 (n = 110) ^c	p Value	2012 (n = 87) ^b	2016 (n = 93) ^b	2019 (n = 96) ^b	p Value	2012 (n = 22) ^b	2016 (n = 43) ^b	2019 (n = 50) ^b	p Value
EEG staff FT	6.0 (4.0, 8.0)	6.0 (4.0, 9.0)	8.0 (4.2, 12.8)	<0.01 ^c	7.5 (5.0, 11.0)	9.0 (5.0, 14.0)	10.0 (6.0, 16.0)	0.01 ^c	10.0 (7.2, 14.5)	10.0 (6.0, 13.5)	12.0 (8.0, 17.5)	0.10
EEG staff REEGT	3.0 (2.0, 5.0)	4.0 (2.0, 6.0)	5.0 (3.0, 7.0)	<0.01 ^c	5.0 (2.0, 7.0)	5.0 (3.0, 8.0)	5.0 (3.0, 9.0)	0.05	4.0 (3.0, 7.2)	6.0 (3.0, 8.5)	7.5 (4.0, 11.8)	0.01 ^c
EEG staff CLTM	0.0 (0.0, 0.0)	0.0 (0.0, 1.0)	0.0 (0.0, 1.0)	<0.01 ^c	0.0 (0.0, 0.0)	0.0 (0.0, 1.0)	1.0 (0.0, 2.0)	<0.01 ^c	0.0 (0.0, 0.0)	0.0 (0.0, 1.0)	1.0 (0.0, 2.0)	<0.01 ^c
Number of epileptologists ^a	3.0 (2.0, 4.0)	3.0 (2.0, 5.0)	4.0 (2.2, 6.0)	0.02 ^c	5.0 (3.0, 7.0)	6.0 (4.0, 9.0)	6.0 (4.0, 9.0)	<0.01 ^c	4.0 (3.0, 7.0)	5.0 (3.0, 7.0)	5.0 (4.0, 8.0)	0.09
Epileptologists ≥10, n (%) ^b	1.0 (1.9)	—	—	—	8.0 (9.2)	—	—	—	1.0 (4.5)	—	—	—
EMU beds, all	5.0 (4.0, 8.0)	5.0 (4.0, 7.0)	6.0 (4.0, 8.0)	0.17	8.0 (6.0, 11.0)	8.0 (6.0, 12.0)	9.0 (6.0, 12.0)	0.47	7.5 (6.0, 9.8)	8.0 (4.0, 8.5)	8.0 (6.0, 10.0)	0.28
EMU admissions, all	200.0 (115.0, 291.5)	181.0 (112.5, 277.0)	177.0 (110.0, 280.5)	0.59	329.5 (198.5, 557.5)	331.0 (183.0, 484.0)	331.0 (203.0, 495.0)	0.66	504.5 (381.5, 589.2)	525.0 (316.0, 659.5)	635.5 (345.0, 810.5)	0.22
Average LOS	4.2 (3.4, 5.0)	4.0 (3.3, 4.5)	4.0 (3.3, 4.5)	0.18	4.0 (3.0, 4.5)	3.5 (3.0, 4.0)	3.3 (2.9, 4.0)	<0.01 ^c	2.5 (2.0, 3.2)	2.0 (1.8, 3.0)	2.4 (1.8, 3.0)	0.31

Abbreviations: CLTM = certified long-term monitoring; EMU = epilepsy monitoring unit; FT = full time; LOS = length of stay; REEGT = registered EEG technologist.

^a Due to the design limitation, the 2012 survey had a predefined choice, but did not record the actual number of epileptologists when there are more than 10.

^b Statistics presented: median (interquartile range).

^c Significant.

Results

Reporting centers increased over time from 161 in 2012 to 256 in 2019. Annual report completion rate was 100% across all years. The range of data missingness for all variables was <10% (eTable 1, links.lww.com/WNL/B678).

Center Demographics and Services

NAEC member centers grew across all levels and populations during the study period (Figure 1). The largest increase occurred in adult- or pediatric-only centers, rather than those serving both children and adults. Summary findings for center personnel, facility size, and characteristics are detailed by center level (Table 1) and patient population (Table 2).

Median center size (epilepsy monitoring unit [EMU] beds and admissions) did not change. However, aggregate access to epilepsy care per population did increase, with most of the increase driven by increased numbers of level 3 and pediatric centers. Per 1 million people, EMU admissions increased from 2012 to 2019 from 11.7 to 30.3 (+159%) in level 3 centers and 193 to 259 (+34%) in level 4 centers. By population served per 1 million people, EMU admissions changed from 37.6 to 75.4 (+100%) in adult centers, 129.2 to 121.3 (−6%) in combined centers, and 38.1 to 93.1 (+144%) in pediatric centers. Length of stay decreased in level 4 centers and those that served combined adult/pediatric populations.

Growth was positive in EMU beds per 1 million people (level 3: 0.3 to 0.9 [+200%]; level 4: 3.6 to 4.3 [+47%]; adult centers: 0.9 to 2.1 [+133%]; combined centers: 2.5 to 2.8 [+12%]; pediatric centers: 0.5 to 1.3 [+160%]). Number of epileptologists per 1 million people also increased from 2012 to 2019 (level 3: 0.2 to 0.5 [+150%]; level 4: 2.0 to 4.1 [+105%]; adult centers: 0.6 to 1.6 [+167%]; combined centers: 1.3 to 2.1 [+62%]; pediatric centers: 0.3 to 1.0 [+233%]).

Growth in median center staff EEG technologists occurred in adult and combined adult/pediatric centers (Table 2). Centers hired more registered EEG technologists (REEGTs) in adult and pediatric centers and certified long-term EEG technologists in combined and pediatric centers. Total certified EEG technologists accounted for more than half of technologists at centers.

Access to testing modalities and specialized services broadly expanded across both levels (eTable 2, links.lww.com/WNL/B678) and all populations (eTable 3). Greatest growth areas included alternative/complementary medicine (adult centers 1.9%–32.9%; combined centers 0%–51.1%; pediatric centers 0%–36%), ketogenic diet in level 3 and adult centers (level 3 28%–48.3%; adult centers 36.5%–54.5%), and genetic testing or counseling (level 3 64%–79.3%; level 4 72.8%–90.9%; pediatric centers 63.6%–100%). Access to MEG, which was only included in 2016 and 2019 survey data, was relatively stable (level 3 0%–8.6%; level 4 23.9%–19.2%; adult centers

11.4%–12.7%; combined centers 18.3%–12.5%; pediatric centers 39.5%–34%).

Center Procedures

Procedure data were analyzed across all years by aggregate volumes and population rates (Table 3) and by center level median volume and utilization rate by level (Table 4) and patient population (Table 5). Most surgeries were performed at level 4 centers (eFigure 1, links.lww.com/WNL/B678).

Over the study period, the total volume of temporal lobectomies (5.9%), extratemporal resections (11.9%), and hemispherectomies/otomies (13.1%) increased slower than center growth (59%). Corpus callosotomies (–12.8%) and VNS implantations (–2.4%) decreased during the study timeframe. In contrast, growth in laser ablations (61%), RNS implantations (114%), and intracranial monitoring without resection (152%) outpaced overall center growth.

Median volumes of some procedure types changed over time between center levels. Median number of temporal lobectomies

decreased in level 4 centers from 9 to 6 ($p = 0.01$), but changes in extratemporal resections in level 4 centers did not reach significance (5.5–4; $p = 0.08$). Level 4 centers performed fewer VNS implantations (13–9; $p \leq 0.01$). Level 4 centers performed more intracranial monitoring without resection (2–4.5; $p \leq 0.01$), while level 3 centers performed less (3–1; $p = 0.05$).

Procedures across target populations changed as well. Corpus callosotomies decreased in adult centers from 15 to 12 ($p = 0.04$). Pediatric centers had a decrease in temporal lobectomies from 10 to 4.5 ($p = 0.04$) and corpus callosotomies from 4 to 2 ($p = 0.04$). Median intracranial monitoring without resection procedures increased in adult centers from 2 to 5 ($p \leq 0.01$) and pediatric centers from 1.5 to 4.5 ($p \leq 0.01$).

Procedure utilization changed over time. LiTT and RNS were most often utilized by level 4 centers and increased substantially from 40.1% to 58.4% ($p \leq 0.01$) and 39.9%–65.4% ($p \leq 0.01$), respectively. LiTT utilization also increased in combined centers from 35.2% to 53.9% ($p = 0.02$). There

Table 3 Aggregate Procedure Counts and Rate per Population per Year

Characteristic	Procedure volume			% Change			p Value
	2012 (n = 161) ^a	2016 (n = 224) ^a	2019 (n = 256) ^a	2012 to 2016	2016 to 2019	2012 to 2019	
Temporal lobectomy	1,424	1,541	1,508	8.2	–2.1	5.9	<0.01 ^c
Temporal lobectomy per 1M population	4.54	4.77	4.59	5.1	–3.8	1.1	<0.01 ^c
Extratemporal resection	784	1,016	877	29.6	–13.7	11.9	0.06
Extratemporal resection per 1M population	2.50	3.14	2.67	25.6	–15.0	6.8	<0.01 ^c
Corpus callosotomy	179	170	156	–5.0	–8.2	–12.8	0.54
Corpus callosotomy per 1M population	0.57	0.53	0.48	–7.0	–9.4	–15.8	<0.01 ^c
VNS implantation	2,732	2,080	2,667	–23.9	28.2	–2.4	<0.01 ^c
VNS implantation per 1M population	8.70	6.44	8.13	–26.0	26.2	–6.6	<0.01 ^c
Hemispherotomy ^b	—	168	190	—	13.1	—	0.85
Hemispherotomy per 1M population ^b	—	0.52	0.58	—	11.5	—	0.05 ^c
Laser ablation ^b	—	438	705	—	61.0	—	0.18
Laser ablation per 1M population ^b	—	1.36	2.15	—	58.1	—	0.88
RNS implantation ^b	—	264	566	—	114.4	—	0.96
RNS implantation per 1M population ^b	—	0.82	1.72	—	109.8	—	0.12
Total intracranial monitoring	1,746	1,817	2,309	4.1	27.1	32.2	0.26
Total intracranial monitoring per 1M population	5.56	5.62	7.03	1.1	25.1	26.4	0.03 ^c
Intracranial electrodes, no resection	406	506	1,022	24.6	102.0	151.7	<0.01 ^c
Intracranial electrodes, no resection per 1M population	1.29	1.57	3.11	21.7	98.1	141.1	0.24

Abbreviations: RNS = responsive neurostimulation; VNS = vagus nerve stimulation.

^a Statistics presented: sum.

^b p Values calculated using Jonckheere-Terpstra test or Mood median test for noted variables.

^c Significant.

Table 4 Procedure Utilization (%) and Median Volume by Center Level Over Time for Centers With at Least 1 Procedure

Characteristic	Level 3			p Value	Level 4			p Value
	2012 (n = 25) ^a	2016 (n = 40) ^a	2019 (n = 58) ^a		2012 (n = 136) ^a	2016 (n = 184) ^a	2019 (n = 198) ^a	
Temporal lobectomy, %	40.0	33.3	17.6	0.08	91.7	93.9	92.1	0.85
Temporal lobectomy	4.5 (3.0, 6.0)	2.0 (1.0, 2.8)	3.0 (2.0, 5.0)	0.20	9.0 (4.0, 15.0)	6.0 (3.0, 12.0)	6.0 (3.0, 11.0)	<0.01 ^d
Extratemporal resection, %	10.0	14.3	10.0	0.91	79.1	79.4	75.3	0.84
Extratemporal resection	1.5 (1.2, 1.8)	3.0 (2.5, 4.0)	1.0 (1.0, 2.0)	0.61	5.5 (3.0, 9.8)	4.0 (2.0, 9.0)	4.0 (2.0, 8.0)	0.08
Corpus callosotomy, %	5.0	3.4	2.0	0.28	48.8	35.0	33.7	0.09
Corpus callosotomy	5.0 (5.0, 5.0)	4.0 (4.0, 4.0)	2.0 (2.0, 2.0)	0.33	2.0 (1.0, 2.8)	2.0 (1.0, 3.0)	2.0 (1.0, 3.0)	0.61
VNS implantation, %	72.7	69.7	77.4	0.85	98.5	93.3	94.9	0.78
VNS implantation	8.5 (4.5, 10.0)	8.0 (2.0, 14.0)	6.0 (3.0, 11.0)	0.51	13.0 (6.2, 24.0)	8.5 (4.0, 16.0)	9.0 (5.0, 17.0)	<0.01 ^d
Hemispherotomy, % ^{b,c}	—	0.0	0.0	—	—	34.5	32.6	0.79
Hemispherotomy ^{b,c}	—	—	—	—	—	2.0 (1.0, 3.0)	2.0 (1.0, 4.0)	0.85
Laser ablation, % ^{b,c}	—	3.6	10.0	—	—	40.1	58.4	<0.01 ^d
Laser ablation ^{b,c}	—	1.0 (1.0, 1.0)	2.0 (2.0, 3.0)	—	—	3.0 (1.0, 9.0)	5.0 (2.0, 8.0)	0.13
RNS implantation, % ^{b,c}	—	3.4	6.0	—	—	39.9	65.4	<0.01 ^d
RNS implantation ^{b,c}	—	1.0 (1.0, 1.0)	1.0 (1.0, 2.0)	—	—	3.0 (1.0, 5.0)	3.0 (1.0, 6.0)	0.88
Total intracranial monitoring, %	24.0	16.7	17.6	0.74	94.9	90.1	94.3	0.82
Total intracranial monitoring	3.0 (2.2, 5.2)	1.0 (1.0, 5.0)	1.0 (1.0, 2.0)	0.05	11.0 (6.0, 17.0)	8.0 (4.0, 14.0)	8.0 (5.0, 17.0)	0.37
Intracranial electrodes, no resection, %	12.0	7.1	11.8	0.94	69.1	62.6	81.2	0.16
Intracranial electrodes, no resection	1.0 (1.0, 4.0)	1.0 (1.0, 1.0)	1.0 (1.0, 1.0)	0.73	2.0 (1.0, 5.0)	3.0 (1.0, 5.0)	4.5 (2.0, 8.0)	<0.01 ^d

Abbreviations: RNS = responsive neurostimulation; VNS = vagus nerve stimulation.

^a Statistics presented: median (interquartile range).

^b p Values calculated using χ^2 test or Mood median test for noted variables.

^c Test could not be conducted because there were fewer than 2 procedures performed in a group.

^d Significant.

were no significant changes in median LiTT procedures, though they trended up in pediatric centers from 2 to 4 ($p = 0.11$). RNS utilization increased in combined centers from 41.6% to 66.7% ($p \leq 0.01$) and pediatric centers from 7.5% to 36.7% ($p \leq 0.01$) with no change in procedure medians across categories.

Discussion

Over the study period, access to specialty epilepsy care in the United States improved as the NAEC implemented accreditation criteria and expanded from 161 to 256 centers. Member centers increased available epilepsy services and hired more epileptologists and EEG technologists. Surgical case complexity increased while aggregate surgical volume remained stable or declined slightly across most procedure types, with a corresponding decline in cases per center. This study highlights several changing trends in epilepsy management in the United States, identifies important data gaps, and underscores future opportunities to study and improve quality care.

Implementation of accreditation criteria emphasizing a broad range of services delivered by expert medical personnel increased center accountability even as center availability also increased. Center growth was greatest among adult- or pediatric-focused centers rather than combined adult/pediatric centers. This is likely due in part to a change in NAEC requirements for center accreditation resulting in hospitals with combined adult/pediatric centers applying for separate accreditation.

Aggregate increases in EMU beds, EEG staff, and epileptologists followed similar trends. Centers hired more epileptologists and EEG technologists, while median EMU beds and admissions per center were stable. Growth in median number of epileptologists per center parallels an increase in case complexity,¹⁷ increasing ICU continuous EEG volume,¹⁸ and outpatient volume. Growth in EEG technologists reflects similar trends. Technologists are increasingly called upon to monitor and screen continuous video-EEG studies.¹⁹ Most are obtaining REEGT credentials after meeting professional competencies, a

Table 5 Procedure Utilization (%) and Median Volume by Center Population Over Time for Centers With at Least 1 Procedure

Characteristic	Adult				Adult/pediatric				Pediatric			
	2012 (n = 52) ^a	2016 (n = 88) ^a	2019 (n = 110) ¹	p Value	2012 (n = 87) ^a	2016 (n = 93) ^a	2019 (n = 96) ^a	p Value	2012 (n = 22) ^a	2016 (n = 43) ^a	2019 (n = 50) ^a	p Value
Temporal lobectomy, %	83.7	82.7	66.3	0.34	88.0	86.5	82.6	0.92	75.0	87.8	85.7	0.74
Temporal lobectomy	8.0 (4.0, 12.0)	6.0 (3.0, 11.0)	6.0 (3.0, 11.0)	0.15	9.0 (4.0, 15.0)	6.0 (3.0, 12.0)	6.5 (3.0, 12.0)	0.15	10.0 (5.5, 11.5)	5.0 (3.0, 10.2)	4.5 (2.2, 7.8)	0.04 ^c
Extratemporal rxn, %	53.1	63.7	49.0	0.68	77.5	68.2	64.0	0.35	80.0	90.0	83.7	0.95
Extratemporal rxn	4.0 (2.0, 7.0)	4.0 (1.5, 7.5)	3.0 (2.0, 5.0)	0.34	5.0 (3.0, 8.8)	3.0 (1.0, 7.2)	4.0 (2.0, 8.0)	0.12	10.0 (7.5, 16.8)	5.5 (3.8, 13.2)	7.0 (2.0, 11.0)	0.11
Corpus callosotomy, %	31.2	17.9	11.8	0.04 ^c	39.2	30.7	29.2	0.09	85.0	55.0	55.1	0.30
Corpus callosotomy	1.0 (1.0, 2.0)	1.0 (1.0, 2.8)	1.5 (1.0, 2.0)	0.90	2.0 (1.0, 2.0)	2.0 (1.0, 2.5)	1.0 (1.0, 3.0)	0.91	4.0 (2.0, 7.0)	3.0 (2.0, 4.5)	2.0 (1.0, 4.0)	0.04 ^c
VNS implantation, %	94.0	82.9	88.7	0.93	95.3	92.1	92.5	0.74	95.5	97.6	93.9	0.97
VNS implantation	10.0 (5.0, 15.0)	7.0 (4.0, 10.0)	7.0 (4.0, 12.0)	0.10	14.0 (6.0, 24.0)	10.0 (5.0, 16.0)	9.5 (5.0, 19.8)	0.05	17.0 (12.0, 36.0)	10.0 (5.0, 18.0)	11.0 (5.0, 16.8)	0.09
Hemispherotomy, % ^b	—	10.3	6.9	0.60	—	21.6	20.7	>0.99	—	85.0	73.5	0.29
Hemispherotomy ^b	—	1.0 (1.0, 2.2)	1.0 (1.0, 2.5)	>0.99	—	2.0 (1.0, 3.0)	2.0 (1.0, 2.8)	>0.99	—	2.0 (1.0, 4.0)	2.0 (1.8, 5.0)	0.64
Laser ablation, % ^b	—	31.2	38.2	0.41	—	35.2	53.9	0.02 ^c	—	42.5	59.2	0.18
Laser ablation ^b	—	5.5 (1.0, 9.2)	6.0 (4.0, 8.0)	>0.99	—	3.0 (2.0, 9.5)	3.0 (2.0, 6.0)	>0.99	—	2.0 (1.0, 4.0)	4.0 (3.0, 9.0)	0.11
RNS implantation, % ^b	—	41.0	49.0	0.36	—	41.6	66.7	<0.01 ^c	—	7.5	36.7	<0.01 ^c
RNS implantation ^b	—	3.0 (1.0, 4.2)	4.0 (2.0, 6.8)	0.07	—	3.0 (2.0, 5.0)	3.0 (1.0, 6.0)	0.83	—	2.0 (2.0, 5.5)	2.0 (1.0, 3.8)	>0.99
Total IC monitoring, %	76.9	75.3	69.9	0.98	85.1	83.1	82.6	0.78	95.5	80.5	87.8	0.94
Total IC monitoring	7.5 (4.0, 14.0)	7.0 (4.0, 12.0)	7.0 (3.0, 15.2)	0.94	10.5 (6.0, 16.0)	7.0 (3.2, 14.0)	7.5 (4.0, 17.2)	0.37	14.0 (7.0, 23.0)	9.0 (5.0, 12.0)	11.0 (4.5, 16.0)	0.17
IC electrodes, no rxn, %	55.8	57.7	61.2	0.29	59.8	56.2	71.4	0.29	72.7	47.5	69.4	0.73
IC electrodes, no rxn	2.0 (1.0, 4.0)	3.0 (1.0, 6.0)	5.0 (2.0, 8.0)	<0.01 ^c	2.5 (1.0, 6.0)	3.0 (2.0, 5.8)	4.0 (2.0, 8.0)	0.15	1.5 (1.0, 2.2)	2.0 (1.0, 3.5)	4.5 (3.0, 6.8)	<0.01 ^c

Abbreviations: IC = intracranial; rxn = resection.

^a Statistics presented: median (interquartile range).^b p Values calculated using χ^2 test or Mood median test for noted variables.^c Significant.

qualifying examination, and continuing education requirements. However, few hold Certification in Long-Term Monitoring credentials, which may present an opportunity to further improve seizure detection and patient care.²⁰

Availability of supplemental services across NAEC-accredited centers increased during the study period. Alternative or complementary medicine is now available at nearly half of centers, where previous availability was sparse. Ketogenic diet availability has grown in level 3 centers and adult centers. Genetic testing and counseling have been increasingly utilized for epilepsy diagnosis and management across level 3 and level 4 centers. Access to advanced diagnostic and therapeutic options increased over the study period even as more epilepsy centers achieved NAEC accreditation.

Despite an increase in access to epilepsy centers, aggregate surgical volumes did not uniformly increase across all procedures. Temporal lobectomies, extratemporal resections, and hemispherectomies/otomies did not keep pace with center growth, and corpus callosotomy and VNS implantation volumes decreased over the study period. In contrast, LiTT, RNS implantation, and intracranial monitoring without resection grew substantially in aggregate, although median center volumes did not. The sole procedure type with increasing median volumes across centers was intracranial monitoring without resection.

The trends in surgical procedures reflect a shift in epilepsy surgery toward greater surgical case complexity and use of newer technology. Stereo EEG is increasingly utilized across the United States and may carry a better safety and tolerability profile compared to subdural grids,²¹ which likely lowers the threshold for performing intracranial monitoring.²² Increasing use of LiTT may have displaced some temporal lobectomies, although the increase of LiTT cases far exceeded the decline in temporal lobectomies from 2016 to 2019.^{12,23-25} However, despite introduction of new techniques and technologies, the overall rate of epilepsy surgery has not kept pace with the growth in NAEC centers.^{26,27}

The current NAEC accreditation process emphasizes structural and process measures of health care quality. Accreditation criteria for all NAEC members include staffing centers with well-trained personnel, establishing safety and quality protocols, and providing a breadth of key services. Board-certified epileptologists and surgeons with specific epilepsy experience are accreditation criteria for level 4 centers and level 3 centers that perform surgery.

Assigning a process measure of minimum surgical volume for epilepsy surgery centers is an ongoing topic of interest. Several evidence-based recommendations from International League Against Epilepsy and other working groups have proposed minimum procedure volumes for epilepsy centers.^{7,28,29} These recommendations are, in part, based on a higher complication rate in disconnection or resection surgeries at centers with fewer than 15 such procedures per

year.^{23,30} However, NAEC does not require a minimal surgical volume for epilepsy centers, as reliance on volume measures alone can also result in unintended consequences, such as centers performing unnecessary surgery or certain types of surgery simply to attain accreditation status.

Patient outcome data across epilepsy centers are urgently needed to develop outcome measures and better incentivize quality care. Data across epilepsy center networks could be collected as part of the NAEC accreditation process, with the goal of progressively expanding the set of data collected and increasing the standard of care. Indeed, NAEC is helping fund a practice-based learning health network focused on epilepsy care that may provide data and infrastructure to improve outcomes.³¹ Addressing these data gaps will better enable the study of seizure-related outcomes and quality of life across large care delivery networks.

The findings are limited primarily by how data were acquired through the NAEC accreditation annual reports. Data acquisition methods changed during the study period and rely primarily on self-reporting of administrative data. The response rate was 100% with a very low range of missingness since the annual data are required for NAEC accreditation. Our data are in line with previous studies while providing novel data regarding center characteristics. Although NAEC member centers do not provide the entirety of epilepsy care in the United States, they likely represent most of the specialized evaluation and procedures for those with DRE. For instance, the Veterans Affairs health care system is a large provider within the United States, yet performed 5 resections for epilepsy in 2019 and implanted 4 neurostimulators.³² Therefore, our analysis likely reflects accurate data regarding major trends and the current state in epilepsy care in the United States.

Specialty epilepsy care in the United States has significantly changed over the study period. Additional studies are required to better understand the influence of epilepsy center characteristics on testing, treatment, and outcomes. Epilepsy centers are key to standardizing access and delivery of epilepsy care across health care networks.

Acknowledgment

The authors thank the former NAEC board members and the medical directors of NAEC-accredited epilepsy centers that submit data annually to NAEC and who supported the strengthening of NAEC's accreditation program to improve the quality of epilepsy care in the United States.

Study Funding

This study was supported by Award 45141-0001-0321 from Nationwide Children's Hospital and Award 810712-1221-00 from the National Association of Epilepsy Centers.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

Publication History

Received by *Neurology* June 25, 2021. Accepted in final form November 4, 2021.

Appendix Authors

Name	Location	Contribution
Adam P. Ostendorf, MD	Nationwide Children's Hospital, Columbus, OH	Designed and conceptualized study; data acquisition; analysis and interpretation of data; study supervision; drafted and revised manuscript
Stephanie M. Ahrens, DO	Nationwide Children's Hospital, Columbus, OH	Designed and conceptualized study; analysis and interpretation of data; drafted and revised manuscript
Fred A. Lado, MD, PhD	Hofstra Northwell Comprehensive Epilepsy Center, Great Neck, NY	Study concept and design; data acquisition; revised the manuscript for intellectual content
Susan T. Arnold, MD	Comprehensive Epilepsy Center, Children's Medical Center Dallas, TX	Study concept and design; data acquisition; revised the manuscript for intellectual content
Shasha Bai, PhD	Emory University School of Medicine, Atlanta, GA	Performed biostatistical analysis; revised the manuscript for intellectual content
Meriem L. Bensalem-Owen, MD	University of Kentucky Comprehensive Epilepsy Center, Lexington	Study concept and design; data acquisition; revised the manuscript for intellectual content
Kevin E. Chapman, MD	Barrow Neurologic Institute at Phoenix Children's Hospital, AZ	Study concept and design; data acquisition; revised the manuscript for intellectual content
Dave F. Clark, MD	UT Health Austin Pediatric Neurosciences at Dell Children's Comprehensive Pediatric Epilepsy Program, Austin, TX	Study concept and design; data acquisition; revised the manuscript for intellectual content
Mariah Eisner, MS	Nationwide Children's Hospital, Columbus, OH	Performed statistical analysis; revised the manuscript for intellectual content
Nathan B. Fountain, MD	F.E. Drefuss Comprehensive Epilepsy Program, University of Virginia, Charlottesville	Study concept and design; data acquisition; revised the manuscript for intellectual content
Johanna M. Gray, MPA	National Association of Epilepsy Centers, Washington, DC	Data acquisition; revised the manuscript for intellectual content
Jennifer L. Hopp, MD	University of Maryland Epilepsy Center, Baltimore	Study concept and design; data acquisition; revised the manuscript for intellectual content
Ellen Riker, MHA	National Association of Epilepsy Centers, Washington, DC	Data acquisition; revised the manuscript for intellectual content
Stephan U. Shuele, MD, MPH	Northwestern Medicine Comprehensive Epilepsy Center, Chicago, IL	Study concept and design; data acquisition; revised the manuscript for intellectual content

Appendix (continued)

Name	Location	Contribution
Barbara V. Small, BA	National Association of Epilepsy Centers, Washington, DC	Data acquisition; revised the manuscript for intellectual content
Susan T. Herman, MD	Barrow Neurologic Institute Comprehensive Epilepsy Center, Phoenix, AZ	Designed and conceptualized study; data acquisition and analysis; supervised the study; revised manuscript for intellectual content

References

- Zack MM, Kobau R. National and state estimates of the numbers of adults and children with active epilepsy: United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2017;66(31):821-825.
- Kwan P, Brodie MJ. Early identification of refractory epilepsy. *N Engl J Med*. 2000;342(5):314-319.
- Ostendorf AP, Gedela S. Effect of epilepsy on families, communities, and society. *Semin Pediatr Neurol*. 2017;24(4):340-347.
- Wilner AN, Sharma BK, Thompson A, Soucy A, Krueger A. Diagnoses, procedures, drug utilization, comorbidities, and cost of health care for people with epilepsy in 2012. *Epilepsy Behav*. 2014;41:83-90.
- Dwivedi R, Ramanujam B, Chandra PS, et al. Surgery for drug-resistant epilepsy in children. *N Engl J Med*. 2017;377(17):1639-1647.
- Wiebe S, Blume WT, Girvin JP, Eliasziw M. Effectiveness and efficiency of surgery for temporal lobe epilepsy study group: a randomized, controlled trial of surgery for temporal-lobe epilepsy. *N Engl J Med*. 2001;345:311-318.
- Rosenow F, Bast T, Czech T, et al. Revised version of quality guidelines for presurgical epilepsy evaluation and surgical epilepsy therapy issued by the Austrian, German, and Swiss working group on presurgical epilepsy diagnosis and operative epilepsy treatment. *Epilepsia*. 2016;57(8):1215-1220.
- Labiner DM, Bagic AI, Herman ST, et al. Essential services, personnel, and facilities in specialized epilepsy centers: revised 2010 guidelines: guidelines for specialized epilepsy centers. *Epilepsia*. 2010;51:2322-2333.
- Patel AD, Baca C, Franklin G, et al. Quality improvement in neurology: epilepsy quality measurement set 2017 update. *Neurology*. 2018;91(18):829-836.
- National Clinical Guideline Centre (UK). *The Epilepsies: The Diagnosis and Management of the Epilepsies in Adults and Children in Primary and Secondary Care: Pharmacological Update of Clinical Guideline 20* [online].ncbin.nlm.nih.gov/books/NBK247130/. Royal College of Physicians; 2012. Accessed February 23, 2021.
- Fountain NB, Van Ness PC, Swain-Eng R, Tonn S, Bever CT. American Academy of Neurology Epilepsy Measure Development Panel and the American Medical Association-Convended Physician Consortium for Performance Improvement Independent Measure Development Process. Quality improvement in neurology: AAN epilepsy quality measures: report of the quality measurement and reporting subcommittee of the American Academy of neurology. *Neurology*. 2011;76(1):94-99.
- Kaiboriboon K, Malkhachroum AM, Zrik A, et al. Epilepsy surgery in the United States: analysis of data from the National Association of Epilepsy Centers. *Epilepsy Res*. 2015;116:105-109.
- Jonckheere AR. A distribution-free k-sample test against ordered alternatives. *Biometrika*. 1954;41:133.
- Cochran WG. Some methods for strengthening the common χ^2 tests. *Biometrics*. 1954;10:417.
- Armitage P. Tests for linear trends in proportions and frequencies. *Biometrics*. 1955;11:375.
- Mood AM. On the asymptotic efficiency of certain nonparametric two-sample tests. *Ann Math Statist*. 1954;25:514-522.
- Cukiert A, Rydenhag B, Harkness W, Cross JH, Gaillard WD; task force for pediatric epilepsy surgery for the ILAE commissions of pediatrics and surgical therapies. Technical aspects of pediatric epilepsy surgery: report of a multicenter, multinational web-based survey by the ILAE task force on pediatric epilepsy surgery. *Epilepsia*. 2016;57:194-200.
- Ney JP, van der Goes DN, Nuwer MR, Nelson L, Echer MA. Continuous and routine EEG in intensive care: utilization and outcomes, United States 2005-2009. *Neurology*. 2013;81(23):2002-2008.
- Gavvala J, Abend N, LaRoche S, et al. Continuous EEG monitoring: a survey of neurophysiologists and neurointensivists. *Epilepsia*. 2014;55(11):1864-1871.
- Ahrens S, Twanow JD, Vidaurre J, Gedela S, Moore-Clingenpeel M, Ostendorf AP. Electroencephalography technologist inter-rater agreement and interpretation of pediatric critical care electroencephalography. *Pediatr Neurol*. 2021;115:66-71.
- Willems LM, Reif PS, Spyranis A, et al. Invasive EEG-electrodes in presurgical evaluation of epilepsies: systematic analysis of implantation-, video-EEG-monitoring- and explantation-related complications, and review of literature. *Epilepsy Behav*. 2019;91:30-37.
- Gavvala J, Zafar M, Sinha SR, Kalamangalam G, Schuele S. Stereotactic EEG practices: a survey of United States tertiary referral epilepsy centers. *J Clin Neurophysiol*. Epub 2020 Nov 10.

23. Kwon CS, Blank L, Mu L, Jetté N. Trends in lobectomy/amygdalohippocampectomy over time and the impact of hospital surgical volume on hospitalization outcomes: a population-based study. *Epilepsia*. 2020;61(10):2173-2182.
24. Barba C, Specchio N, Guerrini R, et al. Increasing volume and complexity of pediatric epilepsy surgery with stable seizure outcome between 2008 and 2014: a nationwide multicenter study. *Epilepsy Behav*. 2017;75:151-157.
25. Englot DJ, Ouyang D, Garcia PA, Barbaro NM, Chang EF. Epilepsy surgery trends in the United States, 1990-2008. *Neurology*. 2012;78(16):1200-1206.
26. Pestana Knight EM, Schiltz NK, Bakaki PM, Koroukian SM, Lhatoo SD, Kaiboriboon K. Increasing utilization of pediatric epilepsy surgery in the United States between 1997 and 2009. *Epilepsia*. 2015;56:375-381.
27. Cloppenborg T, May TW, Blümcke I, et al. Trends in epilepsy surgery: stable surgical numbers despite increasing presurgical volumes. *J Neurol Neurosurg Psychiatry*. 2016; 87(12):1322-1329.
28. Gaillard WD, Jette N, Arnold ST, et al. Establishing criteria for pediatric epilepsy surgery center levels of care: report from the ILAE Pediatric Epilepsy Surgery Task Force. *Epilepsia*. 2020;61(12):2629-2642.
29. Binnie CD, Polkey CE; International League Against Epilepsy. Commission on neurosurgery of the international league against epilepsy (ILAE) 1993-1997: recommended standards. *Epilepsia*. 2000;41(10):1346-1349.
30. Englot DJ, Ouyang D, Wang DD, Rolston JD, Garcia PA, Chang EF. Relationship between hospital surgical volume, lobectomy rates, and adverse perioperative events at US epilepsy centers. *J Neurosurg*. 2013;118(1):169-174.
31. Donahue MA, Herman ST, Dass D, et al. Establishing a learning healthcare system to improve health outcomes for people with epilepsy. *Epilepsy Behav*. 2021;117:107805.
32. Kelly P. *Epilepsy Centers of Excellence. Annual Report Fiscal Year 2019 [online]*. epilepsy.va.gov/docs/FY19AnnualReport-VHAEpilepsyCentersofExcellence.pdf. US Department of Veterans Affairs; 2019. Accessed February 18, 2021.