

How I Teach: Ultrasound-guided Peripheral Venous Access

AUTHORS: Matthew Gorgone¹, Timothy P. O'Connor², and Stephanie I. Maximous¹

¹Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Medicine, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania; and ²University of Rochester Medical Center, Department of Emergency Medicine, Rochester, New York

ABSTRACT

Ultrasound-guided peripheral intravenous (IV) placement is often required for patients with difficult IV access and is associated with a reduction in central line placement. Despite the importance, there is no standardized technical approach, and there is limited ability to attain mastery through simulation. We describe our step-by-step approach for teaching ultrasound-guided IV placement at the bedside using short-axis dynamic guidance, with emphasis on advancing the needle and catheter device almost entirely into the vessel before threading the catheter. Our teaching approach allows the opportunity for trainees to maximize the learning potential of a single insertion experience, which includes focused preprocedure hands-on practice, instruction with real-time feedback at the bedside, and a post-procedure debrief with reinforcement of concepts.

Intravenous (IV) access is essential in hospitalized patients, and poor IV access is a common reason for central line placement. Peripheral IVs are associated with a lower infection risk and fewer placement complications (1, 2). Using ultrasound can reduce the number of attempts required to place peripheral IVs in patients with one or more failed IV attempts, no visually identifiable vein, or previous difficult IV placement (3, 4). Ultrasound-guided peripheral IVs (USPIVs)

allow clinicians to avoid central lines, remove existing central lines, improve patient satisfaction, and safely administer vasoconstrictive agents (4–10). Novice members of the team are often called to place IVs after multiple expert staff have failed, highlighting the importance of this skill.

WHO ARE THE LEARNERS?

Our technique is aimed at teaching anyone learning USPIV placement, including but not

This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern.

Correspondence and requests for reprints should be addressed to Matthew Gorgone, D.O., F.A.C.P., Division of Pulmonary, Allergy, and Critical Care Medicine, NW628 UPMC Montefiore Hospital, 3459 Fifth Avenue, Pittsburgh, PA 15213. E-mail: matthew.j.gorgone@gmail.com.

This article has a related editorial.

This article has a data supplement, which is accessible from this issue's table of contents at www.atsjournals.org.

ATS Scholar Vol 3, Iss 4, pp 598–609, 2022
Copyright © 2022 by the American Thoracic Society
DOI: 10.34197/ats-scholar.2022-0029HT

limited to students, postgraduate trainees, and nurses. There are many techniques described to place USPIVs, including Seldinger technique, long axis, short axis, static, and dynamic guidance (11, 12). The published literature on teaching USPIV placement often includes didactic lectures followed by practicing on models, although these studies focus on teaching medical students, technicians, nurses, or emergency medical services workers and are 2–8 hours in length (13–16). Many training programs do not have the bandwidth or experienced faculty to teach this skill in a workshop format to learners in the intensive care unit (ICU) and on general medicine floors. We present a method of teaching USPIVs that was developed by one of the authors' experiences teaching residents, hospitalists, and advanced practice providers, while building, directing, and then handing off a bedside procedure team (17). Our method has been used across multiple institutions for many levels of learners, providing a broadly applicable approach in a setting where mastery through simulation may not be feasible.

WHAT IS THE SETTING?

The teaching is most commonly performed in the inpatient setting in the emergency department, medical floors, or ICU, typically after failed attempts by members of the healthcare team using conventional IV placement methods.

WHAT IS THE APPROACH?

Given the limitations in time and resources for teaching this core skill, we use a “just in time training” (JITT) strategy to efficiently prepare the learner for the combined educational and procedural experience they are about to undertake. Procedural JITT facilitates practice of a skill immediately before performance of it in a live clinical encounter to improve patient safety and trainee learning and confidence. As investigators discovered in

a qualitative analysis of JITT for pediatric lumbar puncture training, learners found JITT to be useful for reviewing anatomic landmarks, rehearsing steps, asking questions, and troubleshooting based on prior experience or areas of misunderstanding (18). The nature of this on-demand teaching ensures its relevance to the trainee's immediate learning needs and retention, given the temporal proximity of the learning to the actual skills performance.

Our approach mirrors the “briefing, intraoperative teaching, debriefing” model, a practical way to teach this relatively low-stakes procedure (19). As part of briefing, we start by asking the learner about quantitative experience and comfort level, recognizing the limitations of self-assessment. This encouraged self-reflection and assessment is known as metacognition; it has been shown to improve learning and skills retention and has been used successfully in the operating room setting (19–21). This conversation frames the discussion, allowing for more advanced troubleshooting techniques to be highlighted if the level of skill warrants. We set the expectation by telling the learner that we are going to explain the technique as if they have never placed an USPIV before, ensuring that the approach is clear when we move to the bedside. We highlight that although low stakes procedurally, USPIV placement is a skill that requires significant experience to master and that we will be giving real-time feedback and instruction at the bedside to support them through a successful IV insertion to encourage an open dialogue between the learner and instructor. This establishes the learning environment and psychologic safety, which has been described in many educational settings to be beneficial for learning (22–24). Specifically, the psychological safety that we aim to establish emphasizes the trainee's ability to fully engage in the learning experience without a sense of needing to preserve

their image or be concerned about judgement. Because the procedure is relatively low risk and the learning is happening in real time, it is crucial to establish an environment where the learner can ask questions openly, disclose weaknesses, try earnestly, make mistakes, and be primed to receive honest constructive feedback from a supportive educator. This allows them to focus on deeply learning a skill rather than worrying about meeting certain expectations or being judged (25).

During the briefing, we conduct a mini-simulation, allowing the trainee to practice surveying vascular anatomy and gain familiarity with the IV apparatus. We use mental imagery, cued by the description of the technique and supported by explanations and pictures of what the learner should expect to see based on where the ultrasound beam is directed; this has been described in surgical literature and is associated with improved confidence and technical skill in some studies (26–31).

During the intraoperative teaching section, we support the learner as they perform what they just simulated and imagined. We use cueing to highlight the key steps, facilitating real-time deliberate practice by the trainee. This component of the training is a form of guided discovery, where an expert provides verbal and even hands-on guidance during the learning experience (32). The actual procedural performance is a live form of deliberate practice education, where the learner is attentively guided to focus on repetitively practicing very specific aspects of a skill or procedure; the educator provides rigorous assessment and real-time feedback to the trainee that informs their repeated tries under direct supervision. Namely, during this USPIV procedure, dynamic needle tip positioning (DNTP) is the key skill that the learner is made to concentrate on, hone, and repeat throughout

the procedure to both successfully place the IV and demonstrate skill development as assessed by the supervisor (33).

In the debriefing portion, we focus on positive feedback to reinforce what was done correctly and encourage self-reflection to engage the learner on areas of improvement (34). Effective debriefing should result in the learner formulating objectives to guide future practice that can then be assessed during their next USPIV encounter. By allowing the learner to restate their main take-away lessons from the experience, we harness the learner's metacognition and obtain a commitment for their further practice. The educator also uses this opportunity to reinforce the most salient learning objectives and to ensure the learner understands how to correct any errors made during the procedure.

WHAT IS THE CONTENT?

Technique

Before the bedside. After gathering supplies (including extra IV catheters; see Figure E1 in the data supplement for suggested list), we orient the trainee to the ultrasound machine. This entails observing the trainee holding the linear array probe, ensuring they are anchoring their hand to assist with stabilization. With less-experienced trainees, or those who do not anchor their hand while holding the ultrasound probe, we find the “afternoon tea technique” to be an effective way to communicate how to lightly grip the probe while anchoring on the patient with their fifth digit (Figure 1) (35). We then have the trainee apply a tourniquet to the instructor's upper arm and survey the vessels. Scanning the vessels on the instructor's arm allows for reiteration of the ultrasound grip technique, confirms the trainee can identify relevant anatomy, helps the trainee practice orienting the




Figure 1. Ultrasound probe grip technique. The image on the left labeled “Wrong” displays the ultrasound probe being held away from the patient’s arm and without an anchoring point against the patient. The resultant display on the ultrasound screen is more likely to be like “probe heavy” or “probe moderate,” where the circular vessel is deformed into an oval as a result of the pressure applied by the probe. The image labeled “Correct” shows the probe being held close to the patient’s arm with the operator’s fifth digit anchored proximally on the arm, allowing stabilization and preventing excessive pressure from being applied, resulting in the ultrasound display labeled “probe light.”

probe perpendicular to the course of a vessel while performing serial compression, and allows the optimal gain and depth to be set on the machine.

We then move to practicing holding the IV insertion apparatus. The specific technique requires adjustment based on available catheters. Our hospital currently carries the Introcan Safety 64 mm 18- and 20-gauge catheters, which allows the operator to hold the catheter between their index and middle finger, using their thumb to help control the angle of the needle (Figure 2). Insyte BD IV catheters (48 mm, 18 and 20 gauge) are another common style, requiring alternative hand positioning (Figure 2). Having an extra catheter in the supplies allows the trainee to practice holding and manipulating the catheter before going to the bedside.

At this point, we review the concept of DNTP and highlight how to confirm proper needle position on the ultrasound

display (11). We explain that DNTP is a short-axis technique in which the probe is moved just past the tip of the needle and then held in place while the needle is then inserted until it enters the plane of the ultrasound beam and appears on the screen as a hyperechoic dot. These movements are repeated to “walk” the needle tip down to the vessel. We reference Figure 3 to show the appearance of a crescent-shaped vein representing the anterior vessel wall being tented in by the needle and an “empty” anechoic vessel a few millimeters proximal to confirm the location of the tip of the needle.

The final step before transitioning to the bedside is a broad overview of the specific steps for placement, with more details provided at the bedside in real time or in the debrief after the procedure. We review the following script verbally:

After vessel selection, the needle is inserted at the center of the ultrasound probe, nearly touching




Figure 2. How to hold intravenous (IV) insertion device. (A) The IV catheter being held between the index and middle finger, with the thumb placed on the back to aid in angulation. The fifth digit is extended and used to provide stabilization during IV insertion. (B) Another style of IV catheter in which the grip positioning requires the catheter to be held between the middle finger and thumb in a flexed position, with the index finger placed on the top of the catheter for additional stabilization. The fifth digit is extended and used to provide stabilization during IV insertion.

the edge, at a steep angle, and DNTP is used to guide the needle to the vessel. Once the needle is positioned in the center of the vessel, the angle of the needle is decreased to parallel the course of the vein, using the needle tip as the pivot point, and then DNTP is resumed until the needle is inserted entirely, or at a minimum, 2–3 cm.

We particularly stress that the operator should not be looking for a flash of blood as an indication the IV is in the vessel and

that we will use the ultrasound to ensure the IV is properly placed. At this point, questions are answered, and we proceed to the bedside.

At the bedside.

1. **Vascular survey and target vessel selection.** The learner applies the tourniquet and surveys the vessels and surrounding structures. We cue the operator to track the proposed target




Figure 3. Confirming vessel wall tenting. (A) The vessel is being deformed by the catheter and needle. This creates a crescent-shaped anechoic lumen. (B) It can be confirmed that it is the tip of the needle that is tenting in the vessel wall by fanning the ultrasound just proximal to the needle tip, where the vessel resumes its round shape. Note that in B, the vessel is oval shaped, suggesting the operator is applying too much pressure with the probe and is compressing the vessel. The red dotted lines in the center image correlate with the path of the ultrasound beam in A and B.

vessel with serial compression every centimeter to confirm there are no venous clots that may preclude successful cannulation. Ideal vessels for placement include those <1.6 cm below the skin, located medial or lateral to surrounding arteries and nerves, and >3 mm in diameter (36). Once a target is chosen, we have the operator clean the skin and the ultrasound probe.

2. Long IV insertion with steep approach and dynamic guidance.

The operator first centers the vessel on the ultrasound screen and demonstrates with proximal and distal fanning that the probe is perpendicular to the course of the vessel. We then remind them to insert the needle at a 60- to 80-degree angle; if the vessel is <5 mm deep, we tell them to insert the IV at a 30- to 45-degree angle to avoid puncturing the vessel or injuring surrounding structures while passing through the skin. We are deliberate about giving feedback on the direction of gaze throughout this step: the operator's eyes should be initially focused on the screen until the vessel is centered and the probe is stabilized, then eyes shift to the ultrasound probe to watch the needle pass through the skin, then eyes shift back to the ultrasound screen to find the needle tip. The operator then uses DNTP to follow the needle tip to

the target vessel. As the supervisors, we provide active feedback throughout this process on the size of the movements made by the operator and ensure they do not inadvertently decrease the angle of the needle before reaching the vessel.

3. Tent in vessel wall and lower angle of needle to get parallel to the blood vessel. Once the target vessel is reached, we instruct the operator to insert the needle until it appears to be at the center of the vessel, decrease the angle using the needle tip as the pivot point to parallel the course of the vessel, and then continue inserting using DNTP (Figure 4). To confirm the needle is tenting in the vessel wall, we have the operator fan the ultrasound just proximal to where the tip of the needle is located and then back to the needle tip (Figure 3). This gives the supervisor confidence that the operator has the needle inserted at the intended depth.

4. Anterior wall puncture. It is critical for the operator to maintain insertion pressure on the needle while dynamically tracking to prevent the catheter from rebounding before it punctures the anterior wall of the vessel. The supervisor should take note of how much of the IV catheter has been inserted into the skin. If there is consistent observation of a




Figure 4. Summary of ultrasound intravenous insertion. (A) The needle is inserted into the center of the vessel at a steep angle. Note the catheter or needle tip may not puncture the vessel wall. (B) The angle of the needle is lowered to parallel the course of the vessel while maintaining the tip of the needle in the center of the vessel. (C) The needle is inserted further into the vessel. Note the tip of the needle may pierce the vessel wall, but the catheter is not within the vessel yet, and attempting to thread the catheter at this point would likely result in the catheter threading into the subcutaneous tissue. (D) As the needle is inserted farther, the needle and catheter pierce the vessel wall.

similar length of exposed catheter, we instruct the operator to anchor the IV apparatus against the skin before each advancement of the ultrasound. As the IV is inserted, a tactile “pop” may be appreciated by the operator, often occurring between 1 and 2 cm after initial contact with the vessel wall (Figure 4).

5. Continue DNTP until IV is mostly or entirely inserted before attempting to thread catheter. We advocate tracking of the needle tip through the lumen of the vessel until the catheter is fully inserted, or the operator can ensure that there are at least 2–3 cm of catheter in the vessel. If the needle tip is inserted off-center, the operator is directed to ultrasound over the needle tip and slowly pivot the needle until the hyperechoic dot is in the center of the vessel before continuing dynamic tracking. Repeating this process throughout insertion allows the operator to gain appreciation for the magnitude of motion needed to reposition the needle. Once fully inserted, the ultrasound probe is placed aside, and the ultrasound hand is used to thread the remaining IV catheter into the blood vessel and to stabilize the catheter as the needle is removed.

6. Confirm placement using flush test. To confirm placement, in addition to imaging the catheter in short and long axis, we attach a saline flush, image proximal to the tip of the catheter, provide gentle compression to ensure the structure is venous, and then briskly flush 10 ml saline. The vessel will expand or become bright as the saline traverses the ultrasound beam, confirming venous placement (37–40).

Post-procedure debrief. Regardless of the success, we believe debriefing allows for technique reinforcement. Feedback post-procedure has consistently been

demonstrated to improve effectiveness of learning (19, 34, 41, 42). We ask the operator what went well and what could be done differently next time, or areas for improvement, which helps gauge degree of insight to frame the subsequent discussion.

WHAT CAN BE CHALLENGING?

Here are two commonly experienced challenges encountered and how we approach ongoing skill development:

1. The learner is having trouble with DNTP. DNTP results in the needle alternating between being in and out of plane. There is a tendency for learners not to want to “lose” the tip of the needle and instead maintain the ultrasound beam on what they believe is the needle tip throughout the insertion process but in fact is the needle shaft. We remind operators that if they visualize the needle but advance it before repositioning the ultrasound probe, the true tip of the needle will lie beyond the path of the ultrasound beam. This concept is reinforced during the post-procedure debrief by holding up the probe and using a pen to act as the needle, making it clearer that the tip is being moved past the plane of the ultrasound beam.

If the learner is moving the ultrasound too far with each movement, we will often hold the top of the ultrasound probe and deliberately fan the ultrasound while verbally calling out what is seen on the ultrasound screen. This allows the learner to gain an appreciation for the magnitude of each micromovement needed to move the needle tip in and out of plane without “taking over” the procedure. We review Figure 5 in the post-procedure debrief to reinforce that the tip of the needle is pin-point in size and that micromovements should be used.




Figure 5. Images comparing ultrasound appearance of (A and B) needle shaft to (C) needle tip. The hyperechoic structure in the center of the vessel is the intravenous catheter insertion device. The probe location and representative image are shown below. (A) The posterior wall of the vessel is interrupted because of posterior acoustic shadowing, suggesting the probe is positioned over the shaft of the needle, not the tip. (B) The needle is very bright and larger than how you would expect the tip to appear. The probe is positioned over the bevel of the needle, not directly at the tip. (C) Visualizing of this pinpoint structure is the tip of the needle.

2. The learner expresses they are unable to see or find their needle.

When the needle tip is within the subcutaneous tissue, it can be challenging to visualize the tip itself, because it is hyperechoic and may appear like surrounding tissue. To find the needle, we encourage the trainee to gently bounce the needle without advancing it and slowly slide the ultrasound probe from the point the needle enters the skin toward the tip of the needle until the tissue is no longer shifting, which is often just past the tip of the needle (Video E1). In our experience, unless the needle is passing through muscle, this maneuver is rarely painful. During this maneuver, the supervisor should point to the specific location on the ultrasound screen where they are seeing movement, allowing the trainee to begin to appreciate the subtle changes associated with needle movement. During the insertion process there is a tendency for inexperienced learners to

pick the probe entirely off the skin. We discourage this for two reasons: 1) when they pick the probe up, the tissue moves, which can alter the position of the needle relative to the target vessel; and 2) slow, deliberate sliding and fanning of the probe are necessary for mastery of DNTP. Thus, leaving the probe in contact with the skin allows the operator to gain more experience over the course of a single procedure attempt.

Below are a series of questions that are often asked by learners, with suggested answers and explanations.

The Patient Moves When I Try to Puncture Their Skin. How Can This Be Minimized?

We often use ice placed over the target vessel 1–2 minutes before sterile preparation or sterile ethyl chloride spray immediately before venipuncture to desensitize the superficial skin layer. Although counterintuitive to the heat typically applied to

superficial veins to induce vasodilation, we find most deep vessels will not constrict significantly, and even in those that do the decrease in pain experienced results in less movement during insertion, although this has not been studied.

Why Not Just Use Pythagorean Theorem to Determine Where to Insert the Needle?

Although the Pythagorean theorem method has been described, we often observe errors in location of needle insertion (usually too far back from the center of the probe), resulting in the needle reaching the vessel before it crosses the ultrasound beam (43). In addition, the Pythagorean theorem relies on a 45-degree approach; we recommend taking a steeper approach to minimize catheter length in the subcutaneous tissue (44, 45).

Why Should I Not Use the “Flash” of Blood (Blood Visualized within the IV Apparatus) as an Indication I Can Thread the Catheter?

Blood within the IV often indicates the needle tip within the vessel but not the catheter itself, and attempting to thread the catheter off at this point often results in pushing the needle out of the vein, with catheter insertion into the surrounding subcutaneous tissue rather than the lumen of the vein (Figure 4). We find that learners have greater success when not observing for a flash.

Why Should I Continue to Advance the Catheter in the Vein after I Puncture the Anterior Wall?

Once the needle has punctured the anterior wall of the vessel, continuing to insert the needle into the vein helps in two ways:

- 1) When you remove the ultrasound probe from the skin, the subcutaneous tissue is no longer compressed and can result in the needle dislodging from the vessel as it shifts

back to its resting position. The farther the needle is inserted into the vein beyond the puncture point, the more it is “speared” and held in place when the probe is removed. 2) The farther the needle is inserted in the vein, the less likely it is to slide out of the vessel if the needle is inadvertently, even minimally, retracted before the IV is threaded all the way in.

The needle tip is being followed within the lumen of the vessel, so it is usually not uncomfortable for patients while advancing farther. In addition to the described benefits, it allows more deliberate practice with DNTP during a single procedure attempt.

How Many USPIVs Will I Need to Place before I Am Competent?

The available literature, which focuses mostly on nurses and emergency medical services personnel, suggests four attempts for a success rate of 70% and 15–26 attempts for success rate of 88% (13).

WHY IS THIS THE APPROACH?

We present our method of teaching USPIV placement at the bedside, using a short-axis dynamic ultrasound technique, with a particular focus on a steep initial approach and advancing the needle and catheter device almost entirely into the vessel before attempting to thread the catheter. This variation combines principles of the JITT approach with a three-step briefing, intra-procedural teaching, debriefing model for procedural skills teaching (46, 47). Briefly practicing components, particularly ultrasound and catheter holding technique, before proceeding to the bedside creates the opportunity for technique reinforcement at the bedside, while allowing the learner to focus on more-advanced components of the skill set required to independently place USPIVs. The insertion technique itself

creates a similar opportunity for deliberate practice of the micro-skill surrounding needle localization and advancement, maximizing the learning potential of each experience. Post-procedure debriefing allows revisiting teaching points that were delivered in real time and the ability to demonstrate alternative approaches or techniques that may be used during future USPIV placements.

Success using this method of instruction has been demonstrated across multiple institutions in both the ICU population and hospital ward patients in whom landmark

IV placement has been unsuccessful. This teaching approach promotes careful attention to needle location and fine motor ultrasound skill and can be accomplished in real time at the bedside.

Acknowledgment:

Evan Makkas completed the illustration in Figures 3 and 4. Dr. Brian McNichols helped develop the insertion technique described in the paper.

Author disclosures are available with the text of this article at www.atsjournals.org.

REFERENCES

1. Maki DG, Kluger DM, Crnich CJ. The risk of bloodstream infection in adults with different intravascular devices: a systematic review of 200 published prospective studies. *Mayo Clin Proc* 2006; 81:1159–1171.
2. Au A, Rotte M, Gryzbowski R, Ku B, Fields J. Decrease in central venous catheter placement and complications due to utilization of ultrasound-guided peripheral intravenous catheters [abstract]. *Ann Emerg Med* 2011;58:S230.
3. Gorski L, Hadaway L, Hagle ME, McGoldrick M, Orr M, Doellman D. Supplement to infusion therapy standards of practice. *J Infus Nurs* 2016;39:S51–S55.
4. Bridey C, Thilly N, Lefevre T, Maire-Richard A, Morel M, Levy B, *et al*. Ultrasound-guided versus landmark approach for peripheral intravenous access by critical care nurses: a randomised controlled study. *BMJ Open* 2018;8:e020220.
5. Gregg SC, Murthi SB, Sisley AC, Stein DM, Scalea TM. Ultrasound-guided peripheral intravenous access in the intensive care unit. *J Crit Care* 2010;25:514–519.
6. Galen BT, Southern WN. Ultrasound-guided peripheral intravenous catheters to reduce central venous catheter use on the inpatient medical ward. *Qual Manag Health Care* 2018;27:30–32.
7. Schoenfeld E, Shokoohi H, Boniface K. Ultrasound-guided peripheral intravenous access in the emergency department: patient-centered survey. *West J Emerg Med* 2011;12:475–477.
8. Costantino TG, Parikh AK, Satz WA, Fojtik JP. Ultrasonography-guided peripheral intravenous access versus traditional approaches in patients with difficult intravenous access. *Ann Emerg Med* 2005;46:456–461.
9. Cardenas-Garcia J, Schaub KF, Belchikov YG, Narasimhan M, Koenig SJ, Mayo PH. Safety of peripheral intravenous administration of vasoactive medication. *J Hosp Med* 2015;10:581–585.
10. Groetzinger LM, Williams J, Svec S, Donahoe MP, Lamberty PE, Barbash IJ. Peripherally infused norepinephrine to avoid central venous catheter placement in a medical intensive care unit: a pilot study. *Ann Pharmacother* 2022;56:773–781.
11. Clemmesen L, Knudsen L, Sloth E, Bendtsen T. Dynamic needle tip positioning: ultrasound guidance for peripheral vascular access. A randomized, controlled and blinded study in phantoms performed by ultrasound novices. *Ultraschall Med* 2012;33:E321–E325.

12. Sandhu NPS, Sidhu DS. Mid-arm approach to basilic and cephalic vein cannulation using ultrasound guidance. *Br J Anaesth* 2004;93:292–294.
13. Stolz LA, Cappa AR, Minckler MR, Stolz U, Wyatt RG, Binger CW, *et al.* Prospective evaluation of the learning curve for ultrasound-guided peripheral intravenous catheter placement. *J Vasc Access* 2016;17:366–370.
14. Gardecki J, Hughes LP, Zakaria S, Lewiss RE, Goodsell K, Risler Z, *et al.* Use of the color Doppler twinkle artifact for teaching ultrasound guided peripheral vascular access. *J Vasc Access* 2021;22:692–696.
15. Oliveira L, Lawrence M. Ultrasound-guided peripheral intravenous access program for emergency physicians, nurses, and corpsmen (technicians) at a military hospital. *Mil Med* 2016;181:272–276.
16. Schoenfeld E, Boniface K, Shokoohi H. ED technicians can successfully place ultrasound-guided intravenous catheters in patients with poor vascular access. *Am J Emerg Med* 2011;29:496–501.
17. Gorgone M, McNichols B, Lang VJ, Novak W, O'Connor AB. The procedure coordinator: a resident-driven initiative to increase opportunity for inpatient procedures. *J Grad Med Educ* 2018;10:583–586.
18. Kamdar G, Kessler DO, Tilt L, Srivastava G, Khanna K, Chang TP, *et al.* Qualitative evaluation of just-in-time simulation-based learning: the learners' perspective. *Simul Healthc* 2013;8:43–48.
19. Roberts NK, Williams RG, Kim MJ, Dunnington GL. The briefing, intraoperative teaching, debriefing model for teaching in the operating room. *J Am Coll Surg* 2009;208:299–303.
20. Hong WH, Vadivelu J, Daniel EGS, Sim JH. Thinking about thinking: changes in first-year medical students' metacognition and its relation to performance. *Med Educ Online* 2015;20:27561.
21. National Research Council. How people learn: brain, mind, experience, and school: Expanded edition. 2000; Washington, DC: The National Academies Press.
22. Appelbaum NP, Santen SA, Aboff BM, Vega R, Munoz JL, Hemphill RR. Psychological safety and support: assessing resident perceptions of the clinical learning environment. *J Grad Med Educ* 2018;10:651–656.
23. Ramana Feeser V, Zemore Z, Appelbaum N, Santen SA, Moll J, Aboff B, *et al.* Analysis of the emergency medicine clinical learning environment. *AEM Educ Train* 2019;3:286–290.
24. Wakatsuki S, Tanaka P, Vinagre R, Marty A, Thomsen JL, Macario A. What makes for good anesthesia teaching by faculty in the operating room? The perspective of anesthesiology residents. *Cureus* 2018;10:e2563.
25. te Tsuei SH, Lee D, Ho C, Regehr G, Nimmon L. Exploring the construct of psychological safety in medical education. *Acad Med* 2019;94:S28–S35.
26. Geoffrion R, Gebhart J, Dooley Y, Bent A, Dandolu V, Meeks R, *et al.* The mind's scalpel in surgical education: a randomised controlled trial of mental imagery. *BJOG* 2012;119:1040–1048.
27. Komesu Y, Urwitz-Lane R, Ozel B, Lukban J, Kahn M, Muir T, *et al.* Does mental imagery prior to cystoscopy make a difference? A randomized controlled trial. *Am J Obstet Gynecol* 2009;201:218.e1–218.e9.
28. Souiki T, Benzagmout M, Alami B, Ibn Majdoub K, Toughrai I, Mazaz K, *et al.* Impact of mental imagery on enhancing surgical skills learning in novice's surgeons: a pilot study. *BMC Med Educ* 2021;21:545.
29. Lim G, Krohner RG, Metro DG, Rosario BL, Jeong JH, Sakai T. Low-fidelity haptic simulation versus mental imagery training for epidural anesthesia technical achievement in novice anesthesiology residents: a randomized comparative study. *Anesth Analg* 2016;122:1516–1523.

30. Saab SS, Bastek J, Dayaratna S, Hutton E, Salva CR. Development and validation of a mental practice tool for total abdominal hysterectomy. *J Surg Educ* 2017;74:216–221.
31. Arora S, Aggarwal R, Sevdalis N, Moran A, Sirimanna P, Kneebone R, *et al.* Development and validation of mental practice as a training strategy for laparoscopic surgery. *Surg Endosc* 2010;24:179–187.
32. Mayer RE. Should there be a three-strikes rule against pure discovery learning? The case for guided methods of instruction. *Am Psychol* 2004;59:14–19.
33. Ericsson KA. Deliberate practice and the acquisition and maintenance of expert performance in medicine and related domains. *Acad Med* 2004;79:S70–S81.
34. McSparron JI, Ricotta DN, Moskowitz A, Volpicelli FM, Roberts DH, Schwartzstein RM, *et al.* The PrOSTE: identifying key components of effective procedural teaching. *Ann Am Thorac Soc* 2015;12:230–234.
35. McMenamin L, Wolstenhulme S, Hunt M, Nuttall S, Weerasinghe A. Ultrasound probe grip: the afternoon tea technique. *J Intensive Care Soc* 2017;18:258–260.
36. Fields JM, Dean AJ, Todman RW, Au AK, Anderson KL, Ku BS, *et al.* The effect of vessel depth, diameter, and location on ultrasound-guided peripheral intravenous catheter longevity. *Am J Emerg Med* 2012;30:1134–1140.
37. Weekes AJ, Johnson DA, Keller SM, Efuno B, Carey C, Rozario NL, *et al.* Central vascular catheter placement evaluation using saline flush and bedside echocardiography. *Acad Emerg Med* 2014;21:65–72.
38. Liu YT, Bahl A. Evaluation of proper above-the-diaphragm central venous catheter placement: the saline flush test. *Am J Emerg Med* 2011;29:842.e1–842.e3.
39. Gekle R, Dubensky L, Haddad S, Bramante R, Cirilli A, Catlin T, *et al.* Saline flush test: can bedside sonography replace conventional radiography for confirmation of above-the-diaphragm central venous catheter placement? *J Ultrasound Med* 2015;34:1295–1299.
40. Horowitz R, Gossett JG, Bailitz J, Wax D, Pierce MC. The FLUSH study: flush the line and ultrasound the heart: ultrasonographic confirmation of central femoral venous line placement. *Ann Emerg Med* 2014;63:678–683.
41. Kelm DJ, Ridgeway JL, Ratelle JT, Sawatsky AP, Halvorsen AJ, Niven AS, *et al.* Characteristics of effective teachers of invasive bedside procedures: a multi-institutional qualitative study. *Chest* 2020; 158:2047–2057.
42. Burgess A, van Diggele C, Roberts C, Mellis C. Tips for teaching procedural skills. *BMC Med Educ* 2020;20:458.
43. Piton G, Capellier G, Winiszewski H. Ultrasound-guided vessel puncture: calling for Pythagoras' help. *Crit Care* 2018;22:292.
44. Pandurangadu AV, Tucker J, Brackney AR, Bahl A. Ultrasound-guided intravenous catheter survival impacted by amount of catheter residing in the vein. *Emerg Med J* 2018;35:550–555.
45. Bahl A, Hang B, Brackney A, Joseph S, Karabon P, Mohammad A, *et al.* Standard long IV catheters versus extended dwell catheters: a randomized comparison of ultrasound-guided catheter survival. *Am J Emerg Med* 2019;37:715–721.
46. Peyton JWR. Teaching & learning in medical practice. Heronsgate Rickmansworth, Herts: Manticore Europe Ltd., 1998.
47. Giacomino K, Caliesch R, Sattelmayer KM. The effectiveness of the Peyton's 4-step teaching approach on skill acquisition of procedures in health professions education: a systematic review and meta-analysis with integrated meta-regression. *PeerJ* 2020;8:e10129.