




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Meal preparation and consumption before and during the COVID-19 pandemic: The relationship with cooking skills of Brazilian university students

Talissa Dezanetti ^{a,*}, Ricardo Teixeira Quinaud ^b, Martin Caraher ^c, Manuela Mika Jomori ^a

^a Department of Nutrition, Federal University of Santa Catarina (UFSC). Health Sciences Centre. Reitor João David Ferreira Lima Campus, 88040-900, Florianópolis, Santa Catarina, Brazil

^b Physical Education Postgraduate Programme, Federal University of Santa Catarina (UFSC). Health Sciences Centre. Reitor João David Ferreira Lima Campus, 88040-900, Florianópolis, Santa Catarina, Brazil

^c Centre for Food Policy, City, University of London, Northampton Square, EC1V 0HB, London, UK

ARTICLE INFO

Keywords:

Meal preparation
Food consumption
Quarantine
Social distance
Food skills
Culinary skills

ABSTRACT

Objective: This study aimed to analyze meal preparation and the place of its consumption by university students before and during the COVID-19 pandemic, according to their individual characteristics and cooking skills. **Methods:** Data were collected with 1919 Brazilian university students using the Brazilian Cooking Skills Questionnaire (BCSQ) and questions about gender, living arrangement, knowing how to cook and location of meal consumption. Bayesian multilevel ordinal regression models were used to estimate the probabilities of meal preparation and local location of its consumption by Brazilian university students before and during the COVID-19 pandemic, according to their individual characteristics and cooking skills. **Results:** Most of participants showed reported a high level of cooking skills (70.7%). Also, they reported a decrease in ordering delivery of take-away food and eating fast-food, while increasing homemade meals with fresh ingredients, ultra-processed food or both during the pandemic compared to the period before the quarantine. Lastly, they increased their cooking frequency during the pandemic, regardless of their cooking skills level. However, the group with high cooking skills had a greater increase in the use of fresh ingredients. **Conclusions:** The study showed that the pandemic was an opportunity for university students to increase their cooking skills and frequency of cooking at home with a variety of foods. Particular attention should be given to those with low cooking skills and that use ultra-processed foods, promoting interventions to disseminate information and with behavior change components to teach about healthy cooking to increase cooking knowledge and skills, focusing on preparation of fresh ingredients.

1. Introduction

Cooking skills are defined as a combination of the confidence, attitude, and knowledge of individuals in performing cooking tasks. These tasks involve planning menus, shopping, and preparing unprocessed (fresh foods), minimally processed (e.g., polish in rice, fermentation in cheese and yogurt), processed (e.g., canned foods with added salt or sugar, industrialized bread), or ultra-processed foods (Jomori et al.,

2018). Ultra-processing is related to techniques which transform the original food, adding excessive amount of preservatives, salt, sugar and fat, to help with preservation, storage and distribution, generally, making the food ready to eat (Dietary Guidelines for the Brazilian, 2014). High levels of the use cooking skills are associated with an increase in the consumption of fresh fruits and vegetables, which may be related to healthy eating habits among adults and adolescents (Caraher et al., 1999; Laska et al., 2012; Utter et al., 2018; Wolfson & Bleich,

Abbreviations: BCSQ, Brazilian Cooking Skills Questionnaire; CA, Cooking Attitude scale; CB, Cooking Behavior scale; SEC, Cooking Self-Efficacy scale; SEFVS, Self-Efficacy for Using Fruits, Vegetables, and Seasonings scale; SEPC, Produce Consumption Self-Efficacy scale; SETC, Self-efficacy in Using Basic Cooking Techniques scale; UFSC, Federal University of Sana Catarina.

* Corresponding author.

E-mail addresses: talissadezanetti@hotmail.com (T. Dezanetti), ricardoquinaud@gmail.com (R.T. Quinaud), m.caraher@city.ac.uk (M. Caraher), manuela.jomori@ufsc.br (M.M. Jomori).

<https://doi.org/10.1016/j.appet.2022.106036>

Received 9 November 2021; Received in revised form 27 March 2022; Accepted 2 April 2022

Available online 14 April 2022

0195-6663/© 2022 Elsevier Ltd. All rights reserved.

2015).

Processes involved in individual meal preparation (planning, transportation, shopping, preparing, and cleanup) were described as being an 'overwhelming responsibility' (Murray et al., 2016) aside from other barriers to cook among university students, leading them to have low cooking skills and low meal preparation frequency (Bernardo et al., 2017; de Borba et al., 2021; Murray et al., 2016; Sprake et al., 2018; Wilson et al., 2017), low consumption of fruits and vegetables, and high intake of fast foods, snacks, deep-fried foods, refined grains, sweets, carbonated, and other sugar-sweetened beverages (Bernardo et al., 2017).

Low cooking skills and low frequency of meal preparation or behaviors are related to poor diets, with low consumption of fruits and vegetables, high consumption of ultra-processed foods, meals away from home, take-way and fast food, related to high energy-dense food intake in these populations (Bernardo et al., 2017; Bezerra et al., 2020; de Borba et al., 2021; Graham et al., 2013; Knol et al., 2019; Lavelle et al., 2016; Santos et al., 2015; Sprake et al., 2018; Utter et al., 2018; Vilela et al., 2014; Wilson et al., 2017). Additionally, factors related to greater access, convenience and consumption of ready-to-eat foods (most of them are ultra-processed foods) can be identified, such as difficulties in accessing healthy foods and fresh fruits and vegetables, lack of time, money, kitchen facilities, knowledge, and confidence to cook, and their living arrangements (Graham et al., 2013; Jones et al., 2014; Knol et al., 2019; Murray et al., 2016; Pulz et al., 2017; Sprake et al., 2018; Utter et al., 2018; Wilson et al., 2017).

Recommendations to promote healthy eating practices, encouraging meal preparation and the development of cooking skills are found in the United Kingdom (Tsouros et al., 1998; Dooris & Doherty, 2010; Community Food & Health Scotland, 2013), Northern Ireland (Food Standards Agency, 2012), Canada (Chenhall, 2010), in the United States of America (USDA, 2013) and in Brazil (Brasil, 2014). The Health Promotion Universities Network (REDUPS) that operate in collaboration with the Pan American Health Organization (OPAS/OMS) (REDUPS, 2013) has a commitment to health promotion in the university environment (Dooris & Doherty, 2010; Oliveira, 2017; REDUPS, 2013; Soares et al., 2015; Tsouros et al., 1998; WHO, 2015, pp. 23–25). In Brazil, the accreditation of universities at RIUPS is being implemented (Oliveira, 2017). One of the strategies to promote healthy eating practices among university students to achieve the Brazilian nutritional recommendations could be to develop or improve their cooking skills including preparing their own meals often (Bernardo et al., 2017; Wilson et al., 2017) or offer healthy foods at canteens and cafeterias (Pulz et al., 2017).

One of the concerns in promoting these policies is based on the barriers to cooking that students face ranging from access to food to the application of skills, as mentioned before (Jones et al., 2014; Wilson et al., 2017). Due to the COVID-19 pandemic, a suspension of classes in person and academic activities at universities have been put in place. As a result, university students have been affected by food insecurity, probably because many of them have lost their part-time jobs and have not been registered by student assistance programs, as well as the inaccessibility to university cafeterias that were closed during the pandemic. This required the planning, purchasing, and preparation of their own food, when they were the responsible to prepare it, or it is supposed somehow their involvement on food preparation at home (Gerritsen et al., 2020; Owens et al., 2020; Restrepo & Zeballos, 2020; Reyes-Olavarría et al., 2020).

In this context, long periods at home require knowledge and skills to plan and prepare meals or any culinary involvement. Young people tended to increase their consumption of ultra-processed foods and reduce their consumption of fresh foods during the pandemic (Jribi et al., 2020; Owens et al., 2020; Ruiz-Roso et al., 2020). On the other hand, purchases of ready-made meals, such as instant 'TV dinner', were reduced during the pandemic by Italian individuals, although ultra-processed food's consumption (e.g., chocolate, ice-cream, desserts,

and salty snacks) has been increased (Scarmozzino & Visioli, 2020). Moreover, studies have shown that working and studying from home can increase the frequency of home cooking and homemade-recipes, reduce the food waste, and encourage attention to food behaviors (Deschasaux-Tanguy et al., 2020; Di Renzo et al., 2020; Gerritsen et al., 2020; Restrepo & Zeballos, 2020; Reyes-Olavarría et al., 2020; United Nations, 2015).

It is important to consider social and individual characteristics that influence the cooking skills and meal preparation. A systematic review involving 38 papers about the relationship of social determinants and home cooking showed that the main determinants included gender (women and girls are more likely to be involved in cooking than men and boys), greater available time to cook and employment (those who have restrictions in time or working more tend to cook less than who have greater available time), close personal relationships (those who live with a partner or children were more likely to cook) and culture and ethnic background (Mills et al., 2017). Short and Gatley make the point that wider structural and economic determinants in the food environment influence both the acquisition and use of cooking skills (Gatley et al., 2014; Short, 2006).

To improve university students' diets, it is necessary to increase their consumption of fresh foods, such as fruits and vegetables, rather than ultra-processed foods (Graham et al., 2013; Knol et al., 2019; Larson et al., 2006; Laska et al., 2012; Sprake et al., 2018; Utter et al., 2018; Wilson et al., 2017). However, it is important to consider the difficulty in preparing fresh foods, as they require the use of basic pre-preparation cooking techniques to scratch cook, as well as individual factors (i.g. attitude, confidence and knowledge to cook) (Jomori et al., 2018).

Researchers have identified the need for increases culinary skills and their usage among young people to improve their diet during this period when they are at home for long periods of time (Jribi et al., 2020; Owens et al., 2020; Ribeiro et al., 2020; Ruiz-Roso et al., 2020). However, to our knowledge, none of these studies have evaluated meal consumption and preparation related to their level of culinary skills and healthy eating practices during the pandemic. Therefore, it is an opportunity to identify the students' level of culinary skills and relate them to their characteristics and the food environment during the COVID-19 pandemic (Fulkeron et al., 2019; Owens et al., 2020; Rathi et al., 2018; Ribeiro et al., 2020; Ruiz-Roso et al., 2020).

In this context, this study aimed to estimate the probabilities of meal preparation and the place of consumption by university students before and during the COVID-19 pandemic, according to their individual characteristics and cooking skills.

2. Methods

2.1. Study design and participants

This descriptive cross-sectional study was conducted from June to August 2020, with undergraduate students from the Federal University of Rio Grande do Sul (UFRGS) and Federal University of Santa Catarina (UFSC), Brazil. Based on the total number of students enrolled at UFSC (n = 30,000) and UFRGS (n = 31,000), a minimum of 830 and 731 students, respectively, were calculated to be required bearing in mind a possible 10% loss, 2.0 effect, and 5% random error (Jomori et al., 2017).

The eligibility criteria were the enrollment in any undergraduate course at this institution. Those enrolled in postgraduate courses or those who did not answer all the questions provided in the questionnaire were excluded from analysis.

This study was approved by the Human Research Ethics Committee of the Federal University of Santa Catarina (UFSC) and Federal University of Rio Grande do Sul (UFRGS) under the approval number 09427219.5.3001.0121. Informed consent was obtained online before data was collected and were assured that all data would be used only for research purposes.

2.2. Measurements

The Brazilian Cooking Skills and Healthy Eating Questionnaire (BCSQ) was used to assess cooking skill levels. The BCSQ is an adapted and validated questionnaire for the Brazilian population (Jomori et al., 2017, 2021, 2022). The questionnaire was shortened to 36 items and was distributed across seven factors, incorporating all items and structure from the U.S. version, as shown in supplementary material.

Cooking ability was derived from the Likert's ratings of Cooking Attitude (CA), Cooking Behavior (CB), Produce Consumption Self-Efficacy (SEPC), Cooking Self-Efficacy (SEC), and Using Fruits, Vegetables, and Seasonings Self-Efficacy (SEFVS) from Brazilian Cooking Skills and Healthy Eating Questionnaire scored between 1 and 5 points. The Availability and Accessibility of Fruits and Vegetables (AAFV) is dichotomous scale, scored from 0 to 8 points. The sum of these scales' scores was classified among low (20–43), medium (44–73) and high cooking skills (74–100). This classification is based on the theoretical arguments as opposed to the 'mathematical' ones, since the concept of cooking skills related to healthy eating aspects showed very complex associations (Wolfson & Bleich, 2015; Lavelle et al., 2016; Jomori et al., 2018). A team of experts was surveyed to validate the classification associated with Jomori et al. (2022)'s study. This classification considered the sensitivity of the questionnaire to the variability of the answers of participants, according to their demographic characteristics in the validation process in 2015 (Jomori et al., 2022), as well as the literature review (Jomori et al., 2018). Moreover, it was based on their difficulty to achieve higher levels of cooking skills, considering the questionnaire identifies barriers to cook by young adults. So, larger ranges for medium and high than low cooking skills (74–100) were established (Jomori et al., 2022).

Cooking knowledge was calculated based on responses where the correct answer was scored 1 point. A total score of six or higher was characterized as having high cooking knowledge (Jomori et al., 2022).

2.3. Recruitment and data collection

The participants were recruited between June and July 2020 through university e-mail distribution lists and social media related to undergraduate courses, containing messages with a link and a quick response code (QRcode) to easily access an online questionnaire.

The participants voluntarily completed the online questionnaire. The total number of answers and distribution per course was observed and controlled until the established sample size was achieved. For those courses that had no or few answers, the questionnaire was resent through e-mail asking to the course coordinators asking them to send the study details to their students in order to have a heterogeneous sample between the different study areas. When sufficient numbers of participants from these courses were achieved, data collection was stopped. The data collected included the following variables: gender (male or female); living arrangement (alone, colleagues, parents, or partner); frequency (never, 1 to 2 times a month, once a week, several times a week, or daily) of meals prepared with fresh ingredients (e.g. whole lettuce, meat); meals prepared with ultra-processed food (artificial seasonings such as chicken bouillon in tablets or granulated, canned corn, frozen lasagna and hamburger, etc); meals prepared with fresh ingredients combined with ultra-processed food; consumption (fast food or delivered food); self-reported cooking knowledge (yes or no); availability and accessibility of fruits and vegetables (low, medium, or high), cooking skills (low, medium, or high); and cooking knowledge (low or high).

2.4. Data analysis

Bayesian multilevel ordinal regression models were used to estimate the probabilities of meal preparation and local consumption by Brazilian university students before and during the COVID-19 pandemic,

according to their individual characteristics and cooking skills. Bayesian methods consider parameters as random variables. It estimates the probability distribution based on the data available and the prior distribution information that measures the uncertainty about parameters (McElreath, 2015). Taking into account the different sources of inferential uncertainty, Bayesian methods allow for combining the known information before seeing the data (i.e., the prior uncertainty concerning a parameter or hypothesis expressed as a probability distribution) with what is learned from the observed data (i.e., the likelihood of the data conditioned on the parameter or hypothesis) to update knowledge expressed as the posterior distribution (Lee & Wagenmakers, 2013; Kennedy & Gelman, 2021).

Bayesian data analysis reallocates credibility across the possibilities. It allows the combination of prior knowledge with the observed data, resulting in what is called posterior distribution (Kruschke & Liddell, 2018), as well as reducing bias and variance for posterior estimates (Gao et al., 2020). Additionally, the use of multilevel regression models advances over traditional regressions (i.e., considers data structure at just one level) by considering the nested data structure and the different sources of variation (i.e., within and between groups) (Gelman & Hill, 2007).

The use of ordinal models also lies on the structure of the questionnaire, which our dependent variables are peoples' responses on ordinal categories. Although these variables are not metric, researchers commonly analyze them as metric responses, which can cause misinterpretation of the results (Liddell & Kruschke, 2017). Additionally, Bayesian multilevel ordinal models has been recommended as a better option for analyzing rating, which allow for unequal distances between responses (Verissimo, 2021).

The responses about university students' food consumption and place of consumption of the main meal were estimated across time by gender (female and male), age group (<24 years old and >25 years old), self-reported cooking knowledge (yes or no), living arrangements (alone, colleagues, parents, or partner), cooking ability (low, medium, or high), and cooking knowledge (low or high). One question on the CA scale had a reversed score. Models were fitted using "time" (before and during COVID-19) as a fixed effect (population-level effect). To regularize the estimations, weakly informative prior distributions, normal prior (0, 10) for population-level effects, and normal priors (0, 1) for group-level effects were used. Additionally, to guarantee the Markov's convergence, two chains were run for 4000 iterations with a warm-up length of 1000 iterations. The analysis was performed using the BRMS package (Burkner, 2017) in R (R Core Team, 2018).

Considering there was no previous knowledge in the literature that could guide our understanding of how "time" (before and during COVID) could influence people's meal preparation and consumption, we set larger prior for population-level effects in order not to influence the results and lead to misinterpretation. In relation to prior from group-level effects, the variables used are well known by the literature (de Borja et al., 2021; Jomori et al., 2022; Mills et al., 2017; Wilson et al., 2017). However, we have used weakly informative prior to do not placing too much probability in any particular interval and to stabilize posteriors and regularize parameter estimate. Lastly, Bayesian methods require higher computing power than traditional (frequentists) methods and ordinal regressions, in the present case an ordinal cumulative model, require even a much higher computing power to perform the analysis without disturbing the model and not getting errors in the iterations. Thus, setting a larger prior for population-level effects leads to faster sampling, because priors can be vectorized as in this case, and improve model convergence.

It is important to note, when interpreting the results, that estimates are expressed in standard deviation units, which can be interpreted as standardized effect sizes. Additionally, reference categories are fixed as one and interpretations of standardized effect sizes are based on these categories. The literature recognizes that the interpretation of such effects is not immediately obvious. Thus, results are plotted to have a

natural metric interpretation (Burkner & Vuerre, 2019).

3. Results

A total of 1919 students were eligible for the present study out of the 2061 total students who responded to the online survey. Participants were aged 23.9 (± 6.8) years on average, and the majority were female (73.5%), living with parents (48.9%), and believed that they knew how to cook (92.0%). The demographic and cooking characteristics are presented in Table 1.

Table 2 shows the cooking skills and healthy eating characteristics, such as the level of availability of fruits and vegetables, cooking skills, and cooking knowledge. Most participants reported high availability of fruits and vegetables (73.0%), high levels of cooking skills (70.7%), and approximately half of them had high scores for cooking knowledge (50.6%).

Multilevel ordinal regression models were plotted (Figs. 1–5), and the estimates and confidence intervals are presented in Supplementary Table 1. The response categories are presented as 1 (never), 2 (1–2 times a month), 3 (once a week), 4 (several times a week), and 5 (daily). In relation to the consumption of fast food (Fig. 1), university students decreased their probability of consumption during the pandemic (standard deviation [SD] = -1.13 , 95% confidence interval [CI] = $[-1.21, -1.06]$) compared to consumption before the pandemic, with a high probability of responding “never” during the pandemic. We did not find substantial variation between groups for each variable at each time point for consumption of fast food. In Fig. 2 (delivered food), university students did not show substantial variation in their responses between time points ($SD = 0.11$, 95% CI = $[0.04, 0.18]$) or between groups, although there was a small increase during the pandemic.

Fig. 3 shows the use of fresh ingredients to cook before and during the pandemic. Overall, university students substantially increased the use of fresh ingredients during the pandemic ($SD = 0.83$, 95% CI = $[0.75, 0.90]$) compared to before the pandemic. Students with high cooking ability increased the probability of using fresh ingredients during the pandemic. The probability of answering “never” was almost zero among these participants whereas responding “daily” substantially increased ($SD = 1.50$, 95% CI = $[-0.61, 3.62]$). People with a medium level of cooking ability showed a substantial increase in the probability of using fresh ingredients “several times per week” and “daily” during the pandemic when compared with before the pandemic. Those with low cooking ability had a slightly increased probability of using fresh ingredients during the pandemic (1–2 times a month, once a week, several times a week, and daily), even though this increase was lower than in students with medium and high cooking abilities (Fig. 3). The probability variation of the response of “daily” increased for both groups that responded that they knew or did not know how to cook.

Regarding the use of ultra-processed foods (Fig. 4), university students increased the probability of using this type of food during the

Table 1
Demographic and cooking characteristics of the students from two Brazilian universities (n = 1919).

Variables	N	SD or %
Age (years)	23.9	6.8
Gender		
-Female	1410	73.5
-Male	509	26.5
Living Arrangement		
-Alone	348	18.1
-With Parents	939	48.9
-With Partner	248	12.9
-With Colleagues	384	20.0
Do you believe that you know how to cook?		
- Yes	1766	92.0
-No	153	8.0

SD: standard deviation.

Table 2
Level of availability of fruits and vegetables, cooking skills, and cooking knowledge of Brazilian university students (2020).

Measures	N	%
Availability and Accessibility of Fruits and Vegetables (AAFV)		
-Low	110	5.7
-Medium	408	21.3
-High	1401	73.0
Cooking skills		
-Low	14	0.7
-Medium	548	28.6
-High	1357	70.7
Cooking knowledge		
-Low	947	49.3
-High	972	50.6

pandemic ($SD = 0.37$, 95% CI = $[0.30, 0.43]$) when compared to before the pandemic. Students with low, medium, and high cooking skills had a decreased probability of responding “never” and an increased probability of responding “daily” ($SD = 1.49$, 95% CI = $[-0.19, 3.11]$). The probability of using combined fresh ingredients with ultra-processed foods (Fig. 5) substantially increased during the pandemic ($SD = 0.57$, 95% CI = $[0.50, 0.64]$) compared to before the pandemic. University students with high cooking ability presented a higher probability of responding “daily” ($SD = 1.28$, 95% CI = $[-0.57, 3.07]$) compared to the other groups. Additionally, the probability of answering “never” to this behavior decreased for all groups (low, medium, and high cooking skills). Participants who reported that they did or did not know how to cook had an increased probability of responding “daily” and a decreased probability of responding “never” although with no substantial variation between the two groups.

4. Discussion

This study assessed the characteristics of meal preparation and consumption by Brazilian university students before and during the COVID-19 pandemic according to their cooking skill level and individual characteristics. Overall, the sample showed higher availability and accessibility of fruits and vegetables at home, higher scores in their cooking skills, a decrease in the consumption of fast food and an increase in homemade meals made with fresh ingredients, ultra-processed food, or both during the pandemic compared to the period before the pandemic.

The high availability and accessibility of fruit and vegetable scores showed during the pandemic (Table 2) is related to the fact that living at home may have resulted in others taking responsibility for shopping and purchasing of foods. Thus, an access barrier to food supply may have been averted and account for the reported high availability. Studies have shown that high availability and accessibility of fruits and vegetables are consistently positively correlated with fruit and vegetable intake. Neumark-Sztainer et al. (2003) carried out a study with 3957 teenagers and found that the availability of fruits and vegetables at home was one of the factors most related to the consumption of these foods. A study by Kratt et al. (2000) investigated the availability of fruits and vegetables as a moderating variable for fruit and vegetable consumption relationships and how these relationships might change with varying levels of fruit and vegetable availability. The authors found that homes with more fruits and vegetables had a larger and stronger set of motivating factors for the consumption of these foods by parents and children compared to those with low availability. This suggests that participants of the present study not only have a high availability of fruits and vegetables, but also may have a high intake of fruits and vegetables once they also showed an increase in the probability of using fresh ingredients to cook (Fig. 3), independent of their cooking skill level.

Utter et al. showed in a longitudinal study carried out through a questionnaire with young adults (18–23 years, n = 1158), and with




Fig. 1. Consumption of fast food before and during COVID-19 according to gender (A), age (B), whether they know how to cook (C), living arrangements (D), cooking ability (E), and cooking knowledge (F). Response categories are presented as 1 (never), 2 (1–2 times a month), 3 (once a week), 4 (several times a week), and 5 (daily). 95% confidence intervals.

results from the same questionnaire applied a decade later (30–35 years), that a quarter of their sample had very adequate cooking skills as young adults (18–23 years). Utter and colleagues found that having cooking skills led these participants to better eating behaviors through greater involvement in cooking and a lower consumption of fast food 10 years later in adulthood (Utter et al., 2018). The present study sample

reported high scores for cooking skills during COVID-19 (Table 2), suggesting that, after the pandemic period, this group are more likely to experience long-term benefits, such as to prepare their meals with fresh ingredients and less to consume fast food.

On the other hand, approximately half of our sample achieved a high level of cooking knowledge (Table 2), and half of these recorded high




Fig. 2. Consumption of delivered food before and during COVID-19 in relation to gender (A), age (B), whether they know how to cook (C), living arrangements (D), cooking ability (E), and cooking knowledge (F). Response categories are presented as 1 (never), 2 (1–2 times a month), 3 (once a week), 4 (several times a week), and 5 (daily). Error bars indicate 95% confidence intervals.

frequency cooking practices. Cooking knowledge involves perceptual skills (judging flavors, combining and replacing ingredients with a minimum number of errors at the end of preparation, and adjusting the time to perform culinary tasks), conceptual skills (predicting results,

knowing how to adapt ingredients, planning menus, knowledge of culinary terms and techniques, and using appropriate equipment, utensils, and ingredients), and academic knowledge (knowledge about nutrition, food hygiene, and food trends). This knowledge is part of




Fig. 3. Use of fresh ingredients to cook before and during COVID-19 according to gender (A), age (B), whether they know how to cook (C), living arrangements (D), cooking ability (E), and cooking knowledge (F). Response categories are presented as 1 (never), 2 (1–2 times a month), 3 (once a week), 4 (several times a week), and 5 (daily). Error bars indicate 95% confidence intervals.




Fig. 4. Use of ultra-processed food to cook before and during COVID-19 according to gender (A), age (B), whether they know how to cook (C), living arrangements (D), cooking ability (E), and cooking knowledge (F). Response categories are presented as 1 (never), 2 (1–2 times a month), 3 (once a week), 4 (several times a week), and 5 (daily). Error bars indicate 95% confidence intervals.

person-centered cooking skills; therefore, they can have an impact on confidence, attitude, and individual cooking behavior (Jomori et al., 2018).

The lack of culinary knowledge may not necessarily influence the practice and food choices of the participants, but the lack of culinary knowledge can impact on confidence and culinary attitudes, leading to less healthy choices. Murray et al. (2016) carried out a focus group with

college students who did not live on campus, and thus were not part of the campus meal system, but lived independently of their parents (n = 24) and found no evidence that culinary knowledge at different levels had a direct impact on food choices and culinary practices of these students. Those living at home with others may be less likely to be the sole food preparers and may in fact benefit from the activities of others in the household.




Fig. 5. Use of fresh ingredients and ultra-processed food to cook before and during COVID-19 according to gender (A), age (B), whether they know how to cook (C), living arrangements (D), cooking ability (E), and cooking knowledge (F). Response categories are presented as 1 (never), 2 (1–2 times a month), 3 (once a week), 4 (several times a week), and 5 (daily). Error bars indicate 95% confidence intervals.

Long stay-at-home time during the COVID-19 pandemic may lead to a decrease in the probability of fast-food consumption by the participants of our study in all the variables analyzed, showing that it was independent of their level of cooking skills and knowledge. The closure of restaurants, social distancing by imposing safety standards, and fear of the disease could be the factors that justify the decrease in the consumption of fast-food and food delivered food (Figs. 1 and 2), as a

structural determinant. One study carried out in Brazil with people over 18 years old showed that people were more likely to go to restaurants where they could see safety standards were in place such as mask wearing, social distancing and eating in well ventilated places or outside during the pandemic (Piton Hakim et al., 2021). Likewise, Husain and Ashkanani (2020) carried out an online questionnaire with adults (18–73 years old) in Kuwait and found that 49% of their sample were

likely to consume fast food 1–2 times per week before the pandemic, while up to 82% reported not consuming fast food during the pandemic. [Błaszczuk-Bębenek et al. \(2020\)](#) also found a reduction on the daily servings of fast-food consumption in Polish adults during lockdown, where during the pandemic the percentage of participants who do not consume fast food daily increased to 41,7% from 26% before the pandemic.

On the other hand, the present study showed that the long stay-at-home period increased the probability of cooking homemade meals, as observed in other studies worldwide during the pandemic ([Deschasaux-Tanguy et al., 2020](#); [Di Renzo et al., 2020](#); [Gerritsen et al., 2020](#); [Reyes-Olavarria et al., 2020](#)). In Chile, [Reyes-Olavarria et al. \(2020\)](#) found that almost 60% of the adult participants in their study increased home cooking during the pandemic compared to before the pandemic. [Di Renzo et al. \(2020\)](#) showed that their Italian sample aged between 12 and 86 years had increased homemade recipes during the pandemic. In New Zealand, [Gerritsen et al. \(2020\)](#) showed that adult participants had increased home cooking and baking from scratch during quarantine. A study was conducted in France with 37,252 adults from a web-based cohort-filled lockdown-specific questionnaire in April–May 2020. They verified that 40% of the participants had increased home-made meals during the pandemic ([Deschasaux-Tanguy et al., 2020](#)). These studies concluded that even though the pandemic resulted in a small overall shift towards an unhealthy diet, they also created an opportunity for some people to improve their cooking and nutritional behaviors, beginning with home cooking practices, as also shown in the present study. Over a period of extended lockdown there are reports of changing culinary practices, so reports of increased cooking from scratch were verified in the early stages but as lockdown continues some of this initial impetus is lost ([Cummins et al., 2020](#)).

Some of this is also influenced by the fact that although eating out may have been difficult due to restrictions the catering industry adapted quickly to use online ordering and offer home delivery of fast food. A study by [Scarmozzino and Visioli \(2020\)](#) in Italy with 1932 participants assessed the effects of COVID-19-induced confinement policies on self-reported food consumption through an online questionnaire. They found a reduction of nearly 50% in the purchase of ready meals during the pandemic. Nevertheless, the data of these studies showed that, even in the middle of a pandemic, people continued to have access to fresh ingredients, going out to purchase them frequently. These attitudes, such as prioritizing fresh products, preparing homemade meals, and limiting ultra-processed foods, were encouraged by the World Health Organization during the pandemic ([WHO, 2020](#)). On the other hand, in some food insecurity situations during the pandemic context ([Owens et al., 2020](#)), for example, the inaccessibility to fresh foods can lead people to buy cheaper and nutritionally poor foods, most of them ultra-processed foods.

The results of the present study reflect some of the above with people with the highest level of cooking skills reporting that they had already used fresh ingredients more frequently than those with medium or low cooking skills before the pandemic and had a proportional increase during the pandemic among the high, medium, and low cooking skill groups. People who self-reported that they knew how to cook in our study also claimed they used to use more fresh ingredients than those who reported that they did not know how to cook before the pandemic, even though both groups increased the use of this kind of food during the pandemic ([Fig. 3](#)). [De Borba et al. \(2021\)](#) analyzed self-efficacy in cooking and consuming fruits and vegetables among 766 first-year students from a university in southern Brazil. In their study, most participants reported that they knew how to cook (72%), and among them, the majority were confident or extremely confident in cooking from basic ingredients, including fresh vegetables.

The fact that the participants of the present study were in the period of quarantine, where social distancing was imposed and the participants were staying at home for longer periods of time, suggests that they looked for ways to increase their access and use of fresh ingredients to

cook. This may lead to the belief that they are cooking healthier and more frequently during the pandemic when compared to before the imposition of social distancing.

At the same time, the frequency of using ultra-processed foods to cook was also increased by participants during the pandemic in all variables. However, this increase was not as significant as the increase in fresh ingredients ([Fig. 4](#)). The group with low cooking skills was the group with the highest probability of never using ultra-processed foods before and during the pandemic and the smallest probability of using several times per week and daily when compared to the medium and high cooking skills groups. The reason for this could be that they did not cook at all before or during the pandemic, even though the consumption of ultra-processed food requires lower cooking skills than fresh ingredients ([Caraher & Lang, 1999](#); [Chenhall, 2010](#)). Conversely, [Wolfson and Bleich \(2015\)](#) showed that individuals who had a high cooking frequency had a lower frequency of meals taken from fast-food or frozen meals/pizzas per week in the past 30 days, when compared to people with low cooking frequency.

On the other hand, the use of a combination of fresh ingredients and ultra-processed foods had the highest increase during the pandemic among people who had high cooking skills, followed by the medium cooking skills group. Although the group with low cooking skills also increased the consumption of this combination of foods, the probability was lower than in the medium and high cooking skills groups, and they maintained the highest probability of never using combined ingredients before and during the pandemic. This can be justified by the fact that they might not have enough confidence to use some techniques, plan meals, use a large variety of ingredients, and cook from scratch, depending on the type of ultra-processed food ([Jomori et al., 2018](#)). In addition, both students who self-reported knowing and not knowing how to cook reported a higher probability of using combined ingredients during the pandemic when compared to the period before, being higher in the group who reported knowing how to cook ([Fig. 4](#)).

In a study carried out by [Murray et al. \(2016\)](#) with 24 students through focus groups, the participants remarked that they would eat more healthily if they had the knowledge and information to prepare healthy foods. This shows that interventions can be important to this group to develop cooking knowledge, especially regarding the use of fresh ingredients and cooking skills, resulting in the improvement of their diet behaviors. [Seabrook et al. \(2019\)](#) and [Bernardo et al. \(2018\)](#) showed that students who had taken a food and nutrition course, and culinary intervention had more cooking skills than those who had not taken the course. Higher cooking skills acquisition is supported when university students take a culinary intervention and remains in place for long time ([Bernardo et al., 2018](#)).

Cooking knowledge can be related to cooking skills and practices, which are linked to healthy diets. Study of [Utter et al. \(2018\)](#) found that having cooking skills led the participants to better eating behaviors, greater involvement in cooking, and a lower consumption of fast-food in adulthood. [Wolfson and Bleich \(2015\)](#) found a significant association between the habit of making dinner at home with better diet quality and lower consumption of energy, carbohydrates, fat, and sugar, fewer energy from meals consumed outside home, or frozen or ready-to-eat meals. Similarly, [Hartmann et al. \(2013\)](#) showed that cooking skills correlated positively with weekly vegetable consumption, but negatively with weekly convenience food consumption frequency. Thus, there might be a high probability that university students with high levels of cooking skills are more likely to use fresh ingredients to improve their diet compared to those with low levels of cooking skills.

The results of the present study indicate the demand for stimulating and promoting the use of fresh ingredients, teaching how to prepare and cook from scratch, in order to reduce the use of ultra-processed food and to increase the availability and consumption of fruits and vegetables by university students, mainly those with medium and lower cooking skills. The fact that students with low cooking skills have lower probabilities of increasing their frequency of cooking any kind of food leads to the belief

that they are not encouraged to cook, faced by several barriers reported in the literature (Graham et al., 2013; Jones et al., 2014; Knol et al., 2019; Murray et al., 2016; Pulz et al., 2017; Sprake et al., 2018; Utter et al., 2018; Wilson et al., 2017). This highlights the need for strategies focused on cooking skills and excluding these barriers, especially because in Brazil, there are no public policies that promote healthy eating among university students or guidelines for this group. Thus, policies and interventions need to be focused on solving this aspect, independently of the pandemic situation.

5. Conclusion

This study demonstrated that most of the students showed a high level of cooking skill and high availability and accessibility of fruits and vegetables. However, only half of the participants showed high cooking knowledge, even though most believed that they knew how to cook. This suggests that more attention should be given to those who have low cooking knowledge, because the lack of cooking knowledge may affect their confidence in cooking, and in this way, affect their food choices. In addition, the participants increased their cooking frequency during the pandemic, regardless of their cooking skills. However, the group with low cooking skills had only a slight increase in the use of fresh ingredients, possibly because lack sufficient knowledge and confidence to cook from scratch using ingredients that require more cooking skills. Based on these results, interventions to teach, develop and improve the cooking skills to promote healthy eating practices in university students are highly recommended to increase their cooking practices, focusing on the preparation of fresh ingredients and offering options to increase positive cooking attitudes, such as easy and convenient cooking techniques or ways to use fresh foods. Further follow-up studies are required to assess whether these students will retain their cooking knowledge, high cooking skills, and great availability and accessibility of fruits and vegetables after college.

6. Limitations and strengths

The strengths of the study included the large sample size, multilevel analysis, and use of the validated BCSQ, which presents several variables related to healthy eating practices. Moreover, the relationship found between cooking skills and location of meal preparation and consumption before and during the pandemic was able to indicate which situations or university students' profiles need support to improve their cooking skills, cooking knowledge, and diet.

On the other hand, this study was conducted in only two universities in southern Brazil. Even if they receive students from all over the country, this research does not represent all the countries and could be conducted in other Brazilian regions.

Furthermore, our sample included a higher percentage of females (73.48%). This can be justified by the fact that women are usually more concerned about health or diet issues. Recall bias may also be a limitation. However, it was important to analyze this scenario during the pandemic to plan interventions with this population that, in Brazil, does not have a specific public policy.

Moreover, the study did not evaluate measures before and during the pandemic, at different times, but is a cross-sectional study, depending on the record of students. Further analysis is recommended for the follow-up.

Ethical statement

This study was approved by the Human Research Ethics Committee of the Federal University of Santa Catarina (UFSC) and Federal University of Rio Grande do Sul (UFRGS) under the approval number 09427219.5.3001.0121. Participants provided online informed consent prior to all analyses and were assured that all data would be used only for research purposes.

Author contributions

Talissa Dezanetti: Acquisition of data, Interpretation of data, Investigation, Writing as well as for the choice of the Journal.

Ricardo Teixeira Quinaud: The design of the study, data analysis, interpretation of results and manuscript review.

Martin Caraher: contributed for the design, literature, interpretation of results and manuscript review.

Manuela Mika Jomori: Design of the study, Supervision, Interpretation the data, Project administration, Writing and Review, Funding acquisition, and for drafting the manuscript.

Funding support

Funding was supported by National Council for Scientific and Technological Development – CNPq (*Conselho Nacional de Desenvolvimento Científico e Tecnológico*), [grant number 424989/2018–2].

Declaration of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank the contributions of the undergraduate students Clara Nogueira Pacheco, Fernanda Riscado de Souza, and Patrícia Linhares da Cunha from Federal University of Santa Catarina who carried out the data collection, all the university students that participated in our research, the undergraduate course coordinators and students' organizations that contributed to the recruitment of participants. We also acknowledge the contributions of Professor Vanuska Lima da Silva, Federal University of Rio Grande do Sul, regarding the design of the study. We acknowledge the National Council for Scientific and Technological Development – CNPq (*Conselho Nacional de Desenvolvimento Científico e Tecnológico*) from Brazil by the financial support of the research and the scholarship. Finally, we thank the support by from Federal University of Santa Catarina to support internship to students.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.appet.2022.106036>.

References

- Bernardo, G. L., Jomori, M. M., Fernandes, A. C., Colussi, C. F., Condrasky, M. D., & Proença, R. P. da C. (2018). Positive impact of a cooking skills intervention among Brazilian university students: Six months follow-up of a randomized controlled trial. *Appetite*, *130*, 247–255. <https://doi.org/10.1016/j.appet.2018.08.014>
- Bernardo, G. L., Jomori, M. M., Fernandes, A. C., & Proença, R. P. da C. (2017). Food intake of university students. In *Revista de Nutricao* (Vol. 30, pp. 847–865). *Revista de Nutricao*. <https://doi.org/10.1590/1678-98652017000600016>. Issue 6.
- Bezerra, I. N., Medeiros, H. B. N., De Moura Souza, A., & Sichiari, R. (2020). Contribution of away-from-home food to the energy and nutrient intake among Brazilian adolescents. *Public Health Nutrition*, 2–9. <https://doi.org/10.1017/S1368980020001573>
- Błaszczuk-Bębenek, E., Jagielski, P., Boleślawska, I., Jagielska, A., Nitsch-Osuch, A., & Kawalec, P. (2020). Nutrition behaviors in polish adults before and during COVID-19 lockdown. *Nutrients*, *12*(10), 1–16. <https://doi.org/10.3390/nu12103084>
- de Borba, T. P., da Silva, M. V., Jomori, M. M., Bernardo, G. L., Fernandes, A. C., Proença, R. P. da C., Rockenbach, G., & Uggioni, P. L. (2021). Self-efficacy in cooking and consuming fruits and vegetables among Brazilian university students: The relationship with sociodemographic characteristics. *British Food Journal*, ahead-of-p (ahead-of-print). <https://doi.org/10.1108/bfj-04-2020-0311>
- Bürkner, P. (2017). brms: An R package for Bayesian multilevel models using stan. *Journal of Statistical Software*, *80*(1–28).
- Bürkner P., Vuorre, M. Ordinal Regression Models in Psychology: A Tutorial. *Advances in Methods and Practices in Psychological Science*, *2*(1), 77–101. <https://doi.org/10.1177/251524591882319>.

- Caraher, M., Dixon, P., Carr-Hill, R., & Lang, T. (1999). The state of cooking in England: The relationship of cooking skills to food choice. *British Food Journal*, 101(8), 590–609. <https://doi.org/10.1108/00070799910288289>
- Caraher, M., & Lang, T. (1999). Can't cook, won't cook: A review of cooking skills and their relevance to health promotion. *International Journal of Health Promotion and Education*, 37(3), 89–100. <https://doi.org/10.1080/14635240.1999.10806104>
- Chenhall, C. (2010). Improving cooking and food preparation skills: A synthesis of the evidence to inform program and policy development. *Healthy living issue group (HLIG) of the pan-Canadian public health Network*. <http://www.healthcanada.gc.ca>, 1–39.
- Community Food and Health Scotland. (2013). The impact of cooking courses on families: A summary of a research study comparing three different approaches. <http://www.communityfoodandhealth.org.uk/wpcontent/uploads/2013/04/CFHS-im-pact-cooking-courses-families.pdf>.
- Cummins, S., Berger, N., Cornelsen, L., Eling, J., Er, V., Greener, R., Kalbus, A., Karapici, A., Law, C., & Nnlovu, D. (2020). COVID-19: Impact on the Urban Food Retail System and Dietary Inequalities in the UK. *Cities & Health, [S.L.]*, 1–4. <https://doi.org/10.1080/23748834.2020.1785167>
- Deschasaux-Tanguy, M., Druesse-Pecollo, N., Esseddik, Y., de Edelenyi, F. S., Allès, B., Andreeva, V. A., Baudry, J., Charreire, H., Deschamps, V., Egnell, M., Fezeu, L. K., Galan, P., Julia, C., Kesse-Guyot, E., Latino-Martel, P., Oppert, J. M., Péneau, S., Verdout, C., Hercberg, S., & Touvier, M. (2020). Diet and physical activity during the COVID-19 lockdown period (March-May 2020): Results from the French NutriNet-Santé cohort study. *MedRxiv*. <https://doi.org/10.1101/2020.06.04.20121855>. May.
- Di Renzo, L., Gualtieri, P., Pivari, F., Soldati, L., Attinà, A., Cinelli, G., Cinelli, G., Leggeri, C., Caparelli, G., Barrea, L., Scerbo, F., Esposito, E., & De Lorenzo, A. (2020). Eating habits and lifestyle changes during COVID-19 lockdown: An Italian survey. *Journal of Translational Medicine*, 18(1). <https://doi.org/10.1186/s12967-020-02399-5>
- Dooris, M., & Doherty, S. (2010). Healthy universities - time for action: A qualitative research study exploring the potential for a national programme. *Health Promotion International*, 25(1), 94–106. <https://doi.org/10.1093/heapro/daq015>
- Food Standards Agency. (2012). Fresher food – a student survival guide to food. <http://webarchive.nationalarchives.gov.uk/20121205153510/http://www%0Aw.food.gov.uk/northern-ireland/nutritionni/niyoungpeople/fresherfood/>.
- Fulkerson, J. A., Telke, S., Larson, N., Berge, J., Sherwood, N. E., & Neumark-Sztainer, D. (2019). A healthful home food environment: Is it possible amidst household chaos and parental stress? *Appetite*, 142(January), Article 104391. <https://doi.org/10.1016/j.appet.2019.104391>
- Gao, Y., Kennedy, L., Simpson, D., & Gelman, A. (2020). Improving multilevel regression and poststratification with structured priors. *Bayesian Analysis*, 26. <https://doi.org/10.1214/20-BA1223>
- Gatley, A., Caraher, M., & Lang, T. (2014). A qualitative, cross cultural examination of attitudes and behaviour in relation to cooking habits in France and Britain. *Appetite*, 75.
- Gelman, A., & Hill, J. (2007). *Data analysis using regression and multilevel/hierarchical models*. Cambridge: Cambridge University Press.
- Gerritsen, S., Egli, V., Roy, R., Haszard, J., Backer, C. De, Teunissen, L., Cuyck, I., Decorte, P., Pabian, S. P., Van Royen, K., & Te Morenga, L. (2020). Seven weeks of home-cooked meals: Changes to New Zealanders' grocery shopping, cooking and eating during the COVID-19 lockdown. *Journal of the Royal Society of New Zealand*, 1–19. <https://doi.org/10.1080/03036758.2020.1841010>, 0(0).
- Graham, D. J., Pelletier, J. E., Neumark-Sztainer, D., Lust, K., & Laska, M. N. (2013). Perceived social-ecological factors associated with fruit and vegetable purchasing, preparation, and consumption among young adults. *Journal of the Academy of Nutrition and Dietetics*, 113(10), 1366–1374. <https://doi.org/10.1016/j.jand.2013.06.348>
- Hartmann, C., Dohle, S., & Siegrist, M. (2013). Importance of cooking skills for balanced food choices. *Appetite*, 65, 125–131. <https://doi.org/10.1016/j.appet.2013.01.016>
- Husain, W., & Ashkanani, F. (2020). Does COVID-19 change dietary habits and lifestyle behaviours in Kuwait: A community-based cross-sectional study. *Environmental Health and Preventive Medicine*, 25(1). <https://doi.org/10.1186/s12199-020-00901-5>
- Jomori, M., Caraher, M., Bernardo, G. L., Uggioni, P. L., Echevarria-Guanilo, M. E., Condrasky, M., & Proença, R. P. da C. (2021). How was the cooking skills and healthy eating evaluation questionnaire culturally adapted to Brazil? *Ciência & Saúde Coletiva*, 26(6), 2379–2393. <https://doi.org/10.1590/1413-81232021266.22102019>
- Jomori, M. M., de Vasconcelos, F. de A. G., Bernardo, G. L., Uggioni, P. L., & Proença, R. P. da C. (2018). The concept of cooking skills: A review with contributions to the scientific debate. *Revista de Nutrição*, 31(1), 119–135. <https://doi.org/10.1590/1678-98652018000100010>
- Jomori, M., Proença, R., Bernardo, G., & Fernandes, A. C. (2017). Construct validity of Brazilian cooking skills and healthy eating questionnaire by the known-groups method. *Article in British Food Journal*, 119(5), 1003–1016. <https://doi.org/10.1108/BFJ-10-2016-0448>
- Jomori, M., Quinaud, R. T., Condrasky, M., & Caraher, M. (2022). Brazilian Cooking Skills Questionnaire evaluation of using/cooking and consumption of fruits and vegetables. *Nutrition*, 95. <https://doi.org/10.1016/j.nut.2021.111557>
- Jones, S. A., Walter, J., Soliah, L. A., & Phifer, J. T. (2014). Perceived motivators to home food preparation: Focus group findings. *Journal of the Academy of Nutrition and Dietetics*, 114(10), 1552–1556. <https://doi.org/10.1016/j.jand.2014.05.003>
- Jribi, S., Ben Ismail, H., Doggui, D., & Debbabi, H. (2020). COVID-19 virus outbreak lockdown: What impacts on household food wastage? *Environment, Development and Sustainability*, 22(5), 3939–3955. <https://doi.org/10.1007/s10668-020-00740-y>
- Kennedy, L., & Gelman, A. (2021). Know your population and know your model: Using model-based regression and poststratification to generalize findings beyond the observed sample. *Psychological Methods*, 26(5), 547–558. <https://doi.org/10.1037/met0000362>
- Knol, L. L., Robb, C. A., McKinley, E. M., & Wood, M. (2019). Very low food security status is related to lower cooking self-efficacy and less frequent food preparation behaviors among college students. *Journal of Nutrition Education and Behavior*, 51(3), 357–363. <https://doi.org/10.1016/j.jneb.2018.10.009>
- Kratt, P., Reynolds, K., & Shewchuk, R. (2000). The role of availability as a moderator of family fruit and vegetable consumption. *Health Education & Behavior*, 27(4), 471–482. <https://doi.org/10.1177/109019810002700409>
- Kruschke, J. K., & Liddell, T. M. (2018). Bayesian data analysis for newcomers. *Psychonomic Bulletin & Review*, 25(1), 155–177. <https://doi.org/10.3758/s13423-017-1272-1>
- Larson, N. I., Perry, C. L., Story, M., & Neumark-Sztainer, D. (2006). Food preparation by young adults is associated with better diet quality. *Journal of the American Dietetic Association*, 106(12), 2001–2007. <https://doi.org/10.1016/j.jada.2006.09.008>
- Laska, M. N., Larson, N. I., Neumark-Sztainer, D., & Story, M. (2012). Does involvement in food preparation track from adolescence to young adulthood and is it associated with better dietary quality? Findings from a 10-year longitudinal study. *Public Health Nutrition*, 15(7), 1150–1158. <https://doi.org/10.1017/S1368980011003004>
- Lavelle, F., McGowan, L., Spence, M., Caraher, M., Raats, M. M., Hollywood, L., McDowell, D., McCloat, A., Mooney, E., & Dean, M. (2016). Barriers and facilitators to cooking from 'scratch' using basic or raw ingredients: A qualitative interview study. *Appetite*, 107, 383–391. <https://doi.org/10.1016/j.appet.2016.08.115>
- Lee, M. D., & Wagenmakers, E. J. (2013). *Bayesian cognitive modeling: A practical course*. Cambridge University Press.
- Liddell, T., & Kruschke, J. K. (2017). Analyzing ordinal data with metric models: What could possibly go wrong? *Journal of Experimental Social Psychology*, 79, 328–348.
- McElreath, R. (2015). *Statistical rethinking: A Bayesian course with examples in R and Stan* (p. 469). Chapman & Hall/CRC Press. xvii.
- Mills, S., White, M., Brown, H., Wrieden, W., Kwasnicka, D., Halligan, J., Robalino, S., & Adams, J. (2017). Health and social determinants and outcomes of home cooking: A systematic review of observational studies. *Appetite*, 111, 116–134.
- Murray, D. W., Mahadevan, M., Gatto, K., O'Connor, K., Fissinger, A., Bailey, D., & Cassara, E. (2016). Culinary efficacy: An exploratory study of skills, confidence, and healthy cooking competencies among university students. *Perspectives in Public Health*, 136(3), 143–151. <https://doi.org/10.1177/17579139156000195>
- Neumark-Sztainer, D., Wall, M., Perry, C., & Story, M. (2003). Correlates of fruit and vegetable intake among adolescents: Findings from Project EAT. *Preventive Medicine*, 37(3), 198–208. [https://doi.org/10.1016/S0091-7435\(03\)00114-2](https://doi.org/10.1016/S0091-7435(03)00114-2)
- Oliveira, C. de S. (2017). *A universidade promotora da Saúde: uma revisão de literatura*.
- Owens, M. R., Brito-silva, F., Kirkland, T., Moore, C. E., Davis, K. E., Patterson, M. A., Miketinas, D. C., & Tucker, W. J. (2020). Prevalence and social determinants of food. *Nutrients*, 12(9), 1–18.
- Piton Hakim, M., D'Avoglio Zanetta, L., & Thimoteo da Cunha, D. (2021). Should I stay, or should I go? Consumers' perceived risk and intention to visit restaurants during the COVID-19 pandemic in Brazil. *Food Research International*, 141, Article 110152. <https://doi.org/10.1016/j.foodres.2021.110152>
- Pulz, I. S., Martins, P. A., Feldman, C., & Veiros, M. B. (2017). Are campus food environments healthy? A novel perspective for qualitatively evaluating the nutritional quality of food sold at foodservice facilities at a Brazilian university. *Perspectives in Public Health*, 137(2), 122–135. <https://doi.org/10.1177/1757913916636414>
- R Core Team. (2018). *R: A language and environment for statistical computing*.
- Rathi, N., Riddell, L., & Worsley, A. (2018). Indian adolescents' perceptions of the home food environment. *BMC Public Health*, 18(1), 1–7. <https://doi.org/10.1186/s12889-018-5083-8>
- REDUPS. (2013). Guía para la autoevaluación y reconocimiento de instituciones de educación superior promotoras de la salud. <https://deportes.udem.cl/wp-content/uploads/2016/11/09-Guía-para-la-Autoevaluación-y-Reconocimiento-de-Instituciones-de-Educación-Superior-Promotoras-de-la-Salud-Red-Chilena-de-Universidades-Promotoras-de-la-Salud.pdf>, 34.
- Restrepo, B. J., & Zeballos, E. (2020). The effect of working from home on major time allocations with a focus on food-related activities. *Review of Economics of the Household*, 18(4), 1165–1187. <https://doi.org/10.1007/s11150-020-09497-9>
- Reyes-Olavarría, D., Latorre-Román, P.Á., Guzmán-Guzmán, I. P., Jerez-Mayorga, D., Caamaño-Navarrete, F., & Delgado-Floody, P. (2020). Positive and negative changes in food habits, physical activity patterns, and weight status during COVID-19 confinement: Associated factors in the Chilean population. *International Journal of Environmental Research and Public Health*, 17(15), 5431. <https://doi.org/10.3390/ijerph17155431>
- Ribeiro, K. D. D. S., Garcia, L. R. S., Dامتو, J. F. D. S., Assunção, D. G. F., & Maciel, B. L. L. (2020). COVID-19 and nutrition: The need for initiatives to promote healthy eating and prevent obesity in childhood. *Childhood Obesity*, 16(4), 235–237. <https://doi.org/10.1089/chi.2020.0121>
- Ruiz-Roso, M. B., Knott-Torcal, C., Matilla-Escalante, D. C., Garcimartín, A., Sampedro-Nuñez, M. A., Dávalos, A., & Marazuela, M. (2020). COVID-19 lockdown and changes of the dietary pattern and physical activity habits in a cohort of patients with type 2 diabetes mellitus. *Nutrients*, 12(8). <https://doi.org/10.3390/nu12082327>
- Santos, S., Vilela, S., Padrão, P., & Caraher, M. (2015). Sex-related dietary changes of Portuguese university students after migration to London, UK. *Nutrition and Dietetics*, 72, 340–346.
- Dietary Guidelines for the Brazilian. (2014). In M. da Saúde, & Brasil (Eds.) (2nd ed.). Scarmozzino, F., & Visioli, F. (2020). Covid-19 and the subsequent lockdown modified dietary habits of almost half the population in an Italian sample. *Foods*, 9(5), 675. <https://doi.org/10.3390/foods9050675>

- Seabrook, J. A., Dworatzek, P. D. N., & Matthews, J. I. (2019). Predictors of food skills in university students. *Canadian Journal of Dietetic Practice and Research*, 80(4), 205–208. <https://doi.org/10.3148/cjdp-2019-011>
- Short, F. (2006). *Kitchen secrets: The meaning of cooking in everyday life*. New York; Oxford: Berg.
- Soares, A. M., Pereira, A. M. S., & Canavaro, J. M. A. P. (2015). *Promoção da Saúde nas Instituições de Ensino Superior Portuguesas: Reflexões e Desafios* (Vols. 115–137). Revista Portuguesa de Pedagogia. https://doi.org/10.14195/1647-8614_49-2_6
- Sprake, E. F., Russell, J. M., Cecil, J. E., Cooper, R. J., Grabowski, P., Pourshahidi, L. K., & Barker, M. E. (2018). Dietary patterns of university students in the UK: A cross-sectional study. *Nutrition Journal*, 17(1), 1–17. <https://doi.org/10.1186/s12937-018-0398-y>
- Tsouros, A., Dowding, G., Thompson, J., & Dooris, M. (1998). Health promoting universities. <http://www.ncbi.nlm.nih.gov/pubmed/18447035>.
- United Nations. (2015). Transforming our world: The 2030 agenda for sustainable development. <https://doi.org/10.1163/157180910X12665776638740>, 35.
- USDA. (2013). *My Plate on campus Toolkit* (issue march). <http://choosemyplate.gov/MyPlateOnCampus/downloads/MyPlateOnCampusToolkit.pdf>.
- Utter, J., Larson, N., Laska, M. N., Winkler, M., & Neumark-Sztainer, D. (2018). Self-Perceived cooking skills in emerging adulthood predict better dietary behaviors and intake 10 Years later: A longitudinal study. *Journal of Nutrition Education and Behavior*, 50(5), 494–500. <https://doi.org/10.1016/j.jneb.2018.01.021>
- Veríssimo, J. (2021). Analysis of rating scales: A pervasive problem in bilingualism research and a solution with Bayesian ordinal models. *Bilingualism: Language and Cognition*, 24(5), 842–848. <https://doi.org/10.1017/S1366728921000316>
- Vilela, S., Santos, S., Padrao, P., & Caraher, M. (2014). Length of migration and eating habits of Portuguese university students living in London, United Kingdom. *Ecology of Food and Nutrition*, 53, 419–435.
- WHO, W. H. O. (2015). *Global School health initiatives: Achieving health and education outcomes report of a meeting*. Geneva: World Health Organization. November <http://apps.who.int/iris/bitstream/10665/259813/1/WHO-NMH-PND-17-7-eng.pdf?ua=1>.
- WHO. (2020). Food and nutrition tips during self-quarantine. <https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/publications-and-technical-guidance/noncommunicable-diseases/food-and-nutrition-tips-during-self-quarantine?fbclid=IwAR0IxmHZqgX>.
- Wilson, C. K., Matthews, J. I., Seabrook, J. A., & Dworatzek, P. D. N. (2017). Self-reported food skills of university students. *Appetite*, 108, 270–276. <https://doi.org/10.1016/j.appet.2016.10.011>, 2017.
- Wolfson, J. A., & Bleich, S. N. (2015). Is cooking at home associated with better diet quality or weight-loss intention? *Public Health Nutrition*, 18(8), 1397–1406. <https://doi.org/10.1017/S1368980014001943>
- Talissa Dezanetti** is an undergraduate student of Nutrition of Federal University of Santa Catarina and a scholar supported by National Council for Scientific and Technological Development – CNPq (*Conselho Nacional de Desenvolvimento Científico e Tecnológico*) on the research's project “*Nutrition is in the kitchen! Cooking skills and Healthy eating in the university*”.
- Ricardo Teixeira Quinaud** is a PhD student of Physical Education Postgraduate Program of Federal University of Santa Catarina, member of the research project “*Nutrition is in the kitchen! Cooking skills and Healthy eating in the university*”.
- Martin Caraher** is Professor Emeritus in Food and Health Policy at Centre for Food Policy at City, University London. He is expert in the cooking skills' thematic within public health framework and a member of the research's project “*Nutrition is in the kitchen! Cooking skills and Healthy eating in the university*”.
- Manuela Mika Jomori** is Professor of Department of Nutrition of Federal University of Santa Catarina and Visiting Fellow at Centre for Food Policy, City, University of London. Author of the Brazilian Cooking Skills and Healthy Eating Questionnaire and coordinator of the research's project “*Nutrition is in the kitchen! Cooking skills and Healthy eating in the university*” in Brazil.