

Research Article

Open Access

Alessandro Borello, Alessia Ferrarese*, Roberto Passera, Alessandra Surace, Silvia Marola, Claudio Buccelli, Massimo Niola, Pierpaolo Di Lorenzo, Maurizio Amato, Lorenza Di Domenico, Mario Solej, Valter Martino

Use of a simplified consent form to facilitate patient understanding of informed consent for laparoscopic cholecystectomy

DOI 10.1515/med-2016-0092

received October 10, 2016; accepted October 31, 2016

Abstract: Background. Surgical informed consent forms can be complicated for patients to read and understand. We created a consent form with key information presented in bulleted texts and diagrams combined in a graphical format to facilitate the understanding of information during the verbal consent discussion.

Methods. This prospective, randomized study involved 70 adult patients awaiting cholecystectomy for gallstones. Consent was obtained after standard verbal explanation using either a graphically formatted (study group, n=33) or a standard text document (control group, n=37). Comprehension was evaluated with a 9-item multiple-choice questionnaire administered before surgery and factors affecting comprehension were analyzed.

Results. Comparison of questionnaire scores showed no effect of age, sex, time between consent and surgery, or

document format on understanding of informed consent. Educational level was the only predictor of comprehension.

Conclusions. Simplified surgical consent documents meet the goals of health literacy and informed consent. Educational level appears to be a strong predictor of understanding.

Keywords: Informed consent; Laparoscopic cholecystectomy

1 Introduction

Informed consent refers to a voluntary decision and an authorization by a patient that a health care provider can proceed with diagnostic or therapeutic interventions as planned during the consent discussion. An integral component of the consent process, the consent discussion, is structured around the essential points of the procedure, associated side effects, reasonably predictable risks, benefits, and outcomes, as well as other treatment options [1].

Informed consent is now ethically and legally enshrined as the right of patients to receive adequate information to make decisions about proposed treatment and currently available alternative options in medical care [2]. This is a relatively recent concept in some areas of medicine, and reflects the gradual shift away from the paternalistic approach predominant up until the early 1960s when doctors believed that, by virtue of their superior medical knowledge, they could disregard patients' choices and preferences or act against them if they ran counter to clinical indications. Surveys assessing health literacy in adults have found, however, that many people

*Corresponding author: **Alessia Ferrarese**, Department of Oncology, University of Turin, Section of General Surgery, San Luigi Gonzaga University Hospital, Regione Gonzole 10, 10043 Orbassano, Turin, Italy, Phone: +39 0119026224, Fax: +39 0119026523, E-mail: alessia.ferrarese@gmail.com

Alessandro Borello, Alessandra Surace, Silvia Marola, Mario Solej, Valter Martino, University of Turin, Department of Oncology, School of Medicine, San Luigi Gonzaga University Hospital, Section of General Surgery, Orbassano, Turin, Italy

Roberto Passera, Città della Salute e della Scienza Hospital, Section of Nuclear Medicine, Turin, Italy

Claudio Buccelli, Massimo Niola, Pierpaolo Di Lorenzo, Department of Advanced Biomedical Sciences, Naples, Italy, University "Federico II" of Naples. Via Sergio Pansini 5, 80131, Naples, Italy

Maurizio Amato, Department of Neuroscience, Naples, Italy. Fellow in General Surgery at University of Naples, Naples, Italy

Lorenza Di Domenico, Department of Neuroscience, Naples, Italy. Fellow in Anesthesiology at University of Naples, Naples, Italy

have a poor grasp of basic health concepts, medical information and instructions.

Other studies have shown that patients often have trouble to follow an explanation of diagnostic and therapeutic procedures discussed during the consent consultation [3] and are able to recall little of what was explained [4]. Because surgical informed consent forms are complex for patients to read and understand, we created a two-page consent document with key information presented in bulleted texts and diagrams combined in a graphical format to facilitate understanding during the verbal consent process for laparoscopic cholecystectomy. Factors potentially affecting comprehension, as assessed with a feedback questionnaire, were analyzed.

2 Material and methods

2.1 Study sample

In this prospective randomized study, 100 consecutive patients awaiting elective laparoscopic cholecystectomy for gallstones on a day surgery basis were enrolled between 1 April 2013 and 1 April 2015. Exclusion criteria were: incapacity to read and understand the informed consent form unaided, emergency surgery, need for other surgical procedures in addition to elective laparoscopic cholecystectomy, refusal to participate in the study, non-Italian speakers or patients noted to have difficulty understanding or communicating in Italian, severe mental disorders, and incapacity to provide adequate informed consent. A single physician and member of the medical team collected the consent forms from all enrolled patients before hospital admission for surgery.

Of these 100 patients, 18 were excluded because they had undergone emergency surgery during the time between informed consent collection and the scheduled date of elective surgery; 12 others were excluded because they met one of the exclusion criteria listed above. The 70 remaining patients were assigned to one of two groups according to a computer-generated sequence (available at www.randomization.com). Of these 70 participants, 33 (47%) were men and 37 were women (52.9%). The median age was 55 years (range, 24-80); 37 (52.9%) patients were below and 33 (47.1%) were above the median age. The study group included 33 (47.1%) patients and the control group 37 (52.9%). The demographic characteristics were similar in both groups (Table 1).

2.2 Informed consent discussion

The informed consent discussion during surgical preadmission assessment covered the essential points regarding the pathology and its natural history, clinical course and common complications associated with gallstones, the reason for laparoscopic surgery and related risks and benefits, details about the surgical procedure, the potential surgical risks and reasonably predictable complications, the postoperative course, the diet to be followed during the first month after surgery, and aftercare of the surgical wound. Consent was obtained after the exposure to standard verbal explanation with either a simplified informed consent form (study group) or a standard text document (control group) routinely used in our unit for this purpose. The simplified consent form featured a clear layout and visual design elements, including ample margins and white space, short bulleted lists with topic headings, both medical and lay terms, anatomical diagrams and figures with descriptions and captions, and institutional logo (Fig. 1). Neither informed consent form has been tested for readability standards.

2.3 Feedback questionnaire

Comprehension was evaluated with a non-validated, 9-item multiple-choice questionnaire administered the day before surgery. The self-report multiple-choice questionnaire (each of the 9 questions had only one correct answer from 2-4 choices) examined understanding of informed consent regarding the operation itself (questions 1 to 4), potential surgical complications (questions 5 and 6), and length of hospital stay and postoperative care instructions (questions 7 to 9). An additional final question (item 10 had only a yes/no response) investigated patient satisfaction with the clarity of the presentation and the amount of information given (Fig. 2). Physicians that obtained the consent forms also collected completed questionnaires.

Patient age, sex, educational level, time between consent and questionnaire collection, and questionnaire scores were entered into a database for analysis. Eleven (15.7%) patients had completed elementary school, 24 (34.3%) middle school, 10 (14.3%) technical-vocational school, 19 (27.1) high school, and 6 (8.6%) college level education. The average time between consent and collection of the feedback questionnaire was 23.4 days (median 12 days, range 0 to 152); the time interval was less than 15 days for 39 (55.7%) patients and more than 16 days for the remaining 31 (44.3%).

Regione Piemonte 10 - 10043 Orbassano TORINO
 Centrale tel. +39 011 902621 - fax +39 011 9026502
 e-mail: cs@hsapijgpiemonte.it
 Cod. Fisc. 98871000102 - P.I. 02959440119


S.C.D.U. Chirurgia Generale 2
 Direttore: Prof. Mario Nanno
 Tel. 011 9026224-9026369
 Fax: 011 9026383
 e-mail: chirurgia.universitaria@hsapijgpiemonte.it

INFORMED CONSENT FORM
Laparoscopic cholecystectomy for cholelithiasis

Name _____ Surname _____
 Date of birth _____ Birth place _____
 Qualification _____ Date of informed consent _____

THE OPERATION

- Duration: 1-3 h.
- General anesthesia.
- Pneumoperitoneum with CO₂.
- Incisions for trocar: Umbilical incision (10mm) for telescope, epigastric and left flank (5mm) for working port, median incision (10mm) for left working port.
- Identification and complete removal of the gallbladder.
- Reduction of pneumoperitoneum and closure of the breach.




Regione Piemonte 10 - 10043 Orbassano TORINO
 Centrale tel. +39 011 902621 - fax +39 011 9026502
 e-mail: cs@hsapijgpiemonte.it
 Cod. Fisc. 98871000102 - P.I. 02959440119

-Drain placed if needed.

COMMON SURGICAL COMPLICATIONS

- Conversion to right subcostal laparotomy incision (2-7% of cases)
- Biliary duct injury (0.4-0.6% of cases) or biliary fistulas (0.5% of cases)
- Vascular injury (0.3% of cases) or postoperative bleeding (0.5% of cases)
- Intestinal injury (0.09% of cases)
- Infections (4% of cases)
- Other complications (related to anesthesia, related to the patient comorbidity, etc.).




HOSPITALIZATION DETAILS

- Postoperative length of stay: 24-48h
- Thrombosis prophylaxis (compression stockings and LMWH)


POST-OPERATIVE INSTRUCTION

- Return for outpatient visit after 7-10 days
- Relative rest for the first 2 weeks
- Light low-fat diet for the first 20-30 days
- Unrestricted diet after 60 days

DATE _____
 PATIENT SIGNATURE _____ OPERATOR SIGNATURE _____



REGIONE PIEMONTE
 www.regione.piemonte.it



Non c'è cura senza cuore
 Istituto Sanitario Regione del Piemonte

Figure 1: Simplified informed consent form

Table 1: Patient characteristics.

Characteristic	Number of patients (n=70)	Study Group (n=33)	Control Group (n=37)
Sex			
Male, n° (%)	33 (47.1)	15 (45.4)	18 (48.6)
Female n° (%)	37 (52.9)	18 (54.6)	19 (51.4)
Age – average (yrs)	54	54	55
Median	55	55	55
Range	24-80	24-78	35-80
≤55	37 (52.9)	16 (48.5)	21 (56.7)
>56	33 (47.1)	17 (51.5)	16 (43.3)
Educational level			
Elementary n° (%)	11 (15.7)	4 (12.1)	7 (18.9)
Middle school, n° (%)	24 (34.3)	11 (33.3)	13 (35.1)
Technical-vocational, n° (%)	10 (14.3)	5 (15.1)	5 (13.5)
High school, n° (%)	19 (27.1)	11 (33.3)	8 (21.6)
College, n° (%)	6 (8.6)	2 (6.2)	4 (10.8)
Time between consent and surgery (days)			
Average	23.4	23.4	22.9
Median	12	12	12
Range	0-152	0-152	0-115
Δ Time ≤ 15, n° (%)	39 (55.7)	19 (57.6)	20 (54.1)
Δ Time > 16, n° (%)	31 (44.3)	14 (42.4)	17 (45.9)

CENTRO 011 9026224 - TEL 011 9026224
e-mail: info@sanluigi.piemonte.it
Cod. Fisc. 96801020010 - P.I. 0295840016

S.C.D.U. Chirurgia Generale 2
Direttore: Prof. Mario Nano
Tel.: 011.9026224-9026369
Fax: 011.9026383
e-mail: chirurgia_universitaria@sanluigi.piemonte.it

Comprehension TEST
Laparoscopic cholecystectomy for cholelithiasis

Name _____ Surname _____
Date of birth _____ Birth place _____
Qualification _____ Date of informed consent _____

Question 1. Surgery is performed using a/an:
-Laparoscopic approach
-Open subcostal incision

Question 2. Laparoscopic surgery involves:
-Placing a subcostal incision to expose the gallbladder and remove it
-Creating minimally invasive accesses to the abdomen for the instruments

Question 3. Which type of anesthesia is given?
-General
-Local
-Spinal

Question 4. What will be removed?
-The gallbladder
-Only the gallstones

Question 5. If technical or anatomical problems arise, can the procedure be converted from laparoscopic to subcostal incision?
-Yes
-No

Question 6. What kind of complications may occur?
-Vascular injury
-Intestinal injury
-Biliary duct injury
-All of the above

Question 7. How long is the average length of stay in the hospital?
-24-48 hours
-3-5 days
-7-10 days

Question 8. What kind of diet should be followed after the operation?
No restrictions
-A light diet for the first 10-15 days
-A light diet for the first 3-6 months
-A restricted diet indefinitely

Question 9. What kind of food should be avoided the first days after the operation?
-Sugar
-Protein
-Fat

Question 10. Was the explanation of the operation and possible complications sufficiently clear?
-Yes
-No

Date _____
Operator Signature _____
Patient signature _____

REGIONE PIEMONTE
Non c'è cura senza cuore
REGIONE PIEMONTE
Non c'è cura senza cuore

Figure 2: Feedback questionnaire

2.4 Statistical analysis

Univariate descriptive analysis was applied to categorical and continuous variables, followed by bivariate descriptive analysis of the categorical variables, inferential analysis of the categorical variables (two-sided Monte Carlo and Fisher's exact test), and a multivariate binary logistic regression model (odds ratio with 95% confidence intervals). Statistical significance was set at $P \leq 0.05$.

3 Results

The correct response rate was 100% on all 9 questionnaire items for 20 out of 70 patients: 11/33 (33.3%) in the study group and 9/37 (24.3%) in the control group; 23 responded incorrectly to 1 item (11 [33.3%] in the study group vs. 12 [32.4%] in the control group); 17 responded incorrectly to 2 items (7 [21.2%] in the study group vs. 10 [27.1%] in the control group); 8 responded incorrectly to 3 items (3 [9.1%] in the study group vs. 5 [13.5%] in the control group); 2 responded incorrectly to 4 items (1 [3.1%] in the study group vs. 1 [2.7%] in the control group). The percentage of correct responses to the first 5 items investigating understanding of the operation was generally high in both

groups: 97% and 100% to item 1, 96.9% and 97.3% to item 2, 93.9% and 94.6% to item 3, 93.9% and 100% to item 4, and 90.9% and 100% to item 5 in the study and the control group, respectively.

By comparison, the correct response rate was lowest for item 6 (probability of surgical complications), with 57.6% and 62.1% in the study and the control group, respectively, responding incorrectly. The correct response rate improved on items 7 to 9 (length of hospital stay and postoperative care instructions): 78.8% vs. 67.6%, 90.9% vs. 75.7%, and 100% vs. 89.2% in the study and the control group, respectively. All patients stated that they were satisfied with the way they were informed and the clarity of the information provided irrespective of whether they had been exposed to verbal consent with the standard text or with the simplified graphically enhanced document (Table 2).

Univariate and bivariate analysis of the categorical variables (age, sex, educational level, time between consent and surgery) showed that none of these were statistically significant as a predictor of understanding consent nor did exposure to the standard text or graphically formatted consent document predict better understanding (Table 3). Multivariate binary logistic regression (odds ratio with 95% confidence intervals) of these variables showed that, independent of type of exposure,

Table 2: Feedback questionnaire items and number and percentage of patients who responded correctly or incorrectly.

	N° of patients (n=70)	Study Group (n=33)	Control Group (n=37)
Question 1. Surgery is performed using a/an:			
Laparoscopic approach			
Open subcostal incision			
Correct, n(%)	69 (98.6)	32 (97)	37 (100)
Incorrect, n° (%)	1 (1.4)	1 (3)	0 (0)
Question 2. Laparoscopic surgery involves:			
Placing a subcostal incision to expose the gallbladder and remove it			
Creating minimally invasive accesses to the abdomen for the instruments			
Correct, n° (%)	68 (97.1)	32 (96.9)	36 (97.3)
Incorrect, n° (%)	2 (2.9)	1 (3.1)	1 (2.7)
Question 3. Which type of anesthesia is given?			
General			
Local			
Spinal			
Correct, n° (%)	66 (94.3)	31 (93.9)	35 (94.6)
Incorrect, n° (%)	4 (5.7)	2 (6.1)	2 (5.4)
Question 4. What will be removed?			
The gallbladder			
Only the gallstones			
Correct, n° (%)	68 (97.1)	31 (93.9)	37 (100)
Incorrect, n° (%)	2 (2.9)	2(6.1)	0 (0)
Question 5. If technical or anatomical problems arise, can the procedure be converted from laparoscopic to subcostal incision?			
Yes			
No			
Correct, n° (%)	67 (95.7)	30 (90.9)	37 (100)
Incorrect, n° (%)	3 (4.3)	3 (9.1)	0 (0)
Question 6. What kind of complications may occur?			
Vascular injury			
Intestinal injury			
Biliary duct injury			
All of the above			
Correct, n° (%)	28 (40.0)	14 (42.4)	14 (37.8)
Incorrect, n° (%)	42 (60.0)	19 (57.6)	23 (62.1)
Question 7. How long is the average length of stay in the hospital?			
24-48 hours			
3-5 days			
7-10 days			
Correct, n° (%)	51 (72.9)	26 (78.8)	25 (67.6)
Incorrect, n° (%)	19 (27.1)	7 (21.2)	12 (32.4)
Question 8. What kind of diet should be followed after the operation?			
No restrictions			
A light diet for the first 10-15 days			
A light diet for the first 3-6 months			
A restricted diet indefinitely			
Correct, n° (%)	58 (82.9)	30 (90.9)	28 (75.7)
Incorrect, n° (%)	12 (17.1)	3 (9.1)	9 (24.3)

Table 2: Feedback questionnaire items and number and percentage of patients who responded correctly or incorrectly.

Question 9. What kind of food should be avoided the first days after the operation?			
Sugar			
Protein			
Fat			
Correct, n° (%)	66 (94.3)	33 (100)	33 (89.2)
Incorrect, n° (%)	4 (5.7)	0 (0)	4 (10.8)
Question 10. Was the explanation of the operation and possible complications sufficiently clear?			
Yes	70 (100.0)	33 (100)	37 (100)
No	0 (0.0)	0 (0)	0 (0)
Number of errors			
0, n° (%)	20 (28.6)	11 (33.3)	9 (24.3)
1, n° (%)	23 (32.9)	11 (33.3)	12 (32.4)
2, n° (%)	17 (24.3)	7 (21.2)	10 (27.1)
3, n° (%)	8 (11.4)	3 (9.1)	5 (13.5)
4, n° (%)	2 (2.9)	1 (3.1)	1 (2.7)

Table 3: Results of interferential analysis of the categorical variables (two-sided Monte Carlo and Fisher's exact test).

Characteristic	Number of patients n=70 (%)	Study Group n=33 (%)	Control Group n=37 (%)	Univariate Analysis Chi-Square Tests p-value
Gender				0.381
Male	33 (47.1)	15 (45.4)	18 (48.6)	
Female	37 (52.9)	18 (54.6)	19 (51.4)	
Age (yrs)				0.260
≤55	37 (52.9)	16 (48.5)	21 (56.7)	
>56	33 (47.1)	17 (51.5)	16 (43.3)	
Educational level				0.285
Elementary	11 (15.7)	4 (12.1)	7 (18.9)	
Middle school	24 (34.3)	11 (33.3)	13 (35.1)	
Technical-vocational	10 (14.3)	5 (15.1)	5 (13.5)	
High school	19 (27.1)	11 (33.3)	8 (21.6)	
College	6 (8.6)	2 (6.2)	4 (10.8)	
Time between consent and surgery (days)				0.567
Δ Time ≤ 15	39 (55.7)	19 (57.6)	20 (54.1)	
Δ Time > 16	31 (44.3)	14 (42.4)	17 (45.9)	
Type of consent	70 (100)	33(100)	37(100)	0.905

Table 4: Results of multivariate binary logistic regression analysis comparing the factors affecting the probability of making errors: 0 errors (20 patients) vs. ≥ 1 error (50 patients).

Variable	Odds Ratio (95% confidence interval)	P value
Age (yrs) >56 vs. ≤ 55	1.58 (0.4-6.22)	0.515
Study group vs. Control group	0.71 (0.22-2.27)	0.560
Female vs. Male	0.46 (0.14-1.52)	0.203
Δ Time between consent and feedback questionnaire (days) >16 vs. ≤ 15	0.79 (0.25-2.56)	0.698
Educational level		0.048
Middle vs. Elementary	0.5 (0.05-5.08)	0.558
Technical-vocational vs. Elementary	0.4 (0.03-5.25)	0.485
High vs. Elementary	0.09 (0.01-0.85)	0.035
College vs. Elementary	0.1 (0.01-1.35)	0.043

educational level had a statistically significant effect on reducing the risk of responding incorrectly by the patients in both groups who had completed high school or college ($P = 0.035$ and $P = 0.043$, respectively) (Table 4).

4 Discussion

Like the trust that underlies the doctor-patient relationship, informed consent can be viewed as both an event and a process [5]. It begins with the preoperative encounter and continues through to the postoperative evaluation of treatment outcomes and patient satisfaction.

In a questionnaire study, Akkad *et al.* [6] found, however, that patients often had a limited understanding of the legal implications of signing an informed consent form and that they did not recognize the fact that informed consent works in their interest: 46% stated that informed consent serves to protect the doctor's interest. The authors concluded that the current informed consent process is inadequate and its ethical validity and credibility are questionable.

Health literacy surveys have underlined the need to provide medical information in plain language in order to bridge the knowledge gap between doctor and patient. Kusec *et al.* [7] reported that the complexity of presenting concepts could be graded according to patient characteristics by aiming for a balanced choice between medical and lay terms adjusted to the patient's educational level, for example. By the same token, not all patients have a desire of being fully informed about clinical aspects, therapy, common complications, and treatment outcomes.

Degerliyurt *et al.* [8] found a wide variability in the amount of information about therapy and potential complications that patients want to know: 57% of the patients in their study stated they wanted to be fully informed about all the complications associated with their surgical treatment, 33% wanted to know only the most common complications, and 10% did not want to be informed. Despite these discrepancies, the authors recommended that patients receive a standard informed consent form.

In a systematic review of the literature, Falagas *et al.* [9] found that less than one third of the studies they reviewed reported that patients had adequately understood of the components of the informed consent process. Although the majority of the patients felt that the amount of information they had received was adequate, the actual degree of understanding was suboptimal.

There are now various methods and strategies to simplify consent forms and improve comprehension with the aid of especially designed written or multimedia material being used during the informed consent process in clinical trials. The general agreement is that the use of such materials may increase patient understanding of informed consent [10-16]. Multimedia presentations have been effectively implemented in education to increase learning, interest, memory, comprehension, and satisfaction. Hung *et al.* [17] reported that the use of multimedia disclosure of informed consent does not significantly increase hospital costs. Not all authors agree that multimedia use necessarily improves patients' informed consent understanding, however. Clark *et al.* [18] reported that PowerPoint presentations illustrating laparoscopic cholecystectomy did not increase the degree of patient understanding. Other authors have suggested that a consent discussion follow-

ing a checklist of essential concepts is sufficient to obtain good understanding of consent [19] or that repeat back of concepts and repetition of the points not clearly understood can also help to improve understanding [20]. What has been consistently shown is that an increased understanding of the surgical procedure can increase satisfaction with treatment received and reduce perioperative anxiety [21-24] despite of the inherent difficulties of surgical procedures [25-44].

In our study, comparison of feedback questionnaire scores revealed no statistically significant difference in informed consent understanding between the two groups. This finding contrasts with previously published data [10-16]. We agree with Clark et al. [18] and Kondziolka [19] who demonstrated that the addition of educational material to the standard verbal consent process does not improve patient understanding of the risks and benefits associated with surgery. Like Falagas et al. [8], we noted that although the majority of patients felt that the amount of information received was adequate, their actual understanding was low: only 28.6% of the patients responded correctly to all 9 questionnaire items, though both groups stated they were satisfied with the clarity with which the information was presented.

Laparoscopic surgery seems to give greater difficulties in intervention explanation perhaps because of technology that can not be understood by all types of patients . [45-57]. Further to this, since the literature is scarce on the effects of simplifying informed consent forms in the clinical setting, we analyzed several variables that might correlate with patient understanding. Age, sex, and time between consent and questionnaire administration were not found to be statistically significant predictors of increased understanding. The only factor that had a statistically significant effect on reducing the risk of responding incorrectly to a questionnaire item was educational level ($P = 0.048$ on multivariate analysis). This was particularly evident for the patients with a high school ($P = 0.035$ on multivariate analysis) or college education ($P = 0.043$ on multivariate analysis) in both groups. This finding is in line with observations by Kusec et al. [7] and suggests that the complexity of medical information can be appropriately graded in patients with a higher educational level.

That said, we believe that, irrespective of exposure to a standard verbal consent procedures or to a consent process with a simplified, graphically formatted consent document, understanding informed consent is influenced by a patient's health literacy. Voluntariness and interest in their health condition may help patients retain the concepts discussed with their surgeon during the consent process and prompt them to ask for the information they

feel is important when faced with decisions about their health.

Potential and real conflicts of interest: All authors have no conflict of interest to declare.

Sources of funding: All authors have no source of funding to declare.

References

- [1] Faden RR, Beauchamp TL. A history and theory of informed consent. Oxford: Oxford University Press, 1986
- [2] Ferrarese A., Pozzi G., Borghi F., Pellegrino L., Di Lorenzo P., Amato B., Santangelo M., Niola M., Martino V., Capasso E., Informed consent in robotic surgery: quality of information and patient perception. *Open Med.* 2016; 11 (1): 279-285
- [3] Chate RAC. An audit of the level of knowledge and understanding of informed consent amongst consultant orthodontists in England, Wales and Northern Ireland. *Br Dent J* 2008; 205:12
- [4] Scheer AS, O'Connor AM, Chan BP et al. The myth of informed consent in rectal cancer surgery: what do patients retain?. *Dis Colon Rectum* 2012; 55(9):970-975
- [5] Bernat JL, Peterson LM. Patient-Centered Informed Consent in Surgical Practice. *Arch Surg* 2006; 141:86-92
- [6] Akkad A, Jackson C, Kenyon S et al. Patients' perceptions of written consent: questionnaire study. *BMJ* 2006; 333:528
- [7] Kusec S, Oreskovic S, Skegro M. Improving comprehension of informed consent. *Patient Educ Couns* 2006; 60:294-300
- [8] Degerliyurt K, Laskin DM. Informed Consent: What Do Patients Really Want to Know? *J Oral Maxillofac Surg* 2010; 68:1849-52
- [9] Falagas ME, Korbila IP. Informed consent: how much and what do patients understand? *Am J Surg* 2009; 198:420-443
- [10] Bollschweiler E, Apitzsch J. Improving Informed Consent of Surgical Patients Using a Multimedia-Based Program? Results of a Prospective Randomized Multicenter Study of Patients Before Cholecystectomy. *Ann Surg* 2008; 248:2
- [11] Tompsett E. Can video aids increase the validity of patient consent? *Aust N Z J Obstet Gynaecol* 2012; 32:680-682
- [12] Ibrahim T, Meng Ong S. The new consent form: is it any better?. *Ann R Coll Surg Engl* 2004; 86:206-209
- [13] Wang C, Ammon P, Beischer AD. The use of multimedia as an adjunct to the informed consent process for Morton's neuroma resection surgery. *Foot Ankle Int* 2014; 35(10):1037-1044
- [14] Batuyong ED, Jowett AJ, Wickramasinghe N et al. Using multimedia to enhance the consent process for bunion correction surgery. *ANZ J Surg* 2014; 84(4):249-254
- [15] Sherlock A, Brownie S. Patients' recollection and understanding of informed consent: a literature review. *ANZ J Surg* 2014; 84(4):207-210
- [16] Nehme J, El-Khani U, Chow A, et al. The use of multimedia consent programs or surgical procedures: a systematic review. *Surg Innov* 2013; 20(1):13-23

- [17] Hung S, Huang K. An empirical study of the effectiveness of multimedia disclosure of informed consent: A technology mediated learning perspective. *Inf Manage* 2011; 48:135-144
- [18] Clark S, Mangram A. The Informed Consent: A Study of the Efficacy of Informed Consents and the Associated Role of Language Barriers. *J Surg Educ* 2011; 1931-7204
- [19] Kondziolka DS. Improving the informed consent process for surgery. *Neurosurgery* 2006; 58:1184-1189
- [20] Fink AS, Prochazka AV, Henderson WG, et al. Enhancement of surgical informed consent by addition of repeat back: a multicenter, randomized controlled clinical trial. *Ann Surg* 2010; 252(1):27-36
- [21] Mulsow JJ, Feeley TM, Tierney S. Beyond consent-improving understanding in surgical patients. *Am J Surg* 2012; 203(1):112-120
- [22] Betti S, Sironi A, Saino G, et al. Effect of the informed consent process on anxiety and comprehension of patients undergoing esophageal and gastrointestinal surgery. *J Gastrointest Surg* 2011; 15(6):922-927
- [23] Moseley TH, Wiggins MN, O'Sullivan P. Effects of presentation method on the understanding of informed consent. *Br J Ophthalmol* 2006; 90:990-993
- [24] Fink AS, Prochazka AV, Henderson WG, et al. Predictors of comprehension during surgical informed consent. *J Am Coll Surg* 2010; 210(6):919-926
- [25] Azoulay D, Castaing D, Dennison A et al. Transjugular intrahepatic portosystemic shunt worsens the hyperdynamic circulatory state of the cirrhotic patient: preliminary report of a prospective study. *Hepatology* 1994 Jan; 19(1):129-132
- [26] Pozzi G, Ferrarese A, Busso M et al. Percutaneous drainage and sclerosis of mesenteric cysts: literature overview and report of an innovative approach. *Int J Surg* 2014; 12 Suppl 2:S90-93
- [27] Ferrarese A, Marola S, Surace A et al. Fibrin glue versus stapler fixation in laparoscopic transabdominal inguinal hernia repair: a single center 5-year experience and analysis of the results in the elderly. *Int J Surg* 2014; 12 Suppl 2:S94-98
- [28] Surace A, Marola S, Benvenga R, et al. Difficult abdominal access in laparoscopic cholecystectomy in elderly patients: our experience and literature review. *Int J Surg* 2014; 12 Suppl 2:S1-3
- [29] Polistena A., Di Lorenzo P., Sanguinetti A., Buccelli C., Conzo G., Conti A., Niola M., Avenia N. Medicolegal implications of surgical errors and complications in neck surgery: a review based on the Italian current legislation. *Open Med.* 2016; 11 (1): 298-306
- [30] Gentile V, Ferrarese A, Marola S, et al. Perioperative and postoperative outcomes of perforated diverticulitis Hinchey II and III: open Hartmann's procedure vs. laparoscopic lavage and drainage in the elderly. *Int J Surg* 2014; 12 Suppl 2:S86-89
- [31] Martino V, Ferrarese A, Borello A et al. An unusual evolution of a case of Klippel-Trenaunay Syndrome. *Open Med.* 2015; 10:498-501
- [32] Sandrucci S, Garrone C, Mobiglia C et al. Evaluation of the toxicity induced in rat by the intra-arterial cytostatic infusion and by hepatic dearterialization associated with systemic cytostatic therapy. *Bollettino e Memorie della Societa Piemontese di Chirurgia* 1989; 59(2):65-77
- [33] Muzio S, Cassini P, Martino V et al. Transcystic videolaparoscopy for choledocholithiasis with holmium: YAG laser lithotripsy. A case report. *Chir Ital* 2008 Jan-Feb; 60(1):119-123
- [34] Serra R, Grande R, Butrico L et al. Effects of a new nutraceutical substance on clinical and molecular parameters in patients with chronic venous ulceration. *Int Wound J* 2016 Feb; 13(1):88-96
- [35] Surace A, Ferrarese A, Benvenga R et al. ACTH-secreting neuroendocrine pancreatic tumor: a case report. *Int J Surg* 2014; 12 Suppl 1:S222-224
- [36] Ferrarese A, Borello A, Gentile V, et al. Meso-pancreatectomy for pancreatic neuroendocrine tumor. *Int J Surg* 2014; 12 Suppl 1:S123-125
- [37] Serra R, Gallelli L, Conti A, et al. The effects of sulodexide on both clinical and molecular parameters in patients with mixed arterial and venous ulcers of lower limbs. *Drug Des Devel Ther* 2014 May; 13;8:519-527
- [38] Priola AM, Priola SM, Volpicelli G, et al. Accuracy of 64-row multidetector CT in the diagnosis of surgically treated acute abdomen. *Clin Imaging* 2013 Sep-Oct; 37(5):902-907
- [39] Ferrarese A, Enrico S, Solej M et al. Laparoscopic management of non-midline incisional hernia: A multicentric study. *Int J Surg* 2016 Jun 21; pii: S1743-9191(16)30181-9. doi: 0.1016/j.ijisu.2016.06.023. [Epub ahead of print]
- [40] Surace A, Ferrarese A, Marola S et al. Endorectal ultrasound in the diagnosis of rectal cancer: accuracy and criticisms. *Int J Surg* 2014; 12 Suppl 2:S99-102
- [41] Berti S, Ferrarese A, Feleppa C, et al. Laparoscopic perspectives for distal biliary obstruction. *Int J Surg* 2015 Sep; 21 Suppl 1:S64-67
- [42] Surace A, Ferrarese A, Marola S et al. Abdominal compartment syndrome and open abdomen management with negative pressure devices. *Ann Ital Chir.* 2015 Jan-Feb; 86(1):46-50
- [43] Ferrarese A, Falcone A, Solej M et al. Surgeon's clinical valuation and accuracy of ultrasound in the diagnosis of acute appendicitis: A comparison with intraoperative evaluation. Five years experience
- [44] Marola S, Ferrarese A, Solej M et al. Management of venous ulcers: State of the art. *Int J Surg* 2016 Jun 21; in press
- [45] Allaix ME, Giraudo G, Ferrarese A, et al. 10-Year Oncologic Outcomes After Laparoscopic or Open Total Mesorectal Excision for Rectal Cancer. *World J Surg* 2016 Jul 14; in press
- [46] Eretta C, Ferrarese A, Olcese S, et al. Celiac axis compression syndrome: laparoscopic approach in a strange case of Chronic abdominal pain in 71 years old man. *Open Med* 2016; 11: In press
- [47] Nano M, Martino V, Ferrarese A, Falcone A. A brief history of laparoscopy. *G Chir* 2012; (33); 3:53-57
- [48] Martino V, Ferrarese A, Bindi M, et al. Abnormal right hepatic artery injury resulting in right hepatic atrophy: diagnosed by laparoscopic cholecystectomy. *Open Med* 2015; 10:535-537
- [49] Ferrarese A, Pozzi G, Borghi F, et al. Malfunctions of robotic system in surgery: role and responsibility of surgeon in legal point of view. *Open Med* 2016; 11: In press
- [50] Ferrarese A, Pozzi G, Borghi F, et al. Informed consent in robotic surgery: quality of information and patient perception. *Open Med* 2016; 11: In press
- [51] Ferrarese A, Solej M, Enrico S, et al. Elective and emergency laparoscopic cholecystectomy in the elderly: our experience. *BMC Surg* 2013; 13 Suppl 2:S21
- [52] Ferrarese A, Solej M, Enrico S, et al. Diagnosis of incidental gallbladder cancer after laparoscopic cholecystectomy: our experience. *BMC Surg* 2013; 13 Suppl 2:S20

- [53] Ferrarese A, Enrico S, Solej M, et al. Transabdominal pre-peritoneal mesh in inguinal hernia repair in elderly: end point of our experience. *BMC Surg* 2013; 13 Suppl 2:S24
- [54] Ferrarese A, Martino V, Enrico S et al. Laparoscopic repair of wound defects in the elderly: our experience of 5 years. *BMC Surg* 2013; 13 Suppl 2:S23
- [55] Ferrarese A, Martino V, Enrico S, et al. Laparoscopic appendectomy in the elderly: our experience. *BMC Surg* 2013; 13 Suppl 2:S22
- [56] Solej M, Martino V, Mao P, et al. Early versus delayed laparoscopic cholecystectomy for acute cholecystitis. *Minerva Chir* 2012 Oct; 67(5):381-387
- [57] Ferrarese A, Martino V, Falcone A, et al. Diverticoli duodenali perforati: caso clinico e breve review della letteratura. *Chirurgia* 2014 Aprile; 27(2):129-131