

Incomplete Immune Reconstitution and Traditional Chinese Medicine in Patients with HIV/AIDS: Challenges and Perspectives

Xue Ding^{1,*}, Leilei Fan^{2,*}, Liran Xu³, Xiuxia Ma⁴, Pengfei Meng⁴, Jie Li⁴, Jiahe Li³, Jingyu Yue⁴

¹Department of Medicine, The First Affiliated Hospital of Henan University of Chinese Medicine, Zhengzhou, People's Republic of China;

²Department of Cardiovascular, The First People's Hospital of Zhengzhou, Zhengzhou, People's Republic of China; ³Department of the First Clinical Medical College, Henan University of Chinese Medicine, Zhengzhou, People's Republic of China; ⁴Department of AIDS Clinical Research Center, The First Affiliated Hospital of Henan University of Chinese Medicine, Zhengzhou, People's Republic of China

*These authors contributed equally to this work

Correspondence: Jingyu Yue, Department of AIDS Clinical Research Center, The First Affiliated Hospital of Henan University of Chinese Medicine, 19 Renmin Road, Zhengzhou, People's Republic of China, Email yuejingyu@aliyun.com

Abstract: Antiretroviral therapy can reduce human immunodeficiency virus (HIV) load to undetectable levels and restore CD4+ T cells to rebuild immune function in patients with HIV. However, some patients fail to achieve immune reconstitution despite treatment. Traditional Chinese medicine is an important branch of complementary and alternative medicine for the treatment of HIV infection, and a growing number of studies has demonstrated that traditional Chinese medicine can increase CD4+ T cell counts in patients, thereby promoting immune reconstitution, ameliorating symptoms and signs, and improving quality of life. Here, we review pathogenesis in immunological non-responders and research into their treatment with traditional Chinese medicine. Furthermore, we summarize potential future research directions, including elucidation of how traditional Chinese medicine can regulate CD4+ T cells to reduce opportunistic infections and improve quality of life in immunological non-responders.

Keywords: immunological non-responders, drug mechanism, traditional Chinese medicine, future research directions

Introduction

Despite the introduction of effective antiretroviral therapy (ART), which successfully controls human immunodeficiency virus (HIV) replication, 10–40% of people living with HIV (PLWH) fail to recover their CD4+ T lymphocyte counts. Those who experience this incomplete immune reconstitution are termed “immunological non-responders” or “immune non-responders” (INR).^{1,2} Low levels of CD4+ T cell recovery after ART are significantly associated with the occurrence of non-AIDS-related events.³ INR have an elevated risk of non-AIDS-related morbidity and mortality compared with PLWH who achieve complete immune reconstitution.⁴

Traditional Chinese medicine (TCM), which typically utilizes plants with complex components in specific proportions and dosages, is often used to treat infectious diseases. Tu Youyou isolated artemisinin from *Artemisia annua* L., which effectively suppressed the spread of malaria. The ability of TCM to effectively improve CD4+ T cell counts with surprisingly long-term effects when used as a means to treat AIDS—known as “Yi Du” in TCM—is garnering increasing attention.

Numerous single herbs and TCM formulae capable of enhancing CD4+ T cell numbers and immunological function have been listed in medical texts.⁵ Multi-target treatment with TCM can effectively improve the effect of ART. A body of clinical data supports the increasing use of TCM in the treatment of INR.⁶ However, the main challenges to expanding its use are the needs to further improve the efficacy of TCM for the treatment of INR and to understand the mechanisms of action. We will discuss the advantages of integrating TCM and ART to gain in-depth understanding of the mechanisms of TCM in immune restoration and the prevention of acquired immunodeficiency syndrome (AIDS)-related events.

Potential Mechanism Underlying Incomplete Immune Reconstitution

The underlying mechanisms for incomplete immune recovery in INR are complex and may be multifactorial—changes in the bone marrow (BM) microenvironment, decreased thymic output, disturbed cytokine secretion, aberrant immune activation, residual viral replication, gut damage and associated microbial translocation, and specific genetic or metabolic characteristics can all lead to abnormal CD4⁺ T cell numbers and function in patients with AIDS.^{7–10} The low CD4⁺ T cell counts in INR appear to be mainly caused by an imbalance in the production and destruction of CD4⁺ T cells, such that damaged CD4⁺ T cells cannot be effectively replenished.¹¹

BM Microenvironment

CD4⁺ T cells are derived from hematopoietic cells in the BM, and both multilineage and lineage-restricted hematopoietic progenitors are decreased in the BM after HIV infection, leading to a decrease in the number of CD4⁺ T cells.^{12–14} HIV-infected BM cannot provide the optimal milieu for the growth of hematopoietic colonies. Moreover, infected BM secretes TNF- α , which induces apoptosis through the Fas ligand–FAS receptor pathway in the BM and inhibits hematopoiesis.¹⁵ BM cells in INR secrete high levels of TNF- α and show significant upregulation of Fas ligand and the receptor Fas. In addition, the interstitial layer of the BM of INR shows abnormal morphological features similar to those observed in BM obtained from PLWH without ART.^{16,17} INR widely lack colony formation by BM cells and display altered early lymphoid progenitors at the level of BM CD34⁺ precursors; these mechanisms may explain the cause of CD4⁺ T cell lymphopenia in INR.¹⁷

Thymic Output

The thymus is responsible for T cell development, including the production of CD4⁺ T cells, and thymic development is a primary mechanism of CD4⁺ T cell reconstitution.¹⁸ Evaluation of the ability of the thymus to export mature T lymphocytes often involves the quantification of T cell receptor excision circles in cells referred to as recent thymic emigrants (RTE).¹⁹ The thymus is most active in early life, and its function and thymogenic activity are limited in adulthood.²⁰ However, when an individual is infected with HIV-1, the thymus is reactivated and releases RTEs to replenish CD4⁺ T cell counts.²¹ Previous studies have shown that INR exhibit lower levels of RTEs than immunological responders, and a lower proportion of RTEs among CD4⁺ T cells, which indicates that thymic output is significantly lower in INR.²² This decrease in thymic activity is critical for maintaining the immunosuppressed state.²³

Cytokine Disorders

Cytokines—especially the common gamma chain cytokines, including interleukin (IL)-2, IL-4, IL-7, IL-9, IL-15, and IL-21—are important players in immune reconstruction. In particular, IL-2 and IL-7 contribute to T cell recovery owing to their abilities to reverse HIV-1-specific CD4⁺ T cell anergy and activate resting CD4⁺ T cells; their loss is associated with the progressive depletion of CD4⁺ T cells.^{24,25} Most IL-2 is produced by activated T cells, specifically CD4⁺ cells, and it acts on these cells in an autocrine or paracrine manner.²⁶ A central component of T cell homeostasis is signaling through the IL-7 receptor α chain (IL-7R α , CD127).²⁷ Loss of expression of the IL-7R α chain on CD4⁺ T cells has been associated with a decrease in CD4⁺ T cell count.²⁸ T helper 17 (Th17) cells secrete IL-17; they are preferentially targeted by HIV-1 infection, and Th17 cell depletion is associated with disease progression.²⁹ Therefore, cytokine disorders are not conducive to the recovery of immune function in INR.

Pyroptosis

Pyroptotic cells secrete their intracellular contents and pro-inflammatory cytokines, including IL-1 β and IL-18.³⁰ During HIV infection, pyroptosis enhances the host's ability to rapidly eradicate infection and inhibit HIV replication through the production of pro-inflammatory cytokines and endogenous danger signals. However, programmed cell death caused by HIV-induced inflammatory responses leads to the massive loss of CD4⁺ T cells, which may promote and sustain immunological non-responsiveness.³¹ The gp120 receptor, the only viral protein exposed on the surface of HIV, binds to CD4 on the surface of host cells, promotes the expression

of Fas by CD4+ cells, and increases intracellular Ca^{2+} , leading to cell apoptosis.^{32,33} Thus, pyroptosis may play a key role in the inability of the immune systems of INR to mount a protective response to HIV and associated therapies.

Immune Activation

During HIV-1 infection, persistent immune activation commonly occurs in INR, stimulated by the virus and non-AIDS-related events, which causes CD4+ T cell depletion.³⁴ Driving factors for HIV-related immune activation include the persistence of HIV reservoirs, intestinal microbial translocation, depletion of regulatory T (Treg) cells, and coinfection with other viruses.³⁴ Conversely, immune overactivation can maintain virus reservoirs, accelerate the depletion of Treg cells, aggravate mucosal barrier damage, and decrease CD4+ T cell numbers.³⁵ Overactivation of T cells, which is associated with decreases in CD4+ T cell counts, is an important mechanism of action in INR.

Intestinal Barrier and Microbiota

The intestinal lamina propria is the main site of HIV infection and replication. HIV replication through the intestinal barrier leads to intestinal flora disturbance and induces intestinal mucosal CD4+ T cell apoptosis.^{36,37} Depletion of CD4+ T cells decreases the numbers of important Treg cells in the intestinal mucosa, resulting in mucosal immune and functional deficits, thereby accelerating the progression of HIV.³⁸ INR exhibit lower CD4+ T cell counts, and more severe intestinal epithelial cell damage and tight junction protein changes, than immunological responders, indicating more severe intestinal damage and mucosal immune deficiency.³⁹ The gut microbiota is critical for maintaining immune homeostasis, and HIV infection can cause significant changes in the gut microbiota.^{40,41} Studies have demonstrated significant differences in the gut microbiota of INR and responders, with the families Succinivibrionaceae and Erysipelotrichaceae (known for their protective effects against intestinal inflammation) significantly enriched in responders, but not detected in INR.⁴² Therefore, it is particularly important to study the association between intestinal immune barrier function and microbial imbalance in INR.

Th17 and Treg Cells

T lymphocytes can differentiate into various lineages, including Treg and Th17 cells, which mediate inflammation and immune tolerance. Treg cell numbers in INR negatively correlate with Naive CD4+T cells compared with those in responders.⁴³ A balance between Th17 and Treg cells is necessary to facilitate protection against pathogens while maintaining immune stability.⁴⁴ An obvious imbalance between Th17 and Treg cells is observed in the peripheral blood and intestinal mucosa during HIV infection, which is related to the progression of AIDS.⁴⁵ Studies have reported that the Th17:Treg ratio is significantly lower in patients with low CD4+ T lymphocyte counts and lower still in INR.⁴⁶ Therefore, maintaining or reinstating the Th17:Treg ratio may be an important strategy for successful immune reconstruction.

TCM and INR

Since INR is an neglected group associated with increased morbidity and mortality due to immunodeficiency and immune activation. TCM has the effect of multi-target and bidirectional regulation, which can promote the immune reconstitution of INR.

Mechanism of TCM in INR

The pathogenesis of INR is complex, and individual patients have different pathogenesis, so the therapy should be individualized according to different pathogenesis. We explain the various mechanisms of Chinese herbal medicines in treating INR and provide a reference for the advancement of TCM in treating INR (Figure 1).

Reduced the T Cell Activation, Pyroptosis and Promote Immune Reconstruction

INR is unable to achieve adequate immunologic recovery, in which immune activation plays a crucial role. Continuous immune activation affects the proliferation of CD4+T cells and accelerates cell proptosis. The current study results confirmed that TCM can reduce the T cell activation and proptosis, promote the recovery of CD4 + T cells and immune function in INR




Figure 1 Summary of the potential mechanism underlying incomplete immune reconstitution and the role of traditional Chinese medicine in immunological non-responders.

patients. Several studies on TCM in the treatment of INR have been completed, demonstrating the potential advantages of treating INR. *Tripterygium wilfordii* Hook F (TwHF) is a Chinese herbal medicine, its main bioactive component triptolide (TPL) has been extracted, and modified into a novel compound (5R)-5-hydroxytriptolide (LLDT-8) with immunosuppressive activity and reduced toxicity. The application of TwHF pill and LLDT-8 has been clinically proved efficacy drugs in reduced the T cell activation, promoted the circulating CD4⁺ T cell counts and alleviated inflammation among INR.^{47,48} Combined transcriptional, proteomic and network pharmacology approach reveals that TPL could inhibit the interferon (IFN) response and the phosphorylation of signal transducer and activator of transcription 1 (STAT1).⁴⁹ Artesunate, as a semi-synthetic artemisinin derivative of sesquiterpene lactone, is widely used in clinical immune-related diseases treatment due to its immunomodulatory activity and reduced toxicity. Considering its immunomodulatory properties, researchers applied Artesunate in treating INR. The application of Artesunate in 50mg once daily or 50mg twice a day for 48 weeks successfully improved the initial T cell count level, promoted the circulating CD4⁺ T cell counts and immune reconstitution among INR.⁵⁰ They also launched a multi-centered, double-blinded clinical trial to validate these findings clinical effects of Artesunate among 269 INR.⁵¹ Subsequent studies demonstrated that Artesunate could reduce the early apoptosis of CD4⁺ T lymphocytes and increase the abundance of *Verrucomicrobiota*.⁵²

Regulate Intestinal Barrier, Microbes and the Treg/Th17 Balance

Ectopic intestinal flora and intestinal mucosa damage lead to the persistence of abnormal immune activation, which affects the immune reconstruction in INR. The diversity of intestinal flora in patients with poor immune reconstruction is low. Explore the immune reconstruction by regulating intestinal barrier and microbes is of great significance for

improving immune function. Yiaikang capsule is composed of Ginseng, Astragalus membranaceus, Atractylodes rhizoma, et al, which has definite curative effect in the treatment of AIDS. Yiaikang Capsule can increase the CD4+T cell count, regulate Treg/Th17 balance, and enhance the abundance of probiotics in the intestinal flora among 18 INR.⁵³ Yiaikang Capsule inhibited the expression of tight junction protein ZO-1, maintained the normal tight junction, reduced the permeability of intestinal mucosal barrier induced by IFN- γ , and maintained the integrity of intestinal mucosal barrier.⁵⁴ Shenling Guben granule is composed of Ganoderma lucidum, American ginseng, Knotweed, etc. which can enhance the immune function of INR, increase the CD4+T lymphocyte count, and promote the growth of Sarteria.

Regulate Cytokines Disorders

Abnormal immune activation leads to a high inflammatory response and releases interleukin (IL), including IL-1, IL-6, IL-8, IL-10, tumor necrosis factor (TNF) and other cytokines and chemokines, which increases the difficulty of immune reconstitution. Toll-like receptors can promote the maturation of immune cells by promoting the synthesis and release of cytokines. Yiqi Jianpi granules are composed of Astragalus, Poria cocos, Wolfberry fructus, etc., which can increase CD4+ cell count by affecting the activation of Toll-like receptors and up-regulating the expression of IL-2 in INR.^{55,56} Therefore, TCM can regulate the balance of pro-inflammatory factors and anti-inflammatory factors, alleviate inflammatory response, and promote immune reconstruction.

Increased CD4+ T Cell Counts

The number of CD4+CD45RA+T cells produced by the thymus is considered to be an effective predictor of INR, with the number of CD45RA+T cells in immune responders significantly higher than that in non-responders.⁵⁷ CD4+CD45RA+ and CD4+CD45RO+T cells express two important subtypes of CD45 that are closely related to the regeneration of thymic function and immune responsiveness; their expression and abundance can reflect changes in cellular immunity.^{58,59} Clinical research supports the ability of TCM to improve CD4+T lymphocyte counts and CD4+CD45RA+T cell numbers in INR and promote immune reconstitution and improve patient quality of life⁶⁰ (Table 1).

Table 1 The Link Between Chinese Herbal Medicine and Immunological NonResponders

Chinese Herb Medicine	Sample	Research Design	Effect on Immune Reconstitution	Ref.
Inigorating spleen and tonifying kidney method	32	Randomized, double-blind, controlled	The proportion of CD4+CD45RA+T cells and the ratio value of CD4+CD45RA+T/CD4+CD45RO+T cells \uparrow the proportion of CD4+CD45RO+T \downarrow	[61,62]
Artesunate	269	Multicenter, prospective clinical trial	CD4+ T lymphocytes count, CD45RA+ T subsets count, The ratio of CD4+ Ki67+ /CD4+ \uparrow , IR \uparrow	[63–65]
Aikeqing Granule	70	Randomized, controlled	CD4+T lymphocyte count \uparrow , effective rate of immune reconstitution \uparrow ,QoL \uparrow	[66,67]
Fuzheng Kangdu Wan	180	Pretherapy and post-treatment	CD4+T lymphocyte count \uparrow , immune reconstitution \uparrow , clinical symptoms and immune function \uparrow	[68–70]
Wenshen Jianpi Granule	60	Randomized, double-blind, controlled	CD4+ T lymphocytes count, CD45RA+ T cells \uparrow , immune reconstitution \uparrow	[71,72]
Bushenfang	41	Randomized, controlled	CD4+T lymphocyte counts \uparrow , CD4+/CD8+ \uparrow , IR \uparrow , clinical symptoms \uparrow	[73]
Yiaikang Capsules	83	Pretherapy and post-treatment	CD4+T lymphocyte counts \uparrow , CD4+/CD8+ \uparrow , IR \uparrow , clinical symptoms \uparrow , quality of life \uparrow	[74,75]

(Continued)

Table 1 (Continued).

Chinese Herb Medicine	Sample	Research Design	Effect on Immune Reconstitution	Ref.
Jianpi Yiqi Formula	62	Randomized, controlled	CD4+T lymphocyte counts↑, IR ↑	[76]
Diwu Yanggan Capsule	57	Randomized, double-blind, controlled	CD45RA+↑, CD4+T↑, IR ↑	[77]
Astragalus membranaceus	30	Randomized, controlled	CD4+T lymphocyte counts↑	[78,79]
Yiqi Jianpi Granules	22	Pretherapy and post-treatment	CD4+T lymphocyte counts↑	[80]
Ganoderma lucidum	92	Retrospective study	CD4+T lymphocyte counts↑, Th/Ts ↑	[81]
Mianyi NO.2	264	Randomized, double-blind, controlled	CD4+ T lymphocyte counts, CD4CD45RA+, CD4CD45RO +↑, clinical symptoms ↑	[82]
Tripterygium Wilfordii Hook F	33	Pretherapy and post-treatment	CD4 + T cell counts and recovery↑, T cell activation↓	[47,49]

TCM Interventions Against Opportunistic Infections

The incidence of immunological non-responsiveness is 10%–40%, and affected patients are at significantly higher risk for opportunistic infections (OIs).⁸³ The prevention and treatment of OIs is important to improve the quality of life of PLWH. Clinical studies on OIs show that TCM is efficacious for reducing OIs—such as AIDS-associated lung infections, AIDS-related diarrhea, AIDS-associated fevers, AIDS-associated skin injuries, and AIDS-associated oral lesions—and improving quality of life⁸⁴ (Table 2). TCM has preventative and curative effects for OIs, but some OIs, such as tuberculosis, lymphoma, and Kaposi's sarcoma, still require effective treatments.

Potential Future Directions

Elucidate the Potential Mechanisms Through Which TCM Regulates the CD4+ T Cells of INR

Changes in the number and function of CD4+ T cells are critical to the progression of AIDS. Although TCM therapy can effectively increase CD4+ T cell counts, and represents a promising approach to treating INR, its mechanisms of

Table 2 Chinese Herbal Medicine and Opportunistic Infections

Opportunistic infections	Chinese Herb Medicine	Sample	Research Design	Consequence	Potential Mechanisms	Ref.
AIDS-associated lung infections	Qingfei Peiyuan Granules	141	Randomized, double-blind, controlled	Cough, wheezing, chest tightness, phlegm↓, QoL↑	Regulates T lymphocyte subsets, Th1/Th2, TLR/MyD88, and IL-6, PCT↓,	[85–88]
	Fuzheng Qingfei Decoction	106	Randomized, controlled	Cough↓, phlegm↓, fever↓	CD 4+T↑	[89]
	Tanreqing injection	100	Randomized, controlled	Temperature↓, hemameba↓, neutrophile granulocyte↓	CD 4+T↑, CD8+↑, natural killer cell↑	[90]

(Continued)

Table 2 (Continued).

Opportunistic infections	Chinese Herb Medicine	Sample	Research Design	Consequence	Potential Mechanisms	Ref.
AIDS-related diarrhea	Xieli Kang Capsule	130	Randomized, controlled	Diarrhea, abdominal distension and pain↓	Regulate TNF- α , secreted Ig A \uparrow , IL-17 \uparrow , CD40L \downarrow ,	[91–93]
	Jianpi Wenyang Jiedu granule	40	Randomized, controlled	Diarrhea \downarrow , Stomachache \downarrow , Bloating \downarrow , anorexia \downarrow	Acidaminococcus \downarrow , Oscillospira \downarrow , Dorea \uparrow , Coprococcus_3 \uparrow , NF- κ B \downarrow , IL-1 β \downarrow , TNF- α \downarrow , IL-6 \downarrow	[94]
	Guchang Zhixie Pills	86	Randomized, controlled	Diarrhea \downarrow , Stomachache \downarrow ,	NR	[95]
AIDS-associated fevers	Jiebiao Qingli granules, Shenqi Jiere capsules and Chaihopedayuan mixture	270	Randomized, controlled	Fever \downarrow	IL-1 \downarrow , IL-6 \downarrow , TNF- α \downarrow , hemameba \downarrow , C-reactive protein \downarrow ,	[96]
	Modified minor radix bupleuri decoction	90	Randomized, controlled	Fever \downarrow	NR	[97]
	Buzhong Yiqi decoction	40	Randomized, controlled	Fever \downarrow	NR	[98]
	Lianhua Qingwen Decoction	120	Randomized, controlled	Fever \downarrow Fatigue \downarrow	NR	[99]
AIDS-associated skin injuries	Modified Longdan Xiegan Decoction	149	Randomized, double-blind, controlled	Pruritus and skin lesions \downarrow , QoL \uparrow	The eosinophil count and serum IgE \downarrow	[100]
	Kangai Baosheng Pills, Fuzheng Kangdu Pills	34	Randomized, controlled	Rash \downarrow Ulceration \downarrow	NR	[101]
	Luhui Pikang Liquid	58	Pretherapy and post-treatment	Pruritus \downarrow , rash \downarrow , ulceration \downarrow	NR	[102]
AIDS-associated oral lesions	Licorice heart-draining decoction, Xiaolu Granules	78	Randomized, controlled	Ulcer pain and area \downarrow , QoL \uparrow	Regulation of oral flora	[103,104]
	Chinese herbal gargle	80	Randomized, controlled	Leukoplakia \downarrow , Ulceration \downarrow , QoL \uparrow	Regulating oral candida	[105]

Abbreviations: AIDS, acquired immunodeficiency syndrome; QoL, quality of life, NR, not reported.

action in immune recovery remain to be elucidated. A synergistic approach utilizing single-cell sequencing, transcriptomics, proteomics, and metabolomics can effectively analyze the biological processes and molecular interactions involved in the development of disease. We believe a similar synergistic approach applied to the analysis of blood, gut, and stool samples in the context of TCM may reveal key information. Priority should be given to the gut microbiota because some studies have shown that the gut microbiota is closely related to CD4 $^+$ T cell numbers and immune reconstitution during the resolution of poor immune reconstitution upon treatment with TCM.^{74,106–108} Chen et al best illustrated this approach when they explored the mechanism of Yiaikang capsules in the treatment of INR using serum metabolomics. They compared the metabolomics results of 58 patients treated with Yiaikang capsules before and after treatment, and concluded that Yiaikang capsules increased CD4 $^+$ T cell counts and the CD4 $^+$:CD8 $^+$ T cell ratio in INR. Metabolites such as 1-phosphatidylyl-D-mercaptinositol, PC(18:3 (9Z,12Z,15Z)/20:1 (11Z)), PE(14:0/P-18:0),




Figure 2 Mechanism outline of traditional Chinese medicine on treating infectious diseases in incomplete immune reconstitution.

Abbreviation: TCM, traditional Chinese medicine.

quinolinic acid, deoxycorticosterone, and selenocysteine are closely related to the therapeutic effects of Yiaikang capsules. This metabolic pathway may be a potential therapeutic target of Yiaikang capsules in the treatment of INR.¹⁰⁹ Researchers who study TCM to prevent and treat immunological non-responsiveness are gaining insight into potential research targets. Moreover, in-depth analysis of gene upregulation and/or other changes in genes during TCM treatment of INR may provide a clearer understanding of the role of TCM in the recovery of CD4⁺ T cells and immune responses in INR.

Reduce OIs and Improve Quality of Life

Declines in CD4⁺ T lymphocyte counts can increase the risk of OIs, and infected hosts are more susceptible to a variety of OIs, usually caused by fungi, bacteria, viruses, and parasites that are typically controlled by the immune system. OIs are the leading cause of death in PLWH, and the key to AIDS treatment is to accurately evaluate the changes in a patient's condition and reduce the incidence of poor prognosis. Therefore, it is essential to find effective treatments for HIV OIs. TCM treatment is holistic and individual; it has obvious advantages for the clinical treatment of low immunity accompanied by OIs, and has achieved clinical efficacy.¹¹⁰ We believe that early intervention with integrated TCM and Western medicine for the prevention and treatment of OIs will improve clinical efficacy and reduce the recurrence rate and mortality (Figure 2).

Conclusion

INR is an neglected group, we explored the pathogenesis of INR, but it is also associated with older age, male, low level of pre-ART CD4⁺ T lymphocytes, ethnic factors, etc. There is no method to fully restore the CD4⁺ T lymphocyte count. TCM therapy has potential advantages in INR, and its mechanism is related to the reduction of immune activation, pyroptosis and the regulation of cytokine disorders, intestinal barrier and microbiota, Th17 and treg17. TCM can regulate intestinal microecology, and intestinal flora is closely related to the number of CD4⁺ T cells and immune reconstitution, so it should be given priority. Comprehensive analysis of single-cell RNA sequencing, transcriptomics, genomics and proteomics can provide an in-depth understanding of INR. Analyzing the role of TCM therapy in INR using the multi-omics method can accelerate the research of TCM promoting immune reconstitution. In the TCM therapy of INR, it is of great significance to elucidate the possible mechanism of TCM regulating INR CD4⁺ T cells and reducing OIs and improving quality of life. There are limitations in the TCM therapy of INR. First, it will be necessary to screen and analyze the active ingredients of TCM products and their therapeutic targets. Second, combined with serum pharmacology, systems biology and bioinformatics technology must be applied to verify the research results. As this area of research continues to evolve, we hope that TCM will use these technologies to overcome its own limitations to achieve a more comprehensive treatment of complex diseases. We recommend that researchers use a synergistic omics approach to study the key problems restricting clinical efficacy in INR and to develop Chinese herbs with clear mechanisms and precise efficacy.

Funding

This work was supported by Henan Province Special Project of Traditional Chinese Medicine Scientific research (2021JDZX2078, 2022JDZX154, 2023ZY2204, 2023ZXZX1057), National Natural Science Foundation of China, Youth Science Foundation Project (82104559), National Administration of Traditional Chinese Medicine of China, 2022 Zhang Zhongjing Inheritance and innovation project (GZY-KJS-2022-041-1), Zhengzhou Medical and health science and technology Innovation guidance Program (2024Y LZDJH134), Henan Province Key Research and Development and promotion Project (232102311222).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Yang X, Su B, Zhang X, Liu Y, Wu H, Zhang T. Incomplete immune reconstitution in HIV/AIDS patients on antiretroviral therapy: challenges of immunological non-responders. *J Leukocyte Biol.* 2020;107(4):597–612. doi:10.1002/jlb.4mr1019-189r
2. Rb-Silva R, Goios A, Kelly C, et al. Definition of immunological nonresponse to antiretroviral therapy: a systematic review. *J Acquir Immune Defic Syndr.* 2019;82(5):452–461. doi:10.1097/qai.0000000000002157
3. Hsu DC, Sereti I, Ananworanich J. Serious Non-AIDS events: immunopathogenesis and interventional strategies. *AIDS Res Ther.* 2013;10(1):29. doi:10.1186/1742-6405-10-29
4. Kelley CF, Kitchen CM, Hunt PW, et al. Incomplete peripheral CD4+ cell count restoration in HIV-infected patients receiving long-term antiretroviral treatment. *Clin Infect Dis.* 2009;48(6):787–794. doi:10.1086/597093
5. Wang J, Liang B, Zhang X, et al. An 84-month observational study of the changes in CD4 T-lymphocyte cell count of 110 HIV/AIDS patients treated with traditional Chinese medicine. *Front Med.* 2014;8(3):362–367. doi:10.1007/s11684-014-0363-x
6. Tao Z, Huang XJ, Liu Y, et al. Efficacy of integrated traditional Chinese Medicine and anti-retroviral therapy on immunological nonresponse in patients with human immunodeficiency virus/acquired immunodeficiency syndrome: a meta-analysis of randomized controlled trial. *J Trad Chin Med.* 2021;41(5):669–676. doi:10.19852/j.cnki.jtcm.2021.05.002
7. Rodríguez-Gallego E, Gómez J, Pacheco YM, et al. A baseline metabolomic signature is associated with immunological CD4+ T-cell recovery after 36 months of antiretroviral therapy in HIV-infected patients. *AIDS.* 2018;32(5):565–573. doi:10.1097/qad.0000000000001730
8. Gunda DW, Kilonzo SB, Kamugisha E, Rauya EZ, Mpondo BC. Prevalence and risk factors of poor immune recovery among adult HIV patients attending care and treatment centre in northwestern Tanzania following the use of highly active antiretroviral therapy: a retrospective study. *BMC Res Notes.* 2017;10(1):197. doi:10.1186/s13104-017-2521-0
9. Yan J, Ouyang J, Isnard S, et al. Alcohol use and abuse conspires with HIV infection to aggravate intestinal dysbiosis and increase microbial translocation in people living with HIV: a review. *Front Immunol.* 2021;12:741658. doi:10.3389/fimmu.2021.741658
10. Guo XY, Guo YT, Wang ZR, et al. Severe intestinal barrier damage in HIV-infected immunological non-responders. *Heliyon.* 2023;9(10):e20790. doi:10.1016/j.heliyon.2023.e20790
11. Gaardbo JC, Hartling HJ, Gerstoft J, Nielsen SD. Incomplete immune recovery in HIV infection: mechanisms, relevance for clinical care, and possible solutions. *Clinic Develop Immunol.* 2012;2012:670957. doi:10.1155/2012/670957
12. Hoang TN, Harper JL, Pino M, et al. Bone marrow-derived CD4(+) T cells are depleted in simian immunodeficiency virus-infected macaques and contribute to the size of the replication-competent reservoir. *J Virol.* 2019;93(1). doi:10.1128/jvi.01344-18
13. van Elsacker-Neile AM, Kroon FP, van der Ende ME, Salimans MM, Spaan WJ, Kroes AC. Prevalence of parvovirus B19 infection in patients infected with human immunodeficiency virus. *Clin Infect Dis.* 1996;23(6):1255–1260. doi:10.1093/clinids/23.6.1255
14. Koka PS, Reddy ST. Cytopenias in HIV infection: mechanisms and alleviation of hematopoietic inhibition. *Curr HIV Res.* 2004;2(3):275–282. doi:10.2174/1570162043351282
15. Maciejewski J, Selleri C, Anderson S, Young NS. Fas antigen expression on CD34+ human marrow cells is induced by interferon gamma and tumor necrosis factor alpha and potentiates cytokine-mediated hematopoietic suppression in vitro. *Blood.* 1995;85(11):3183–3190. doi:10.1182/blood.V85.11.3183.bloodjournal85113183
16. Isgrò A, Aiuti A, Mezzaroma I, et al. Improvement of interleukin 2 production, clonogenic capability and restoration of stromal cell function in human immunodeficiency virus-type-1 patients after highly active antiretroviral therapy. *Br J Haematol.* 2002;118(3):864–874. doi:10.1046/j.1365-2141.2002.03680.x
17. Badolato R. Immunological nonresponse to highly active antiretroviral therapy in HIV-infected subjects: is the bone marrow impairment causing CD4 lymphopenia? *Clin Infect Dis.* 2008;46(12):1911–1912. doi:10.1086/588481
18. Corbeau P, Reynes J. Immune reconstitution under antiretroviral therapy: the new challenge in HIV-1 infection. *Blood.* 2011;117(21):5582–5590. doi:10.1182/blood-2010-12-322453
19. Cunningham CA, Helm EY, Fink PJ. Reinterpreting recent thymic emigrant function: defective or adaptive? *Current Opin Immunol.* 2018;51:1–6. doi:10.1016/j.coi.2017.12.006
20. Kolte L. Thymic function in HIV-infection. *Danish Med J.* 2013;60(4):B4622.
21. Santos Guedes MC D, Carvalho-Silva WHV, Andrade-Santos JL, Brelaz-de-Castro MCA, Souto FO, Guimarães RL. Thymic exhaustion and increased immune activation are the main mechanisms involved in impaired immunological recovery of HIV-positive patients under ART. *Viruses.* 2023;15(2). doi:10.3390/v15020440
22. Xiao Q, Yan L, Han J, et al. Metabolism-dependent ferroptosis promotes mitochondrial dysfunction and inflammation in CD4(+) T lymphocytes in HIV-infected immune non-responders. *EBioMedicine.* 2022;86:104382. doi:10.1016/j.ebiom.2022.104382

23. Carvalho-Silva WH V, Andrade-Santos JL, Santos Guedes MC D, Guimarães RL. Genetics and immunological recovery with antiretroviral treatment for HIV. *Pharmacogenomics*. 2020;21(14):979–983. doi:10.2217/pgs-2020-0083
24. Gu Y, X X, FY KCM, Ostrowski MA. The role of cytokines which signal through the common gamma chain cytokine receptor in the reversal of HIV specific CD4(+) and CD8(+) T cell anergy. *PLoS One*. 2007;2(3):e300. doi:10.1371/journal.pone.0000300
25. Zeng M, Southern PJ, Reilly CS, et al. Lymphoid tissue damage in HIV-1 infection depletes naïve T cells and limits T cell reconstitution after antiretroviral therapy. *PLoS Pathogens*. 2012;8(1):e1002437. doi:10.1371/journal.ppat.1002437
26. Létourneau S, van Leeuwen EM, Krieg C, et al. IL-2/anti-IL-2 antibody complexes show strong biological activity by avoiding interaction with IL-2 receptor alpha subunit CD25. *Proc Natl Acad Sci USA*. 2010;107(5):2171–2176. doi:10.1073/pnas.0909384107
27. Fry TJ, Mackall CL. The many faces of IL-7: from lymphopoiesis to peripheral T cell maintenance. *J Immunol*. 2005;174(11):6571–6576. doi:10.4049/jimmunol.174.11.6571
28. Kiazzyk SA, Fowke KR. Loss of CD127 expression links immune activation and CD4(+) T cell loss in HIV infection. *Trends Microbiol*. 2008;16(12):567–573. doi:10.1016/j.tim.2008.08.011
29. Brechley JM, Paiardini M, Knox KS, et al. Differential Th17 CD4 T-cell depletion in pathogenic and nonpathogenic lentiviral infections. *Blood*. 2008;112(7):2826–2835. doi:10.1182/blood-2008-05-159301
30. Wang X, Mbondji-Wonje C, Zhao J, Hewlett I. IL-1 β and IL-18 inhibition of HIV-1 replication in Jurkat cells and PBMCs. *Biochem. Biophys. Res. Commun*. 2016;473(4):926–930. doi:10.1016/j.bbrc.2016.03.153
31. Doitsh G, Greene WC. Dissecting how CD4 T cells are lost during HIV infection. *Cell Host Microbe*. 2016;19(3):280–291. doi:10.1016/j.chom.2016.02.012
32. Pancera M, Majeed S, Ban YE, et al. Structure of HIV-1 gp120 with gp41-interactive region reveals layered envelope architecture and basis of conformational mobility. *Proc Natl Acad Sci USA*. 2010;107(3):1166–1171. doi:10.1073/pnas.0911004107
33. Xia C, Zhang X, Harypursat V, Ouyang J, Chen Y. The role of pyroptosis in incomplete immune reconstitution among people living with HIV: Potential therapeutic targets. *Pharmacol Res*. 2023;197:106969. doi:10.1016/j.phrs.2023.106969
34. Lv T, Cao W, Li T. HIV-related immune activation and inflammation: current understanding and strategies. *J Immunol Res*. 2021;2021:7316456. doi:10.1155/2021/7316456
35. Somsouk M, Estes JD, Deleage C, et al. Gut epithelial barrier and systemic inflammation during chronic HIV infection. *AIDS*. 2015;29(1):43–51. doi:10.1097/qad.0000000000000511
36. Godfrey C, Bremer A, Alba D, et al. Obesity and fat metabolism in human immunodeficiency virus-infected individuals: immunopathogenic mechanisms and clinical implications. *J Infect Dis*. 2019;220(3):420–431. doi:10.1093/infdis/jiz118
37. Alzahrani J, Hussain T, Simar D, et al. Inflammatory and immunometabolic consequences of gut dysfunction in HIV: parallels with IBD and implications for reservoir persistence and non-AIDS comorbidities. *EBioMedicine*. 2019;46:522–531. doi:10.1016/j.ebiom.2019.07.027
38. Meyer-Myklestad MH, Medhus AW, Lorvik KB, et al. Human immunodeficiency virus-infected immunological nonresponders have colon-restricted gut mucosal immune dysfunction. *J Infect Dis*. 2022;225(4):661–674. doi:10.1093/infdis/jiaa714
39. Tincati C, Merlini E, Braidotti P, et al. Impaired gut junctional complexes feature late-treated individuals with suboptimal CD4+ T-cell recovery upon virologically suppressive combination antiretroviral therapy. *AIDS*. 2016;30(7):991–1003. doi:10.1097/qad.0000000000001015
40. Pickard JM, Zeng MY, Caruso R, Núñez G. Gut microbiota: role in pathogen colonization, immune responses, and inflammatory disease. *Immunol Rev*. 2017;279(1):70–89. doi:10.1111/imr.12567
41. Vujkovic-Cvijin I, Somsouk M. HIV and the gut microbiota: composition, consequences, and avenues for amelioration. *Current HIV/AIDS Reports*. 2019;16(3):204–213. doi:10.1007/s11904-019-00441-w
42. Xie Y, Sun J, Wei L, et al. Altered gut microbiota correlate with different immune responses to HAART in HIV-infected individuals. *BMC Microbiol*. 2021;21(1):11. doi:10.1186/s12866-020-02074-1
43. Méndez-Lagares G, Pozo-Balado MM, Genebat M, García Pergañeda A, Leal M, Pacheco YM. Severe immune dysregulation affects CD4⁺CD25(hi)FoxP3⁺ regulatory T cells in HIV-infected patients with low-level CD4 T-cell repopulation despite suppressive highly active antiretroviral therapy. *J Infect Dis*. 2012;205(10):1501–1509. doi:10.1093/infdis/jis230
44. Guo YT, Guo XY, Fan LN, et al. The imbalance between intestinal Th17 and Treg cells is associated with an incomplete immune reconstitution during long-term antiretroviral therapy in patients with HIV. *Viral Immunol*. 2023;36(5):331–342. doi:10.1089/vim.2023.0017
45. Xin C, Mingfeng Z, Chengpeng Z, et al. To explore the effect of traditional Chinese medicine dual external treatment combined with HAART on poor immune reconstitution of AIDS from the balance of Th17/Treg. *J Chin J AIDS STD*. 2019;25(08):776–778. doi:10.13419/j.cnki.aids.2019.08.04
46. Man Z, Baoling R, Tao S, Hua L. Relationship between Th17/Treg imbalance and disease progression and monocyte activation in HIV-1 patients with intravenous drug abuse. *J Chin J Viral Dis*. 2013;3(05):330–334. doi:10.16505/j.2095-0136.2013.05.009
47. Li T, Xie J, Li Y, et al. Tripterygium wilfordii Hook F extract in cART-treated HIV patients with poor immune response: a pilot study to assess its immunomodulatory effects and safety. *HIV Clin Trials*. 2015;16(2):49–56. doi:10.1179/1528433614z.0000000005
48. Cao W, Liu X, Han Y, et al. (5R)-5-hydroxytriptolide for HIV immunological non-responders receiving ART: a randomized, double-blinded, placebo-controlled Phase II study. *Lancet Region Health Western Pacific*. 2023;34:100724. doi:10.1016/j.lanwpc.2023.100724
49. Liu X, Lin L, Lv T, et al. Combined multi-omics and network pharmacology approach reveals the role of Tripterygium Wilfordii Hook F in treating HIV immunological non-responders. *Phytomedicine*. 2022;101:154103. doi:10.1016/j.phymed.2022.154103
50. Tao Z, Dong J, Guo H, et al. The clinical effects and safety of artesunate in immunologic non-responders of ART-treated HIV/AIDS patients: a pilot study. *Chinese Journal of AIDS & STD*. 2021;27(09):921–925. doi:10.13419/j.cnki.aids.2021.09.03
51. Jipeng D, Zhuang T, Huijun G, et al. Efficacy characteristics of artesunate for immune dysfunction after antiretroviral therapy in HIV/AIDS. *China J Trad Chinese Med Pharm*. 2022;37(07):4170–4175.
52. X Q, W X, Z H, W J, T X. Effect of artesunate on immune function and intestinal flora in HIV/AIDS patients with poor immune reconstitution after HAART. *Acta Chin Med Pharmacol*. 2022;50(12):54–59. doi:10.19664/j.cnki.1002-2392.220274
53. L Y, L Z, S F, L Z, L P, G H. Effects of Yiaikang Capsules on intestinal microflora and immune function in HIV/AIDS patients with poor immune reconstitution of lung and spleen qi deficiency syndrome. *China J Tradition Chinese Med Pharm*. 2022;37(05):2729–2733.
54. Feng S, L Q, Q J, et al. Discussion on the protective effect of yiaikang capsule on intestinal mucosal barrier injury of HIV/AIDS based on permeability and tight junction. *J Emergency Traditional Chin Med*. 2018;27(05):769–772.

55. Z W, W J, G G, et al. Effects of TLR4 and TLR9 in immunological unresponsiveness AIDS patients treated with Yiqi Jianpi Granules combined with HAART. *China J Tradition Chinese Med Pharm.* 2019;34(09):4426–4428.
56. Z W, W J, L Y, X Q, Z F. Effects of Immune 2 Granule Combined with HAART therapy treatment on IL-2 and IL-10 mRNA level of HIV/AIDS with incomplete immune recovery. *J Basic Chin Med.* 2017;23(06):824–825+836. doi:10.19945/j.cnki.issn.1006-3250.2017.06.029
57. Shete A, Dhayarkar S, Sangale S, et al. Incomplete functional T-cell reconstitution in immunological non-responders at one year after initiation of antiretroviral therapy possibly predisposes them to infectious diseases. *Inter J Infect Dis.* 2019;81:114–122. doi:10.1016/j.ijid.2019.01.017
58. Yanmei J, Ming S, Ruonan X, Jianguo Z, Fusheng W. Study on clinical immune basis and translational application of AIDS. *J Chin J AIDS & STD.* 2019;25(09):877–881. doi:10.13419/j.cnki.aids.2019.09.01
59. Bin Z, Zhongjing S. Advances in CD45 isoforms in T-cell receptor signaling. *J Anat Res.* 2021;43(01):82–85.
60. Liu ZB, Yang JP, Xu LR. Effectiveness and safety of traditional Chinese medicine in treating acquired immune deficiency syndrome: 2004-2014. *Infect Diseases Poverty.* 2015;4:59. doi:10.1186/s40249-015-0093-6
61. Yinglin W, Jianzhong L, Mingzhong X, et al. Clinical effect of the method of invigorating spleen and tonifying kidney on acquired immunodeficiency syndrome patients. *J China Medical Herald.* 2022;19(02):112–116.
62. Weihua L, Chuanyun L, Huiping Y, Xin Z, Dantong Z, Xiuhui L. Effect of spleen-invigorating and kidney-invigorating Chinese medicine on HIV-specific T cell immune function. *J Chin J Integrat Trad Western Med.* 2014;34(07):812–815.
63. Jipeng D, Zhuang T, Huijun G, et al. Efficacy of artesunate for AIDS patients with incomplete immune reconstitution after antiretroviral therapy. *J China J Trad Chin Med Pharm.* 2022;37(07):4170–4175.
64. Qihua X, Xue W, Haipeng Z, Jian W, Xinghua T. Effects of artesunate on immune function and gut microbiota in HIV/AIDS patients with poor immune reconstitution after HAART. *J Acta Chin Med Pharmacol.* 2022;50(12):54–59. doi:10.19664/j.cnki.1002-2392.220274
65. Jiantao C, Siyi X, Zifeng L, Qianfeng L, Qin T, Linchun F. Regulating effects of artesunate on SIV-infected Chinese RhesusMacaque T lymphocyte activation. *J Acta Chin Med.* 2017;32(11):2029–2034. doi:10.16368/j.issn.1674-8999.2017.11.532
66. Siyi X, Jiantao C, Yuanfen Z, Linchun F. Effects of Aikeqing on T Lymphocyte subsets of SIV/SAIDS. *J Chin Archi Trad Chin Med.* 2018;36(01):173–176. doi:10.13193/j.issn.1673-7717.2018.01.042
67. Ke C, Linchun F, Xinghua T, Haipeng Z, Shuyun C. Effect of Aikeqing Granule combined with HAART on CD4 counts of HIV/AIDS patients with incomplete immune reconstitution. *J Chin J AIDS & STD.* 2017;23(08):688–690. doi:10.13419/j.cnki.aids.2017.08.04
68. Yuqi Y, Guangcheng Q, Yunzhong L, Yaping P, Li W, Xiandong W. Clinical efficacy analysis of Fuzheng anti-poison pill combined with HAART in the treatment of AIDS immune reconstitution insufficiency. *J Guiding J Trad Chin Med Pharm.* 2020;26(10):66–68. doi:10.13862/j.cnki.cn43-1446/r.2020.10.013
69. Yuqi Y, Guangcheng Q, Zhongyun L, Yaping P, Li W, Xiandong W. Clinical efficacy analysis of Fuzheng anti-poison pill combined with HAART in the treatment of AIDS immune reconstitution insufficiency. *J Guiding J Trad Chin Med Pharm.* 2020;26(10):66–68. doi:10.13862/j.cnki.cn43-1446/r.2020.10.013
70. Yuqi Y, Hongmei W, Nali S, Zhengzheng H, Jun S, Yi C. Intervention of traditional Chinese medicine treatment on the expression levels of PD-1 and Tim-3 on T cells in HIV-infected patients. *J Yunnan J Trad Chin Med Materia Medica.* 2016;37(09):76–78. doi:10.16254/j.cnki.53-1120/r.2016.09.035
71. Qingfei X, Ying L, Wen Z, et al. Effect of Wenshen Jianpi Granule on T lymphocyte subsets in AIDS patients with immune reconstitution insufficiency after HAART. *J Basic Chin Med.* 2019;25(10):1391–1392+1455. doi:10.19945/j.cnki.issn.1006-3250.2019.10.023
72. Yingyu C, Song C, Linchun F, et al. Effect of Chuankezhi injection on balance between T helper type 17 and regulatory T Cells in Chinese Rhesus Macaques infected with Simian immunodeficiency virus. *J Guangzhou Univ Trad Chin Med.* 2014;31(04):566–572+577. doi:10.13359/j.cnki.gzxbtcm.2014.04.017
73. Qin Z. Expression levels of HLA-DR, CD38 and oxidized lipoprotein in AIDS patients with incomplete immune reconstitution and intervention of invigorating spleen and tonifying kidney prescription. *Master.* 2020. doi:10.27465/d.cnki.gzzyc.2020.000142.
74. Ya-nan L, Zhi-bin L, Feng S, Zhen L, Peng-yu L, Hui-jun G. Effects of Yiaikang Capsules on intestinal microflora and immune function in HIV/AIDS patients with poor immune reconstitution of lung and spleen qi deficiency syndrome. *J China J Trad Chin Med Pharm.* 2022;37(05):2729–2733.
75. Lihua C, Feng S, Qianlei X, et al. Explore the mechanism of Yiaikang capsule in treating AIDS patients with poor immune reconstitution based on serum metabolomics. *J China J Trad Chin Med Pharm.* 2024;39(05):2651–2656.
76. Quan Z, Xiangqun L, Kui X, et al. Clinical study of Jianpi Yiqi decoction combined with HAART to promote immune reconstitution in AIDS patients with spleen deficiency and dampness syndrome. *J Clin J Chin Med.* 2018;10(30):102–105.
77. Ze L, Kuo RL, Yao C, et al. Curative effect of Diwu Yanggan Capsule combined with HAART on HIV/AIDS patients with poor immune reconstitution. *J Chinese J AIDS STD.* 2022;28(10):1127–1131. doi:10.13419/j.cnki.aids.2022.10.03
78. Fauci SR, Jamieson BD, Chin AC, et al. Telomerase-based pharmacologic enhancement of antiviral function of human CD8+ T lymphocytes. *J Immunol.* 2008;181(10):7400–7406. doi:10.4049/jimmunol.181.10.7400
79. Peng L, Genhui M, Sillonite. Effect of Astragalus membranaceus on CD4+T lymphocyte count in AIDS patients without immune response. *J Chinese J Exp Clin Infect Dis.* 2014;8(02):108–110.
80. Wen Z, Jian W, Guo-jian G, et al. Effects of TLR4 and TLR9 in immunological unresponsiveness AIDS patients treated with Yiqi Jianpi Granules combined with HAART. *J China J Trad Chin Med Pharm.* 2019;34(09):4426–4428.
81. Yuping L, Qihua X, Siyan C, Xinghua T. Retrospective study of 92 patients with HIV/AIDS treated with Ganoderma lucidum combined with highly effective anti-retroviral therapy. *J Chin J AIDS & STD.* 2020;26(02):120–124. doi:10.13419/j.cnki.aids.2020.02.03
82. Yanli T. Clinical and mechanism study on the intervention of immunological reconstitution in AIDS patients by immune-2 prescription. *Doctor.* 2012. https://kns.cnki.net/kcms2/article/abstract?v=MBBTPQIn9ZKHTbUx22CUOw3sbawkNmJpXc6dbkxsFarmFb15vAxB2Ov971myXY64sagJry0_Bqx9OdcXpaga1v-8i2PUd35PEXgP_nSnMeIYc6Iw5wJBXn_IncanPCEO2-oPz60yinDZS7v0Gv15w25GWERvTCqpE052J-TlhNpE0-2CP1zQxXZ2LsQfMZjTHIXNlnBOQ6Ho=&uniplatform=NZKPT&language=CHS.
83. Acquired Immunodeficiency Syndrome Professional Group SoLD, Chinese Medical Association, Control CCfD, Prevention. Chinese guidelines for diagnosis and treatment of human immunodeficiency virus infection/acquired immunodeficiency syndrome (2024 edition). *J Chin J Infect Dis.* 2024;05:257–284.
84. Zhouyan W, Xiyuan S, Liran X, et al. Research progress of traditional Chinese medicine in treating common opportunistic infections of AIDS. *J Chin J AIDS & STD.* 2023;29(06):723–726. doi:10.13419/j.cnki.aids.2023.06.24

85. Xiuxia M, Liran X, Zhipan Z, et al. Effect of Qingfei Peiyuan capsules on patients with phlegm-heat obstructing lung infection of AIDS. *J China J Trad Chin Med Pharm.* 2014;29(06):2068–2070.
86. Xiuxia M, Liran X, Zhipan Z, et al. Effect of Qingfei Peiyuan capsules on main symptoms of 141 AIDS patients with pulmonary infection of phlegm-heat obstructing lung. *J Modern Trad Chin Med Materia Medica-World Sci Technol.* 2014;16(05):1127–1132.
87. Xiuxia M, Feng S, Weifeng C, et al. Effect of Qingfei Peiyuan Granule on Th1/Th2 in AIDS patients with phlegm-heat obstructing lung infection. *J China J Trad Chin Med Pharm.* 2021;36(06):3721–3724.
88. Xiuxia M, Feng S, Weifeng C, et al. Effect of combined treatment of Chinese and Western medicine on TLR/MyD88 signaling pathway in patients with AIDS pulmonary infection. *J Liaoning J Trad Chin Med.* 2022;49(09):9–12. doi:10.13192/j.issn.1000-1719.2022.09.003
89. Jianhui Y, Xiaotao Z, Zengjia X. Observation on clinical effect of Fuzheng Qingfei Decoction on AIDS complicated with pulmonary infection. *J Heilongjiang J Trad Chin Med.* 2019;48(03):91–92.
90. Jinmi L, Jiong L. Clinical effect analysis of Tanreqing in AIDS complicated with pulmonary infection. *J Global Trad Chin Med.* 2013;6(S2):190–191.
91. Shaoqun D. Clinical study on Xielikang capsule treating AIDS chronic diarrhea. *J Acta Chin Med.* 2014;29(11):1549–1551. doi:10.16368/j.issn.1674-8999.2014.11.025
92. Xiaoping Y, Zhenzhen S. Effect of Xielikang capsule on intestinal microecology of AIDS-related chronic diarrhea. *J Trad Chin Med Res.* 2018;31(01):16–19.
93. Pengyu L, Lihua C, Qianlei X, et al. Discussion on HIV/AIDS diarrhea and Xielikang capsule treatment status and thinking based on intestinal flora. 2020;7.
94. Jingjing L, Xin P, Liyan T, et al. 效Clinical Efficacy of Jianpi Wenyang Jiedu Granules in the Treatment of AIDS-Related Diarrhea of Spleen-Kidney Yang Deficiency Based on 16S rDNA. *J Pharmacol Clin Chin Materia Medica.* 1–14. doi:10.13412/j.cnki.zyyl.20240320.012
95. Zhangwei Z, Yuling L, Jingfeng Y. Clinical analysis of Guchang Zhixie pills for treating AIDS diarrhea with spleen and stomach weakness. *J Syst Med.* 2017;2(13):125–127. doi:10.19368/j.cnki.2096-1782.2017.13.125
96. Guanpeng H. Clinical intervention study on the commonly witnessed type of AIDS fever. *Master.* 2015. https://kns.cnki.net/kcms2/article/abstract?v=JgtjNxUAsgfavMFEbWSsRQU3zpDhR2iwPf5e58azrLZ6BNC2jZtbaB1Yj6vw7lZulvRmfSid_Wb7M9hrAOX5jMRU5Xo-MhDdy5qSc4NjqwRRZhkp3V0NfzwJzVClkqE5cMCFxrSI3ksJ0UedXREiilBROWRPYVqInwkeh6hFD2u2O4UapvrDpLv2ymmm8JeqeKPK eM72U=&uniplatform=NZKPT&language=CHS.
97. Subing Z. Analysis of clinical effect of Xiaobuplehuhu decoction on AIDS fever. *J Health Way.* 2017;16(10):199–200.
98. Qiong F, Qingke D, Jian L, Jiantao G. Observation on therapeutic effect of Buzhong Yiqi Decoction on 40 cases of AIDS fever. *J Forum Trad Chin Med.* 2014;29(01):38–39. doi:10.13913/j.cnki.41-1110/r.2014.01.022
99. Baoliang W, Bin G. Lianhua Qingwen decoction was added to treat 60 cases of AIDS fever. *J Trad Chin Med Res.* 2013;26(12):19–21.
100. Yujin Z, Junwen W, Jiangrong W, et al. Study on the curative effect and mechanism of Gentian Xiegan Decoction in treating the intrinsic syndrome of itchy damp-heat of AIDS skin. *J Hunan Univ Chin Med.* 2022;42(04):680–684.
101. Xiaoling S, Jinbao N. 34 cases of AIDS skin mucosal lesions treated by traditional Chinese medicine. *J Yunnan J Trad Chin Med Materia Medica.* 2011;32(10):49. doi:10.16254/j.cnki.53-1120/r.2011.10.011
102. Baoping Z. Treatment of 58 cases of AIDS accompanied by skin damage with aloe Pikang liquid. *J Liaoning Univ Trad Chin Med.* 2008;09:111–112. doi:10.13194/j.jlunivtm.2008.09.113.zhaobp.059
103. Pengfei M, Tao W, Xiuxia M, Xue D, Haojie W, Liran X. Gancao Xiexin decoction treats AIDS recurrent oral ulcer of spleen and stomach damp-heat syndrome. *J Acta Chin Med.* 2019;34(10):2203–2205. doi:10.16368/j.issn.1674-8999.2019.10.508
104. Pengfei M, Xiuxia M, Xue D, Haojie W, Liran X. Effect of licorice heart-draining decoction on patients with spleen and stomach damp-heat syndrome of recurrent oral ulcer of AIDS. *J China J Trad Chin Med Pharm.* 2019;34(08):3829–3831.
105. Manqin X, Xinya H. Clinical observation of traditional Chinese medicine treating AIDS complicated with oral candida infection. *J Electron J Pract Clin Nurs Sci.* 2018;3(45):131–141.
106. Qihua X, Xue W, Jian ZH, Xinghua W, T. Effects of artesunate on immune function and gut microbiota in HIV/AIDS patients with poor immune reconstitution after HAART. *J Acta Chin Med Pharmacol.* 2022;50(12):54–59. doi:10.19664/j.cnki.1002-2392.220274
107. Yanan L, Feng S, Pengyu L, Zhen L, Lihua C, Huijun G. Research on intestinal microbiota in HIV/AIDS patients with poor immune reconstitution by 16S rDNA. *J Chin J AIDS STD.* 2022;28(07):775–780. doi:10.13419/j.cnki.aids.2022.07.04
108. Qihua X, Siyan C, Yuping L, Jian W, Xinghua T. Preliminary study of the effect of traditional Chinese medicine on gut microbiota of HIV/AIDS with poor immune reconstitution based on 16S rDNA. *J Chin J AIDS & STD.* 2020;26(11):1150–1153+1177. doi:10.13419/j.cnki.aids.2020.11.02
109. Lihua C, Feng S, Qianlei X, et al. Explore the mechanism of Yiaikang capsule in treating AIDS patients with poor immune reconstitution based on serum metabolomics. *J China J Trad Chin Med Pharm.* 2024;39(05):2651–2656.
110. Zhouyan W, Xiyuan S, Liran X, et al. Traditional Chinese medicine treatment of HIV/AIDS research progress of common opportunistic infections. *J Chin J AIDS & STD.* 2023;29(06):723–726. doi:10.13419/j.cnki.aids.2023.06.24

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>