

CORRECTION

Correction: The Association between Polyclonal Combined Serum Free Light Chain Concentration and Mortality in Individuals with Early Chronic Kidney Disease

Lakhvir K. Assi, Natasha McIntyre, Simon Fraser, Scott Harris, Colin A. Hutchison, Chris W. McIntyre, Paul Cockwell, Maarten W. Taal

There is an error in the units in [Fig 4](#), “Correlation of cFLC and eGFR.” Please see the corrected [Fig 4](#) here.




Fig 4. Correlation of cFLC and eGFR. There was a significant inverse association with cFLC (combined serum free light chains) and estimated glomerular filtration rate (eGFR) ($\rho = -0.49$, $P < 0.0001$). Spearman rank correlation was performed.

doi:10.1371/journal.pone.0141404.g001



OPEN ACCESS

Citation: Assi LK, McIntyre N, Fraser S, Harris S, Hutchison CA, McIntyre CW, et al. (2015) Correction: The Association between Polyclonal Combined Serum Free Light Chain Concentration and Mortality in Individuals with Early Chronic Kidney Disease. PLoS ONE 10(10): e0141404. doi:10.1371/journal.pone.0141404

Published: October 20, 2015

Copyright: © 2015 Assi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Reference

1. Assi LK, McIntyre N, Fraser S, Harris S, Hutchison CA, McIntyre CW, et al. (2015) The Association between Polyclonal Combined Serum Free Light Chain Concentration and Mortality in Individuals with Early Chronic Kidney Disease. PLoS ONE 10(7): e0129980. doi:[10.1371/journal.pone.0129980](https://doi.org/10.1371/journal.pone.0129980) PMID: [26132658](https://pubmed.ncbi.nlm.nih.gov/26132658/)