



Published in final edited form as:

Mucosal Immunol. 2015 July ; 8(4): 852–862. doi:10.1038/mi.2014.116.

Lipoxin B₄ promotes the resolution of allergic inflammation in the upper and lower airways of mice

L. Karra¹, O. Haworth², R. Priluck², B.D. Levy^{2,#}, and F. Levi-Schaffer^{1,#}

¹The Institute for Drug Research, Faculty of Medicine, The Hebrew University of Jerusalem, Israel

²Pulmonary and Critical Care Medicine Division, Department of Internal Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

Abstract

Chronic mucosal inflammation is the hallmark of important and common airway diseases, such as allergic rhinitis and asthma. Lipoxin A₄ (LXA₄) is an endogenous pro-resolving mediator for mucosal inflammation that decreases allergic and asthmatic responses. Lipoxin B₄ (LXB₄) is a structurally distinct member of the lipoxin family that signals in a manner distinct from LXA₄. LXB₄ is generated by mucosal tissues, but its actions in allergic inflammation are unknown. Here, we used murine models of allergic rhinitis and asthma to investigate LXB₄'s activity in mucosal inflammation. In the upper airway, LXB₄ significantly decreased nasal mucosal leukocytes and degranulation of mast cells and eosinophils. In the lower airway, LXB₄ significantly decreased airway inflammation, mucus metaplasia and hyper-responsiveness. Inhibition of mast cell degranulation *in vivo* by LXB₄ was more potent than dexamethasone, and these agents displayed unique profiles for cytokine regulation; however, their overall anti-inflammatory actions were comparable. LXB₄ decreased eotaxin-dependent eosinophil chemotaxis, IgE-mediated mast cell degranulation and expression of type 2 cytokine receptors. Together, these findings indicate that LXB₄ carries cell type selective and mucosal protective actions that broaden the lipoxin family's therapeutic potential for upper and lower airway catabasis.

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Corresponding authors: Bruce D. Levy, Pulmonary and Critical Care Medicine, Brigham and Women's Hospital, Harvard Institutes of Medicine Bldg. (HIM855), Boston, MA, USA 02115 Tel: 617-525-5407, Fax: 617-525-5413, blevy@partners.org, Francesca Levi-Schaffer, The Institute for Drug Research, Faculty of Medicine, The Hebrew University of Jerusalem, Israel, 91120, POB. 12065, Tel: +972-2-6757512, Fax: +972-2-6758144, Francescal@ekmd.huji.ac.il.

[#]Equal contributions by these authors

Disclosure/Conflict of Interest:

None of the authors declare any financial competing interests.

Supplementary Material is linked to the online version of the paper at <http://www.nature.com/mi>.

The PDF file contains additional explanation on nasal mucosa extraction as well as flow cytometry staining on nasal mucosal cells and LXB₄ validation methodology.

AUTHOR CONTRIBUTIONS

L.K. planned and performed experiments, collected and analyzed data, and wrote the manuscript; O.H. planned and performed experiments, collected and analyzed data, and edited the manuscript; R.P. performed experiments, B.D.L and F.L.S co- designed the study, performed experiments, analyzed data, and wrote the manuscript.

INTRODUCTION

Allergic airway diseases, such as rhinitis and asthma, are common and adversely impact patients function and quality of life¹⁻². Allergic rhinitis (AR) and atopic asthma can present with chronic mucosal inflammatory processes, and have been referred to as a “united airways” disease³⁻⁴. Despite distinct clinical manifestations, these diseases have immunological similarities. For example, eosinophil tissue infiltration and IgE-mediated mast cell (MC) activation through FcεRI bound allergen are major factors in the pathogenesis of AR and atopic asthma⁵. Current treatments for AR and asthma principally target symptom relief and anti-inflammation with glucocorticosteroids (GC). Since GC are associated with heterogeneous therapeutic responses and can have serious side effects, there is still a need for new treatment strategies to control AR and asthma. Importantly, no curative therapy is available to resolve allergic airway inflammation.

Recently, research focused on the physiological termination of inflammatory responses has led to the identification of specialized mediators involved in the natural resolution of inflammation in health, including lipoxins (i.e., lipoxin A₄ (LXA₄), lipoxin B₄ (LXB₄))⁶. These specialized pro-resolving mediators are lipoxygenase-derived eicosanoids formed from arachidonic acid by transcellular metabolism during cell–cell interactions (reviewed in⁶). Lipoxins are generated by the upper respiratory tract, including by nasal polyps⁷, and are present in nasal lavage fluids from aspirin- challenged patients with aspirin-exacerbated respiratory disease⁸. Lipoxins are also present in the lower airways in broncho-alveolar lavage fluid (BALF) from patients with asthma⁹. Of note, severe asthma is associated with decreased LXA₄ generation⁹. Administration of a stable analog of LXA₄ blocks allergic airway inflammation, mucus metaplasia and hyper-reactivity to methacholine¹⁰. LXA₄ is a potent inhibitor of granulocyte recruitment and activation, including eosinophils¹⁰. LXB₄ is a structurally distinct product of arachidonic acid metabolism. LXB₄ regulates neutrophil activation¹¹; however, its actions in allergic inflammation and on the functional responses of mucosal effector cells (i.e. MC and eosinophils), has yet to be established. Here we provide the first evidence that LXB₄ mediates anti- inflammatory and pro-resolving actions for allergic airway responses in murine models of upper and lower airway mucosal inflammation.

RESULTS

LXB₄ reduces leukocyte infiltration and mucus secretion in the nasal mucosa

Mice (BALB/c) were systemically sensitized with OVA/Alum and then challenged with OVA intranasally to target the allergic responses to the nasal mucosa (Figure 1a, see Methods). Twenty four hours after the last nasal allergen challenge, mice were given LXB₄ (as validated in Supplementary Figure S1) (100 ng, i.v), dexamethasone (100 ng) or vehicle control for 3 consecutive days (protocol day 28, 29, 30) (Figure 1a). Mice were sacrificed 24 hours later (protocol day 31). Histopathological assessment of the nasal cavity in mice exposed to vehicle, revealed mucosal allergic inflammation with mucin secretion and leukocyte infiltration (Figure 1b). Relative to vehicle, LXB₄ reduced inflammatory infiltrates and mucin secretion in the lumen (Figure 1c). Equivalent amounts of dexamethasone also decreased the mucosal inflammation, but to a lesser degree than LXB₄

(Figure 1d). BALF that was collected from these mice at the time of nasal tissue excision did not reveal significant increases in eosinophils or other leukocytes (data not shown).

LXB₄ has pro-resolving actions on select nasal mucosa cell populations

Leukocyte infiltration and total nasal mucosal cell numbers were significantly decreased by LXB₄ compared to vehicle (Figure 2a). Differential cell count performed on Wright-Giemsa stained cytopins of nasal mucosa cells indicated a significant decrease in eosinophils numbers with LXB₄ and dexamethasone (Figure 2b). The numbers of MC still containing granules significantly increased with LXB₄ while dexamethasone gave no significant changes from vehicle (Figure 2c).

Mast cell and eosinophil degranulation is decreased by LXB₄

Examination of the cytopins following Wright-Giemsa staining revealed striking evidence of changes in leukocyte granule contents (Figure 3). Mice (BALB/c) exposed to vehicle had highly degranulated eosinophils and MC with evidence for released MC granules (Figure 3a). In sharp contrast, nasal mucosal MC from LXB₄ treated mice were larger and retained most of their granule contents, indicative of a less activated state (Figure 3b). Eosinophils also appeared less degranulated (Figure 3b). Nasal mucosal MC following dexamethasone treatment showed cellular changes consistent with partial activation and degranulation with cells generally smaller in size than with LXB₄ (Figure 3c). Next, the apparent decrease in eosinophils number and degranulation was validated by flow cytometry analysis where Siglec-F⁺ cells (Supplementary Figure S2a) and CD107a⁺ cells (Supplementary Figure S2b) were significantly decreased by LXB₄ (3.50%±0.87 Siglec-F⁺ cells; 19.54%±1.74 CD107a⁺ cells, p<0.05) compared to vehicle (8.51%± 1.86 Siglec-F⁺ cells; 27.15%±1.706 CD107a⁺ cells). In comparison, dexamethasone decreased Siglec-F expressing cells (4.59%±0.79, p<0.05, Supplementary Figure S2a), but had no significant effect on cell degranulation (CD107a⁺, p=0.531, Supplementary Figure S2b).

LXB₄ decreases serum OVA-specific IgE levels and select cytokines/chemokines

Serum titers of OVA-specific IgE (BALB/c mice) were significantly decreased with LXB₄ (0.57±0.10 pg/ml, p<0.01) and dexamethasone (0.56±0.08 pg/ml, p<0.01) relative to vehicle (1.54±0.29 pg/ml) (Figure 4a). Additionally, LXB₄ significantly decreased serum levels of several mediators associated with IgE production and allergic responses (Figure 4b-k), including IL-4 (Figure 4b), IL-9 (Figure 4c), RANTES (CCL5) (Figure 4d), granulocyte macrophage colony-stimulating factor (GM-CSF; Figure 4e), vascular endothelial growth factor (VEGF; Figure 4f) and IL-12 (Figure 4h). Monocyte chemoattractant protein-1 (MCP-1) was also decreased by LXB₄, but did not reach statistical significance with this sample size (Figure 4g; p=0.053). Dexamethasone did not share LXB₄'s broad immunoregulatory actions at this time point, but did decrease serum IL-4, IL-9 and GM-CSF (Figure 4b, 4c, 4e). Of interest, LXB₄ gave modest increases in keratinocyte chemoattractant (KC) (Figure 4i; p<0.01) and IL-13 (Figure 4j, p<0.05) levels. The increased serum KC level was not associated with increased tissue neutrophils in the nasal mucosa. Levels of IL-5 were not significantly influenced by either LXB₄ or dexamethasone (Figure 4k). Serum levels of IL-17 were also detected and significantly down-regulated by LXB₄ (Figure 4l, p<0.01), but not by dexamethasone (Figure 4l).

LXB₄ decreases bone marrow-derived MC degranulation and bone marrow- derived eosinophil chemotaxis

To quantitatively determine if LXB₄ carried direct actions for cell functional responses, IgE-sensitized bone marrow (BM)-derived mast cells (BMMCs) from naive mice were exposed to LXB₄ and compared to LXA₄, dexamethasone or vehicle. Cell activation for degranulation was initiated in a non-antigen specific manner with an IgG (Fab)2' antibody as in ¹², see Methods. BMMC degranulation, monitored by the percentage of β-hexosaminidase release, was significantly decreased with LXB₄ (19.90%±0.86 of total release relative to vehicle control cells: 39.11%±0.83, p<0.0001) (Figure 5a). LXA₄ also decreased BMMC degranulation to 23.97%±1.64 of total release (p<0.01), and dexamethasone (10–1000nM) exhibited concentration-dependent inhibition of BMMC degranulation that relative to vehicle (38.07±0.4917) was similar in magnitude to LXB₄ (100nM dexamethasone: 33.93±0.5301, p<0.01; 1000nM dexamethasone: 27.05±0.3910, p<0.0001) (Figure 5b). To determine if these cellular responses were strain-specific, BALB/c and FVB cell responses were determined. Both BALB/c (Supplementary Figure S3a-b) and FVB-derived (Supplementary Figure S3c-d) BMMC responded similarly to IgE-mediated activation and degranulation, which was inhibited by LXB₄ and LXA₄ in a concentration-dependent manner. For eosinophil responses, murine BM-derived eosinophils were exposed to LXB₄, and for comparison LXA₄ or vehicle, prior to eotaxin exposure, to determine their impact on chemotaxis (see Methods). Relative to vehicle, LXB₄ and LXA₄ significantly decreased eotaxin-mediated chemotaxis (Figure 5c).

LXB₄ and LXA₄ decrease cytokine release from BMMC following IgE-mediated activation

Based on the serum cytokine regulation of AR mice by LXB₄, the direct actions of lipoxins on BMMC cytokine release was next determined. BMMC were exposed to LXB₄, LXA₄ or vehicle prior to activation. After 18 hrs, IgE-mediated activation of BMMC led to a significant TNF-α release that was significantly inhibited by LXB₄ and LXA₄ in a concentration-dependent manner (Figure 6a-d). LXB₄ also decreased FCεRI expression (Supplementary Figure S4a-b) without affecting cell viability (Supplementary Figure S4c).

LXB₄ regulates allergic lung inflammation

To determine if the actions of LXB₄ in the lower airways were similar to the upper airways, we first investigated the anti-inflammatory activity of LXB₄ (Figure 7I. Prevention). OVA sensitized animals (FVB) received LXB₄ (100ng, 1000ng) or vehicle (i.v.) 30 minutes before OVA aerosol challenge on protocol days 14–17 and were then sacrificed 24 hours later on day 18– a time of peak allergic inflammation. LXB₄ led to a significant dose-dependent decrease in total BALF cell numbers (Figure 7I. a), BALF eosinophils (Figure 7I.b) and BALF lymphocyte numbers (Figure 7I. d) but no significant change in macrophage numbers was seen (Figure 7I. c).

Given these anti-inflammatory actions, LXB₄'s pro-resolving actions in the lower airways were next determined by administration after the allergic airways responses were fully established (Figure 7II. Treatment). Twenty four hours after the last OVA aerosol challenge, LXB₄ (100ng, i.v.) was given (protocol day 18) and repeated on days 19 and 20. Mice were sacrificed 24 hours later (day 21) during the resolution phase of this model. Relative to

vehicle, LXB₄ significantly accelerated the resolution of the allergic lung inflammation with decreased BALF total cell number (Figure 7II. e) and decreased eosinophils, macrophage and lymphocyte numbers (Figure 7II. f-h). The resolution interval (R_i) was calculated for BALF total cells (approximately 6 days with vehicle) and LXB₄ shortened the R_i by over 30% to ~4 days (Figure 7II. i).

LXB₄ promotes the resolution of Airway Hyper-responsiveness and lung inflammation

In addition to airway inflammation, airway hyperresponsiveness (AHR) is a clinical hallmark of asthma¹, therefore the impact of LXB₄ on methacholine-triggered increases in lung resistance was determined. Relative to vehicle, mice (FVB) exposed to 100ng LXB₄ treatment (as in Figure 7II.) had significantly less AHR to methacholine (Figure 8a). In addition, histological analysis of lung sections from animals not subjected to AHR testing revealed that LXB₄ led to significant reductions in leukocyte infiltration and epithelial cell activation (Figure 8b-c). Mice treated with LXB₄ (100 ng) also exhibited a faster resolution of mucus metaplasia, as evidenced by decreased PAS positive airway epithelial cells compared to vehicle (Figure 8d). A significant decrease in BALF Th2 cytokines (Figure 8e-g) was observed, particularly for IL-4 (Figure 8e vehicle: 27.68±12.98; LXB₄:15.51±7.40, p<0.05) and IL-13 (Figure 8f vehicle: 3.21±1.145; LXB₄: 0.845±0.483, p<0.05). Given the importance of IL-13 signaling in airway mucus metaplasia and airway hyperreactivity to methacholine¹³⁻¹⁶, we next determined the influence of LXB₄ on IL-13 receptors (Figure 8h-j). Lung expression of the shared IL-4 receptor decreased during resolution (day 21) relative to the peak of inflammation (day 18), and LXB₄ further accelerated this decrease in lung IL-4 receptor expression (Figure 8h). LXB₄ administration also markedly decreased the expression of IL-13 specific receptors IL-13Rα1 and IL-13Rα2 (Figure 8i-j).

DISCUSSION

Allergic rhinitis and asthma are common allergic conditions with high morbidity. Allergen avoidance is often difficult and impractical and there is no curative therapy for the resultant allergic inflammation. Thus, new therapeutic approaches are needed. To this end, LXB₄ (~5µg/kg) displayed pro-resolving actions for allergic mucosal inflammation in murine models of AR and asthma. LXB₄ decreased mucosal leukocyte infiltration, OVA-specific IgE, MC and eosinophils degranulation, mucus metaplasia and AHR to methacholine.

LXB₄ mediated direct actions on pivotal cellular effectors for allergy, namely MC and eosinophils (as shown on cell functional responses *in vitro*), as well as indirect actions on these cells (regulation of levels of inflammatory mediators). Distinct from their important role during the induction of allergic airway responses, the levels of type 2 cytokines during resolution and their response to LXB₄ varied. In resolving upper and lower airways, LXB₄ potently decreased serum IL-4 and IgE levels. Because significant changes were not present in IL-5 levels, LXB₄ regulation of eosinophils function appears to have resulted from a direct action on these cells as well as from the reduced levels of GM-CSF. Surprisingly, LXB₄ increased serum IL-13 as the upper airways resolved. In contrast, BALF IL-13 was decreased by LXB₄, suggesting local and distinct regulatory mechanisms. Although IL-13 can initiate allergic inflammation, it can also play a role in limiting inflammation. Of note,

IL-13 has the capacity to induce the expression of 15-lipoxygenase, which catalyzes the formation of both LXB₄ and LXA₄⁶.

IL-4 plays important roles in isotype switching and IgE production¹⁷⁻¹⁸, and LXB₄ inhibited both IL-4 and serum IgE levels. Given the differential actions on IL-13 levels, expression of its receptors was determined. Lung expression of IL-4R, IL-13Rα1 and IL-13Rα2 were down regulated during resolution and further decreased by LXB₄. IL-4R deficient mice display weak Th2 responses¹⁹. IL-13Rα1 deficiency has an established role in allergic airway responses²⁰. Thus, LXB₄ regulation of IL-4R and IL-13Rα1 expression would serve to decrease mucus metaplasia and airway hyper-reactivity. No LXB₄-mediated increases in expression of the IL-13 decoy receptor IL-13Rα2 were apparent. In asthma, regulation of IgE and IL-13 has beneficial actions in some asthmatic patients²¹⁻²², so together these findings suggest therapeutic potential for LXB₄ or its stable analogs²³.

The increase of serum KC with LXB₄ was unexpected and distinct from LXA₄²⁴. In addition to its well defined role as a neutrophil chemoattractant, KC can modulate ciliary function of murine sinonasal epithelial cells²⁵⁻²⁶, suggesting that LXB₄ might enhance mucociliary clearance via this mechanism. In contrast to KC, serum IL-17 levels were low and significantly decreased by LXB₄. Compared to the GC dexamethasone, LXB₄'s actions were of similar potency for decreasing inflammatory cell activation and more potent for regulating degranulation *in vivo*. Although consistently decreased by LXB₄, there was donor-to-donor variability for the magnitude of stimulus-driven BMMC degranulation, suggesting context dependent responses for mast cells and their regulation by LXB₄. Direct cellular actions for LXB₄ were also identified *in vitro* for inhibition of BMMC TNF-α production and BM-derived eosinophil chemotaxis. Together, these lines of evidence support potent anti-allergic and pro-resolving actions for LXB₄ on MC and eosinophils that were relevant for both upper and lower airways.

Notably, LXA₄ blocks histamine release from MC during interactions with epithelial cells²⁷ and decreases neutrophil degranulation of azurophilic granules²⁸. In conjunction with LXB₄'s actions here *in vitro*, the lipoxins (LXA₄ and LXB₄) carry MC and granulocyte stabilizing activity. Apart from its direct regulation of MC and eosinophils, LXB₄ blocks T-cell activation, cytokine release²⁹ and neutrophil chemotaxis¹¹. Indirect mechanisms are also likely in the accelerated resolution of allergic lung inflammation, as levels of several inflammatory mediators, including GM-CSF, a pro-survival cytokine for eosinophils, were decreased by LXB₄. Of interest, LXA₄ has a counter-regulatory effect on GM-CSF signaling in a human eosinophil cell line (EoL-1) and peripheral blood eosinophils³⁰. LXA₄ also inhibits human eosinophils recruitment *in vitro*³¹ and murine eosinophils *in vivo*¹⁰, and can induce natural killer cells to promote human eosinophils apoptosis in asthma in order to enhance their non-phlogistic clearance from inflamed tissues³². Together, these findings highlight several cellular mechanisms for lipoxin regulation of allergic airway inflammation.

LXB₄ and LXA₄ are generated by nasal polyps where a mucosal epithelial 15-lipoxygenase plays an important role in their biosynthesis. LXA₄ is present in nasal secretions¹⁰ and airway samples from human asthmatic patients^{9,33}. Eosinophilic donors generate LXA₄ preferentially to LXB₄³⁴, however, this is cell type specific as human platelet 12-

lipoxygenase converts leukotriene A₄ to LXA₄ and LXB₄ in an equimolar ratio³⁵, and IL-4 exposed alveolar macrophage 15-lipoxygenase generates LXB₄ preferentially to LXA₄³⁶. The nanogram doses of LXB₄ used here are within physiological concentrations generated during inflammatory responses and were very potent (~5µg/kg) in comparison to a similar dose of dexamethasone. LXA₄ interacts with several receptors, including the G-protein coupled receptor ALX/FPR2. While LXB₄ also displays pharmacological properties of a G-protein coupled receptor agonist, it binds to a receptor distinct from ALX/FPR2 that has yet to be identified. In conclusion, LXB₄ displayed pro-resolving actions for mucosal allergic responses in the upper and lower airways. LXB₄ is generated in the airway and the present results are the first demonstration of an airway protective effect for LXB₄ or that any specialized pro-resolving mediator is active in the nasal mucosa. With the development of LXB₄ stable analogs and the limited treatment options currently available, these findings suggest a potential new pro-resolving therapeutic strategy for allergic conditions.

Materials & Methods

Materials

Lipoxin A₄ and Lipoxin B₄ were from Cayman Chemical (Ann Arbor, Michigan, USA) and dexamethasone was from MP Biomedicals LLC (Solon-Ohio, USA). The lipoxins were stored at -80°C under a blanket of N₂ and their integrity was monitored by UV-Vis spectrophotometry and HPLC (Supplementary Figure S1 Online).

Allergic upper and lower airway models

The allergic upper airway model was designed to reflect AR and was a modification of a previously published protocol³⁷ (Figure 1). Briefly, BALB/c female mice (Harlan Laboratories), aged 7–8 weeks (20–23gr) were sensitized on day 0 and day 7 by intraperitoneal injection (i.p.) of 200µl chicken ovalbumin (OVA; 1mg/ml) (Amresco, Ohio, USA) and aluminum hydroxide (alum; 50 mg/ml) (Acros Organics, New Jersey, USA) in PBS. Starting on day 14, awake mice were challenged twice a day by intranasal (i.n.) instillation of 20µl OVA (25mg/ml) daily until day 27. For the allergic lower airway model, a previously published protocol with OVA was used to reflect atopic asthma³⁸. Briefly, FVB male mice aged 5–7 weeks (Charles River Laboratories) were housed in isolation cages under viral antibody-free conditions. Mice were fed a standard diet (Laboratory Rodent Diet 5001, PMI Nutrition International). Mice were sensitized with i.p. injections of OVA (10µg) and 1mg alum in 0.2ml saline on days 0 and 7. Mice were then challenged on days 14–17 with an aerosol of 6% OVA (25min).

Treatment protocols

Mice were given either intravenous (i.v.) LXB₄ (100ng or 1000ng), vehicle (0.1% ethanol) or dexamethasone (100ng). To determine pro-resolving actions, treatments were given after the completion of allergen challenge at peak allergic airway inflammation time: days 28, 29, 30 in the upper airway model (Figure 1a) or days 18, 19 and 20 in the lower airway model (Figure 7II). Calculation of the resolution interval (R_i) (i.e. time interval for the maximum number of cells to decrease by 50%) was then evaluated as in³⁹. To determine anti-inflammatory actions, LXB₄ (100ng) or vehicle were given i.v. on protocol days 14, 15, 16

and 17, 30 minutes prior to OVA aerosol challenge (Figure 7I). Twenty four hours after the last i.v. treatment, mice were euthanized for the collection of airway and systemic specimens. Mouse experiments were approved by the Animal Ethics committee, The Hebrew University, Jerusalem, Israel, and the Harvard Medical Area IRB, Boston, MA, USA (Protocol #03618).

Nasal tissue preparation of cell suspensions and histopathology

As in ³⁷, extensive dissection was required to obtain the nasal mucosa cell suspensions (see supplemental methods in the Online Data Supplement). Total viable cell numbers were enumerated by Trypan blue exclusion. Nasal tissue sections were prepared and stained for H&E.

Flow cytometry

Single cell suspensions from nasal mucosa were resuspended in PBS/0.1% bovine serum albumin (BSA) seeded in a 96-well plate at 10^6 cells/ml, blocked with 5% goat serum and stained with Phycoerythrin conjugated anti-CD107a (eBioscience, San Diego, California, USA) and anti-Siglec-F (BD biosciences) for the detection of total degranulation⁴⁰ and eosinophils, respectively. For *in vitro* assessment of BMMC expression of FCεRI, cells were stained with anti-FCεRI (eBioscience, San Diego, California, USA) and cell viability was determined by propidium iodide staining (Sigma Aldrich, Rehovot, Israel). Data were analyzed by a FACSCalibur and CellQuest software (Becton Dickinson, San Jose California, USA).

LXA₄/LXB₄ activity on mouse bone marrow derived mast cells (BMMC) and eosinophils

BMMC and BM eosinophils were obtained as previously described ⁴¹⁻⁴². BMMC from BALB/c and FVB mice (1×10^6 /ml) were sensitized overnight with Rat Myeloma IgE (1 μg/ml; Molecular Probes, Invitrogen, Carlsbad, California, USA). Cells were washed, resuspended in Tyrode's buffer and pre-incubated in a 96-well plate (2×10^5 /well) with LXB₄, LXA₄ or vehicle at the indicated concentrations for 30 min at 37°C. Cells were then activated by the addition of a goat anti-rat IgG (Fab)₂' antibody (10 μg/ml)(Jackson ImmunoResearch Laboratories Inc, PA, USA) (30 min, 37°C) in Tyrode's buffer. Degranulation was evaluated by a colorimetric enzymatic β-hexosaminidase release assay⁴³. For TNF-α release, supernatants were collected 18 hrs following BMMC activation for evaluation of mTNF-α quantified by ELISA (Peprotech Asia, Israel). For chemotaxis, BM eosinophils were similarly pre-treated and then seeded into the upper chamber of a 24 transwell plate (2×10^5 /well/100 μl) (Corning) with a 5.0 μm pore size polycarbonate membrane. Cells were incubated (37°C, 3 hours) to allow migration across the membrane in response to medium only or medium with murine eotaxin (R&D biosystems, Minneapolis, Minnesota, USA) in the lower wells (50 ng/ml/600 μl). Migrated BM eosinophils were assessed by aspirating a fixed volume from the lower wells and counting the cells by flow cytometer (20 seconds, duplicate determinations).

Cytospin preparation

Nasal mucosa and BALF cells were resuspended in PBS/2% BSA, enumerated by a hemacytometer and cytocentrifuged (1000 rpm, 4 mins, r.t.) followed by staining with Modified Wright-Giemsa (Sigma Aldrich, Rehovot, Israel). Total cells, MC and eosinophils numbers were counted.

Evaluation of serum cytokines/chemokines and OVA specific IgE-levels

Blood samples were collected from the inferior vena cava and sera were stored (-80°C) until measurement of OVA-specific IgE levels by ELISA (USCN life, China) and select cytokines/chemokines with Quantibody® Mouse cytokine array 1 (Ray Biotech, Inc. Norcross, Georgia, USA).

BALF and Lung sample preparation and evaluation

Bronchoalveolar lavage (BAL) was performed with 2×1 ml aliquots of PBS with 0.6 mM EDTA. Lungs were fixed at 25cm H₂O in 10% buffered formalin and paraffin embedded for hematoxylin and eosin (H&E) and Periodic Acid Schiff (PAS) staining (Sigma). Cell-free BALF (centrifuged at 2000g for 10 min) were coded and select cytokines/chemokines measured by bead array (Pierce Biotechnology, Inc., Rockford, Illinois, USA).

RNA isolation and Real Time PCR

Lungs were obtained and snap frozen. RNA was extracted with TRIzol (Invitrogen) and reverse transcribed. The cDNA was used as a template for the amplification of selected genes. The difference between the Ct value for the gene of interest and the respective Ct value for the control gene was then calculated (ΔCt). The fold change was calculated as $2^{-\Delta\text{Ct}}$. Primers were purchased from Invitrogen: mouse IL-13R α 1: right CGTGCAGAGATTTTCGACAG, left AGCTGTTGGTGCTGCTACTG, IL13R α 2: right GGCAAAGAAGTAACAAAAGGAAT, left CGAATGGAGTGAAGAGGAATG, IL-4R: right AAGCACGCAGATCCAAAATC, left GTGGAGCCTGAACTCGCA, GAPDH: right TTGATGGCAACAATCTCCAC, left CGTCCCGTAGACAAAATGGT, β -actin: right ATGGAGGGGAATACAGCCC, left TTCTTTGCAGCTCCTTCGTT.

Statistical analysis

All parameters were analyzed using GraphPad Prism 5 software (GraphPad Software, La Jolla, California). Data are expressed as means \pm SEMs. Statistical comparisons were performed using 1-way ANOVA, followed by the Tukey post test. When only 2 groups were compared statistical differences were determined using the unpaired Student *t*-test. A *p* value of less than 0.05 was considered significant.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Professor Charles N. Serhan for helpful discussions on this project and critical appraisal of the manuscript, as well as Nour Karra and Jennifer Colby for assisting with sample preparations.

This research was supported in part by the US National Institutes of Health grants AI068084, HL68669 and P01-GM095467 (BDL) and by the Aimwell Charitable Fund, UK (FLS).

References

1. Fanta CH. Asthma. *N. Engl. J. Med.* 2009; 360:1002–1014. [PubMed: 19264689]
2. Plaut M, Valentine MD. Clinical practice. Allergic rhinitis. *N. Engl. J. Med.* 2005; 353:1934–1944. [PubMed: 16267324]
3. KleinJan A, et al. United airways: circulating Th2 effector cells in an allergic rhinitis model are responsible for promoting lower airways inflammation. *Clin. Exp. Allergy.* 2010; 40:494–504. [PubMed: 19968652]
4. Palm NW, Rosenstein RK, Medzhitov R. Allergic host defences. *Nature.* 2012; 484:465–472. [PubMed: 22538607]
5. Locksley RM. Asthma and allergic inflammation. *Cell.* 2010; 140:777–783. [PubMed: 20303868]
6. Levy BD, Vachier I, Serhan CN. Resolution of inflammation in asthma. *Clin. Chest. Med.* 2012; 33:559–570. [PubMed: 22929102]
7. Edenius C, Kumlin M, Bjork T, Anggard A, Lindgren JA. Lipoxin formation in human nasal polyps and bronchial tissue. *FEBS Lett.* 1990; 272:25–28. [PubMed: 2172016]
8. Levy BD, et al. Agonist-induced lipoxin A4 generation: detection by a novel lipoxin A4-ELISA. *Lipids.* 1993; 28:1047–1053. [PubMed: 8121246]
9. Planaguma A, et al. Airway lipoxin A4 generation and lipoxin A4 receptor expression are decreased in severe asthma. *Am. J. Respir. Crit. Care Med.* 2008; 178:574–582. [PubMed: 18583575]
10. Levy BD, et al. Multi-pronged inhibition of airway hyper-responsiveness and inflammation by lipoxin A(4). *Nat. Med.* 2002; 8:1018–1023. [PubMed: 12172542]
11. Lee TH, et al. Lipoxin A4 and lipoxin B4 inhibit chemotactic responses of human neutrophils stimulated by leukotriene B4 and N-formyl-L-methionyl-L-leucyl-L-phenylalanine. *Clin. Sci.* 1989; 77:195–203. [PubMed: 2548801]
12. Lu-Kuo JM, Joyal DM, Austen KF, Katz HR. gp49B1 inhibits IgE-initiated mast cell activation through both immunoreceptor tyrosine-based inhibitory motifs, recruitment of src homology 2 domain-containing phosphatase-1, and suppression of early and late calcium mobilization. *J. Biol. Chem.* 1999; 274:5791–5796. [PubMed: 10026201]
13. Fallon PG, Emson CL, Smith P, McKenzie AN. IL-13 overexpression predisposes to anaphylaxis following antigen sensitization. *J. Immunol.* 2001; 166:2712–2716. [PubMed: 11160336]
14. Wills-Karp M, et al. Interleukin-13: central mediator of allergic asthma. *Science.* 1998; 282:2258–2261. [PubMed: 9856949]
15. Kuperman DA, et al. Direct effects of interleukin-13 on epithelial cells cause airway hyperreactivity and mucus overproduction in asthma. *Nat. Med.* 2002; 8:885, 889. [PubMed: 12091879]
16. Zhu Z, et al. Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *J. Clin. Invest.* 1999; 103:779–788. [PubMed: 10079098]
17. Gilmour J, Lavender P. Control of IL-4 expression in T helper 1 and 2 cells. *Immunology.* 2008; 124:437–444. [PubMed: 18479352]
18. Kashiwada M, et al. IL-4-induced transcription factor NFIL3/E4BP4 controls IgE class switching. *Proc. Natl. Acad. Sci. USA.* 2010; 107:821–826. [PubMed: 20080759]
19. Kopf M, et al. Disruption of the murine IL-4 gene blocks Th2 cytokine responses. *Nature.* 1993; 362:245–248. [PubMed: 8384701]

20. Ramalingam TR, et al. Unique functions of the type II interleukin 4 receptor identified in mice lacking the interleukin 13 receptor alpha1 chain. *Nat. Immunol.* 2008; 9:25–33. [PubMed: 18066066]
21. Busse WW, et al. Randomized trial of omalizumab (anti-IgE) for asthma in inner-city children. *N. Engl. J. Med.* 2011; 364:1005–1015. [PubMed: 21410369]
22. Wenzel S, et al. Dupilumab in persistent asthma with elevated eosinophil levels. *N. Engl. J. Med.* 2013; 368:2455–2466. [PubMed: 23688323]
23. Maddox JF, et al. Lipoxin B4 regulates human monocyte/neutrophil adherence and motility: design of stable lipoxin B4 analogs with increased biologic activity. *FASEB J.* 1998; 12:487–494. [PubMed: 9535221]
24. Gronert K, et al. A role for the mouse 12/15-lipoxygenase pathway in promoting epithelial wound healing and host defense. *J. Biol. Chem.* 2005; 280:15267–15278. [PubMed: 15708862]
25. Shen JC, Chen B, Cohen NA. Keratinocyte chemoattractant (interleukin-8) regulation of sinonasal cilia function in a murine model. *Int. Forum Allergy Rhinol.* 2012; 2:75–79. [PubMed: 22311845]
26. Shen JC, Cope E, Chen B, Leid JG, Cohen NA. Regulation of murine sinonasal cilia function by microbial secreted factors. *Int. Forum Allergy Rhinol.* 2012; 2:104–110. [PubMed: 22253045]
27. Martin N, et al. Primary human airway epithelial cell-dependent inhibition of human lung mast cell degranulation. *PLoS One.* 2012; 7:e43545. [PubMed: 22970103]
28. Gewirtz AT, Fokin VV, Petasis NA, Serhan CN, Madara JL. LXA4, aspirin-triggered 15-epi-LXA4, and their analogs selectively downregulate PMN azurophilic degranulation. *Am. J. Physiol.* 1999; 276:C988–C994. [PubMed: 10199831]
29. Ariel A, Chiang N, Arita M, Petasis NA, Serhan CN. Aspirin-triggered lipoxin A4 and B4 analogs block extracellular signal-regulated kinase-dependent TNF-alpha secretion from human T cells. *J. Immunol.* 2003; 170:6266–6272. [PubMed: 12794159]
30. Starosta V, Pazdrak K, Boldogh I, Svider T, Kurosky A. Lipoxin A4 counterregulates GM-CSF signaling in eosinophilic granulocytes. *J. Immunol.* 2008; 181:8688–8699. [PubMed: 19050289]
31. Soyombo O, Spur BW, Lee TH. Effects of lipoxin A4 on chemotaxis and degranulation of human eosinophils stimulated by platelet-activating factor and N-formyl-L-methionyl-L-leucyl-L-phenylalanine. *Allergy.* 1994; 49:230–234. [PubMed: 8037356]
32. Barnig C, et al. Lipoxin A4 regulates natural killer cell and type 2 innate lymphoid cell activation in asthma. *Sci. Transl. Med.* 2013; 5 174ra126.
33. Kazani S, et al. Exhaled breath condensate eicosanoid levels associate with asthma and its severity. *J. Allergy Clin. Immunol.* 2013; 132:547–553. [PubMed: 23608729]
34. Serhan CN, Hirsch U, Palmblad J, Samuelsson B. Formation of lipoxin A by granulocytes from eosinophilic donors. *FEBS Lett.* 1987; 217:242–246. [PubMed: 3109942]
35. Romano M, et al. Lipoxin synthase activity of human platelet 12-lipoxygenase. *Biochem. J.* 1993; 296:127–133. [PubMed: 8250832]
36. Levy BD, et al. Human alveolar macrophages have 15-lipoxygenase and generate 15(S)-hydroxy-5,8,11-cis-13-trans-eicosatetraenoic acid and lipoxins. *J. Clin. Invest.* 1993; 92:1572–1579. [PubMed: 8376607]
37. Miyahara S, Miyahara N, Takeda K, Joetham A, Gelfand EW. Physiologic assessment of allergic rhinitis in mice: role of the high-affinity IgE receptor (FcepsilonRI). *J. Allergy Clin. Immunol.* 2005; 116:1020–1027. [PubMed: 16275370]
38. Levy BD, et al. Protectin D1 is generated in asthma and dampens airway inflammation and hyperresponsiveness. *J. Immunol.* 2007; 178:496–502. [PubMed: 17182589]
39. Bannenberg GL, et al. Molecular circuits of resolution: formation and actions of resolvins and protectins. *J. Immunol.* 2005; 174:4345–4355. [PubMed: 15778399]
40. Grutzkau A, et al. LAMP-1 and LAMP-2, but not LAMP-3, are reliable markers for activation-induced secretion of human mast cells. *Cytometry A.* 2004; 61:62–68. [PubMed: 15351990]
41. Jensen BM, Beaven MA, Iwaki S, Metcalfe DD, Gilfillan AM. Concurrent inhibition of kit- and FcepsilonRI-mediated signaling: coordinated suppression of mast cell activation. *J. Pharmacol. Exp. Ther.* 2008; 324:128–138. [PubMed: 17925481]

42. Dyer KD, et al. Functionally competent eosinophils differentiated ex vivo in high purity from normal mouse bone marrow. *J. Immunol.* 2008; 181:4004–4009. [PubMed: 18768855]
43. Blank, U.; Rivera, J. Assays for regulated exocytosis of mast cell granules. In: Bonifacino, JS.; Dasso, M.; Harford, JB.; Lippincott-Schwartz, J.; Yamada, KM., editors. *Current Protocols in Cell Biology*. Wiley & Sons: New York; 2006. p. 15.11.1-15.11.18.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 1. LXB₄ reduces leukocyte infiltration and mucus secretion in the nasal mucosa
(a) Days 0, 7: OVA/Alum i.p. sensitization, days 14–27: OVA i.n. challenge, days 29–30: vehicle (0.1%), LXB₄ or dexamethasone (100ng) i.v. injection, day 30: sacrifice. **(b-d)** H&E stained mucosal sections of BALB/c mice injected with vehicle **(b)**, LXB₄ **(c)** or dexamethasone **(d)**. Red arrows- mucus; blue arrows- leukocyte infiltration sm-submucosa; mc-mucosa; b-bone (×20 magnification- left panel; ×40 magnification- right panel).




Fig. 2. LXB₄ has pro-resolving effects on select nasal mucosa cell populations
(a) Total cell number as evaluated in trypan blue exclusion test. **(b-c)** Differential counts of eosinophils **(b)** and granulated mast cells **(c)** in stained cytopins from BALB/c mice (n = 8).




Fig. 3. Fewer degranulated mast cells and eosinophils were evident in nasal mucosa with LXB₄ treatment

(a-c) Images of Wright-Giemsa stained cytopins of vehicle (a), LXB₄ (b) and dexamethasone (c) injected BALB/c mice ($\times 40$). Representative images from 3 mice. Black arrows-mast cells; hollow arrowheads- mast cell granules; red arrows- eosinophils.




Fig. 4. LXB₄ decreases OVA-specific IgE levels and modulates various cytokines in serum of AR mice

(a) OVA-specific IgE levels in vehicle, LXB₄ and dexamethasone injected BALB/c mice (n=9). (b-k) Cytokine expression in serum of vehicle, LXB₄ and dexamethasone injected mice (n=5), p<0.05; p<0.01 two-tailed Student's *t*-test.




Fig. 5. LXB₄ decreases BMMC degranulation and BM eosinophils chemotaxis
 Cells were derived from BALB/c mice bone marrow as described in Methods. **(a-b)** BMMC IgE-dependent activation (β -hexosaminidase release) **(c)** BM eosinophil chemotaxis to eotaxin presented as number of events counted per 20 seconds in the flow cytometer (see Methods) (n=3).




Fig. 6. LXB₄ and LXA₄ decrease TNF-α release from BMMC following IgE-mediated activation BMMC were derived from BALB/c and FVB mice bone marrow and treated as described in Methods. IgE activation was carried out with anti-rat IgG (Fab)₂' (10μg/ml) **(a-b)** TNF-α levels (pg/ml) in BALB/c cell supernatants following incubation with LXB₄ **(a)** and LXA₄ **(b)**. **(c-d)** TNF-α levels (pg/ml) in FVB cell supernatants following incubation with LXB₄ **(c)** and LXA₄ **(d)**




Fig. 7. LXB₄ regulates allergic lung inflammation

FVB Mice were sensitized and aerosol challenged with OVA. LXB₄ (100, 1000ng) or vehicle were injected: **I) prevention**- 30 minutes before challenge (days 14–17). **II) Treatment**- after the last challenge (days 18–20). Total cells (**a, e**), eosinophils (**b, f**), macrophages (**c, g**) and lymphocytes (**d, h**) numbers in BALF. The resolution interval-R_i for vehicle/LXB₄ injected mice (n = 5), p<0.005 one-tailed Student's *t*-test.




Fig. 8. LXB₄ promotes the resolution of Airway Hyper-responsiveness and lung inflammation (a) Determination of airway hyper-responsiveness to metacholine, (n = 5), p<0.005 one-tailed Student's *t*-test (b, c) H&E and (d) PAS staining of FVB mice lungs. Arrow heads point to mucus (magenta positive) Goblet cells (x20). (e-g) IL-4, IL-5 and IL-13 cytokine levels in BALF, (n = 5) p<0.05 paired Student's *t*-test. (h-j) mRNA levels of lung IL-4R, IL-13R α 1 and IL-13R α 2 (n = 5).