

Systematic review on strength training in Parkinson's disease: an unsolved question

Ileana Ramazzina¹
Benedetta Bernazzoli²
Cosimo Costantino¹

¹Department of Biomedical, Biotechnological and Translational Sciences, ²Department of Clinical and Experimental Medicine, Graduate School of Rehabilitation and Physical Medicine, University of Parma, Parma, Italy

Abstract: The purpose of this study was to investigate the effectiveness of strength training, performed against a different resistance from body weight, in improving motor and nonmotor symptoms in patients with Parkinson's disease (PD). The following electronic databases were searched: PubMed, Physiotherapy Evidence Database, Cochrane Central Register of Controlled Trials, Scopus, and Web of Science. The review was conducted and reported in accordance with the PRISMA statement. Thirteen high-quality randomized controlled trials were included. Strength training performed against external resistance is well tolerated and appears to be a suitable physical activity to improve both physical parameters and quality of life parameters of PD subjects. However, although the study intervention included strength training, only a few selected studies assessed the improvement of muscle strength. Despite the encouraging results, it is difficult to establish a correlation between strength training and the improvements made. Our review highlights the lack of common intent in terms of study design and the presence of different primary and secondary outcomes. Accordingly, further studies are needed to support the beneficial effects of different types of strength training in PD subjects and to underline the superiority of strength training in PD patients with respect to other training.

Keywords: Parkinson's disease, strength training, muscle strength

Introduction

Parkinson's disease (PD) is an age-related neurodegenerative disorder characterized by specific motor symptoms such as tremor, slowed movement, rigid muscles, impaired posture, and balance.^{1,2} An increased body of evidence highlights also the presence of nonmotor symptoms like hyposmia, sleep disorders, and gastrointestinal dysfunctions. These symptoms accompany the onset and progression of the pathology, often preceding cardinal motor features of PD.²⁻⁴

PD is characterized by the selective loss of dopaminergic neurons of the *substantia nigra pars compacta*, the main region affected by this pathology. The histopathological hallmark of PD is the development of cytoplasmic inclusions known as Lewy bodies (LB) and Lewy neurites (LN). The latter are insoluble proteinaceous aggregate formed primarily within the body and processes of brain cell neurons, but they are also present in the spinal cord and peripheral nervous system.² LB and LN are mainly composed of alpha-synuclein, an endogenous protein that plays a crucial role in PD. Toxic alpha-synuclein oligomers may impact cells in a number of ways, including the disruption of membranes, mitochondrial depolarization, cytoskeleton changes, impairment of protein clearance pathways, and enhanced oxidative stress. The loss of functions of native alpha-synuclein and the gain of toxic functions following the misfolding or oligomerization process play a pivotal role in the pathogenesis of PD.^{5,6}

Correspondence: Cosimo Costantino
Department of Biomedical,
Biotechnological and Translational
Sciences, University of Parma,
Via Gramsci, 14, 43126 Parma, Italy
Tel +39 521 703 517
Fax +39 521 702 893
Email cosimo.costantino@unipr.it

Although the etiology of PD remains obscure, multifactorial theories have been postulated concerning both environmental^{7,8} and genetic factors.^{9,10} The current therapeutic approaches to PD include levodopa (the most potent drug for controlling PD symptoms), dopamine agonists, catechol-O-methyltransferase inhibitors, and non-dopaminergic agents.^{11,12} Nevertheless, these drugs only slow down the disease progression and are often associated with comorbid problems. In order to improve the quality of life of PD patients, physical activity is considered to be the one of the most important non-pharmacological strategies. There is a strong consensus that physical exercise can 1) improve the management of symptoms,^{13–16} 2) delay disease progression,^{13,16–19} and 3) improve the physiological and structural function of the human brain.^{15,17,20,21} However, a great deal of additional research must be carried out to clarify the effects of physical exercise on brain neuroplasticity. Despite the fact that physical activity improves quality of life and functional independence of PD patients, it cannot reverse PD symptoms.

A lot of recent reviews describe the effect of aerobic exercise in improving motor and nonmotor symptoms in PD patients.^{15,22–24} Less attention has been focused on specific muscle training against an external resistance (resistance/strength training).²⁵ For this reason, the purpose of this study was to conduct a systematic review of published literature concerning the effects of resistance training, performed against a resistance different from the body weight, in PD patients. The primary outcome of this review was to assess the effectiveness of resistance training on muscle strength improvement. The secondary outcome was to shed light on the effects of resistance training in improving physical performance and quality of life of PD patients.

Materials and methods

The review was conducted and reported in accordance with the PRISMA statement (www.prisma-statement.org).

Databases and search strategy

The following electronic databases were searched: PubMed, Physiotherapy Evidence Database (PEDro), Cochrane Central Register of Controlled Trials, Scopus, and Web of Science.

The keywords used were “Parkinson disease and physical exercise”, “Parkinson disease and physical therapy”, “Parkinson disease and training”, “Parkinson disease and strength physical exercise”, “Parkinson disease and strength physical therapy”, “Parkinson disease and strength training”, “Parkinson disease and eccentric training”, “Parkinson disease and resistance physical exercise”, “Parkinson disease and resistance physical therapy”, “Parkinson disease and

resistance training”, “Parkinson disease and aquatic training” and “Parkinson disease and aquatic exercise”.

A manual search of reference lists of selected papers and reviews on the topics was performed to identify additional relevant articles. To identify gray literature, a search was conducted in Google and Google Scholar using the aforementioned keywords. The electronic databases were investigated until February 2016.

Selection criteria for studies

Strength training was defined as an intervention in which participants exercised a muscle or group of muscles against an external resistance.²⁵ For this study, we considered as external resistance cycle ergometer, weight machine, elastic band, punching bag, and water. In further analysis, we included articles in which the effect of strength training in subjects affected by PD was evaluated and the articles that matched the following inclusion criteria based on PICO (Patient, Intervention, Comparison, Outcome) principles:

- randomized controlled trials related to both sexes;
- stages 1–3 on the Hoehn and Yahr scale;
- study design comparing the effects of strength training versus different exercise protocol;
- study outcomes: muscle strength, physical performance, quality of life;
- training/assessment of subjects during the “on” medication period;
- articles written in English.

Exclusion criteria were as follows:

- observational studies;
- studies with healthy or non-exercise controls;
- studies employing supplementary intervention therapies in addition to strength training;
- studies with tailored exercise programs to meet individual capacity.

Two authors (IR and BB) independently screened the articles by title and abstract against the selection criteria. Articles that were unclear from their title or abstract were reviewed against the selection criteria through the full text. Any discrepancies between authors were resolved through discussion. The second step was to screen all full-text articles that passed the first step.

Conference and symposium abstracts were assessed but deemed unsuitable due to the limited body of data related to the study design and the intervention program.

Data extraction and analysis

Two authors (IR and BB) independently extracted data from the 13 studies that met the inclusion criteria. In agreement

with PICO principles, the data included the following: disease population and disease status, the study design and number of participants, strength training (duration, frequency, intensity of strength training, and specific exercises employed), outcomes, participant retention and dropouts, and adverse effects associated with strength training. Any discrepancies between reviewers were resolved through discussion with the third author (CC).

Assessment of risk of bias

The methodological quality of selected articles was assessed using the PEDro scale checklist. For the purposes of this review, studies were included if they achieved a score ≥ 6 (high-quality study). If the articles' score was not reported in the PEDro database, two researchers (IR and BB) assessed the score independently. The researchers were blinded to each other's quality assessment, and in the event of disagreement, a third opinion was sought (CC).

Results

Overview of the inclusion process and methodological quality assessment

The reviewing process is presented in Figure 1; duplicate articles, review articles, conference proceedings, book

chapters, and articles written in languages other than in English were excluded from the initial records retrieved from different databases. From a total of 511 selected articles, 449 did not meet the inclusion criteria as previously reported in materials and methods section. Of the remaining 62 items, 49 were excluded after quality assessment. A total of 13 articles were included in this review.²⁶⁻³⁸ Two articles showed shared data, so we combined them and considered them as a single item.^{35,36} The PEDro scale score of the articles is reported in Table 1. All the studies included had a PEDro scale score of 6 or higher, indicating low risk of bias. Eligibility criteria were not used to calculate the PEDro scale score because they influenced external validity but not the internal or statistical validity of the trial.

Details of the included articles

The details of the reviewed articles are reported in Table 2. The range of disease severity assessed by the Hoehn and Yahr scale is 1.5–3; the lowest mean age considered is 58.6 ± 5.6 years old²⁸ and the maximum is 75.7 ± 7.2 years old,³² and the number of subjects per study ranged from 22 to 60. The lowest intervention per week considered is reported in the study of Mateos-Toset et al, in which they highlighted the effects of a brief exercise session of hand




Figure 1 Study selection process. Electronic databases searched were PubMed, Physiotherapy Evidence Database (PEDro), Cochrane Central Register of Controlled Trials, Scopus, and Web of Science.

Table 1 Study design quality assessment based on Physiotherapy Evidence Database Scale

Study	1	2	3	4	5	6	7	8	9	10	11	Total score
Arcolin et al ²⁶	X	X		X			X	X		X	X	6
Carvalho et al ^{38,*}	X	X	X	X				X	X	X	X	7
Combs et al ²⁷	X	X	X	X			X		X	X	X	7
Corcos et al ²⁸	X	X	X	X			X	X		X	X	7
Dibble et al ²⁹		X	X	X			X	X	X	X	X	8
Mateos-Toset et al ³⁰	X	X	X	X			X	X		X	X	7
Paul et al ³¹	X	X	X	X			X	X	X	X	X	8
Ridgel et al ^{33,*}	X	X		X			X	X		X	X	6
Schlenstedt et al ³²		X		X			X		X	X	X	6
Shen and Mak ³⁴	X	X	X	X			X	X		X	X	7
Shen and Mak ³⁵	X	X		X			X	X	X	X	X	7
Shen and Mak ³⁶	X	X	X	X			X	X	X	X	X	8
Volpe et al ³⁷	X	X	X	X			X	X	X	X	X	8

Notes: PEDro criteria: 1, eligibility criteria (not used to calculate the PEDro score); 2, random allocation; 3, concealed allocation; 4, baseline comparability; 5, blind subjects; 6, blind therapists; 7, blind assessors; 8, adequate follow-up; 9, intention-to-treat analysis; 10, between-group comparisons; 11, point estimates and variability provided; *PEDro score attributed by the authors.

Abbreviation: PEDro, Physiotherapy Evidence Database.

training with therapeutic putty.³⁰ The longest intervention is reported in the study of Corcos et al, in which they assessed the effects of 2 years of progressive resistance training.²⁸ As regards the length of the training session, the shortest was the single 15-minute hand exercise reported by Mateos-Toset et al³⁰ and the longest was the double 30-minute session of specific training plus one additional 60-minute session of common exercises, reported by Arcolin et al.²⁶ Most of the reviewed articles assessed the post-treatment effects, and only three reported the long-term effects.^{31,32,35,36}

Three out of the twelve studies included, considered as external resistance cycle ergometer,^{26,29,33} three weight machines,^{28,31,38} and two studies considered both weight machines and cuff weights.³⁴⁻³⁶ One study analyzed elastic bands and cuff weights,³² punching bag,²⁷ water,³⁷ or therapeutic putty,³⁰ respectively, as external resistance. In all the studies reviewed, strength training was compared with the PD active control group. The latter performed traditional physiotherapy, treadmill training, and balance training. The focus of training was to maintain and/or improve functional ability of PD patients and improve their quality of life. However, which intervention was the most effective is still being debated.^{16,39,40}

The details of strength training included in the selected articles are reported in Table 3. Most of the studies involved the training of the lower body with cycle ergometer,^{26,29,33} water,³⁷ weight machines,³¹ elastic bands, and cuff weights,³² or both weight machines and cuff weights.³⁴⁻³⁶ Two studies

involved training of both upper and lower limbs with weights/resistance machines.^{28,38} One study focused on manual dexterity and strength.³⁰ Regarding the intervention program with weight machines, the initial intensity settled at a different percentage of one repetition maximum (1RM), ranging from 40% to 70%–80% of 1RM for lower limbs and from 30%–40% to 70%–80% of 1RM for upper limbs. Also increase within the training program varied between the studies.

Primary outcome: muscle strength

Out of the twelve studies that performed strength training, only six^{28-32,38} involved specific tests to verify the improvement of PD subjects' muscle strength (Tables 2 and 3). In particular, three studies assessed the muscle isometric contraction,^{28,29,32} one of which also assessed the maximal voluntary contraction and the rate of force development.³² One study considered peak muscle power,³¹ one-hand grip, and pinch strength,³⁰ and the other performed physical tests and no instrumental evaluation.³⁸ In this review, we have considered only the effects tested at the peak of the medication cycle. As reported in Table 2, a statistically significant improvement was reported between the strength training group and the control group in two studies that used weight machines.^{28,31} Moreover, a statistically significant improvement in manual dexterity was reported after a single-hand exercise session with therapeutic putty.³⁰ On the contrary, Carvalho et al reported a statistical improvement within both the strength- and treadmill-training groups but no statistically significant improvement was highlighted between the groups and with respect to the physiotherapy group.³⁸ Dibble et al also reported a statistically significant improvement within the groups but not between the strength training group and the control group.²⁹ The study reported by Schlenstedt et al showed an improvement in muscle strength both in the strength training group and control group, but the data are not statistically significant.³²

Due to the heterogeneity of data reported in the selected studies, it was not possible to perform a meta-analysis to highlight the effect of resistance training in improving PD muscle strength from baseline to post-intervention.

Secondary outcomes: physical parameters

The studies reviewed have analyzed a variety of outcomes to assess the effects of training in people with PD, including both physical parameters and quality of life parameters, as reported in Table 2. For the secondary outcomes, we also considered the effects tested at the peak of the medication cycle.

Seven studies assessed balance using both self-reported scales and specific physical tests. Mini-Balance Evaluation System Test,²⁶ Berg Balance Scale,^{27,37,38} Activities-Specific Balance Confidence Scale (ABC),^{27,35,37} FAB Scale,³² Choice Stepping Reaction Time,³¹ Single Leg Stand Time,^{31,35} Maximum Balance Range,³¹ Limit of Stability Test,^{34,35} Center of Mass Displacement,^{32,37} and the Latency of Compensatory Postural Response to External Perturbation Test³⁶ were administered. Regardless of intervention type, most of the articles reviewed reported an improvement in balance in agreement with the literature data.^{19,41,42} However, a statistically significant improvement highlighted by the ABC test was reported in the study of Combs et al in the group that carried out traditional exercises compared to the box training group.²⁷ Shen and Mak also reported a statistically significant improvement in terms of balance/limit of stability in repetitive step group training with respect to the strength training group, although there was no statistically significant improvement between groups.^{35,36} It is worth noting that Volpe et al pointed out a statistically significant improvement in the hydrotherapy group with respect to physiotherapy.³⁷

Gait parameters were evaluated in six studies,^{26,27,31,32,34–36} three of which reported an improvement but no statistically significant difference between groups.^{27,34–36} Only one study analyzed aerobic performance with the 2-minute step test and reported a statistically significant improvement in strength- and treadmill-training groups with respect to traditional physiotherapy exercises.³⁸ The same study reported improvement in former two trainings with regard to electroencephalographic activity, while only treadmill training determined a moderate effect in PD subjects' flexibility.³⁸

Only one study assessed the effect of strength training with respect to freezing of gait events using Freezing of Gait Questionnaire, but no group difference was highlighted.³¹ Four studies reported the effect of training on the number of falls.^{31,34–37} Only Volpe et al pointed out an improvement in PD subjects who underwent hydrotherapy.³⁷

Most of the studies selected reported an improvement in Unified Parkinson's Disease Rating Scale, part III (UPDRS-III), but no statistically significant difference between groups.

Regarding the effects of training on physical performance, different tests were administered to PD subjects. The most frequent test was the Timed Up and Go Test.^{26,27,31,33,37,38} Other widely used tests were the 6-Minute Walk Test^{26,27,29} and the 10-Meter Walk Test.^{31,38} Most of the studies reported an improvement in physical performance but only one study evidenced an improvement in PD subjects who underwent strength training.³⁰

Secondary outcome: quality of life parameters

Six of the twelve selected studies assessed the quality of life of PD subjects, with the administration of different questionnaires: UPDRS-II,^{37,38} Parkinson's Disease Questionnaire,^{28,29,32,37} Parkinson's Disease Quality of Life Scale,²⁷ and Beck Depression Inventory.³² Two studies^{29,32} reported no change in outcome, while four studies reported improvement in quality of life.^{27,28,37,38} In particular, two studies highlighted a statistically significant improvement in the strength training group with respect to the control group.^{28,37}

Strengths and limits of this review

The strengths of this review are the selective eligibility criteria and high methodological quality of the articles included. However, we could not answer the question of whether strength training improves motor symptoms and quality of life of PD patients, as we were not able to perform a meta-analysis because data included studies with different designs and different primary and secondary outcomes.

What are the findings?

- Strength training performed against a resistance different from body weight is well tolerated in subjects with mild to moderate PD.
- Strength training improves both physical parameters and quality of life parameters of PD patients.
- To support the beneficial effects of strength training, clinical trials that include specific muscle strength evaluation are required.

Conclusion

Our purpose was to investigate the effect of different types of strength training performed against an external resistance (cycle ergometer, weight machine, elastic band, punching bag, and water) on the improvement of different symptoms of PD patients. In most of the studies selected, the results were positive and strength training appears to be a suitable physical activity to improve both physical parameters and quality of life parameters of PD subjects. Importantly, strength training did not determine a decline in most of the outcomes considered with respect to the other training. Nevertheless, the effect of strength training on balance is uncertain because, on the basis of the items included, data are conflicting. There is also limited evidence of improvements on freezing events. Moreover, the effect on the number of falls is not clear and one study reported no training effect on the majority of outcomes examined.³² These evidences are

Table 2 Details of intervention type and outcomes of the articles reviewed

Study	Intervention type	Number of subjects	Intervention details			Follow-up	Outcomes	
			Intervention (weeks)	Sessions per week	Session lasting (minutes)		Physical parameters	Strength
Arcolin et al ²⁶	Treadmill	13	3	5	2×30+60	Post-treatment		↑*
	Cycle ergometer	16				Post-treatment		↑*
Carvalho et al ³⁸	PKT	9	12	2	40	Post-treatment	↑	=
	Treadmill	5				Post-treatment	↑*	=
	ST (weight machines)	8				Post-treatment	↑*	=
Combs et al ²⁷	Traditional exercise	14 (3)	12	2–3	90	Post-treatment		↑**
	Box	17 (6)				Post-treatment		↑*
Corcos et al ²⁸	mFC	24	104	2	60–90	6 months	↑	
		(1)				12 months	↓	
		(1)				18 months	↓	
		(4)				24 months	↓	
	PREP (weight machines)	24				6 months	↑	
		(1)				12 months	↑**	
		(2)				18 months	↑**	
(1)					24 months	↑**		
Dibble et al ²⁹	Active control group	21 (3)	12	2	60	Post-treatment	↑*	
	Renew (eccentric ergometer)	20 (2)				Post-treatment	↑*	
Mateos-Toset et al ³⁰	Upper limb exercises	30		1	15	Post-treatment	=	
	ST (therapeutic putty)	30				Post-treatment	↑**	
Paul et al ³¹	Control group	20 (2)	12	2	45	Post-treatment	↑	↑/=
	ST (weight machines)	20 (2)				6 months	↑**	↑
		(0)				6 months		
Ridgel et al ³³	Static cycling	25 (2)	1	3	40	Post-treatment		
	Dynamic cycling	25 (1)				Post-treatment		
Schlenstedt et al ³²	BAL	20 (5)	7	2	60	Post-treatment	↑	↑/=
		(4)				4 weeks	=	
	ST (cuff weights, elasticated bands)	20 (3)				Post-treatment	↑	↑*/=
		(3)				4 weeks		=
Shen and Mak ³⁴	RST and visual cues	15 (1)	4	3	60	Post-treatment		
	ST (weight machines and cuff weights)	14				Post-treatment		
Shen and Mak ^{35,36}	BAL (technology-assisted)	26 (4)	12	8	3	60	Post-treatment	
		(1)					3 months	↑*
		(3)					12 months	↑*
	ST (weight machines and cuff weights)	25 (2)		4	5	20	Post-treatment	↑
		(6)					3 months	↑
(6)					12 months	=		
Volpe et al ³⁷	PKT	17	8	5	60	Post-treatment		↑*
	Hydrotherapy	17				Post-treatment		↑**

Notes: ↑, outcomes improvement; ↓, outcomes decline; =, no outcomes change; ↑*, outcomes significant improvement within group; ↑**, outcomes significant improvement between group. Numbers in bracket represent the drop-out events recorded during follow-up.

Abbreviations: ST, strength training; PKT, physiotherapy; mFC, modified fitness counts exercise program; PREP, progressive resistance exercise program; BAL, balance training; RST, repetitive step training; RENEW, Resistance Exercise via Negative Eccentric Work; UPDRS-II, Unified Parkinson's Disease Rating Scale Part II; UPDRS-III, Unified Parkinson's Disease Rating Scale Part III; PDQ-39, Parkinson's Disease Questionnaire; PDQL, Parkinson's Disease Quality of Life Scale; BDI, Beck Depression Inventory; EEG, electroencephalographic activity.

Limit of stability	Gait parameters	Number of falls	Freezing	Physical performance	UPDRS-III	Aerobic performance	Flexibility	Quality of life parameters					
								EEG	UPDRS-II	PDQ-39	PDQL BDI		
↑*				↑*	↑*								
↑*				↑*	↑*								
				↑*/=	=	↓	=	↑	↓				
				↑*	↑*	↑**	↑	↑	↑*				
				↑*	↑*	↑**	=	↑	↑				
↑				↑*									↑*
↑*				↑*									↑*
				↑	↑								
				↑	↑							↑	
				↑	↑							↓	
				↑	↑							↓	
				↑	↑							↓	
				↑	↑							↓	
				↑	↑							↓	
				↑	↑*							↑	
				↑*	↑*							=	
				=								=	
				↑**									
=				=									
		110		↑									
		62		↑									
				↑	↑								
				↑	↑*								
				↑	↑								
				↑*	↑								
				=									
↑**	↑*	0			↑								
=	↑*	1			↑								
↑*	↑*	↑*											
↑*	↑*	↑*											
↑	↑*	↑											
↑*	↑*	↓											
↑	↑*	↓											
↑	↑*	↓											
		↑*		↑*	↑*					↑*	↑*		
		↑**		↑*	↑*					↑*	↑**		

Table 3 Details of strength training of the articles reviewed

Study	Strength training	Strength assessment
Arcolin et al ²⁶	Warm-up: 5 minutes of pedaling at 60 rpm on cycle ergometer Training: 20 minutes of pedaling at a fixed frequency, workload increase maintaining RPE between 11 and 14 Cool-down: 5 minutes	ND
Carvalho et al ³⁸	Warm-up: leg extensions, leg curls, leg presses, chest presses, and low row with minimum load Training: two series of 8–12 maximum repetitions, 90-second rest between exercise sets; intensity 70%–80% of 1RM Cool-down: stretching session	Chair stand test Arm curl test
Combs et al ²⁷	Warm-up: 20 minutes breathing and stretching exercises Training: 45–60 minutes of 3-minute endurance and punching activities, 1-minute rest between exercise sets Cool-down: 15–20 minutes stretching, strengthening, and breathing exercises	ND
Corcos et al ²⁸	Warm-up: 10 minutes of walking and stretching exercises Training: 3 set of 8 repetitions, 6–9 seconds duration of set, pause 2–3 seconds for chest press, later pull downs, reverse fly, double leg press, biceps curl, shoulder press, triceps extension, back extension, knee extension, hip extension, and rotary calf. First week: 30%–40% of 1RM for upper limbs, 50%–60% lower limbs with 5% increase at least for 8 weeks. After 8 weeks, patients performed strength plus speed training: 70%–80% of 1RM, 2 sets of 12 repetitions. Every 8 weeks, patients alternated between strength and strength plus training Cool-down: 10 minutes of walking and stretching exercises	Elbow flexion torque in isometric contraction
Dibble et al ²⁹	Warm-up: 15 minutes stationary bicycling or treadmill Training: 20 minutes of flexibility, balance, and upper/lower extremity concentric resistance training plus 15 minutes of lower extremity eccentric ergometer	Quadriceps KinCom isometric dynamometer
Mateos-Toset et al ³⁰	Training: 15 minutes of hand training with therapeutic putty with a soft medium resistance; exercises include rolling the putty, opening and closing the hands, and exercises involving pinch performance, finger abduction, finger adduction, finger flexion, finger extension, and finger opposition	Jamar dynamometer and pinch meter
Paul et al ³¹	Training: 45 minutes, 3 sets of 8 repetitions using pneumatic resistance equipment. The first set at 40% of 1RM, the second set at 50%, the third set at 60%. 1RM was increased by 5% when patients performed 10 repetitions	Peak muscle power with a pneumatic variable resistance equipment
Ridgel et al ³³	Warm-up: 5 minutes of low resistance pedaling at 40–50 rpm Training: 30 minutes of dynamic cycling with motor output speed between 75 and 85 rpm Cool-down: 5 minutes of low resistance pedaling at 40–50 rpm	ND
Schlenstedt et al ³²	Warm-up: 10 minutes Training: 50 minutes resistance training, 3 sets of 15–20 repetitions of squats, knee extensions, toe/calf raises, hip abductions, and other exercises incrementing resistance with cuff weights, elastic bands, or by therapist; 2-minutes rest between exercise sets	Maximal isometric leg strength, MVC, RFD with a leg press equipped with a force platform
Shen and Mak ³⁴	Training: 2 sets of 15 repetitions for each muscle group at 60% of 1RM using dynamometers and leg-press machines; 1RM was reassessed after 2 weeks of training. In addition, subjects performed rowing exercises, repetitive step on a 6-inch curb, and walking with 1–1.5 kg sandbag strapped to each ankle, 3-minutes each, increasing repetitions within the set duration	ND
Shen and Mak ^{35,36}	Training laboratory based: 2 sets of 15 repetitions for each muscle group at 60% 1RM using dynamometers and leg-press machines; 1RM was reassessed after 2 weeks of training. In addition, subjects performed rowing exercises, repetitive step on a 6-inch curb, and walking with 0.5–1.5 kg sandbag strapped to each ankle, 3 minutes each, increasing repetitions within the set duration Home training: 20 minutes of stepping and walking with 0.5–1.5 kg sandbag strapped to each ankle, increasing repetitions within the 2-week training	ND
Volpe et al ³⁷	Warm up: 10 minutes cardiovascular and stretching exercises Training: 40 minutes of hydrotherapy treatment with perturbation-based balance and strength training Cool-down: 10-minutes	ND

Abbreviations: RPE, rating of perceived exertion; 1RM, one repetition maximum; MVC, maximal voluntary contraction; RFD, rate of force development; ND, not determined.

consistent with other recent reviews that assessed the effect of strength training on PD.^{43–45}

We would like to point out that, despite the fact that the study intervention included strength training, only six

out of twelve articles selected assessed the improvement of muscle strength; moreover, only two studies reported muscle power/strength as primary outcome. Mainly balance, gait parameters, and measure of mobility were assessed. On the

basis of these evidences, it is hard to establish a correlation between strength training and the improvements highlighted. It is worth noting that an improvement in strength correlates positively with physical performance/UPDRS-III results, but the correlation with balance is not well defined.

Despite the encouraging results, our review highlighted the lack of common intent in terms of strength training, control group training, intervention design, and outcomes. Accordingly, further studies are necessary to support the beneficial effects of different types of strength training in PD subjects and to underline the superiority of strength training in people with PD with respect to other training.

Acknowledgment

The authors thank Maurizio Agosti for statistical analysis.

Disclosure

The authors report no conflicts of interest in this work.

References

- Dickson DW. Parkinson's disease and parkinsonism: neuropathology. *Cold Spring Harb Perspect Med*. 2012;2(8):a009258.
- Kalia LV, Lang AE. Parkinson's disease. *Lancet*. 2015;386(9996):896–912.
- Schrag A, Sauerbier A, Chaudhuri KR. New clinical trials for nonmotor manifestations of Parkinson's disease. *Mov Disord*. 2015;30(11):1490–1504.
- Goldman JG, Postuma R. Premotor and nonmotor features of Parkinson's disease. *Curr Opin Neurol*. 2014;27(4):434–441.
- Jones DR, Moussaud S, McLean P. Targeting heat shock proteins to modulate α -synuclein toxicity. *Ther Adv Neurol Disord*. 2014;7(1):33–51.
- Lashuel HA, Overk CR, Oueslati A, Masliah E. The many faces of α -synuclein: from structure and toxicity to therapeutic target. *Nat Rev Neurosci*. 2013;14(1):38–48.
- Bellou V, Belbasis L, Tzoulaki I, Evangelou E, Ioannidis JP. Environmental risk factors and Parkinson's disease: an umbrella review of meta-analyses. *Parkinsonism Relat Disord*. 2016;23:1–9.
- Chin-Chan M, Navarro-Yepes J, Quintanilla-Vega B. Environmental pollutants as risk factors for neurodegenerative disorders: Alzheimer and Parkinson diseases. *Front Cell Neurosci*. 2015;9:124.
- Klein C, Westenberger A. Genetics of Parkinson's disease. *Cold Spring Harb Perspect Med*. 2012;2(1):a008888.
- Verstraeten A, Theuns J, Van Broeckhoven C. Progress in unraveling the genetic etiology of Parkinson disease in a genomic era. *Trends Genet*. 2015;31(3):140–149.
- Jankovic J, Aguilar LG. Current approaches to the treatment of Parkinson's disease. *Neuropsychiatr Dis Treat*. 2008;4(4):743–757.
- Connolly BS, Lang AE. Pharmacological treatment of Parkinson disease: a review. *JAMA*. 2014;311(16):1670–1683.
- Borrione P, Tranchita E, Sansone P, Parisi A. Effects of physical activity in Parkinson's disease: a new tool for rehabilitation. *World J Methodol*. 2014;4(3):133–143.
- Shariffar S, Coronado RA, Romero S, Azari H, Thigpen M. The effects of whole body vibration on mobility and balance in Parkinson disease: a systematic review. *Iran J Med Sci*. 2014;39(4):318–326.
- Paillard T, Rolland Y, de Souto Barreto P. Protective effects of physical exercise in Alzheimer's disease and Parkinson's disease: a narrative review. *J Clin Neurol*. 2015;11(3):212–219.
- Klamroth S, Steib S, Devan S, Pfeifer K. Effects of exercise therapy on postural instability in Parkinson disease: a meta-analysis. *J Neurol Phys Ther*. 2016;40(1):3–14.
- Tuon T, Valvassori SS, Dal Pont GC, et al. Physical training prevents depressive symptoms and a decrease in brain-derived neurotrophic factor in Parkinson's disease. *Brain Res Bull*. 2014;108:106–112.
- Reynolds GO, Otto MW, Ellis TD, Cronin-Golomb A. The therapeutic potential of exercise to improve mood, cognition, and sleep in Parkinson's disease. *Mov Disord*. 2016;31(1):23–38.
- Lamotte G, Rafferty MR, Prodoehl J, et al. Effects of endurance exercise training on the motor and non-motor features of Parkinson's disease: a review. *J Parkinsons Dis*. 2015;5(1):21–41.
- Hirsch MA, Iyer SS, Sanjak M. Exercise-induced neuroplasticity in human Parkinson's disease: what is the evidence telling us? *Parkinsonism Relat Disord*. 2016;22(Suppl 1):S78–S81.
- Petzinger GM, Fisher BE, McEwen S, Beeler JA, Walsh JP, Jakowec MW. Exercise-enhanced neuroplasticity targeting motor and cognitive circuitry in Parkinson's disease. *Lancet Neurol*. 2013;12(7):716–726.
- Shu HF, Yang T, Yu SX, et al. Aerobic exercise for Parkinson's disease: a systematic review and meta-analysis of randomized controlled trials. *PLoS One*. 2014;9(7):e100503.
- Tambosco L, Percebois-Macadré L, Rapin A, Nicomette-Bardel J, Boyer FC. Effort training in Parkinson's disease: a systematic review. *Ann Phys Rehabil Med*. 2014;57(2):79–104.
- Lima LO, Scianni A, Rodrigues-de-Paula F. Progressive resistance exercise improves strength and physical performance in people with mild to moderate Parkinson's disease: a systematic review. *J Physiother*. 2013;59(1):7–13.
- Esco MR. Resistance training for health and fitness. In: Medicine ACoS, editor. *American College of Sports Medicine*. Indianapolis: American College of Sport Medicine; 2013:1–2.
- Arcolin I, Pisano F, Delconte C, et al. Intensive cycle ergometer training improves gait speed and endurance in patients with Parkinson's disease: a comparison with treadmill training. *Restor Neurol Neurosci*. 2015;34(1):125–138.
- Combs SA, Diehl MD, Chrzastowski C, et al. Community-based group exercise for persons with Parkinson disease: a randomized controlled trial. *NeuroRehabilitation*. 2013;32(1):117–124.
- Corcos DM, Robichaud JA, David FJ, et al. A two-year randomized controlled trial of progressive resistance exercise for Parkinson's disease. *Mov Disord*. 2013;28(9):1230–1240.
- Dibble LE, Foreman KB, Addison O, Marcus RL, LaStayo PC. Exercise and medication effects on persons with Parkinson disease across the domains of disability: a randomized clinical trial. *J Neurol Phys Ther*. 2015;39(2):85–92.
- Mateos-Toset S, Cabrera-Martos I, Torres-Sánchez I, Ortiz-Rubio A, González-Jiménez E, Valenza MC. Effects of a single hand-exercise session on manual dexterity and strength in persons with Parkinson disease: a randomized controlled trial. *PMR*. 2016;8(2):115–122.
- Paul SS, Canning CG, Song J, Fung VS, Sherrington C. Leg muscle power is enhanced by training in people with Parkinson's disease: a randomized controlled trial. *Clin Rehabil*. 2014;28(3):275–288.
- Schlenstedt C, Paschen S, Kruse A, Raethjen J, Weisser B, Deuschl G. Resistance versus balance training to improve postural control in Parkinson's disease: a randomized rater blinded controlled study. *PLoS One*. 2015;10(10):e0140584.
- Ridgel AL, Phillips RS, Walter BL, Discenzo FM, Loparo KA. Dynamic high-cadence cycling improves motor symptoms in Parkinson's disease. *Front Neurol*. 2015;6:194.
- Shen X, Mak MK. Repetitive step training with preparatory signals improves stability limits in patients with Parkinson's disease. *J Rehabil Med*. 2012;44(11):944–949.

35. Shen X, Mak MK. Balance and gait training with augmented feedback improves balance confidence in people with Parkinson's disease: a randomized controlled trial. *Neurorehabil Neural Repair*. 2014; 28(6):524–535.
36. Shen X, Mak MK. Technology-assisted balance and gait training reduces falls in patients with Parkinson's disease: a randomized controlled trial with 12-month follow-up. *Neurorehabil Neural Repair*. 2015; 29(2):103–111.
37. Volpe D, Giantin MG, Maestri R, Frazzitta G. Comparing the effects of hydrotherapy and land-based therapy on balance in patients with Parkinson's disease: a randomized controlled pilot study. *Clin Rehabil*. 2014;28(12):1210–1217.
38. Carvalho A, Barbirato D, Araujo N, et al. Comparison of strength training, aerobic training, and additional physical therapy as supplementary treatments for Parkinson's disease: pilot study. *Clin Interv Aging*. 2015; 10:183–191.
39. Tomlinson CL, Herd CP, Clarke CE, et al. Physiotherapy for Parkinson's disease: a comparison of techniques. *Cochrane Database Syst Rev*. 2014; (6):CD002815.
40. Mehrholz J, Kugler J, Storch A, Pohl M, Elsner B, Hirsch K. Treadmill training for patients with Parkinson's disease. *Cochrane Database Syst Rev*. 2015;(8):CD007830.
41. Dibble LE, Addison O, Papa E. The effects of exercise on balance in persons with Parkinson's disease: a systematic review across the disability spectrum. *J Neurol Phys Ther*. 2009;33(1):14–26.
42. Mansfield A, Wong JS, Bryce J, Knorr S, Patterson KK. Does perturbation-based balance training prevent falls? Systematic review and meta-analysis of preliminary randomized controlled trials. *Phys Ther*. 2015; 95(5):700–709.
43. Cruickshank TM, Reyes AR, Ziman MR. A systematic review and meta-analysis of strength training in individuals with multiple sclerosis or Parkinson disease. *Medicine (Baltimore)*. 2015;94(4):e411.
44. Uhrbrand A, Stenager E, Pedersen MS, Dalgas U. Parkinson's disease and intensive exercise therapy – a systematic review and meta-analysis of randomized controlled trials. *J Neurol Sci*. 2015;353(1–2):9–19.
45. Saltychev M, Bärlund E, Paltamaa J, Katajapuu N, Laimi K. Progressive resistance training in Parkinson's disease: a systematic review and meta-analysis. *BMJ Open*. 2016;6(1):e008756.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine,

CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress