

Circadian preferences of birdwatchers in Poland: do “owls” prefer watching night birds, and “larks” prefer daytime ones?

Martyna Frątczak¹, Tim H. Sparks¹, Christoph Randler² and Piotr Tryjanowski¹

¹ Poznan University of Life Sciences, Poznan, Poland

² Eberhard-Karls-Universität Tübingen, Tübingen, Germany

ABSTRACT

Birdwatching is a very popular and increasing leisure activity, and the study and observation of birds is more popular in the morning due to the greater activity among birds at that time of day. The aim of our study was to find out whether there was a relationship between the circadian preference of observers and their favourite bird species and whether it was influenced by such factors as professional status, age and gender. In an e-mail survey we asked a total of 433 Polish ornithologists (professionals) or birdwatchers (non-professionals) for their morningness–eveningness preferences (four categories) and favourite (open choice) bird species and received 143 responses. The temporal (circadian) preferences of respondents declined from early morning (35.7%) to evening/nighttime (11.4%). Circadian preference categories differed significantly by age, with early morning respondents significantly older. These preference categories did not differ significantly in terms of response time to the survey invitation or in the percentage of their favourite birds that were categorised as daytime birds. A total of 204 species were identified as favourite birds of which 34 species were mentioned by five or more respondents, with only two, the common crane *Grus grus* and the Eurasian pygmy owl *Glaucidium passerinum* mentioned by more than 10% of respondents. The white stork *Ciconia ciconia* was more popular with professionals than non-professionals and the swift *Apus apus* less popular. A significant gender × circadian preference interaction was detected for the percentage of favourite birds categorised as daytime birds, with fewer daytime birds among early morning female recorders. The presented results are obviously of a correlative nature, but open the door for further, more advanced study and suggest there may be a need to investigate temporal biases when analysing citizen-based data.

Submitted 31 May 2019
Accepted 31 January 2020
Published 2 March 2020

Corresponding author
Piotr Tryjanowski,
piotr.tryjanowski@gmail.com

Academic editor
Michael Wink

Additional Information and
Declarations can be found on
page 9

DOI 10.7717/peerj.8673

© Copyright
2020 Frątczak et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Animal Behavior, Anthropology, Zoology

Keywords Circadian preference, Environmental attitudes, Morningness–eveningness, Birdwatching

INTRODUCTION

Many factors influence human attitudes to animals, including animal traits such as colour, size, rarity and popularity and human traits such as gender, age, education level and even food preferences (reviewed in: *Prokop & Randler, 2018*). Obviously, both animal and

human perception have co-evolved, and the latter changes due to life experience and education (Prokop, Kubiato & Fančovičová, 2008; Prokop & Randler, 2018). Recently, dealings with animals have not necessarily been linked just to a targeted use (e.g. food source, guard role, beast of burden), but simply watching animals for fun, of which birdwatching is one example, is increasing worldwide (Kronenberg, 2016; Callaghan et al., 2018). Bird species differ in many ecological traits and these can also affect human behaviour, for example searching for information on particular species (Žmihorski et al., 2013; Correia et al., 2016, 2017) and even willingness to pay for their conservation (Raab, Randler & Bogner, 2018). However, people watching birds, and other taxa, differ in many psychological and physical traits and one very marked behavioural difference is their particular morningness–eveningness spectrum of activity (Werner, Geisler & Randler, 2015). In the case of bird observation, this can affect not only intellectual and physical peak performance during the day (Pica et al., 2015; Randler, Faßl & Kalb, 2017; Arrona-Palacios & Díaz-Morales, 2018), but also punctuality (Werner, Geisler & Randler, 2015) and risk-taking (Wang & Chartrand, 2015), which can be useful in finding and watching rare birds, which is a particularly focussed activity of some birdwatchers (Žmihorski, Sparks & Tryjanowski, 2012; Callaghan et al., 2018).

The morningness–eveningness preference may change in relation to age, gender and even seasons (Pica et al., 2015; Randler, Faßl & Kalb, 2017), which also makes this interesting to study in terms of the potential relationship between humans and birds. Birds are a very variable taxonomical group. There are more than 10,000 species, they live in different habitats and they have different seasonal and daily activities mainly related to breeding, wintering and seasonal migrations (Del Hoyo, Elliott & Sargatal, 1992–2011; Jetz et al., 2012), all of which combine to strongly affect people who watch and study birds (Callaghan et al., 2018). Because people have different financial resources, time availability, personal preferences, skills, knowledge and commitment, they may focus on different bird species/taxonomic groups (Lee & Scott, 2004; Prokop, Kubiato & Fančovičová, 2008; Yarwood, Weston & Garnett, 2014). Furthermore, the psychological terminology on circadian preference and morningness–eveningness has always been linked to bird behaviour, as shown by use of the classical phrase ‘owls vs. larks’ (Pica et al., 2015; Putilov, Donskaya & Verevkin, 2015). Furthermore, differences in daily patterns of data collection and a focus on particular bird species may attach a bias to obtained results. Given the current debate on the importance of data quality (La Sorte & Somveille, 2020) this is a very pertinent issue, also in the context of bird monitoring schemes (Snäll et al., 2011).

However, we do not know of a single study that has linked morningness–eveningness to preferences of particular bird species or groups and therefore it is difficult to specify exact testable hypotheses. However, we hazard a guess that people with stronger morningness than eveningness will prefer morning rather than nocturnally active bird species and vice versa. Therefore, based on an e-mail questionnaire to birdwatchers, we attempted to answer whether: (1) if there is a relationship between observer morningness–eveningness and favourite bird species; (2) if there are differences between amateur birdwatchers and professional ornithologists in this human trait; (3) if potential differences are influenced by other factors (age and gender).

METHODS

Data collection

At 00:00 h on the 19 February 2019 the first author (MF) sent an e-mail questionnaire to 433 (356 male, 77 female) Polish ornithologists and advanced birdwatchers and asked for a response by e-mail. Potential participants were chosen by preparing a database of e-mail addresses using information provided in scientific publications, including short notes published in Polish naturalists' journals dedicated to birds. We accessed 28 journals publishing even local records of rare species, unusual behaviour, species lists and even notes on untypical plumage (mainly in Polish. The whole list of accessed journals is available in [Supplemental Material](#)). Participants were asked for up to five favourite bird species (without any suggestion as to local or exotic), their year of birth and their classical circadian general type as used in similar studies ([Randler, Schredl & Göritz, 2017](#)) as follows: (A) definitely morning type; (B) more morning than evening type; (C) more evening than morning type; and (D) definitely evening and night type. Measuring circadian preference in this way could be criticised because it is based on a single question, but [Loureiro & Garcia-Marques \(2015\)](#) showed that results were rather similar, whether based on only a single question or derived from the full scale. [Randler, Schredl & Göritz \(2017\)](#) showed that results were comparable, irrespective of whether a four-item version or a single-item measurement was used; both measures of chronotype revealed identical results. This is also in line with [Loureiro & Garcia-Marques \(2015\)](#). We closed the survey two weeks after the initial e-mail posting. For each respondent we recorded when their reply was sent and used this to calculate the time to respond, expressed in decimal hours.

Statement on ethical approval

Research reported in this manuscript has been conducted in an ethical and responsible manner, in full compliance with all relevant codes of experimentation and legislation.

Data analyses

For further analyses we added information on gender and we divided participants, according to their academic degrees and jobs, into two general categories: professional and non-professional (amateurs). The suggested favourite bird species were divided into two broad categories—nighttime birds (such as owls, rails etc.) and daytime birds (the majority of passerines, cranes, ducks etc.) using data from handbooks (mainly [Del Hoyo, Elliott & Sargatal, 1992–2011](#)). For each respondent the percentage of favourite birds that were daytime birds was calculated.

Because of the rather skew nature of some of the variables it was decided to focus on nonparametric tests. Kruskal Wallis tests, adjusted for ties, were used to compare scale variables to categories and chi squared tests to compare categories with one another. Where expected numbers were so low as to question the validity of chi squared results, Fisher exact tests were used. For testing individual species such tests were restricted to species mentioned by 10 or more respondents. Spearman correlations were used to compare scale variables. Binary logistic regression was used to compare the proportions of

favourite birds identified as daytime birds to year of birth, professional status, gender and time preference category. The model initially included all main effects and two-way interactions, but nonsignificant two-way interactions were dropped one-by-one until only the main effects and any significant interactions remained.

Year of birth was used in analyses, but is presented as age in results. The significance threshold in all analyses was taken as $P = 0.05$. All analysis was carried out in Minitab 18 and SPSS 24.

RESULTS

Description of participants

Complete or near complete responses were received from 143 individuals (33.0%), with response times ranging from 3 min to 14 days (the cut-off point). Response rates were 31.7% for males and 39.0% for females with no significant difference between the two ($\chi^2_1 = 0.73$, $P = 0.394$).

Response times were positively skewed with a median of 10 h and a mean of 44 h. Years of birth were provided by 140 individuals, indicating a mean age of 45 years (range 22–85). The professional status of the 142 birders providing detail was roughly evenly split between non-professionals (51.4%) and professionals (48.6%). As expected from the targeted individuals, the majority of respondents (79%) were male. The circadian preferences of respondents changed from early morning; 50 (35.7%) coded A, 41 (29.3%) coded B, 33 (23.6%) coded C and 16 (11.4%) coded D, 3 people didn't provide a response to this question.

It is worth noting that 11 complaints (mainly about the aims and methods of the study) were also received, which in combination with other responses indicate a response rate of 35.6% of the targeted population. Nine of the complaints came from professional birders and a chi squared test indeed confirmed that professionals were significantly more likely to complain ($\chi^2_1 = 4.70$, $P = 0.030$). There was no significant association between gender and complaining ($\chi^2_1 = 1.18$, $P = 0.277$). Furthermore, those complaining were significantly older ($H = 11.91$, $P = 0.001$, medians 64, 42 years).

Professional birdwatchers were significantly older ($H = 5.89$, $P = 0.015$) than non-professionals (median ages 47 and 41 years respectively), but did not differ significantly in response times to the survey ($H = 0.72$, $P = 0.395$) or in the percentage of their favourite birds that were categorised as daytime birds ($H = 0.29$, $P = 0.589$). There was no significant association between professional status and gender ($\chi^2_1 = 0.76$, $P = 0.384$) or between professional status and circadian preference ($\chi^2_3 = 5.34$, $P = 0.149$).

Female respondents were significantly younger than male respondents ($H = 11.91$, $P = 0.001$, medians of 37 and 45 years respectively), but did not differ significantly in terms of response times ($H = 0.04$, $P = 0.835$) or percentage of their favourite birds that were categorised as daytime birds ($H = 0.12$, $P = 0.726$). There was a suggestion that females were less likely to be category A recorders but this did not quite reach significance (Fisher exact test $P = 0.053$).

Circadian preference categories differed significantly in age, with category A respondents noticeably older ($H = 10.56$, $P = 0.014$, medians 48, 41, 40 and 42 years

respectively for categories A–D). Circadian preference categories did not differ significantly in terms of response time ($H = 3.29$, $P = 0.349$) or percentage of their favourite birds that were categorised as daytime birds ($H = 2.29$, $P = 0.514$).

Favourite bird species

A total of 204 species were identified as favourite birds, of which 184 (90.2%) were classified as daytime birds. A full table of species indicating their daytime/nighttime category is given in [Supplemental Material 1](#). The 34 species mentioned by five or more respondents are listed in [Table 1](#). Only two species, the common crane *Grus Grus* and the Eurasian pygmy owl *Glaucidium passerinum*, were mentioned by more than 10% of respondents. Only the 12 species that were mentioned 10 or more times were subject to further analysis. This additional analysis suggested that the white stork *Ciconia ciconia* was more popular with professionals than non-professionals (13% cf. 3%) and the swift *Apus apus* less popular (3% cf. 12%). No significant gender preferences were detected in this subset of species. There was a significant difference in the circadian preference categories for swift which was not mentioned by any of the category A respondents.

There were no significant Spearman correlations between age, response times and percentage of favourite birds that were categorised as daytime birds (all $P > 0.297$).

Binary logistic regression on the proportion of favourite birds categorised as daytime produced an overall significant model ($\chi^2_9 = 23.47$, $P = 0.005$) with one significant main effect (gender: $\chi^2_1 = 5.09$, $P = 0.024$) and one significant interaction (gender \times circadian preference: $\chi^2_3 = 14.26$, $P = 0.003$). In addition, circadian preference was very close to significance ($\chi^2_3 = 7.80$, $P = 0.0503$). This interaction is shown in [Fig. 1](#) where female respondents in category A reported fewer favourite daytime birds than their male counterparts and in contrast to the pattern in other categories.

DISCUSSION

The number of participants may appear lower than expected in western European countries such as the Netherlands, Germany and especially the UK ([Cocker, 2012](#)). But in reality we asked a large proportion of bird observers that usually participate in national citizen science projects in Poland, such as Common Birds Monitoring Scheme, Winter Bird Counts and BirdFeeder Project. These schemes usually attract only 100–180 people. We used an e-mail based questionnaire, with an overall response rate of 35.6% which may appear low, but is similar to the average rate in other studies using similar methods in psychology ([Cook, Heath & Thompson, 2000](#); [Sheehan, 2001](#)). We also noted with great interest the few complaining e-mails where respondents, mainly professionals, expressed concerns over the aims and methodology of the study. It may seem obvious that a significantly greater rate of complaint would derive from professional ornithologists rather than amateur birdwatchers, because professionals focus much more on science than simply gaining pleasure from watching birds ([Yarwood, Weston & Garnett, 2014](#); [Yarwood, Weston & Symonds, 2019](#) and see also below).

The study also suggests that watching birds is still more popular among men than women, similar to other studies in Poland, but which were based on a much smaller sample

Table 1 The 34 most popular species (those with $n \geq 5$) reported by respondents. In addition to the overall number and percentages, percentages are also given by professional status (NP nonprofessional, P professional), gender (F female, M male) and circadian preference (categories A–D, see main text). Significance (indicated by *) tested using chi squared or Fisher exact tests for the 12 species with $n \geq 10$. Species in grey shading were categorised as nighttime; all others as daytime.

Species	<i>n</i>	%	NP%	P%	F%	M%	A%	B%	C%	D%	
<i>Grus grus</i>	28	19.6	23.3	15.9	23.3	18.6	20.0	29.3	15.2	6.3	
<i>Glaucidium passerinum</i>	16	11.2	12.3	10.1	10.0	11.5	10.0	14.6	9.1	12.5	
<i>Alcedo atthis</i>	14	9.8	11.0	8.7	10.0	9.7	12.0	7.3	9.1	12.5	
<i>Upupa epops</i>	14	9.8	11.0	8.7	10.0	9.7	8.0	12.2	12.1	6.3	
<i>Turdus merula</i>	13	9.1	11.0	7.2	6.7	9.7	14.0	2.4	9.1	12.5	
<i>Aegithalos caudatus</i>	12	8.4	8.2	8.7	16.7	6.2	8.0	9.8	9.1	6.3	
<i>Ciconia ciconia</i>	11	7.7	2.7	13.0	*	3.3	8.8	12.0	4.9	9.1	0.0
<i>Vanellus vanellus</i>	11	7.7	8.2	7.2	6.7	8.0	2.0	9.8	9.1	18.8	
<i>Apus apus</i>	11	7.7	12.3	2.9	*	10.0	7.1	0.0	14.6	12.1	6.3
<i>Merops apiaster</i>	11	7.7	9.6	5.8	10.0	7.1	10.0	2.4	12.1	0.0	
<i>Haliaetus albicilla</i>	10	7.0	8.2	5.8	6.7	7.1	8.0	9.8	0.0	6.3	
<i>Lanius collurio</i>	10	7.0	5.5	8.7	6.7	7.1	10.0	2.4	9.1	6.3	
<i>Strix aluco</i>	9	6.3	9.6	2.9	6.7	6.2	6.0	12.2	3.0	0.0	
<i>Circus pygargus</i>	9	6.3	8.2	4.3	6.7	6.2	6.0	9.8	3.0	6.3	
<i>Corvus monedula</i>	9	6.3	2.7	10.1	10.0	5.3	8.0	7.3	6.1	0.0	
<i>Ciconia nigra</i>	8	5.6	8.2	2.9	6.7	5.3	6.0	7.3	3.0	6.3	
<i>Corvus frugilegus</i>	8	5.6	4.1	7.2	6.7	5.3	10.0	0.0	6.1	6.3	
<i>Corvus corax</i>	8	5.6	8.2	2.9	13.3	3.5	4.0	7.3	0.0	12.5	
<i>Galerida cristata</i>	8	5.6	6.8	4.3	3.3	6.2	8.0	4.9	3.0	6.3	
<i>Erithacus rubecula</i>	7	4.9	6.8	2.9	3.3	5.3	8.0	4.9	3.0	0.0	
<i>Bucephala clangula</i>	7	4.9	4.1	5.8	3.3	5.3	6.0	2.4	6.1	6.3	
<i>Sturnus vulgaris</i>	6	4.2	5.5	2.9	3.3	4.4	8.0	0.0	6.1	0.0	
<i>Parus major</i>	6	4.2	1.4	7.2	3.3	4.4	6.0	0.0	6.1	6.3	
<i>Lyrurus tetrix</i>	6	4.2	5.5	2.9	0.0	5.3	2.0	7.3	6.1	0.0	
<i>Athene noctua</i>	6	4.2	4.1	4.3	10.0	2.7	0.0	0.0	9.1	18.8	
<i>Aquila chrysaetos</i>	6	4.2	8.2	0.0	6.7	3.5	2.0	9.8	0.0	6.3	
<i>Limosa limosa</i>	6	4.2	5.5	2.9	3.3	4.4	2.0	7.3	3.0	6.3	
<i>Tyto alba</i>	5	3.5	4.1	2.9	0.0	4.4	6.0	0.0	6.1	0.0	
<i>Milvus milvus</i>	5	3.5	4.1	2.9	3.3	3.5	6.0	0.0	3.0	6.3	
<i>Tetrao urogallus</i>	5	3.5	2.7	4.3	3.3	3.5	0.0	7.3	6.1	0.0	
<i>Sylvia nisoria</i>	5	3.5	2.7	4.3	0.0	4.4	2.0	4.9	6.1	0.0	
<i>Coracias garrulus</i>	5	3.5	2.7	4.3	3.3	3.5	4.0	2.4	0.0	12.5	
<i>Strix nebulosa</i>	5	3.5	2.7	4.3	6.7	2.7	4.0	0.0	3.0	12.5	
<i>Clanga pomarina</i>	5	3.5	6.8	0.0	3.3	3.5	4.0	2.4	0.0	12.5	

size (Skłodowski & Jurkowska, 2015). However, it is interesting to note that women participating in our study were significantly younger than men, which is a sign of an encouraging (probably global) trend that birdwatching has recently become more popular with women (Cooper & Smith, 2010). The most represented circadian preference was the




Figure 1 The mean percentage \pm SE of favourite birds classified as daytime by gender and circadian preference. [Full-size](#) DOI: 10.7717/peerj.8673/fig-1

‘early morning type,’ a finding similar to that found by [Raab, Randler & Bogner \(2018\)](#). Moreover, those authors even suggested that students with a morningness preference were more likely to be protective of and appreciative towards, nature, including birds. Furthermore, early morning (lark type) people more often participated in voluntary activities, shared information, were open to new ideas, were faster and more punctual ([Song & Stough, 2000](#); [Werner, Geisler & Randler, 2015](#)). On the other hand, if we compare birders with the general Polish population ([Jankowski, 2013](#); [Tęgowska, Sobocińska & Maliszewska, 2014](#)) there is still apparent a visible bias to the early morning type. In a correlative-type study like ours it is impossible to determine if early morning types are more likely to take up an ornithological career, or indeed if they just have to get up early to watch more bird species. This can only be resolved if a representative sample of Polish people was surveyed and matched for gender and age, because there are significant age effects in circadian preference in the adult population ([Randler et al., 2016](#)).

The 143 respondents mentioned 204 species as favourite birds with great variation between respondents, but some patterns are worth noting. Only two species, the common crane and the Eurasian pygmy owl, were mentioned by more than 10% of respondents. We did not ask why people decided on their particular species, but Eurasian pygmy owl was a surprise to us. The common crane is a more obvious choice because it is a big charismatic bird with a characteristic call, an increasing population size and is also represented in Polish culture and art ([Wqs, 2013](#)). Interestingly, the white stork was a more popular choice with professionals than with non-professionals but the swift showed an

opposite pattern. The white stork is a Polish icon, well recognised, easy-to-spot even by beginners and very popular in both scientific studies and the cultural life of Poland (Kronenberg, Andersson & Tryjanowski, 2017). There are long term-data on white storks that permit advanced statistical analyses and so it is often chosen as a study species by ornithologists, although it is not very attractive to classical birdwatchers because it is still common, but is more exciting to visitors to Poland rather than local birdwatchers (Kronenberg, 2016; Kronenberg, Andersson & Tryjanowski, 2017). In choosing study species, ornithologists follow different criteria, based on the probability of obtaining a good sample size, conservation status and even size and behaviour (Yarwood, Weston & Garnett, 2014; Yarwood, Weston & Symonds, 2019). After some years of study, they know much about the biology of the species, including its secrets and are fascinated by the species which then also becomes a favourite bird (personal experience of the authors). It is likely that preferences change over time and are sensitive to mass-media information making species more or less charismatic in recent years (Żmihorski, Sparks & Tryjanowski, 2012; Correia et al., 2016, 2017). In our sample, swift is perhaps an indicator of changes in recent times, because the species is an object of recent debate focused on bird conservation in urban areas and information has been promoted on this species biology, ecology and importance to humans (Luniak & Grzeniewski, 2011). It is interesting that this species was not suggested as a favourite bird by any of the 'early morning' respondents, which fits especially well with the late afternoon and early evening activity of swifts.

A number of other interesting patterns appear evident in Table 1 but failed to achieve either significance or sufficient sample size for testing. A study based on a larger sample would be needed to confirm or refute any of these apparent patterns. Probably a more advanced study should compare species choice with other factors, for example how much attention she/he pays to that particular bird species, how many publications are dedicated to that particular species and even a measure of emotional attachment. Similar studies could be useful for understanding the choice of different outdoor activities between people, as well as a deeper understanding of the variability of other groups of passionate recreationalists (e.g. anglers, hunters, mountain climbers, cf. Wałęga, Wałęga & Graczyk, 2017). Given that much of the data used to estimate species population sizes and geographical distributions and to produce bird guides, originate from recorders of which our respondents were a sample, it is important to know if their behaviour may subconsciously influence their records. Furthermore, a better understanding of the reasons for the popularity of species could help enormously in planning for and funding, their conservation. One important aspect that chronobiology may contribute to bird research is the influence on citizen science programs. Citizen science programs, such as eBird rely heavily on the submissions of many volunteer birdwatchers (Wiersma, 2010). However, the reporting of given species may be significantly influenced if the chronotype of the birdwatchers and the circadian rhythms of specific species do not map. Early morning bird species may go underreported if most birdwatchers are not morning oriented themselves. In a similar manner, a higher proportion of 'owls' in birdwatchers may lead to a better registration of owls in these databases. Of course, these assumptions need further study based on larger samples.

CONCLUSION

Although much is known about birds and the importance of citizen science, only scarce evidence is available about the nature of birdwatchers, both professional and amateur. These data, especially about their morning or evening preference, may severely influence data collection behaviour but, in this study, we were able to show that there is only a weak influence of circadian preference on leisure birdwatching; thus, we assume a minimal or even absent influence of circadian preference on birdwatching.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The authors received no funding for this work.

Competing Interests

Piotr Tryjanowski is an Academic Editor for PeerJ.

Author Contributions

- Martyna Frątczak conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Tim H. Sparks analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Christoph Randler conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Piotr Tryjanowski conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

Raw data is available as a [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8673#supplemental-information>.

REFERENCES

- Arrona-Palacios A, Díaz-Morales JF. 2018.** Morningness–eveningness is not associated with academic performance in the afternoon school shift: preliminary findings. *British Journal of Educational Psychology* **88**(3):480–498 DOI [10.1111/bjep.12196](https://doi.org/10.1111/bjep.12196).
- Callaghan CT, Slater M, Major RE, Morrison M, Martin JM, Kingsford RT. 2018.** Travelling birds generate eco-travellers: the economic potential of vagrant birdwatching. *Human Dimensions of Wildlife* **23**(1):71–82 DOI [10.1080/10871209.2017.1392654](https://doi.org/10.1080/10871209.2017.1392654).
- Cocker M. 2012.** *Birders*. London: Random House.

- Cook C, Heath F, Thompson RL. 2000.** A meta-analysis of response rates in web-or internet-based surveys. *Educational and Psychological Measurement* **60**(6):821–836
DOI [10.1177/00131640021970934](https://doi.org/10.1177/00131640021970934).
- Cooper C, Smith J. 2010.** Gender patterns in bird-related recreation in the USA and UK. *Ecology and Society* **15**(4):4 DOI [10.5751/ES-03603-150404](https://doi.org/10.5751/ES-03603-150404).
- Correia RA, Jepson PR, Malhado AC, Ladle RJ. 2016.** Familiarity breeds content: assessing bird species popularity with culturomics. *PeerJ* **4**(11):e1728 DOI [10.7717/peerj.1728](https://doi.org/10.7717/peerj.1728).
- Correia RA, Jepson P, Malhado AC, Ladle RJ. 2017.** Internet scientific name frequency as an indicator of cultural salience of biodiversity. *Ecological Indicators* **78**:549–555
DOI [10.1016/j.ecolind.2017.03.052](https://doi.org/10.1016/j.ecolind.2017.03.052).
- Del Hoyo J, Elliott A, Sargatal J. 1992–2011.** *Handbook of the birds of the world*. Barcelona: Lynx Edicions.
- Jankowski KS. 2013.** Polish version of the reduced morningness–eveningness questionnaire. *Biological Rhythm Research* **44**(3):427–433 DOI [10.1080/09291016.2012.704791](https://doi.org/10.1080/09291016.2012.704791).
- Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. 2012.** The global diversity of birds in space and time. *Nature* **491**(7424):444–448 DOI [10.1038/nature11631](https://doi.org/10.1038/nature11631).
- Kronenberg J. 2016.** Birdwatchers’ wonderland? Prospects for the development of birdwatching tourism in Poland. *Journal of Ecotourism* **15**(1):78–94 DOI [10.1080/14724049.2016.1142556](https://doi.org/10.1080/14724049.2016.1142556).
- Kronenberg J, Andersson E, Tryjanowski P. 2017.** Connecting the social and the ecological in the focal species concept: case study of White Stork. *Nature Conservation* **22**(1–2):79–105
DOI [10.3897/natureconservation.22.12055](https://doi.org/10.3897/natureconservation.22.12055).
- La Sorte FA, Somveille M. 2020.** Survey completeness of a global citizen-science database of bird occurrence. *Ecography* **43**(1):34–43 DOI [10.1111/ecog.04632](https://doi.org/10.1111/ecog.04632).
- Lee J-H, Scott D. 2004.** Measuring birding specialization: a confirmatory factor analysis. *Leisure Sciences* **26**(3):245–260 DOI [10.1080/01490400490461387](https://doi.org/10.1080/01490400490461387).
- Loureiro F, Garcia-Marques T. 2015.** Morning or evening person? Which type are you? Self-assessment of chronotype. *Personality and Individual Differences* **86**:168–171
DOI [10.1016/j.paid.2015.06.022](https://doi.org/10.1016/j.paid.2015.06.022).
- Luniak M, Grzeniewski M. 2011.** Nest-boxes for the common swift *Apus apus*—experience from Poland. *Ecologia Urbana* **23**:3–5.
- Pica G, Amato C, Pierro A, Kruglanski AW. 2015.** The early bird gets the worm: on locomotors’ preference for morningness. *Personality and Individual Differences* **76**:158–160
DOI [10.1016/j.paid.2014.12.020](https://doi.org/10.1016/j.paid.2014.12.020).
- Prokop P, Kubiato M, Fančovičová J. 2008.** Slovakian pupils’ knowledge of, and attitudes toward birds. *Anthrozoös* **21**(3):221–235.
- Prokop P, Randler C. 2018.** Biological predispositions and individual differences in human attitudes toward animals. In: Alves RRN, Albuquerque UP, eds. *Ethnozoology*. Cambridge: Academic Press, 447–466.
- Putilov AA, Donskaya OG, Verevkin EG. 2015.** How many diurnal types are there? A search for two further bird species. *Personality and Individual Differences* **72**:12–17
DOI [10.1016/j.paid.2014.08.003](https://doi.org/10.1016/j.paid.2014.08.003).
- Raab P, Randler C, Bogner F. 2018.** How young early birds prefer preservation, appreciation and utilization of nature. *Sustainability* **10**(11):4000 DOI [10.3390/su10114000](https://doi.org/10.3390/su10114000).
- Randler C, Faßl C, Kalb N. 2017.** From lark to owl: developmental changes in morningness–eveningness from new-borns to early adulthood. *Scientific Reports* **7**(1):45874
DOI [10.1038/srep45874](https://doi.org/10.1038/srep45874).

- Randler C, Freyth-Weber K, Rahafar A, Jurado AF, Kriegs JO. 2016. Morningness–eveningness in a large sample of German adolescents and adults. *Heliyon* 2(11):e00200 DOI 10.1016/j.heliyon.2016.e00200.
- Randler C, Schredl M, Göritz AS. 2017. Chronotype, sleep behavior, and the big five personality factors. *SAGE Open* 7(3):2158244017728321 DOI 10.1177/2158244017728321.
- Sheehan KB. 2001. E-mail survey response rates: a review. *Journal of Computer-Mediated Communication* 6(2):JCMC621.
- Skłodowski J, Jurkowska A. 2015. Charakterystyka sylwetki i zainteresowań uczestników turystyki birdwatchingowej w Polsce [Profile and interests of bird-watching tourists in Poland]. *Studia i Materiały Centrum Edukacji Przyrodniczo-Leśnej* 17:203–208.
- Snäll T, Kindvall O, Nilsson J, Pärt T. 2011. Evaluating citizen-based presence data for bird monitoring. *Biological Conservation* 144(2):804–810 DOI 10.1016/j.biocon.2010.11.010.
- Song J, Stough C. 2000. The relationship between morningness–eveningness, time-of-day, speed of information processing, and intelligence. *Personality and Individual Differences* 29(6):1179–1190 DOI 10.1016/S0191-8869(00)00002-7.
- Tęgowska E, Sobocińska P, Maliszewska J. 2014. Disruption of circadian rhythmicity—an exploratory study in Poland. *Personality and Individual Differences* 66:98–101 DOI 10.1016/j.paid.2014.03.007.
- Wang L, Chartrand TL. 2015. Morningness–eveningness and risk taking. *Journal of Psychology* 149(4):394–411 DOI 10.1080/00223980.2014.885874.
- Wałęga A, Wałęga G, Graczyk A. 2017. Leisure time and income: some evidence from Poland. *Economic and Environmental Studies* 17(43):555–575 DOI 10.25167/ees.2017.43.7.
- Werner L, Geisler J, Randler C. 2015. Morningness as a personality predictor of punctuality. *Current Psychology* 34(1):130–139 DOI 10.1007/s12144-014-9246-1.
- Wiersma Y. 2010. Birding 2.0: citizen science and effective monitoring in the Web 2.0 World. *Avian Conservation and Ecology* 5(2):1–9 DOI 10.5751/ACE-00427-050213.
- Wąs A. 2013. Symbolika i wymiar dydaktyczny zwierząt w literaturze (na przykładzie twórczości Juliusza Słowackiego) [The symbolism and educational dimension of animals in literature (as seen in Juliusz Słowacki's works)]. *Prace Literackie* 53:107–115.
- Yarwood MR, Weston MA, Garnett ST. 2014. From little things, big things grow; trends and fads in 110 years of Australian ornithology. *Scientometrics* 98(3):2235–2254 DOI 10.1007/s11192-013-1144-z.
- Yarwood MR, Weston MA, Symonds MR. 2019. Biological determinants of research effort on Australian birds: a comparative analysis. *Emu-Austral Ornithology* 119(1):38–44 DOI 10.1080/01584197.2018.1501274.
- Żmihorski M, Dziarska-Pałac J, Sparks TH, Tryjanowski P. 2013. Ecological correlates of the popularity of birds and butterflies in Internet information resources. *Oikos* 122(2):183–190 DOI 10.1111/j.1600-0706.2012.20486.x.
- Żmihorski M, Sparks TH, Tryjanowski P. 2012. The weekend bias in recording rare birds: mechanisms and consequences. *Acta Ornithologica* 47(1):87–94 DOI 10.3161/000164512X653953.