



## Review Article

Korean J Anesthesiol 2020;73(1):8-29  
<https://doi.org/10.4097/kja.19391>  
pISSN 2005-6419 • eISSN 2005-7563

Received: September 25, 2019

Accepted: October 20, 2019

### Corresponding author:

Il-Ok Lee, M.D., Ph.D.

Department of Anesthesiology and Pain  
Medicine, Korea University Guro Hospital,  
Korea University College of Medicine, 148  
Gurodong-ro, Guro-gu, Seoul 08308, Korea  
Tel: 82-2-2626-3234

Fax: 82-2-2626-1438

Email: [iloklee@korea.ac.kr](mailto:iloklee@korea.ac.kr)

ORCID: <https://orcid.org/0000-0001-8062-1496>

# Anesthetic management of geriatric patients

Byung-Gun Lim, Il-Ok Lee

*Department of Anesthesiology and Pain Medicine, Korea University Guro Hospital, Korea University College of Medicine, Seoul, Korea*

The number of elderly patients who frequently access health care services is increasing worldwide. While anesthesiologists are developing the expertise to care for these elderly patients, areas of concern remain. We conducted a comprehensive search of major international databases (PubMed, Embase, and Cochrane) and a Korean database (KoreaMed) to review preoperative considerations, intraoperative management, and postoperative problems when anesthetizing elderly patients. Preoperative preparation of elderly patients included functional assessment to identify preexisting cognitive impairment or cardiopulmonary reserve, depression, frailty, nutrition, polypharmacy, and anticoagulation issues. Intraoperative management included anesthetic mode and pharmacology, monitoring, intravenous fluid or transfusion management, lung-protective ventilation, and prevention of hypothermia. Postoperative checklists included perioperative analgesia, postoperative delirium and cognitive dysfunction, and other complications. A higher level of perioperative care was required for older surgical patients, as multiple chronic diseases often makes them prone to developing postoperative complications, including functional decline and loss of independence. Although the guiding evidence remains poor so far, elderly patients have to be provided optimal perioperative care through close interdisciplinary, interprofessional, and cross-sectional collaboration to minimize unwanted postoperative outcomes. Furthermore, along with adequate anesthetic care, well-planned postoperative care should begin immediately after surgery and extend until discharge.

**Keywords:** Aged; Anesthesia; Frail elderly; Geriatrics; Perioperative care.

## Introduction

The WHO's 'World Report on Aging and Health' revealed significant impairments in the elderly population in Europe, and the number of elderly people is expected to double by 2050 [1]. As the population of a country continues to age, the demands for surgical services increase. Elderly patients often require a higher level of care than younger patients do during the perioperative period, with higher health care costs. Strategies to optimize anesthesia care to reduce complications and improve outcomes in elderly surgical patients will also be of great value to the individual patients and society.

This article reviews the perianesthetic considerations for geriatric patients according to three conceptual categories: preoperative considerations, intraoperative management, and postoperative problems. It also aims to establish a framework to assess the complex issues related to the perioperative care of elderly patients.

The authors performed a comprehensive literature review based on major international databases (PubMed, Embase, Cochrane) and a Korean database (KoreaMed) to identify systematic reviews, meta-analyses, practice guidelines, and clinical trials published in the last 10 years (Appendix 1). The initial search resulted in 1,551 citations and an additional 15 articles were obtained by a manual search through the related references. Of these, the

© The Korean Society of Anesthesiologists, 2020

© This is an open-access article distributed under the terms of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

authors selected 262 publications, which were further narrowed to 172 based on strength of evidence, relevance to geriatric patients, and focused on the perioperative period (Fig. 1).

## Preoperative Management

### Assessment of functional reserve

Comprehensive geriatric assessment (CGA) in the preoperative period include systematic evaluation of comorbidities, functional status, neurocognitive function, sensory impairment, substance abuse, frailty, nutrition, and medication. Preoperative CGA has a positive impact on postoperative outcomes in older patients undergoing elective surgery [2]. A recent Cochrane review including 1,583 hip fracture surgeries in subjects  $\geq 65$  years showed that CGA probably reduces mortality (risk ratio [RR] 0.85, 95% CI 0.68–1.05) and referral to an increased level of care (RR 0.71, 95% CI 0.55–0.92) [2]. However, CGA may make little or no difference in major postoperative complications and delirium rates [3]. Physicians must therefore recommend their patients to undergo all the appropriate preoperative evaluations and interventions such that they can improve their functional reserve and make an in-

formed decision [4,5]. Furthermore, anesthesiologists are strongly encouraged to get involved in national surveys and clinical outcomes research focused on elderly surgical patients [6].

Baseline functional status should first be evaluated in ambulatory patients using a simple screening test, followed by in-depth or full screening of basic and instrumental activities of daily living (ADL) [7,8]. The patient should be evaluated for limitations in gait and mobility using the Time Up and Go (TUG) test [9,10]. In a 2018 prospective cohort study among 131 patients  $\geq 65$  years old who underwent elective major surgery for cancer, those who were dependent in ADLs and had an unfavorable TUG test had a significantly higher 1-year mortality (odds ratio [OR] 4.5, 95% CI 1.21–18.25,  $P = 0.034$ ) than the other patients. As such, functional assessments such as ADL and TUG test as well as mild cognitive impairment are predictors of long-term outcomes in elderly cancer patients [11]. A 2005 prospective study of 120 patients  $\geq 60$  years old who underwent thoracic surgery also showed dependence in ADLs and impaired cognitive conditions as important predictors of postoperative complications [12].

Age-associated organ reserve decline, compounded by chronic diseases, also leads to high incidence of postoperative complications in the elderly patient, including neurologic, pulmonary, car-




Fig. 1. Flow diagram for article search and study inclusion in this review.

diac, and renal comorbidities. More data on the effectiveness of various components of the CGA are needed, but a 2018 meta-analysis showed that comorbidity (Charlson Comorbidity Index  $\geq 3$ ), polypharmacy ( $\geq 5$  drugs/day), and ADL dependency were predictive factors for postoperative complications in gastrointestinal cancer patients [13,14]. A more independent preoperative functional status strongly predicts better postoperative function and shorter recovery period after major abdominal surgery [15]. Using the one million longitudinal health insurance database in Taiwan, another 2017 retrospective study of patients  $\geq 65$  years old who received anesthesia showed that the leading comorbidity contributing to postoperative mortality is chronic renal failure (OR = 2.806), acute MI (OR = 4.58), and intracranial hemorrhage (OR = 3.758) [16].

The American College of Surgeons (ACS) and the American Geriatrics Society (AGS) together recently proposed a framework for routine multidomain preoperative assessment of geriatric patients. The ACS and AGS are also developing best practice guidelines for optimal perioperative care in geriatric surgical patients [17,18]. One component of this collaboration is the initiation of patient preparation (prehabilitation) before hospital admission through measures including, e.g., correction of any deficiency states, optimization of chronic drug treatment, and respiratory training [19].

## Neurocognitive and behavioral assessment

Dementia has a prevalence of 5% to 8% among people  $\geq 65$  years old and may affect more than one third of individuals  $> 85$  years [20]. As the geriatric population grows, the number of people living with dementia is projected to rise dramatically [21]. A 2018 meta-analysis study showed that the mortality of patients with dementia who underwent hip fracture surgery was increased 1.57, 1.97, 1.77, and 1.60-fold after a 30-day, 6-month, 1-year, and  $> 1$ -year follow up, respectively [22]. A 2014 retrospective cohort study of 45,602 aged patients who had a hip fracture also showed that dementia was associated with an increased risk of long-term care admission (hazard ratio [HR] 2.49, 95% CI 2.38–2.61,  $P < 0.001$ ), higher mortality (HR 1.47, 95% CI 1.41–1.52,  $P < 0.001$ ), and poor prognosis [23].

The recent ACS-AGS Guidelines for Optimal Preoperative Assessment of the Geriatric Patient include a routine preoperative neurocognitive assessment before surgery. For patients without a known history of cognitive impairment, “Mini-Cog” [24] is recommended as a short, easily applicable, and well-studied tool. Careful documentation of the patient’s preoperative cognitive status is critical for diagnosing and anticipating common postopera-

tive complications such as postoperative delirium (POD) or cognitive dysfunction (POCD). Preexisting cognitive impairment predicts POD [25,26]. Postoperative cognitive impairment is associated with longer hospital stay, increased mortality, and functional decline. Explanations may be that patients with impaired cognition are less likely to engage in postoperative aggressive pulmonary hygiene and ambulation, causing a high risk of developing postoperative complications such as pneumonia, deep vein thrombosis, stroke, and cerebrovascular accident with neurologic deficit.

## Cardiac evaluation

Diminished cardiac reserve in elderly patients often manifests as exaggerated drops in blood pressure during induction of general anesthesia (GA). Reductions in the responsiveness of beta-receptors caused by a beta-blocked state limits patients’ ability to increase cardiac output and properly respond to blood losses. Baroreceptor dysfunction and reduced responsiveness to angiotensin II further limit responsiveness to hypovolemia. All these factors may be compounded by comorbid myocardial ischemia related to atherosclerosis.

Assessing the metabolic equivalents (METs) of daily activities is a useful way to assess exercise tolerance for patients who may not participate in regular exercise. Geriatric patients should undergo cardiac tests and risk stratification and evidence-based optimization strategies should be applied before surgery [27]. The web-based ACS National Surgical Quality Improvement Program (NSQUIP) Risk Calculator is one of the recommended tools in the 2014 American College of Cardiology/American Heart Association Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery [27]. However, critical knowledge gaps remain for informed decision-making and recommendations targeting older patients. A 2016 study highlighted a critical need for large population-based studies including a broad spectrum of older patients [28]. A 2002 prospective study of 513 patients  $\geq 70$  years old undergoing non-cardiac surgery showed that abnormalities in preoperative electrocardiogram (ECG) are common but non-specific and thus are of limited value in predicting postoperative cardiac complications compared to the presence and severity of comorbidities [29]. Conversely, echocardiography may help provide insights into ventricular function and valve status; it may be considered in patients with significant cardiac comorbidities such as history of myocardial infarction (MI), congestive heart failure (CHF), or valvular heart disease.

Older patients are more vulnerable to perioperative cardiac ad-

verse events. A 2018 retrospective study of 8,441 adult patients who underwent general and vascular surgeries showed that the underlying predictors of cardiac events after surgery included age > 65 years (OR 4.9, 95% CI 3.4–6.9,  $P < 0.01$ ) [30]. In addition, a 2017 study showed one-year mortality in elderly patients undergoing hip fracture repair to be significantly higher in patients with postoperative atrial fibrillation, even in patients receiving anti-arrhythmic therapy [31]. The anesthesia-related cardiac arrest rate is a quality indicator to improve patient safety in the perioperative period. A 2017 systematic review showed that perioperative and anesthesia-related cardiac arrest rates only decreased with a country's high Human Development Index (HDI), with perioperative cardiac arrest being 4-fold higher in geriatric patients in low-HDI countries compared to that in high-HDI countries [32]. Another 2014 study showed that a large majority of cardiac arrests in older patients were caused by not anesthesia-related factors. The major predictors of intraoperative cardiac arrests were poorer physical status as based on assessment guidelines from the American Society of Anesthesiologists (ASA) and the need for emergency surgery. All anesthesia-related cardiac arrests were medication-related or airway-related, highlighting the importance of preventive strategies [33].

## Pulmonary evaluation

Pulmonary function declines with age due to loss of both lung and chest wall compliance and oxygen diffusion capacity, especially in smokers, contributing to decline in oxygen uptake and delivery. Age and functional dependence have been identified as the most reliable risk factors for postoperative pulmonary complications (PPC). A 2006 systematic review showed that patient-related risk factors for PPC included advanced age, ASA class 2 physical status or higher, functional dependence, chronic obstructive pulmonary disease, and CHF. In patients undergoing noncardiac surgery, the rates of PPC are 14% and 15% for ages  $\geq 65$  years and  $\geq 70$  years, respectively [34]. A 2017 prospective study for PPC, which included 1,202 patients showed that significant PPC risk factors, such as emergency (OR 4.47, 95% CI 1.59–12.56), surgical site (OR 2.54, 95% CI 1.67–3.89), and age (OR 1.03, 95% CI, 1.02–1.05) were nonmodifiable [35]. In addition, a 2003 prospective study of 517 patients  $\geq 70$  years old and undergoing noncardiac surgery showed that 31.7 percent of patients died from renal complications (HR 6.07, 95% CI 2.23–16.52,  $P < 0.001$ ), cancer (HR 2.44, 95% CI 1.78–3.38,  $P < 0.001$ ), and pulmonary complications (HR 2.41, 95% CI 1.30–4.48,  $P = 0.005$ ). These studies showed that pulmonary complications were an independent predictor of mortality in elderly patients [36].

## Depression

Senility alone is a high risk factor for depression, and the preoperative psychological burden that patients likely suffer may complicate the situation. ACS-AGS guidelines strongly recommend preoperative depression and substance abuse screening using a simple questionnaire [37]. More than 10% of elderly people have depressive symptoms significant enough to warrant clinical intervention [38]. A 2018 prospective cohort study of 1,035 individuals  $\geq 70$  years old who underwent transcatheter or surgical aortic valve replacement surgery showed that baseline depression (31.5 percent of patients screened positive) was associated with mortality after 1 month (OR 2.20, 95% CI 1.18–4.10) and 12 months (OR 1.532, 95% CI 1.03–2.24). Persistent depression 6 months after the procedure was associated with a 3-fold increase in mortality at 12 months (OR 2.98, 95% CI 1.08–8.20) [39]. Therefore, active multi-disciplinary management of these patients is needed perioperatively.

## Frailty

Frailty is a syndrome of decreased physiologic reserve and resistance to stressors. According to the Cardiovascular Health Study, with 5,317 participants  $\geq 65$  years old, the overall prevalence of frailty in this community was 6.9%, with frailty increasing with age and being greater in women than men [40]. A prospective measure of frailty in 594 patients found that preoperative frailty was associated with an increased risk of postoperative complications (intermediately frail: OR 2.06; 95% CI 1.18–3.60; frail: OR 2.54; 95% CI 1.12–5.77), length of stay (intermediately frail: incidence rate ratio 1.49; 95% CI 1.24–1.80; frail: incidence rate ratio 1.69; 95% CI 1.28–2.23), and discharge to an assisted-living facility after previously living at home (intermediately frail: OR 3.16; 95% CI 1.0–9.99; frail: OR 20.48; 95% CI 5.54–75.68). Frailty independently predicted higher rates of postoperative complications in elderly patients [41]. In addition, a 2019 prospective cohort study of 326 geriatric patients who required emergency general surgery showed that a frail status increased the odds ratio of failure to rescue 3 times when compared with a nonfrail status [42]. A 2014 study including 275 subjects  $\geq 65$  years old showed that the multidimensional frailty score based on CGA is more useful than conventional methods like the ASA classification (area under the receiver operating characteristic curve, 0.821 vs. 0.647,  $P = 0.01$ ) for predicting all-cause mortality rates in geriatric patients undergoing surgery [43]. According to a 2017 meta-analysis, sarcopenia and frailty seem to have significant adverse impacts on postoperative outcomes [44]. Another 2018 meta-analysis

identified potentially modifiable prognostic factors (i.e., frailty, depressive symptoms, and smoking) associated with developing postoperative complications that can be targeted preoperatively to optimize care [45]. These results lend support to the view that intensive management of preoperative modifiable factors can prevent postoperative complications.

There is growing evidence that preoperative frailty in elderly patients is associated with increased adverse outcomes after surgery. Although further studies are needed, frailty evaluation will be a useful preoperative risk-stratification tool in perioperative geriatrics [46]. Geriatric specialists will be able not only to make more extensive assessments but also to implement prior rehabilitation measures. Moreover, anesthesiologists should be aware of their role in patient preparation, maintaining or enhancing patient's functional reserve to facilitate postoperative rehabilitation and discharge back into the society.

## Nutrition

A 2015 meta-analysis showed that perioperative oral nutritional supplementation had a positive effect on serum total protein and led to fewer complications, such as wound, respiratory, and urinary tract infections, but did not have a positive effect on postoperative mortality [47]. Another clinical trial study showed that perioperative taurine supplementation attenuated postoperative oxidative stress in elderly patients with a hip fracture, but did not improve postoperative morbidity and mortality [48]. However, a 2016 review, which included 41 trials with a total of 3,881 participants, showed that oral multinutrient supplements started before or soon after surgery might prevent complications within the first 12 months after a hip fracture although no clear effects on mortality were seen [49].

Oral nutrition and supplementation counteracted the effects of poor appetite and illness. Prolonged preoperative fasting should be avoided (except in cases with an intraabdominal pathology). Yeniy et al. [50] measured preoperative fasting duration with respect to time of the day and its effect on vital parameters and ECG in elderly patients undergoing surgery under spinal anesthesia (SA). The fasting times were far longer than recommended, with 15-hour fasting being related to a transiently increased cardiac stress and mild hypothermia. Malnutrition is a frequent yet often overlooked problem in the surgical field.

## Polypharmacy

Geriatric patients usually takes various medications, especially cardiovascular and central nervous system acting medications.

When compared with younger adults, older people are more likely to have impaired renal function; therefore, it is critical to adjust dosage to prevent adverse effects. The risk of adverse drug reactions increases with the number of drugs taken, leading to more hospital stays [51]. A 2016 analysis of 272 elderly patients with consecutive hip fracture showed that the total number of medications at the time of discharge was predictive of rehospitalization (OR 1.08, 95% CI 1.01–1.17,  $P = 0.030$ ) but not predictive of mortality [52]. Furthermore, a 2018 meta-analysis showed that comorbidity (Charlson Comorbidity Index  $\geq 3$ ) [13], polypharmacy ( $\geq 5$  drugs/day), and ADL dependency were predictive factors for postoperative complications [14]. The ACS/AGS Best Practices Guidelines for the Optimal Preoperative Assessment of the Geriatric Surgical Patients recommended that whenever possible, nonessential medications should be discontinued perioperatively and the addition of new medications should be kept to a minimum [17,53].

For patients at risk of POD, new benzodiazepines [54] and meperidine should be avoided [55,56]. Antihistamine H1 antagonists and strong anticholinergic effective drugs should be cautiously prescribed [54,57]. According to American College of Cardiology/American Heart Association guidelines, beta-blockers are indicated for patients who are already on it or whose known coronary artery disease represents an immediate risk in vascular surgery [58–60]. Beta-blockers should be started days to weeks before elective surgery and titrated to a heart rate of 60 to 80 beats/min in the absence of hypotension. Preoperative statin should be started as soon as possible before surgery for patients who have known vascular disease. For patients undergoing noncardiac surgery and currently taking statins, statins should be continued [60].

Cholinesterase inhibitors (e.g., galantamine, rivastigmine, and donepezil), which are used to slow cognitive decline, was not associated with an increased risk of postoperative respiratory complications among older patients with dementia undergoing hip fracture surgery [61]. Cholinesterase inhibitors may interact with muscle relaxants, thereby prolonging the actions of succinylcholine, reversing the effects of non-depolarizing neuromuscular blocking agents, and leading to larger doses needed to achieve a proper degree of neuromuscular blockade [61,62].

## Anticoagulation therapy

A 2016 population-based cohort study of 154,047 hip fracture patients showed that 33% of them used one or more antithrombotic on admission, with a higher proportion of men and a higher mean age [63]. In most cases, perioperative bridging anticoagulation is no longer recommended when using novel oral anticoagu-

lants and vitamin-K antagonists [64]. However, this recommendation does not apply to patients at high risk of thromboembolism. Although the use of anticoagulants in the elderly is basically similar with that in younger patients, the reduced renal function frequently seen in elderly patients requires consideration.

## Intraoperative Management

### Anesthesia in elderly patients

A 2019 meta-analysis showed that postoperative morbidity and mortality increased with age [65]. However, a 2018 meta-analysis showed that laparoscopic hepatectomy is a feasible and safe alternative to open hepatectomy in elderly patients because of the lower rates of morbidity and favorable postoperative recovery and outcomes [66]. Another 2018 meta-analysis showed that the outcomes of laparoscopic gastrectomy for elderly gastric cancer patients were comparable to those in nonelderly patients. A 2017 study showed that laparoscopic liver resection of colorectal liver metastases in subjects  $\geq 70$  years of age is associated with a significant lower morbidity and a shorter stay [67]. The oncological outcomes were comparable with those in open liver resection even though the benefits of the laparoscopic approach appeared to fade with increasing age. Another recent retrospective study of patients  $\geq 80$  years old who underwent transabdominal preperitoneal repair of groin hernia showed that the incidence of postoperative complication was influenced by poor performance status, lower hemoglobin level, and lower albumin level rather than old age [68]. Therefore, age alone should not preclude laparoscopic gastrectomy in elderly patients [69].

However, whether old age itself is the only criterion for determining the indication of surgery remains controversial. A 2017 meta-analysis that included 18 studies of pancreaticoduodenectomy in patients  $\geq 80$  years old showed a higher 30-day postoperative mortality rate (OR 1.51, 95% CI 1.48–3.31,  $P < 0.001$ ) and length of hospital stay (OR 2.23, 95% CI 1.36–3.10,  $P < 0.001$ ) in this group of patients compared to that in younger patients. The overall postoperative complication rate was high (OR 1.51, 95% CI 1.25–1.83,  $P < 0.001$ ) in aged patients [70]. Another recent prospective study including 165 patients who had pancreaticoduodenectomy showed that the 90-day mortality rate (5.9% in those  $\geq 80$  years old vs. 2% in the younger group,  $P = 0.335$ ) and the postoperative complication rate (64.7% in the elderly vs. 62.8% in the younger group,  $P = 0.83$ ) were similar, although the older patients were far more likely to be discharged to a rehabilitation facility than younger patients were (47.1% vs. 12.8%,  $P < 0.001$ ) [71]. Careful selection of elderly patients and optimal periopera-

tive care, rather than age, should be used to determine whether surgical intervention is indicated. Therefore, the question arises as to whether surgery is associated with a better outcome for the elderly patient when compared with conservative treatment. Ideally, individual parameters should be assessed at an interdisciplinary level, thereby preventing a complication-prone decision-making process based on surgical diagnosis. Once the decision to operate has been made, experienced staffs should be available at all times to anesthetize and operate on the patient as well as to organize appropriate postoperative care.

### Mode of anesthesia

To date, there is insufficient evidence to support a single best anesthetic plan for elderly patients. No difference was found in postoperative morbidity, rates of rehospitalization, in-patient mortality, or hospitalization costs in geriatric patients undergoing regional anesthesia (RA) or GA for hip fracture repair [72,73]. GA and RA are both useful for older non-cardiac patients, but for some procedures, e.g., hip fracture surgery, RA seems the technique of choice. The mode of anesthesia may only play a secondary role in mobility, rehabilitation, and discharge delay. There are no specific recommendations regarding the preferred type of anesthesia for elderly non-cardiac patients [74]. Another 2014 population-based retrospective cohort study that included 6,135 age-matched adult pairs with dementia undergoing hip fracture surgery showed that GA and RA are associated with similar rates of most perioperative adverse events (GA, 11.3%; RA, 10.8%,  $P = 0.44$ ) [75]. The mode of anesthesia also did not have any significant effects on perioperative outcomes (MI, pulmonary complication, stroke, urinary tract infection, and wound infection) after lower extremity amputation in a total of 3,260 geriatric patients [76]. While anesthesia preparation time, start time of surgery, length of surgery, time to sit, and time to walk were shorter in GA, time to fast-track eligibility, phase 1 recovery time, and time to discharge were similar among patients who received SA [77]. Another 2014 retrospective cohort study that included 56,729 patients  $\geq 50$  years old undergoing hip repair surgery found that RA was not associated with a lower 30-day mortality when compared with GA (RA, 5.4%; GA, 5.8%; instrumental variable estimate of risk difference, -1.1%, 95% CI -2.8 to 0.5),  $P = 0.20$ ) but was associated with a modestly shorter length of stay. These findings do not support a mortality benefit for RA in this setting [78].

However, a recent retrospective study that included 16,695 geriatric patients and analyzed mortality within 90 days of undergoing hip fracture repair surgery showed that GA and conversion from RA to GA were associated with a higher risk of mortality

(HR 1.22, 95% CI 1.11–1.35,  $P < 0.001$ ) during hospital stay when compared with RA, but this higher risk did not persist after discharge. Furthermore, GA was associated with a higher risk of all-cause readmission when compared with RA, suggesting that RA may be preferable [79]. Another 2019 retrospective comparative study including 89 patients  $\geq 70$  years old undergoing major lumbar surgery showed that there was an association between epidural anesthesia and superior perioperative outcomes, such as episodes of hypertension and tachycardia during anesthesia and POD, nausea, vomiting, and pain [80]. A 2000 meta-analysis concluded that there were marginal advantages for RA when compared with GA for hip fracture patients in terms of early mortality and risk of deep vein thrombosis [81]. A 2014 review showed that multimodal drug therapy and RA can be effective for perioperative pain management in the elderly, although it did not improve long-term mortality [82].

Several ongoing large-scale randomized controlled trials (RCTs) should provide future guidance on the best anesthesia technique for hip surgery in the elderly [83–85]. Regardless of the type of anesthesia chosen, anesthesia should be carried out by experienced anesthesiologists who are trained to manage the perioperative care of geriatric patients [62,86]. Anesthesiologists should then base the choice of anesthetic on patient preferences, comorbidities, and potential postoperative complications.

A 2018 study of 392 older patients showed that propofol-based GA might decrease the incidence of delayed neurocognitive recovery at one week after surgery when compared with sevoflurane-based GA (propofol 14%; sevoflurane 23.2%; OR 0.577, 95% CI 0.342–0.975,  $P = 0.038$ ) [87]. However, another recent study that included 28 RCTs with 4,507 participants found low-certainty evidence that maintenance with propofol-based total intravenous anesthesia (TIVA) or with inhalational agents may affect the incidence of POD, mortality, or length of hospital stay [88].

In elderly patients undergoing hip fracture repair, continuous spinal anesthesia (CSA) provided fewer episodes of moderate-to-severe hypotension when compared with a single intrathecal injection of 7.5 mg bupivacaine [89]. A prospective cohort study showed that cardiac output and blood pressure decreased significantly after the onset of SA in elderly patients, mainly due to a decrease in stroke volume and not in systemic vascular resistance [90]. A 2014 prospective randomized study of elderly high-risk patients undergoing hip replacement surgery showed that CSA and psoas compartment sciatic nerve block (PCSNB) produced a satisfactory quality of anesthesia. Fewer hemodynamic changes in PCSNB cases when compared with CSA cases were also noted [91]. A 2015 prospective randomized study comparing

midazolam with propofol sedation in hypoalbuminemia (albumin level below 3 g/dl) geriatric patients under SA found that when compared with midazolam, propofol is associated with greater hemodynamic stability, lesser respiratory depression, and faster recovery [92].

## Anesthetic pharmacology

Standard anesthetic doses can cause more profound clinical effects in the elderly, because of differences in pharmacokinetics and pharmacodynamics with the general population. Lower doses are required for propofol, remifentanyl, ropivacaine, and desflurane [93]. Particular care should be taken with hypnotic agents, as the dose required to induce anesthesia is lower but the onset time is prolonged [94]. Depth of anesthesia monitoring is recommended [95,96]. Anesthesiologists should be familiar with potentially inappropriate medications for older patients according to Beers criteria. Thus, for example, elderly people have increased sensitivity to benzodiazepines and decreased metabolism of long-acting agents; in general, all benzodiazepines increase the risk for cognitive impairment, delirium, falls, fractures, and motor vehicle crashes in elderly people [53].

## Neuromuscular blocking and reversal agents

The dose of neuromuscular blocking agents (NMBAs) should hardly be reduced in the elderly for intubation, but their duration of action is often prolonged and difficult to predict along with age-induced changes in pharmacokinetics of long and intermediate-acting NMBAs (especially, aminosteroids including rocuronium and vecuronium), which may cause postoperative residual neuromuscular blockade and associated complications. Therefore, perioperative neuromuscular monitoring including a train-of-four monitoring is strongly recommended [97].

Benzylisoquinoliniums including atracurium and cisatracurium have more reliable durations of action because they depend less on renal and hepatic function for their elimination and can thus be favorably considered for use in the elderly [98]. Neostigmine and pyridostigmine are preferable to edrophonium as NMBA reversal agents because their prolonged duration of action can counterbalance that of NMBAs; however, neostigmine reversal may be ineffective or prolonged and standard doses of sugammadex are required in the elderly [93,98].

## Monitoring

The professional association of anesthesiologists recommends

routinely considering using the following monitoring devices for the elderly, particularly during major or emergency surgery [6].

For intra-arterial blood pressure monitoring, hemoglobin concentration, blood glucose, arterial blood gas testing, and beat-to-beat blood pressure monitoring are recommended. A suitable limit of blood pressure is a fall in systolic blood pressure of more than 20% from pre-anesthesia induction baseline.

For central venous monitoring, catheterization may provide an additional route of venous access after complex surgery when vasoactive drug support is necessary, but its use should be balanced against the possible complication of the procedure. For cardiac output monitoring, there is still limited evidence in the elderly. Because elderly patients have poorly compliant aortas, cardiac output monitoring using Doppler directed at the aorta may provide less accurate information.

For cerebral oxygen saturation, an episode of a decrease in regional cerebral oxygen saturation (rSO<sub>2</sub>) of more than 15% of the baseline value is indicative of cerebral ischemia. Ružman et al. [99] evaluated the changes of rSO<sub>2</sub> measured by near-infrared spectroscopy during elective laparoscopic cholecystectomy under TIVA and the association between patient characteristics and critical decline in rSO<sub>2</sub>. The rSO<sub>2</sub> was significantly lower in patients older than 65 years, suggesting that monitoring of cerebral oxygenation could be an important part of the perioperative care to prevent cerebral hypoxia in older patients. In addition, early evidence suggests that monitoring of cerebral oxygen desaturation and early intervention may reduce POD or POCD [100].

For depth of anesthesia monitoring, processed electroencephalogram (EEG) neuromonitoring including a bispectral index or entropy monitor is recommended to avoid excessive depth of anesthesia, thereby preventing the development of POD [95,96] or POCD [100] in elderly patients. Lastly, perioperative neuromuscular monitoring is strongly recommended to keep the proper degree of neuromuscular blockade and its safe reversal [97,101].

## Fluid management and blood transfusion

Intraoperative fluid optimization may be associated with benefit in geriatric hip fracture patients. A 2015 systematic review showed that goal-directed fluid therapy during hip fracture repair under SA does not result in a significant reduction in length of stay or postoperative complications [102]. Discharge time was similar in the anesthetist-directed fluid therapy group and the pulse-count-guided fluid optimization strategy group, as was total length of stay. A nested meta-analysis of 355 patients found non-significant reduction in early mortality (relative risk [RR] 0.66, 95% CI 0.24–1.79) and in-hospital complications (RR 0.80, 95% CI 0.61–

1.05) when goal-directed intervention fluid therapy was implemented. However, the study provided preliminary evidence that goal-directed fluid therapy may have a mortality reduction benefit. A 2014 systematic review including 734 high-risk patients aged 50 years or older found that the use of a cardiac output-guided hemodynamic therapy algorithm did not reduce a composite outcome of complications or 30-day mortality when compared with the usual standard of care [103].

In high-risk surgical patients, several studies have demonstrated that goal-directed hemodynamic therapy (GDHT) significantly reduced postoperative mortality and morbidity [104,105]. However, several studies found that GDHT was not associated with improved cardiac performance and exerted a statistically uncertain risk reduction in postoperative complications in elderly patients [102,106]. Subsequently, a 2016 RCT in patients aged  $\geq$  70 years old undergoing hip-fracture surgery also indicated that the main GDHT component, the stroke volume maximization by fluid challenges, and traditional pre-anesthesia fluid loading are of questionable value in the elderly [107]. A multicenter RCT involving 807 patients  $\geq$  75 years old and using a stepped wedge cluster design to assess the effectiveness of an optimization strategy involving GDHT, lung-protective ventilation, and depth of anesthesia monitoring for GA on postoperative morbidity and mortality in high-risk elderly patients undergoing high-risk surgeries (the OPTI-AGED study) identified a considerable gap between clinical practice and the relevant guidelines for anesthetic optimization [108]. Implementation of such multimodal optimization strategies varied independently of factors related to the population or type of surgery, and thus, potential benefit should be further addressed in elderly patients [109].

To date, it is still controversial whether red blood cell (RBC) transfusions might increase the risk of infection after hip fracture surgery in geriatric patients. The Transfusion Requirements In Frail Elderly (TRIFE) randomized study with 284 patients showed that a more liberal RBC transfusion strategy (Hb  $<$  11.3 g/dl; 7 mmol/L) was not associated with higher infection risk in subjects undergoing hip fracture surgery compared to the restrictive RBC strategy group (hemoglobin [Hb]  $<$  9.7 g/dl; 6 mmol/L). The rate of infection was 72% in the restrictive RBC strategy group compared with 66% in the liberal group (RR 1.08, 95% CI 0.93–1.27,  $P = 0.29$ ) [110]. A 2015 Cochrane review of 2,722 participants between 81 and 87 years old undergoing hip fracture surgery provided preliminary evidence for similar mortality, functional recovery, or postoperative morbidity when using the thresholds for RBC transfusion in the liberal strategy (aiming to maintain a Hb level usually around 10 g/dl) versus the restrictive strategy (based on symptoms of anemia or a lower Hb concentration usually



around 8 g/dl) [111]. A 2015-nested analysis in the Transfusion Requirements After Cardiac Surgery (TRACS) RCT focusing on 30-day all-cause mortality and severe morbidity showed no difference between liberal and restrictive transfusion strategy inpatients aged less or more than 60 years old. However, there was an increased rate of cardiogenic shock in the elderly patients in the restrictive transfusion strategy group [112]. A 2019 prospective, multicenter, controlled trial randomizing 2,470 elderly ( $\geq 70$  years) patients undergoing non-cardiac surgery is now to evaluate whether a liberal transfusion strategy (defined as the transfusion of a single RBC unit when Hb is  $< 9$  g/dl with a subsequent target range of 9–10.5 g/dl) reduces the major adverse events after non-cardiac surgery compared to a restrictive strategy (transfusion of a single RBC unit when Hb  $< 7.5$  g/dl with a subsequent target range of 7.5–9 g/dl) within 90 days after surgery [113].

Another 2019 RCT showed that salvaged autologous blood leukocyte filtration could improve ventilation, promote oxygenation and gas exchange, and inhibit lung inflammatory and oxidative stress reactions in elderly patients undergoing lumbar spinal surgery [114].

## Lung-protective ventilation

The ACS best practice guidelines for PPC detailed risk factors and strategies to prevent complications. A 2018 expert survey including 362 respondents suggested a care bundle composed of factors before surgery, i.e., supervised exercise programs and inspiratory muscle training; factors during surgery, i.e., low tidal volume ventilation (6–8 ml/kg) with individualized positive end-expiratory pressure (PEEP) of 5–8 cmH<sub>2</sub>O and repeated recruitment maneuvers, use of routine monitoring to avoid hyperoxia, and efforts to limit neuromuscular blockade; and post-operative factors, i.e., deep breathing exercises and elevation of the head of the bed [115].

## Prevention of hypothermia

Elderly patients are vulnerable to perioperative hypothermia, leading to increased morbidity. Especially, during transurethral resection of the prostate (TURP) or bladder tumor (TURB) under SA or GA, or arthroscopic shoulder surgery under GA, it is important to maintain a normal range of body temperature throughout the whole perioperative period. Many studies have therefore been performed to investigate the effects of various active or passive warming devices and methods including a forced-air warming blanket or heated humidifier circuit on perioperative hypothermia or shivering in elderly patients undergoing TURP or TURB under

SA or GA and arthroscopic or open urologic surgeries under GA [116–119]. Jo et al. [116] suggested that a brief period of preoperative forced-air warming did not significantly reduce the incidence of intraoperative hypothermia, but it could significantly reduce its severity in elderly male patients undergoing TURP under SA. Moreover, Zhang et al. [120] reported that use of a forced-air warming system combined with an electric blanket was a more effective method for maintaining body temperature compared to individual devices alone in elderly TURP patients.

Hong et al. [117] reported that warming blanket application for 10 min before induction of anesthesia reduced the incidence of hypothermia as measured one hour after induction compared to one-layer cotton blanket. A prospective observational study showed a heated humidifier was more effective in preventing intraoperative hypothermia in elderly patients undergoing open urologic surgeries than a heat moisture exchanger was [118]. A retrospective study performed in arthroscopic shoulder surgery reported that the incidence of postoperative hypothermia was higher and the associated temperature drop was more prominent in geriatric patients compared to young adult patients and suggested that additional warming methods are needed to prevent perioperative hypothermia in geriatric patients [119]. Altogether, these and other studies demonstrated that various warming strategies could be helpful in keeping the body temperature stable in elderly patients undergoing surgery under GA or RA.

## Postoperative Management

### Postoperative adverse outcomes

Despite the prevalence of preoperative chronic medical conditions, most patients do well postoperatively. Nonetheless, in a prospective cohort study of 544 patients aged 70 and older undergoing non-cardiac surgery, an overall 21 percent of patients developed adverse outcomes and 3.7% died during the in-hospital postoperative period. ASA physical classification, emergency surgery, and intraoperative tachycardia increased the odds of adverse events [121]. Another prospective study of 517 patients  $\geq 70$  years old and undergoing non-cardiac surgery showed that 31.7 percent of patients were deceased at the time of follow-up and a history of cancer (HR 2.44, 95% CI 1.78–3.38,  $P < 0.001$ ), ASA physical status  $> 2$  (HR 2.27, 95% CI 1.61–3.21,  $P < 0.001$ ), neurologic disease (HR 1.59, 95% CI 1.13–2.24,  $P = 0.008$ ), age (HR 1.42 per decade, 95% CI 1.11–1.81,  $P = 0.005$ ), postoperative pulmonary complication (HR 2.41, 95% CI 1.30–4.48,  $P = 0.005$ ), and renal complication (HR 6.07, 95% CI 2.23–16.52,  $P < 0.001$ ) were all significant independent predictors of decreased long-

term survival [122]. Co-morbid conditions, age, and new hospitalization after discharge were important independent predictors of a long-term decrease in quality of life. To improve postoperative long-term quality of life, geriatric surgical patients should be evaluated for their potential pre- and intra-operative risk factors.

## Postoperative transfusion

In a 2016 RCT including 284 frail elderly patients undergoing surgical hip fracture, postoperative transfusion using the liberal hemoglobin target (7 mmol/L, or 11.3 g/dl) improved survival within one year after surgery in the frailest elderly (the nursing home residents) without impairing recovery from physical disabilities and overall quality of life or increasing the risk of infection when compared with the restrictive hemoglobin target (6 mmol/L, or 9.7 g/dl) [123]. According to the Hb thresholds, recovery from physical disabilities in frail elderly hip fracture patients was similar after a restrictive RBC transfusion strategy (Hb < 9.7 g/dl; < 6 mmol/L) and after a liberal strategy (Hb < 11.3 g/dl; < 7 mmol/L). The 90-day mortality rate was higher for the nursing home residents in the restrictive transfusion group (36%) than for those in the liberal group (20%) (HR 2.0, 95% CI 1.1–3.6, P = 0.010). Implementation of a liberal RBC transfusion strategy in nursing home residents has thus the potential to increase survival [124].

## Perioperative analgesia

Elderly patients are often undertreated for pain. Acute pain management in the elderly is challenging, with physiological frailty, medical comorbidities, and cognitive impairment commonly compounding pain assessment and treatment. A 2003 retrospective cohort study including 8,855 subjects aged 16 years and older showed that the risk of respiratory depression after short-term opioids use increased with age, substantially after 60 years of age [125]. A 2014 review suggested that multimodal drug therapy and perioperative regional analgesia can be very effective for perioperative pain management in elderly patients [81]. Paracetamol is safe and considered first-line therapy. Nonsteroidal anti-inflammatory drugs should be used with caution because they can cause gastric and renal damage. Although morphine is effective, cautious administration to elderly patients with poor renal or respiratory function and impaired cognition must be taken into consideration.

RA as part of multimodal perioperative treatment can often reduce postoperative neurological, pulmonary, cardiac, and endocrine complications. RA or analgesia has not been proven to improve long-term morbidity but does benefit immediate postoperative pain control. In addition, multimodal drug therapy utilizes a

variety of nonopioid analgesic medications in order to minimize dosage and prevent adverse effects from opioids while maximizing analgesic benefit [81]. A 2018 review showed that comprehensive pain protocols for elderly hip fracture patients are required, with fascia iliaca blocks as a local anesthesia method of choice [126]. Ultrasound-guided regional anesthesia/analgesia is an important part of anesthesia practice in the elderly population, the growth of which will continue to outpace that of the younger population due to improvements in lifespan worldwide [127]. In a 2016 RCT comparing ultrasound-guided continuous femoral nerve block versus continuous fascia iliaca compartment block in 60 elderly patients undergoing hip replacement surgery, both ultrasound-guided blocks provided effective anesthesia and postoperative analgesia [128]. Moreover, in a 2014 RCT comparing the hemodynamic effects of combined PCSNB with CSA in elderly high-risk patients undergoing hip replacement surgery, PCSNB produced satisfactory-quality anesthesia in elderly high-risk patients with fewer hemodynamic changes compared with CSA [91].

Another RCT showed that after a major abdominal surgery in the elderly patient, patient-controlled analgesia, regardless of the route (epidural or parenteral), was effective. Furthermore, patient-controlled epidural analgesia using local anesthetics and an opioid provided better pain relief and improved mental status and bowel activity when compared with intravenous patient-controlled analgesia [129].

## Postoperative delirium

POD is a common serious postoperative complication, especially in older people, and is associated with increased mortality, morbidity, and health costs. The overall prevalence of delirium in older patients after surgery has been estimated to be 10% [130]. One study in which 144 patients > 50 years were scheduled for an operation requiring a postoperative intensive care unit (ICU) admission showed that 44 percent of patients developed delirium [131]. A recent review of 35 selected articles showed that the incidence of POD was up to 50% [132]. However, a 2015 study including 459 elderly patients found that incident POD was not significantly associated with decreased survival after hip fracture repair and that survival was a function of age at the time of surgery, illness severity based on ASA physical status, and duration of ICU stay after surgery [133].

Among the risk factors for POD, preexisting cognitive impairment and dementia are the strongest predisposing factors [131, 134]. A 2006 study including 333 elderly patients undergoing noncardiac surgery found that 46 percent of patients developed

POD. By multivariate logistic regression, age (OR 2.5, 95% CI 1.5–4.2) and moderate (OR 2.2, 95% CI 1.2–4.0) or severe (OR 3.7, 95% CI 1.5–9.0) preoperative resting pain were important contributing factors [135]. Therefore, adequate opioid-reduced analgesia is of great importance. A 2019 review suggested that a clinical trial on the usefulness of the STOP-BANG questionnaire on obstructive sleep apnea as a preoperative stratification for POD showed no difference between the low-risk group and the intermediate-to-high risk groups in POD incidence, duration of delirium, and length of ICU. However, a higher preoperative risk for obstructive sleep apnea was associated with a 3-fold higher risk for POD and coma [136]. A retrospective study that included 318 elderly patients undergoing total knee arthroplasty also showed that preoperative dementia is the most important risk factor for POD, suggesting that those patients should be thoroughly evaluated and their dementia should be managed preoperatively. Adequate management of intraoperative hypotension and preoperative hemoglobin might also be helpful in reducing the incidence of POD [137].

A 2018 systematic review, which included 104 studies found no evidence of anesthesia types (GA vs. RA) influencing POD [138]. Another 2018 RCT, which included 256 patients  $\geq$  75 years old, found that Xenon anesthesia did not reduce the incidence of POD (Xenon; 9.7%, 95% CI 4.5–14.9, sevoflurane; 13.6%, 95% CI 7.8–19.5,  $P = 0.33$ ) [139]. Meanwhile, a comparison of RA with GA on POD in elderly patients is underway in nine clinical trial centers in China with an expected total enrollment of 1,000 patients [84].

Intraoperative EEG waveform suppression, which often suggests excessive GA, has been associated with POD, prompting research on whether EEG-guided anesthetic administration decreases the incidence of POD. A 2019 systematic review for prevention of POD in elderly patients planned for elective surgery offered preliminary evidence that multicomponent interventions (i.e., comprehensive multidisciplinary care and multimodal interventions), antipsychotics, bispectral index-guidance, and dexmedetomidine treatment can successfully reduce POD incidence in elderly patients undergoing elective non-cardiac surgery [95]. However, recently, a remarkable large-scale RCT of 1,232 adults aged 60 years and older undergoing major surgery showed that EEG-guided anesthetic administration, compared with standard care, did not decrease the incidence of POD. This finding does not support the use of EEG-guided anesthetic administration for decreasing the incidence of POD [140]. Another 2019 RCT with 200 elderly patients undergoing hip fracture repair with SA supplemented with propofol sedation found that heavier intraoperative sedation was not associated with significant differences in mortality or return to pre-fracture ambulation up to one year after surgery [141]. There-

fore, further large-scale and well-designed RCTs are needed to clarify the association between EEG-guided anesthetic administration or depth of anesthesia, and POD incidence.

A 2012 clinical trial that included 171 elderly subjects with hip fracture found that delirium episodes and cognitive decline during hospitalization were common, but inpatient geriatric consultation teams intervention reduced the incidence of POD (control group 53.2%; intervention group 37.2%; OR 1.92, 95% CI 1.04–3.54,  $P = 0.04$ ). However, another study found that geriatric consultation had no effect on the severity or duration of POD episodes [142]. Although the results are not conclusive, a close collaboration with the geriatric team can be useful [143]. A 2017 meta-analysis that included 1,840 elderly patients concluded that comprehensive geriatric care may reduce the incidence of POD (OR 0.71, 95% CI 0.57–0.89,  $P = 0.003$ ) [144].

Two systematic 2019 meta-analyses of RCTs showed that in elderly patients undergoing noncardiac surgery, perioperative administration of dexmedetomidine, compared with placebo, reduced the incidence of POD [145,146]. In contrast, a 2019 double-blinded, multi-center, randomized study that included 164 elderly patients undergoing cardiac surgery reported that dexmedetomidine-based GA resulted in reduced extubation time and postoperative morphine requirements when compared with propofol-based GA, but no significant difference was observed in POD incidence [147].

If possible, drugs that precipitate POD such as opioids, anticholinergics, atropine, sedative hypnotics, and corticosteroids should be avoided in patients at risk, including benzodiazepine [54,148].

## Postoperative cognitive dysfunction

POCD is another frequent neurologic complication occurring in geriatric patients. The type of anesthesia or analgesia and patient inflammatory response may contribute to POCD. A 2014 clinical trial that included 200 elderly patients with mild cognitive impairment showed that there is no difference in the incidence of POCD at 7 days after radical rectal resection under sevoflurane (33.3%) or propofol-based (29.7%) GA, even though sevoflurane had more severe impact on cognitive function than propofol [149]. However, a 2015 clinical trial that included 90 elderly patients scheduled for resection of an esophageal carcinoma showed the incidence of POCD was higher in sevoflurane than propofol anesthesia using the Mini Mental State Examination (MMSE) and the Montreal cognitive assessment (MoCA) scores. Furthermore, elevated plasma concentrations of TNF- $\alpha$ , interleukin (IL)-6, and S-100 $\beta$  protein were found in patients receiving sevoflurane anesthesia throughout the first postoperative week [150]. However, a

2016 RCT that included 80 elderly patients scheduled for a non-cardiac operation found a negative influence by sevoflurane anesthesia on the early (48 hours postoperatively) and late (9 months postoperatively) state compared to propofol anesthesia and no difference in inflammatory markers (IL6, IL10, TNF- $\alpha$ ) between the two anesthesia groups [151]. A 2018 clinical trial that included 120 elderly scheduled for esophageal carcinoma resection showed that POCD incidence was higher in elderly patient receiving sevoflurane anesthesia while dexmedetomidine could alleviate POCD through decreasing TNF- $\alpha$  and IL-6 [152].

A 2018 RCT compared the effect of GA and SA on the occurrence of POCD up to postoperative 30 days in elderly patients undergoing hip fracture surgery and reported that the choice of anesthesia modality did not appear to influence the emergence of PCOD in the elderly patients [153]. Meanwhile, a 2019 RCT with 80 elderly patients undergoing orthopedic surgery concluded that compared with GA, SA can effectively reduce eye opening and language presentation times and also has few negative impacts on the short-term cognitive function and mental status of elderly patients, along with lower incidence of POCD [154].

A 2017 RCT was performed to observe whether combined GA and RA affected perioperative cognitive trajectory compared to only GA in elderly patients with arthroplasty [155]. Postoperative MMSE was significantly higher in the combined anesthesia group as well as significantly improved compared with preoperative MMSE score in both groups. Combined GA and RA protected perioperative cognitive trajectory, providing evidence supporting the combined use of RA and GA in elderly orthopedic patients vulnerable to POCD.

A 2016 meta-analysis including 13 RCTs of GA showed that dexmedetomidine significantly reduced the incidence of POCD (RR 0.59, 95% CI 0.45–2.95) and improved the MMSE score (mean difference [MD] 1.74, 95% CI 0.43–3.05) on the first postoperative day and reduced the incidence of POCD thereafter (MD 2.73, 95% CI 1.33–4.12) [156]. Another 2019 meta-analysis including 26 RCTs found that perioperative dexmedetomidine treatment significantly reduced the incidence of POCD (pooled ORs 0.59, 95% CI 0.45–2.95) and improved MMSE score (standardized mean difference [SMD] 1.74, 95% CI 0.43–3.05) on the first postoperative day and decreased IL-6 (SMD: -1.31, 95% CI: -1.87–-0.75,  $P < 0.001$ ) and TNF- $\alpha$  (SMD: -1.31, 95% CI: -1.87 to -0.75,  $P < 0.001$ ) and TNF- $\alpha$  (SMD: -2.14, 95% CI: -3.14 to -1.14,  $P < 0.001$ ) when compared with saline treatment [157]. A 2016 clinical trial with 134 elderly patients undergoing total knee arthroplasty found that parecoxib sodium decreased POCD incidence and plasma IL-1 $\beta$ , IL-6, and TNF- $\alpha$  levels, suggesting that parecoxib may influence POCD incidence through suppression of

inflammation and pain [158]. Another clinical trial in 2017 that included 152 elderly patients scheduled for shoulder arthroscopy showed that parecoxib sodium pretreatment combined with dexmedetomidine could reduce the incidence of early POCD and yielded higher jugular venous oxygen partial pressure and jugular venous oxygen saturation values at postoperative day one when compared with the control group. This effect may be related to the improvement of postoperative analgesia and cerebral oxygen metabolism [159].

In a prospective randomized double-blinded controlled study, the effect of remifentanyl and fentanyl on POCD and cytokine levels were investigated in elderly patients undergoing major abdominal surgery. The two opioid groups were comparable in terms of POCD incidence; however, IL-6 levels were lower at the seventh day after surgery in the remifentanyl group, suggesting that remifentanyl did not reduce POCD compared to fentanyl [160].

Enhanced inflammation response has been increasingly reported in association with POCD [161]. Glucocorticoid receptor (GR) signal plays a key role in suppression of inflammation. In a 2010 prospective cohort that included 126 elderly patients undergoing hip fracture surgery with GA, plasma cortisol levels and the expression levels of GR and FK506 binding protein 51 (FKBP51) in leukocytes were determined from one day before surgery and to up to seven days after surgery. When compared with non-POCD patients, visual analogue scale (VAS) scores at 12 hours after surgery were higher in POCD patients. No significant difference in expression levels of GR was found between POCD and non-POCD patients, but high expression of FKBP51 in leukocytes and glucocorticoid resistance were associated with POCD in aged patients following hip fracture surgery [162].

A 2017 RCT reported that preoperative oral melatonin supplementation might improve early POCD in elderly patients undergoing hip arthroplasty, suggesting that the restoration of normal circadian function with good sleep quality may be a key factor in preventing or treating POCD [163].

To manage POCD, care bundles and protocols for the perioperative period may improve outcomes in the elderly patient. However, to date, there is no clear strategy to improve POD or POCD. Preventive strategies, early recognition, and management of perioperative risk factors seems to be the best modality to treat POCD until further progress is made [164]. Meanwhile, in Germany, a stepped-wedge cluster randomized (PAWEL) trial for the reduction of delirium and POD risks after elective procedures in adults > 70 years old is now planned with an expected 1,500 patients to enroll. Results of the trial should form the basis of future standards for preventing delirium and POCD in surgical wards [165].

## Prevention of postoperative pulmonary complications

Pulmonary complications increase the risk of mortality after surgery and age is a significant risk predictor of pulmonary complications [34,166]. A large retrospective cohort study that included 8,920 elderly patients with hip fracture repair patients found that cardiac and pulmonary complications were most frequent (8% and 4% of patients, respectively) [167].

Well-documented risk factors for pulmonary complications include atelectasis, pneumonia, and pulmonary thromboembolism, advanced age, poor general health status, current infections, pre-existing cardiopulmonary diseases, hypoalbuminemia, and renal dysfunction. Interventions such as lung expansion maneuvers and thromboprophylaxis are effective in reducing the risk of pulmonary complications [168]. According to a 2006 systematic review that included 20 RCTs and 11 meta-analyses not limited to the elderly patient, lung expansion therapy (i.e., incentive spirometry, deep breathing exercises, and continuous positive airway pressure) reduces postoperative pulmonary risk after abdominal surgery. Well-designed trials are needed to clarify the magnitude of benefit and the comparative effectiveness of different modalities [169].

In a 2019 RCT with 76 elderly patients scheduled for hip joint surgery investigating whether pressure-controlled ventilation-volume guaranteed (PCV-VG) may result in better lung ultrasound score (LUS) by reducing atelectasis in the dependent areas of the lung and minimizing respiratory deterioration after surgery in elderly patients compared with volume-controlled ventilation (VCV), PCV-VG showed better LUS results as well as higher dynamic compliance and lower inspiratory peak pressure compared to VCV [170].

## Prevention of urinary tract infections

In a 2019 retrospective cohort study in which 221 female patients (age  $85.3 \pm 7.0$  years) with a history of hip surgery, urinary retention occurred in 34 out of the 221 cases (15.4%) and was significantly associated with cognitive impairment (OR 4.11, 95% CI 1.53–11.03,  $P = 0.005$ ) and ADL (OR 2.61, 95% CI 1.11–6.18,  $P = 0.029$ ), under adjustment with age and body mass index (BMI). This study demonstrated that cognitive function and ADL were important risk factors for urinary retention, and suggested that the postoperative management of urinary retention is important when considering neurofunctional assistance and nursing care in daily living, especially in elderly female patients undergoing surgery after femoral neck and trochanteric fractures [171]. A 2014 clinical study of risk factors for urinary retention including 72 fe-

male elderly patients undergoing hip surgery showed that the early removal of the urethral catheter (per 1-day indwelling period increase, OR 0.33, 95% CI 0.11–0.96,  $P = 0.04$ ) and preoperative dementia and/or delirium (OR 10.4, 95% CI 1.21–89.2,  $P = 0.03$ ) had significant correlations with postoperative urinary retention. Femoral neck fractures and the surgical procedure used for the hip surgery did not induce damage to the bladder and nerves involved in voiding function, with the voiding function being recovered in all the patients after short-term intermittent catheterization [172]. Older adults are at particular risk for urinary tract infection, and indwelling urinary catheters should not be used as a substitute for adequate nursing care of incontinent patients.

## Conclusion

The main aims of effective perioperative care in elderly patients are to improve the likelihood of them returning to their pre-morbid conditions and maintaining their presence in the community. The perioperative care of elderly patients requires optimization through a multidisciplinary approach that includes risk stratification models. However, the aforementioned approaches are time-consuming and remain a challenge in clinical routine because of limited human resources and lack of funding from healthcare systems. Moreover, the evidence base to inform perioperative care for elderly patients remains poor. More RCTs are needed to clarify the efficacy of GA versus RA for surgery. There are no clear care bundles and protocols, which improve POD or POCD. Preventive strategies and early recognition and management of perioperative risk factors seem to be the best modality until there is further progress in therapeutic interventions. Anesthetic techniques to manage appropriate hemodynamic status during the perioperative period to avoid ischemic complications are required. Anesthesiologists must participate in discussions on the utility of surgery and resuscitation and are strongly encouraged to participate in national surveys and outcomes research. We hope that health care practitioners will use this information to improve their daily practice and that additional research will be undertaken to further improve our future.

## Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

## Author Contributions

Byung-Gun Lim (Conceptualization; Data acquisition and analy-

sis; Investigation; Methodology; Writing—original draft; Writing—review & editing)

Il-Ok Lee (Conceptualization; Data acquisition and analysis; Investigation; Methodology; Supervision; Validation; Writing—original draft; Writing—review & editing)

## ORCID

Byung-Gun Lim, <https://orcid.org/0000-0002-3302-1831>

Il-Ok Lee, <https://orcid.org/0000-0001-8062-1496>

## References

- World report on ageing and health [Internet]. Geneva: World Health Organization; 2005 [cited by 2019 Sep 25]. Available from <https://apps.who.int/iris/handle/10665/186463>.
- Partridge JS, Harari D, Martin FC, Dhesei JK. The impact of pre-operative comprehensive geriatric assessment on postoperative outcomes in older patients undergoing scheduled surgery: a systematic review. *Anaesthesia* 2014; 69 Suppl 1: 8-16.
- Eamer G, Taheri A, Chen SS, Daviduck Q, Chambers T, Shi X, et al. Comprehensive geriatric assessment for older people admitted to a surgical service. *Cochrane Database Syst Rev* 2018; 1: CD012485.
- Wozniak SE, Coleman J, Katlic MR. Optimal preoperative evaluation and perioperative care of the geriatric patient: a surgeon's perspective. *Anesthesiol Clin* 2015; 33: 481-9.
- Knittel JG, Wildes TS. Preoperative assessment of geriatric patients. *Anesthesiol Clin* 2016; 34: 171-83.
- Griffiths R, Beech F, Brown A, Dhesei J, Foo I, Goodall J, et al. Peri-operative care of the elderly 2014: Association of Anaesthetists of Great Britain and Ireland. *Anaesthesia* 2014; 69 Suppl 1: 81-98.
- Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969; 9: 179-86.
- Katz S, Downs TD, Cash HR, Grotz RC. Progress in development of the index of ADL. *Gerontologist* 1970; 10: 20-30.
- Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991; 39: 142-8.
- Panel on Prevention of Falls in Older Persons AGSABGS. Summary of the Updated American Geriatrics Society/British Geriatrics Society clinical practice guideline for prevention of falls in older persons. *J Am Geriatr Soc* 2011; 59: 148-57.
- Schmidt M, Eckardt R, Altmeppen S, Wernecke KD, Spies C. Functional impairment prior to major non-cardiac surgery is associated with mortality within one year in elderly patients with gastrointestinal, gynaecological and urogenital cancer: a prospective observational cohort study. *J Geriatr Oncol* 2018; 9: 53-9.
- Fukuse T, Satoda N, Hijiya K, Fujinaga T. Importance of a comprehensive geriatric assessment in prediction of complications following thoracic surgery in elderly patients. *Chest* 2005; 127: 886-91.
- Robinson TN, Wallace JJ, Wu DS, Wiktor A, Pointer LF, Pfister SM, et al. Accumulated frailty characteristics predict postoperative discharge institutionalization in the geriatric patient. *J Am Coll Surg* 2011; 213: 37-42.
- Xue DD, Cheng Y, Wu M, Zhang Y. Comprehensive geriatric assessment prediction of postoperative complications in gastrointestinal cancer patients: a meta-analysis. *Clin Interv Aging* 2018; 13: 723-36.
- Lawrence VA, Hazuda HP, Cornell JE, Pederson T, Bradshaw PT, Mulrow CD, et al. Functional independence after major abdominal surgery in the elderly. *J Am Coll Surg* 2004; 199: 762-72.
- Chu CL, Chiou HY, Chou WH, Chang PY, Huang YY, Yeh HM. Leading comorbidity associated with 30-day post-anesthetic mortality in geriatric surgical patients in Taiwan: a retrospective study from the health insurance data. *BMC Geriatr* 2017; 17: 245.
- Chow WB, Rosenthal RA, Merkow RP, Ko CY, Esnaola NF; American College of Surgeons National Surgical Quality Improvement Program, et al. Optimal preoperative assessment of the geriatric surgical patient: a best practices guideline from the American College of Surgeons National Surgical Quality Improvement Program and the American Geriatrics Society. *J Am Coll Surg* 2012; 215: 453-66.
- Mohanty S, Rosenthal RA, Russell MM, Neuman MD, Ko CY, Esnaola NF. Optimal perioperative management of the geriatric patient: a best practices guideline from the American College of Surgeons NSQIP and the American Geriatrics Society. *J Am Coll Surg* 2016; 222: 930-47.
- Olotu C, Weimann A, Bahrs C, Schwenk W, Scherer M, Kiefmann R. The perioperative care of older patients. *Dtsch Arztebl Int* 2019; 116: 63-9.
- Lee HB, Ankrom M, Lyketsos CG. Cognitive disorders: delirium, mild cognitive impairment, and dementia. In: *Psychiatry for Primary Care Physicians*. 2nd ed. Edited by Goldman LS, Wise TN, Brody DS: Chicago, American Medical Association Press. 2004, p 197.
- Brookmeyer R, Evans DA, Hebert L, Langa KM, Heeringa SG, Plassman BL, et al. National estimates of the prevalence of Alzheimer's disease in the United States. *Alzheimers Dement* 2011;

- 7: 61-73.
22. Bai J, Zhang P, Liang X, Wu Z, Wang J, Liang Y. Association between dementia and mortality in the elderly patients undergoing hip fracture surgery: a meta-analysis. *J Orthop Surg Res* 2018; 13: 298.
  23. Seitz DP, Gill SS, Gruneir A, Austin PC, Anderson GM, Bell CM, et al. Effects of dementia on postoperative outcomes of older adults with hip fractures: a population-based study. *J Am Med Dir Assoc* 2014; 15: 334-41.
  24. Borson S, Scanlan J, Brush M, Vitaliano P, Dokmak A. The mini-cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry* 2000; 15: 1021-7.
  25. Ansaloni L, Catena F, Chattat R, Fortuna D, Franceschi C, Mascitti P, et al. Risk factors and incidence of postoperative delirium in elderly patients after elective and emergency surgery. *Br J Surg* 2010; 97: 273-80.
  26. Rudolph JL, Jones RN, Levkoff SE, Rockett C, Inouye SK, Sellke FW, et al. Derivation and validation of a preoperative prediction rule for delirium after cardiac surgery. *Circulation* 2009; 119: 229-36.
  27. Fleisher LA, Fleischmann KE, Auerbach AD, Barnason SA, Beckman JA, Bozkurt B, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol* 2014; 64: e77-137.
  28. Rich MW, Chyun DA, Skolnick AH, Alexander KP, Forman DE, Kitzman DW, et al. Knowledge gaps in cardiovascular care of the older adult population: a scientific statement from the American Heart Association, American College of Cardiology, and American Geriatrics Society. *J Am Coll Cardiol* 2016; 67: 2419-40.
  29. Liu LL, Dzankic S, Leung JM. Preoperative electrocardiogram abnormalities do not predict postoperative cardiac complications in geriatric surgical patients. *J Am Geriatr Soc* 2002; 50: 1186-91.
  30. Acheampong D, Guerrier S, Lavarias V, Pechman D, Mills C, Inabnet W, et al. Risk factors contributing to cardiac events following general and vascular surgery. *Ann Med Surg (Lond)* 2018; 33: 16-23.
  31. Leibowitz D, Abitbol C, Alcalai R, Rivkin G, Kandel L. Perioperative atrial fibrillation is associated with increased one-year mortality in elderly patients after repair of hip fracture. *Int J Cardiol* 2017; 227: 58-60.
  32. Braghiroli KS, Braz JRC, Rocha B, El Dib R, Corrente JE, Braz MG, et al. Perioperative and anesthesia-related cardiac arrests in geriatric patients: a systematic review using meta-regression analysis. *Sci Rep* 2017; 7: 2622.
  33. Nunes JC, Braz JR, Oliveira TS, de Carvalho LR, Castiglia YM, Braz LG. Intraoperative and anesthesia-related cardiac arrest and its mortality in older patients: a 15-year survey in a tertiary teaching hospital. *PLoS One* 2014; 9: e104041.
  34. Smetana GW, Lawrence VA, Cornell JE; American College of Physicians. Preoperative pulmonary risk stratification for non-cardiothoracic surgery: systematic review for the American College of Physicians. *Ann Intern Med* 2006; 144: 581-95.
  35. Fernandez-Bustamante A, Frenzl G, Sprung J, Kor DJ, Subramaniam B, Martinez Ruiz R, et al. Postoperative pulmonary complications, early mortality, and hospital stay following noncardiothoracic surgery: a multicenter study by the Perioperative Research Network Investigators. *JAMA Surg* 2017; 152: 157-66.
  36. Manku K, Bacchetti P, Leung JM. Prognostic significance of postoperative in-hospital complications in elderly patients. I. Long-term survival. *Anesth Analg* 2003; 96: 583-9.
  37. Li C, Friedman B, Conwell Y, Fiscella K. Validity of the Patient Health Questionnaire 2 (PHQ-2) in identifying major depression in older people. *J Am Geriatr Soc* 2007; 55: 596-602.
  38. Steffens DC, Fisher GG, Langa KM, Potter GG, Plassman BL. Prevalence of depression among older Americans: the Aging, Demographics and Memory Study. *Int Psychogeriatr* 2009; 21: 879-88.
  39. Drudi LM, Ades M, Turkdogan S, Huynh C, Lauck S, Webb JG, et al. Association of depression with mortality in older adults undergoing transcatheter or surgical aortic valve replacement. *JAMA Cardiol* 2018; 3: 191-7.
  40. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001; 56: M146-56.
  41. Makary MA, Segev DL, Pronovost PJ, Syin D, Bandeen-Roche K, Patel P, et al. Frailty as a predictor of surgical outcomes in older patients. *J Am Coll Surg* 2010; 210: 901-8.
  42. Khan M, Jehan F, Zeeshan M, Kulvatunyou N, Fain MJ, Saljuqi AT, et al. Failure to rescue after emergency general surgery in geriatric patients: does frailty matter? *J Surg Res* 2019; 233: 397-402.
  43. Kim SW, Han HS, Jung HW, Kim KI, Hwang DW, Kang SB, et al. Multidimensional frailty score for the prediction of postoperative mortality risk. *JAMA Surg* 2014; 149: 633-40.
  44. Shen Y, Hao Q, Zhou J, Dong B. The impact of frailty and sarcopenia on postoperative outcomes in older patients undergoing gastrectomy surgery: a systematic review and meta-analysis. *BMC Geriatr* 2017; 17: 188.
  45. Watt J, Tricco AC, Talbot-Hamon C, Pham B, Rios P, Grudniewicz A, et al. Identifying older adults at risk of harm following elective surgery: a systematic review and meta-analysis. *BMC*

- Med 2018; 16: 2.
46. Ko FC. Preoperative frailty evaluation: a promising risk-stratification tool in older adults undergoing general surgery. *Clin Ther* 2019; 41: 387-99.
  47. Liu M, Yang J, Yu X, Huang X, Vaidya S, Huang F, et al. The role of perioperative oral nutritional supplementation in elderly patients after hip surgery. *Clin Interv Aging* 2015; 10: 849-58.
  48. Van Stijn MF, Bruins AA, Vermeulen MA, Witlox J, Teerlink T, School MG, et al. Effect of oral taurine on morbidity and mortality in elderly hip fracture patients: a randomized trial. *Int J Mol Sci* 2015; 16: 12288-306.
  49. Avenell A, Smith TO, Curtain JP, Mak JC, Myint PK. Nutritional supplementation for hip fracture aftercare in older people. *Cochrane Database Syst Rev* 2016; 11: CD001880.
  50. Yeniay O, Tekgul ZT, Okur O, Koroglu N. Unexpectedly prolonged fasting and its consequences on elderly patients undergoing spinal anesthetics. A prospective observational study. *Acta Cir Bras* 2019; 34: e201900309.
  51. Barnett SR. Polypharmacy and perioperative medications in the elderly. *Anesthesiol Clin* 2009; 27: 377-89.
  52. Harstedt M, Rogmark C, Sutton R, Melander O, Fedorowski A. Polypharmacy and adverse outcomes after hip fracture surgery. *J Orthop Surg Res* 2016; 11: 151.
  53. The 2019 American Geriatrics Society Beers Criteria® Update Expert Panel. American Geriatrics Society 2019 Updated AGS Beers Criteria(R) for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc* 2019; 67: 674-94.
  54. Clegg A, Young JB. Which medications to avoid in people at risk of delirium: a systematic review. *Age Ageing* 2011; 40: 23-9.
  55. Fong HK, Sands LP, Leung JM. The role of postoperative analgesia in delirium and cognitive decline in elderly patients: a systematic review. *Anesth Analg* 2006; 102: 1255-66.
  56. Swart LM, van der Zanden V, Spies PE, de Rooij SE, van Munster BC. The comparative risk of delirium with different opioids: a systematic review. *Drugs Aging* 2017; 34: 437-43.
  57. Flinn DR, Diehl KM, Seyfried LS, Malani PN. Prevention, diagnosis, and management of postoperative delirium in older adults. *J Am Coll Surg* 2009; 209: 261-8.
  58. Woolger JM. Preoperative testing and medication management. *Clin Geriatr Med* 2008; 24: 573-83.
  59. Fleischmann KE, Beckman JA, Buller CE, Calkins H, Fleisher LA, Freeman WK, et al. 2009 ACCF/AHA focused update on perioperative beta blockade: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines. *Circulation* 2009; 120: 2123-51.
  60. Biccard BM, Sear JW, Foëx P. Statin therapy: a potentially useful peri-operative intervention in patients with cardiovascular disease. *Anaesthesia* 2005; 60: 1106-14.
  61. Seitz DP, Gill SS, Gruneir A, Austin PC, Anderson G, Reimer CL, et al. Effects of cholinesterase inhibitors on postoperative outcomes of older adults with dementia undergoing hip fracture surgery. *Am J Geriatr Psychiatry* 2011; 19: 803-13.
  62. White SM, Foss NB, Griffiths R. Anaesthetic aspects in the treatment of fragility fracture patients. *Injury* 2018; 49: 1403-8.
  63. Madsen CM, Jantzen C, Lauritzen JB, Abrahamsen B, Jorgensen HL. Temporal trends in the use of antithrombotics at admission. *Acta Orthop* 2016; 87: 368-73.
  64. Douketis JD, Spyropoulos AC, Kaatz S, Becker RC, Caprini JA, Dunn AS, et al. Perioperative bridging anticoagulation in patients with atrial fibrillation. *N Engl J Med* 2015; 373: 823-33.
  65. van Tuil T, Dhaif AA, Te Riele WW, van Ramshorst B, van Santvoort HC. Systematic review and meta-analysis of liver resection for colorectal metastases in elderly patients. *Dig Surg* 2019; 36: 111-23.
  66. Chen K, Pan Y, Maher H, Zhang B, Zheng XY. Laparoscopic hepatectomy for elderly patients: Major findings based on a systematic review and meta-analysis. *Medicine (Baltimore)* 2018; 97: e11703.
  67. Martinez-Cecilia D, Cipriani F, Vishal S, Ratti F, Tranchart H, Barkhatov L, et al. Laparoscopic versus open liver resection for colorectal metastases in elderly and octogenarian patients: a multicenter propensity score based analysis of short-and long-term outcomes. *Ann Surg* 2017; 265: 1192-200.
  68. Egawa N, Nakamura J, Manabe T, Iwasaki H, Noshiro H. Incidence of postoperative complications in transabdominal preperitoneal repair for groin hernia is influenced by poor performance status rather than by old age. *Ann Gastroenterol Surg* 2019; 3: 318-24.
  69. Pan Y, Chen K, Yu WH, Maher H, Wang SH, Zhao HF, et al. Laparoscopic gastrectomy for elderly patients with gastric cancer: A systematic review with meta-analysis. *Medicine (Baltimore)* 2018; 97: e0007.
  70. Kim SY, Weinberg L, Christophi C, Nikfarjam M. The outcomes of pancreaticoduodenectomy in patients aged 80 or older: a systematic review and meta-analysis. *HPB (Oxford)* 2017; 19: 475-82.
  71. Kim SY, Fink MA, Perini M, Houli N, Weinberg L, Muralidharan V, et al. Age 80 years and over is not associated with increased morbidity and mortality following pancreaticoduodenectomy. *ANZ J Surg* 2018; 88: e445-50.
  72. Luger TJ, Kammerlander C, Gosch M, Luger MF, Kammerlander-Knauer U, Roth T, et al. Neuroaxial versus general anaesthesia in geriatric patients for hip fracture surgery: does it matter? *Osteoporos Int* 2010; 21: S555-72.



73. Le-Wendling L, Bihorac A, Baslanti TO, Lucas S, Sadasivan K, Wendling A, et al. Regional anesthesia as compared with general anesthesia for surgery in geriatric patients with hip fracture: does it decrease morbidity, mortality, and health care costs? Results of a single-centered study. *Pain Med* 2012; 13: 948-56.
74. Luger TJ, Kammerlander C, Luger MF, Kammerlander-Knauer U, Gosch M. Mode of anesthesia, mortality and outcome in geriatric patients. *Z Gerontol Geriatr* 2014; 47: 110-24.
75. Seitz DP, Gill SS, Bell CM, Austin PC, Gruneir A, Anderson GM, et al. Postoperative medical complications associated with anesthesia in older adults with dementia. *J Am Geriatr Soc* 2014; 62: 2102-9.
76. Moreira CC, Farber A, Kalish JA, Eslami MH, Didato S, Rybin D, et al. The effect of anesthesia type on major lower extremity amputation in functionally impaired elderly patients. *J Vasc Surg* 2016; 63: 696-701.
77. Ornek D, Metin S, Deren S, Un C, Metin M, Dikmen B, et al. The influence of various anesthesia techniques on postoperative recovery and discharge criteria among geriatric patients. *Clinics (Sao Paulo)* 2010; 65: 941-6.
78. Neuman MD, Rosenbaum PR, Ludwig JM, Zubizarreta JR, Silber JH. Anesthesia technique, mortality, and length of stay after hip fracture surgery. *JAMA* 2014; 311: 2508-17.
79. Desai V, Chan PH, Prentice HA, Zohman GL, Diekmann GR, Maletis GB, et al. Is anesthesia technique associated with a higher risk of mortality or complications within 90 days of surgery for geriatric patients with hip fractures? *Clin Orthop Relat Res* 2018; 476: 1178-88.
80. Li Z, Long H, Huang F, Zhang Y, Xu J, Wang X. Impact of epidural versus general anesthesia on major lumbar surgery in elderly patients. *Clin Spine Surg* 2019; 32: E7-12.
81. Urwin SC, Parker MJ, Griffiths R. General versus regional anaesthesia for hip fracture surgery: a meta-analysis of randomized trials. *Br J Anaesth* 2000; 84: 450-5.
82. Nordquist D, Halaszynski TM. Perioperative multimodal anesthesia using regional techniques in the aging surgical patient. *Pain Res Treat* 2014; 2014: 902174.
83. Neuman MD, Ellenberg SS, Sieber FE, Magaziner JS, Feng R, Carson JL, et al. Regional versus General Anesthesia for Promoting Independence after Hip Fracture (REGAIN): protocol for a pragmatic, international multicentre trial. *BMJ Open* 2016; 6: e013473.
84. Li T, Yeung J, Li J, Zhang Y, Melody T, Gao Y, et al. Comparison of regional with general anaesthesia on postoperative delirium (RAGA-delirium) in the older patients undergoing hip fracture surgery: study protocol for a multicentre randomised controlled trial. *BMJ Open* 2017; 7: e016937.
85. Kowark A, Adam C, Ahrens J, Bajbouj M, Bollheimer C, Borowski M, et al. Improve hip fracture outcome in the elderly patient (iHOPE): a study protocol for a pragmatic, multicentre randomised controlled trial to test the efficacy of spinal versus general anaesthesia. *BMJ Open* 2018; 8: e023609.
86. White SM, Altermatt F, Barry J, Ben-David B, Coburn M, Coluzzi F, et al. International Fragility Fracture Network Delphi consensus statement on the principles of anaesthesia for patients with hip fracture. *Anaesthesia* 2018; 73: 863-74.
87. Zhang Y, Shan GJ, Zhang YX, Cao SJ, Zhu SN, Li HJ, et al. Propofol compared with sevoflurane general anaesthesia is associated with decreased delayed neurocognitive recovery in older adults. *Br J Anaesth* 2018; 121: 595-604.
88. Miller D, Lewis SR, Pritchard MW, Schofield-Robinson OJ, Shelton CL, Alderson P, et al. Intravenous versus inhalational maintenance of anaesthesia for postoperative cognitive outcomes in elderly people undergoing non-cardiac surgery. *Cochrane Database Syst Rev* 2018; 8: CD012317.
89. Minville V, Fourcade O, Grousset D, Chassery C, Nguyen L, Asehnoune K, et al. Spinal anesthesia using single injection small-dose bupivacaine versus continuous catheter injection techniques for surgical repair of hip fracture in elderly patients. *Anesth Analg* 2006; 102: 1559-63.
90. Hofhuizen C, Lemson J, Snoeck M, Scheffer GJ. Spinal anesthesia-induced hypotension is caused by a decrease in stroke volume in elderly patients. *Local Reg Anesth* 2019; 12: 19-26.
91. Aksoy M, Dostbil A, Ince I, Ahiskalioglu A, Alici HA, Aydin A, et al. Continuous spinal anaesthesia versus ultrasound-guided combined psoas compartment-sciatic nerve block for hip replacement surgery in elderly high-risk patients: a prospective randomised study. *BMC Anesthesiol* 2014; 14: 99.
92. Ersoy A, Kara D, Ervatan Z, Çakırgöz M, Kiran Ö. Sedation in hypoalbuminemic geriatric patients under spinal anesthesia in hip surgery. Midazolam or Propofol? *Saudi Med J* 2015; 36: 1191-8.
93. Kruijt Spanjer MR, Bakker NA, Absalom AR. Pharmacology in the elderly and newer anaesthesia drugs. *Best Pract Res Clin Anaesthesiol* 2011; 25: 355-65.
94. Steinmetz J, Rasmussen LS. The elderly and general anesthesia. *Minerva Anesthesiol* 2010; 76: 745-52.
95. Janssen TL, Alberts AR, Hooft L, Mattace-Raso F, Mosk CA, van der Laan L. Prevention of postoperative delirium in elderly patients planned for elective surgery: systematic review and meta-analysis. *Clin Interv Aging* 2019; 14: 1095-117.
96. Koch S, Spies C. Neuromonitoring in the elderly. *Curr Opin Anaesthesiol* 2019; 32: 101-7.
97. Strøm C, Rasmussen LS, Steinmetz J. Practical management of

- anaesthesia in the elderly. *Drugs Aging* 2016; 33: 765-77.
98. Lee LA, Athanassoglou V, Pandit JJ. Neuromuscular blockade in the elderly patient. *J Pain Res* 2016; 9: 437-44.
  99. Ružman T, Mraović B, Šimurina T, Gulam D, Ružman N, Miškulin M. Transcranial cerebral oxymetric monitoring reduces brain hypoxia in obese and elderly patients undergoing general anaesthesia for laparoscopic cholecystectomy. *Surg Laparosc Endosc Percutan Tech* 2017; 27: 248-52.
  100. Ballard C, Jones E, Gauge N, Aarsland D, Nilsen OB, Saxby BK, et al. Optimised anaesthesia to reduce post operative cognitive decline (POCD) in older patients undergoing elective surgery, a randomised controlled trial. *PLoS One* 2012; 7: e37410.
  101. Pietraszewski P, Gaszynski T. Residual neuromuscular block in elderly patients after surgical procedures under general anaesthesia with rocuronium. *Anaesthesiol Intensive Ther* 2013; 45: 77-81.
  102. Moppett IK, Rowlands M, Mannings A, Moran CG, Wiles MD, NOTTS Investigators. LiDCO-based fluid management in patients undergoing hip fracture surgery under spinal anaesthesia: A randomized trial and systematic review. *Br J Anaesth* 2015; 114: 444-59.
  103. Pearse RM, Harrison DA, MacDonald N, Gillies MA, Blunt M, Ackland G, et al. Effect of a perioperative, cardiac output-guided hemodynamic therapy algorithm on outcomes following major gastrointestinal surgery: a randomized clinical trial and systematic review. *JAMA* 2014; 311: 2181-90.
  104. Gurgel ST, do Nascimento P. Maintaining tissue perfusion in high-risk surgical patients: a systematic review of randomized clinical trials. *Anesth Analg* 2011; 112: 1384-91.
  105. Hamilton MA, Cecconi M, Rhodes A. A systematic review and meta-analysis on the use of preemptive hemodynamic intervention to improve postoperative outcomes in moderate and high-risk surgical patients. *Anesth Analg* 2011; 112: 1392-402.
  106. Bartha E, Arfwedson C, Imnell A, Fernlund ME, Andersson LE, Kalman S. Randomized controlled trial of goal-directed haemodynamic treatment in patients with proximal femoral fracture. *Br J Anaesth* 2013; 110: 545-53.
  107. Bartha E, Arfwedson C, Imnell A, Kalman S. Towards individualized perioperative, goal-directed haemodynamic algorithms for patients of advanced age: observations during a randomized controlled trial (NCT01141894). *Br J Anaesth* 2016; 116: 486-92.
  108. Molliex S, Passot S, Futier E, Bonnefoi M, Rancon F, Lemanach Y, et al. Stepped wedge cluster randomised controlled trial to assess the effectiveness of an optimisation strategy for general anaesthesia on postoperative morbidity and mortality in elderly patients (the OPTI-AGED study): A study protocol. *BMJ Open* 2018; 8: e021053.
  109. Molliex S, Passot S, Morel J, Futier E, Lefrant JY, Constantin JM, et al. A multicentre observational study on management of general anaesthesia in elderly patients at high-risk of postoperative adverse outcomes. *Anaesth Crit Care Pain Med* 2019; 38: 15-23.
  110. Gregersen M, Damsgaard EM, Borris LC. Blood transfusion and risk of infection in frail elderly after hip fracture surgery: the TRIFE randomized controlled trial. *Eur J Orthop Surg Traumatol* 2015; 25: 1031-8.
  111. Brunskill SJ, Millette SL, Shokoohi A, Pulford EC, Doree C, Murphy ME, et al. Red blood cell transfusion for people undergoing hip fracture surgery. *Cochrane Database Syst Rev* 2015; 4: CD009699.
  112. Nakamura RE, Vincent JL, Fukushima JT, de Almeida JP, Franco RA, Lee Park C, et al. A liberal strategy of red blood cell transfusion reduces cardiogenic shock in elderly patients undergoing cardiac surgery. *J Thorac Cardiovasc Surg* 2015; 150: 1314-20.
  113. Meybohm P, Lindau S, Treskatsch S, Francis R, Spies C, Velten M, et al. Liberal transfusion strategy to prevent mortality and anaemia-associated, ischaemic events in elderly non-cardiac surgical patients - the study design of the LIBERAL-Trial. *Trials* 2019; 20: 101.
  114. Xu L, Shen J, Sun J, McQuillan PM, Hu Z. The effects of leukocyte filtration on cell salvaged autologous blood transfusion on lung function and lung inflammatory and oxidative stress reactions in elderly patients undergoing lumbar spinal surgery. *J Neurosurg Anesthesiol* 2019; 31: 36-42.
  115. Griffiths SV, Conway DH, POPC-CB Investigators, Sander M, Jammer I, Grocott MPW, et al. What are the optimum components in a care bundle aimed at reducing post-operative pulmonary complications in high-risk patients? *Perioper Med (Lond)* 2018; 7: 7.
  116. Jo YY, Chang YJ, Kim YB, Lee S, Kwak HJ. Effect of preoperative forced-air warming on hypothermia in elderly patients undergoing transurethral resection of the prostate. *Urol J* 2015; 12: 2366-70.
  117. Hong S, Yoo BH, Kim KM, Kim MC, Yon JH, Lee S. The efficacy of warming blanket on reducing intraoperative hypothermia in patients undergoing transurethral resection of bladder tumor under general anesthesia. *Anesth Pain Med* 2016; 11: 404-9.
  118. Seo H, Kim K, Oh EA, Moon YJ, Kim YK, Hwang JH. Effect of electrically heated humidifier on intraoperative core body temperature decrease in elderly patients: a prospective observational study. *Anesth Pain Med* 2016; 11: 211-6.
  119. Chun EH, Lee GY, Kim CH. Postoperative hypothermia in geriatric patients undergoing arthroscopic shoulder surgery.

- Anesth Pain Med 2019; 14: 112-6.
120. Zhang R, Chen X, Xiao Y. The effects of a forced-air warming system plus electric blanket for elderly patients undergoing transurethral resection of the prostate: a randomized controlled trial. *Medicine (Baltimore)* 2018; 97: e13119.
  121. Leung JM, Dzankic S. Relative importance of preoperative health status versus intraoperative factors in predicting postoperative adverse outcomes in geriatric surgical patients. *J Am Geriatr Soc* 2001; 49: 1080-5.
  122. Manku K, Leung JM. Prognostic significance of postoperative in-hospital complications in elderly patients. II. Long-term quality of life. *Anesth Analg* 2003; 96: 590-4.
  123. Gregersen M. Postoperative red blood cell transfusion strategy in frail anemic elderly with hip fracture. a randomized controlled trial. *Dan Med J* 2016; 63: pii: B5221.
  124. Gregersen M, Borris LC, Damsgaard EM. Postoperative blood transfusion strategy in frail, anemic elderly patients with hip fracture: the TRIFE randomized controlled trial. *Acta Orthop* 2015; 86: 363-72.
  125. Cepeda MS, Farrar JT, Baumgarten M, Boston R, Carr DB, Strom BL. Side effects of opioids during short-term administration: effect of age, gender, and race. *Clin Pharmacol Ther* 2003; 74: 102-12.
  126. Scurrah A, Shiner CT, Stevens JA, Faux SG. Regional nerve blockade for early analgesic management of elderly patients with hip fracture - a narrative review. *Anaesthesia* 2018; 73: 769-83.
  127. Lin C, Darling C, Tsui BCH. Practical regional anesthesia guide for elderly patients. *Drugs Aging* 2019; 36: 213-34.
  128. Yu B, He M, Cai GY, Zou TX, Zhang N. Ultrasound-guided continuous femoral nerve block vs continuous fascia iliaca compartment block for hip replacement in the elderly: a randomized controlled clinical trial (CONSORT). *Medicine (Baltimore)* 2016; 95: e5056.
  129. Mann C, Pouzeratte Y, Boccara G, Peccoux C, Vergne C, Brunat G, et al. Comparison of intravenous or epidural patient-controlled analgesia in the elderly after major abdominal surgery. *Anesthesiology* 2000; 92: 433-41.
  130. Rasmussen LS, Moller JT. Central nervous system dysfunction after anesthesia in the geriatric patient. *Anesthesiol Clin North Am* 2000; 18: 59-70.
  131. Robinson TN, Raeburn CD, Tran ZV, Angles EM, Brenner LA, Moss M. Postoperative delirium in the elderly: risk factors and outcomes. *Ann Surg* 2009; 249: 173-8.
  132. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet* 2014; 383: 911-22.
  133. Gottschalk A, Hubbs J, Vikani AR, Gottschalk LB, Sieber FE. The impact of incident postoperative delirium on survival of elderly patients after surgery for hip fracture repair. *Anesth Analg* 2015; 121: 1336-43.
  134. Demeure MJ, Fain MJ. The elderly surgical patient and postoperative delirium. *J Am Coll Surg* 2006; 203: 752-7.
  135. Vaurio LE, Sands LP, Wang Y, Mullen EA, Leung JM. Postoperative delirium: the importance of pain and pain management. *Anesth Analg* 2006; 102: 1267-73.
  136. Bilotta F, Giordano G, Pugliese F. Preoperative stratification for postoperative delirium: obstructive sleep apnea is a predictor, the STOP-BANG is not? *J Thorac Dis* 2019; 11: S202-6.
  137. Korean Statistical Information S, Kim SW, Kim N, Kim J, Kim KM, Lee S. Risk factors for postoperative delirium following total knee arthroplasty in elderly patients. *Anesth Pain Med* 2018; 13: 143-8.
  138. Patel V, Champaneria R, Dretzke J, Yeung J. Effect of regional versus general anaesthesia on postoperative delirium in elderly patients undergoing surgery for hip fracture: a systematic review. *BMJ Open* 2018; 8: e020757.
  139. Coburn M, Sanders RD, Maze M, Nguyen-Pascal ML, Rex S, Garrigues B, et al. The hip fracture surgery in elderly patients (HIPELD) study to evaluate xenon anaesthesia for the prevention of postoperative delirium: a multicentre, randomized clinical trial. *Br J Anaesth* 2018; 120: 127-37.
  140. Wildes TS, Mickle AM, Ben Abdallah A, Maybrier HR, Oberhaus J, Budelier TP, et al. Effect of electroencephalography-guided anesthetic administration on postoperative delirium among older adults undergoing major surgery: the EN-GAGES randomized clinical trial. *JAMA* 2019; 321: 473-83.
  141. Sieber F, Neufeld KJ, Gottschalk A, Bigelow GE, Oh ES, Rosenberg PB, et al. Depth of sedation as an interventional target to reduce postoperative delirium: mortality and functional outcomes of the Strategy to Reduce the Incidence of Postoperative Delirium in Elderly Patients randomised clinical trial. *Br J Anaesth* 2019; 122: 480-9.
  142. Deschodt M, Braes T, Flamaing J, Detroyer E, Broos P, Haentjens P, et al. Preventing delirium in older adults with recent hip fracture through multidisciplinary geriatric consultation. *J Am Geriatr Soc* 2012; 60: 733-9.
  143. Martocchia A, Curto M, Comite F, Scaccianoce S, Girardi P, Ferracuti S, et al. The prevention and treatment of delirium in elderly patients following hip fracture surgery. *Recent Pat CNS Drug Discov* 2015; 10: 55-64.
  144. Wang Y, Tang J, Zhou F, Yang L, Wu J. Comprehensive geriatric care reduces acute perioperative delirium in elderly patients with hip fractures: a meta-analysis. *Medicine (Baltimore)* 2017; 96: e7361.

145. Pan H, Liu C, Ma X, Xu Y, Zhang M, Wang Y. Perioperative dexmedetomidine reduces delirium in elderly patients after non-cardiac surgery: a systematic review and meta-analysis of randomized-controlled trials. *Can J Anaesth* 2019; 66: 1489-500.
146. Zeng H, Li Z, He J, Fu W. Dexmedetomidine for the prevention of postoperative delirium in elderly patients undergoing non-cardiac surgery: a meta-analysis of randomized controlled trials. *PLoS One* 2019; 14: e0218088.
147. Shi C, Jin J, Qiao L, Li T, Ma J, Ma Z. Effect of perioperative administration of dexmedetomidine on delirium after cardiac surgery in elderly patients: a double-blinded, multi-center, randomized study. *Clin Interv Aging* 2019; 14: 571-5.
148. American Geriatrics Society 2012 Beers Criteria Update Expert Panel. American Geriatrics Society updated Beers Criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc* 2012; 60: 616-31.
149. Tang N, Ou C, Liu Y, Zuo Y, Bai Y. Effect of inhalational anaesthetic on postoperative cognitive dysfunction following radical rectal resection in elderly patients with mild cognitive impairment. *J Int Med Res* 2014; 42: 1252-61.
150. Qiao Y, Feng H, Zhao T, Yan H, Zhang H, Zhao X. Postoperative cognitive dysfunction after inhalational anesthesia in elderly patients undergoing major surgery: the influence of anesthetic technique, cerebral injury and systemic inflammation. *BMC Anesthesiol* 2015; 15: 154.
151. Micha G, Tzimas P, Zalonis I, Kotsis K, Papdopoulos G, Arnaoutoglou E. Propofol vs sevoflurane anaesthesia on postoperative cognitive dysfunction in the elderly. A randomized controlled trial. *Acta Anaesthesiol Belg* 2016; 67: 129-37.
152. Zhang H, Wu Z, Zhao X, Qiao Y. Role of dexmedetomidine in reducing the incidence of postoperative cognitive dysfunction caused by sevoflurane inhalation anesthesia in elderly patients with esophageal carcinoma. *J Cancer Res Ther* 2018; 14: 1497-502.
153. Tzimas P, Samara E, Petrou A, Korompilias A, Chalkias A, Papadopoulos G. The influence of anesthetic techniques on postoperative cognitive function in elderly patients undergoing hip fracture surgery: General vs spinal anesthesia. *Injury* 2018; 49: 2221-6.
154. Zhang X, Dong Q, Fang J. Impacts of general and spinal anaesthesia on short-term cognitive function and mental status in elderly patients undergoing orthopaedic surgery. *J Coll Physicians Surg Pak* 2019; 29: 101-4.
155. Chen C, Li M, Wang K, Shen J, Yang L, Bu X, et al. Protective effect of combined general and regional anesthesia on postoperative cognitive function in older arthroplasty patients. *Int J Clin Exp Med* 2017; 10: 15453-8.
156. Zhou C, Zhu Y, Liu Z, Ruan L. Effect of dexmedetomidine on postoperative cognitive dysfunction in elderly patients after general anaesthesia: a meta-analysis. *J Int Med Res* 2016; 44: 1182-90.
157. Yang W, Kong LS, Zhu XX, Wang RX, Liu Y, Chen LR. Effect of dexmedetomidine on postoperative cognitive dysfunction and inflammation in patients after general anaesthesia: a PRISMA-compliant systematic review and meta-analysis. *Medicine (Baltimore)* 2019; 98: e15383.
158. Zhu YZ, Yao R, Zhang Z, Xu H, Wang LW. Parecoxib prevents early postoperative cognitive dysfunction in elderly patients undergoing total knee arthroplasty: a double-blind, randomized clinical consort study. *Medicine (Baltimore)* 2016; 95: e4082.
159. Lu J, Chen G, Zhou H, Zhou Q, Zhu Z, Wu C. Effect of parecoxib sodium pretreatment combined with dexmedetomidine on early postoperative cognitive dysfunction in elderly patients after shoulder arthroscopy: a randomized double blinded controlled trial. *J Clin Anesth* 2017; 41: 30-4.
160. De Cosmo G, Sessa F, Fiorini F, Congedo E. Effect of remifentanyl and fentanyl on postoperative cognitive function and cytokines level in elderly patients undergoing major abdominal surgery. *J Clin Anesth* 2016; 35: 40-6.
161. Luo A, Yan J, Tang X, Zhao Y, Zhou B, Li S. Postoperative cognitive dysfunction in the aged: the collision of neuroinflammation with perioperative neuroinflammation. *Inflammopharmacology* 2019; 27: 27-37.
162. Wang LW, Zhu MJ, Li Y, Wang ST, Zhou MY, Yu YJ, et al. FKBP51 is associated with early postoperative cognitive dysfunction in elderly patients undergoing hip fracture surgery. *Medicine (Baltimore)* 2019; 98: e14037.
163. Fan Y, Yuan L, Ji M, Yang J, Gao D. The effect of melatonin on early postoperative cognitive decline in elderly patients undergoing hip arthroplasty: a randomized controlled trial. *J Clin Anesth* 2017; 39: 77-81.
164. Kotekar N, Shenkar A, Nagaraj R. Postoperative cognitive dysfunction - current preventive strategies. *Clin Interv Aging* 2018; 13: 2267-73.
165. Sánchez A, Thomas C, Deeken F, Wagner S, Klöppel S, Kentischer F, et al. Patient safety, cost-effectiveness, and quality of life: reduction of delirium risk and postoperative cognitive dysfunction after elective procedures in older adults-study protocol for a stepped-wedge cluster randomized trial (PAWEL Study). *Trials* 2019; 20: 71.
166. Qaseem A, Snow V, Fitterman N, Hornbake ER, Lawrence VA, Smetana GW, et al. Risk assessment for and strategies to reduce perioperative pulmonary complications for patients undergoing

- noncardiothoracic surgery: a guideline from the American College of Physicians. *Ann Intern Med* 2006; 144: 575-80.
167. Lawrence VA, Hilsenbeck SG, Noveck H, Poses RM, Carson JL. Medical complications and outcomes after hip fracture repair. *Arch Intern Med* 2002; 162: 2053-7.
168. Lo IL, Siu CW, Tse HF, Lau TW, Leung F, Wong M. Pre-operative pulmonary assessment for patients with hip fracture. *Osteoporos Int* 2010; 21 Suppl 4: S579-86.
169. Lawrence VA, Cornell JE, Smetana GW; American College of Physicians. Strategies to reduce postoperative pulmonary complications after noncardiothoracic surgery: systematic review for the American College of Physicians. *Ann Intern Med* 2006; 144: 596-608.
170. Kim YS, Won YJ, Lee DK, Lim BG, Kim H, Lee IO, et al. Lung ultrasound score-based perioperative assessment of pressure-controlled ventilation-volume guaranteed or volume-controlled ventilation in geriatrics: a prospective randomized controlled trial. *Clin Interv Aging* 2019; 14: 1319-29.
171. Higashikawa T, Shigemoto K, Goshima K, Usuda D, Okuro M, Moriyama M, et al. Urinary retention as a postoperative complication associated with functional decline in elderly female patients with femoral neck and trochanteric fractures: a retrospective study of a patient cohort. *Medicine (Baltimore)* 2019; 98: e16023.
172. Tobu S, Noguchi M, Hashikawa T, Uozumi J. Risk factors of postoperative urinary retention after hip surgery for femoral neck fracture in elderly women. *Geriatr Gerontol Int* 2014; 14: 636-9.

## Appendix

Appendix 1. Search strategies and items found for each database

Items found	Embase	Items found	Cochrane	Items found	KoreaMed	Items found
2960	((geriatric anesthesia/de) AND ('perioperative period'/de OR 'preoperative care'/de OR 'peroperative care'/de OR 'postoperative care'/de)) OR (anesthesia/de AND ('perioperative period'/de OR 'preoperative care'/de OR 'peroperative care'/de OR 'postoperative care'/de) AND (geriatrics/de OR aged/exp)) (anesthesia* OR postanesthesia* OR anesthetic):ti,ab  (perioperative* OR preoperative* OR intraoperative* OR postoperative* OR peri-operative* OR pre-operative* OR intra-operative* OR post-operative*):ti,ab  (geriatric OR geriatrics OR elder OR elders OR elder* OR older OR aged OR senior OR seniors OR gerontolog* OR frail OR frailty OR 'very old' OR (old NEXT/1 (patient OR patients OR inpatient OR inpatients))):ti	827	[mh Anesthesia] AND ([mh ^"Perioperative Care"] OR [mh "Preoperative Care"] OR [mh "Intraoperative Care"] OR [mh "Postoperative Care"]) AND ([mh Geriatrics] OR [mh Aged])  (anesthesia* OR postanesthesia* OR anesthetic):ti,ab  (perioperative* OR preoperative* OR intraoperative* OR postoperative* OR peri-operative* OR pre-operative* OR intra-operative* OR post-operative*):ti,ab  (geriatric OR geriatrics OR elder OR elders OR elder* OR older OR aged OR senior OR seniors OR gerontolog* OR frail OR frailty OR "very old" OR (old NEXT/1 (patient OR patients OR inpatient OR inpatients))):ti	3	(anesthesia* OR anaesthesia* OR anesthetic OR anaesthetic)  (perioperative* OR preoperative* OR intraoperative* OR postoperative* OR peri-operative* OR pre-operative* OR intra-operative* OR post-operative*)  (geriatric OR geriatrics OR elder OR elders OR elder* OR older OR aged OR senior OR seniors OR gerontolog* OR frail OR frailty OR "old patient" OR "old patients" OR "old inpatient" OR "old inpatients" OR "very old")	
1420	#2 AND #3 AND #4	2038	#2 AND #3 AND #4	738	#2 AND #3 AND #4	
4262	#1 OR #5  review:it OR 'meta analysis'/exp OR 'systematic review'/de OR 'multi-center study'/de OR 'observational study'/de OR 'randomized controlled trial'/exp	2768	#1 OR #5  random*:ti,ab	740		
1197	#6 AND #7  Animal/exp NOT (Human/exp AND Animal/exp)  (adolescent/exp OR adult/de OR 'middle aged'/de OR 'young adult'/de OR juvenile/exp) NOT (aged/exp AND (adolescent/exp OR adult/de OR 'middle aged'/de OR 'young adult'/de OR juvenile/exp))	702	#6 AND #7  ([mh Adolescent] OR [mh ^Adult] OR [mh "Middle Aged"] OR [mh "Young Adult"] OR [mh Child] OR [mh Infant]) NOT ([mh Aged] AND ([mh Adolescent] OR [mh ^Adult] OR [mh "Middle Aged"] OR [mh "Young Adult"] OR [mh Child] OR [mh Infant]))	548		
1182	#8 NOT (#9 OR #10)	671	#6 NOT #10	546		361
502	([english]/lim OR [korean]/lim) AND [2009-2019]/py	366	Publication date from 2009/01/01	399	(DP:2009:2019), except case report	284
502		366		399		284
502		183		191		276
1152						