

## Atherosclerotic Cardiovascular Disease in Rheumatoid Arthritis: Impact of Inflammation and Antirheumatic Treatment

Anne Mirjam Kerola <sup>1,2</sup> Silvia Rollefstad <sup>1</sup> and Anne Grete Semb <sup>1</sup>

1. Preventive Cardio-Rheuma Clinic, Division of Rheumatology and Research, Diakonhjemmet Hospital, Oslo, Norway;

2. Department of Rheumatology, Päijät-Häme Joint Authority for Health and Wellbeing, Lahti, Finland

### Abstract

Patients with rheumatoid arthritis (RA) are at approximately 1.5-fold risk of atherosclerotic cardiovascular disease (CVD) compared with the general population, a phenomenon resulting from combined effects of traditional CVD risk factors and systemic inflammation. Rheumatoid synovitis and unstable atherosclerotic plaques share common inflammatory mechanisms, such as expression of proinflammatory cytokines interleukin (IL)-1, tumour necrosis factor (TNF)- $\alpha$  and IL-6. RA patients are undertreated in terms of CVD prevention, and structured CVD prevention programmes are warranted. Alongside management of traditional risk factors, suppressing systemic inflammation with antirheumatic medication is fundamental for the reduction of CVD risk among this high-risk patient group. Many antirheumatic drugs, especially methotrexate, TNF- $\alpha$ -inhibitors and IL-6-inhibitors are associated with reduced risk of CVD in observational studies among RA patients, but randomised controlled trials with hard CVD endpoints are lacking. In patients without rheumatic disease, anti-inflammatory therapies targeting nucleotide-binding oligomerisation domain, leucine-rich repeat and pyrin domain-containing protein 3 inflammasome and the IL-1/IL-6 pathway arise as potential therapies after an atherosclerotic CVD event.

### Keywords

Rheumatoid arthritis, cardiovascular disease, atherosclerosis, inflammation, antirheumatic agents

**Disclosure:** This work was supported by research grants from the Foundation for Research in Rheumatology (FOREUM; AMK) and by grants from the South Eastern Regional Health Authorities of Norway (2016063 for SR and 2013064 for AGS). AMK has received speaker's fees or congress sponsorship from Boehringer-Ingelheim, Pfizer, Celgene, UCB Pharma and Mylan, and attended advisory boards of Pfizer, Gilead and Boehringer-Ingelheim. AGS has received speaker honoraria and/or consulting fees from Sanofi, AbbVie, Novartis, Bayer and Lilly. SR has no conflicts of interest to declare.

**Received:** 21 November 2020 **Accepted:** 19 February 2021 **Citation:** *European Cardiology Review* 2021;16:e18. **DOI:** <https://doi.org/10.15420/ecr.2020.44>

**Correspondence:** Anne Kerola, Preventive Cardio-Rheuma Clinic, Division of Rheumatology and Research, Diakonhjemmet Hospital, Diakonveien 12, 0370 Oslo, Norway. E: [anne.kerola@helsinki.fi](mailto:anne.kerola@helsinki.fi)

**Open Access:** This work is open access under the CC-BY-NC 4.0 License which allows users to copy, redistribute and make derivative works for non-commercial purposes, provided the original work is cited correctly.

Atherosclerotic cardiovascular disease (ASCVD) is an important cause of disability and is the leading cause of death globally.<sup>1</sup> Targeting modifiable cardiovascular risk factors, such as diabetes, hypercholesterolaemia, hypertension, smoking, physical inactivity and obesity, improves CVD outcomes in individuals with established CVD or those at high risk of CVD. Rheumatoid arthritis (RA) is a chronic inflammatory joint disease (IJD) with a prevalence of 0.5–1%. It is characterised by symmetric polyarthritis and, in the majority of RA patients, formation of autoantibodies, such as rheumatoid factor (RF) and anti-citrullinated protein antibodies (ACPA). Similar to diabetes, RA is recognised as an independent risk factor for CVD and the key underlying pathology is systemic inflammation.

During the past few decades, mounting evidence on CVD risk in RA has led to substantial advances in CVD risk management in this patient population.<sup>2</sup> RA patients are still undertreated in terms of primary and secondary CVD prevention.<sup>3,4</sup> Alongside targeting traditional CVD risk factors, suppressing disease activity and inflammation with antirheumatic treatment is expected to reduce CVD burden in these patients. Inflammation plays a major role in atherosclerosis and it has been recognised that among well-treated ASCVD patients without IJD,

low-grade inflammation is a key determinant of residual CVD risk.<sup>5</sup> Groundbreakingly, anti-inflammatory therapies are being studied for the treatment of ASCVD outside the context of IJDs.


In this review, we summarise the evidence on CVD risk in RA with emphasis on inflammation and the impact of antirheumatic treatment on CVD risk. We also discuss the current knowledge on anti-inflammatory therapies for secondary prevention of ASCVD from a rheumatological point of view.

### Epidemiology of CVD in RA

People with RA have an approximate 1.5-fold risk of cardiovascular events and mortality compared to the general population.<sup>6,7</sup> This excess cardiovascular burden is comparable to that of people with diabetes, although it is not as well recognised or structurally managed.<sup>8</sup>

According to a meta-analysis of observational studies, the risk of MI in people with RA compared to the general population is increased by 70% with no differences in the relative risk for men and women.<sup>6</sup> The increase in the risk of coronary heart disease (CHD) seems to occur early on in the

Figure 1: Central Inflammatory Mediators in Rheumatoid Synovitis and Atherosclerosis



NLRP3 inflammasome activates the conversion of pro-IL-1 $\beta$  to biologically active IL-1 $\beta$ . IL-1 $\beta$  promotes production of downstream cytokines TNF- $\alpha$  and IL-6. IL-6 induces hepatic production of acute phase reactants such as CRP. Vascular inflammation in atherosclerosis is usually associated with normal or only slightly increased CRP (low-grade inflammation), whereas active rheumatoid arthritis often results in high-grade systemic inflammation that accelerates atherosclerosis. Antirheumatic drugs that are used to target the central proinflammatory cytokines IL-1, TNF- $\alpha$  and IL-6 are depicted in white arrows, whereas anti-inflammatory therapies that have been investigated in secondary prevention of ASCVD are depicted in grey arrows. ASCVD = atherosclerotic cardiovascular disease; CRP = C-reactive protein; IL = interleukin; LDL = low-density lipoprotein; MMPs = matrix metalloproteinases; NLRP3 = nucleotide-binding oligomerisation domain, leucine-rich repeat and pyrin domain-containing protein 3; oxLDL = oxidised low-density lipoprotein; TNF = tumour necrosis factor.

course of RA, or according to some studies, even before RA is diagnosable.<sup>10</sup> Patients with RA are less likely to report symptoms of angina and more likely to experience unrecognised MI and sudden cardiac death than the general population, suggesting an overrepresentation of vulnerable coronary plaques in patients with RA.<sup>10</sup> After an MI, RA patients have impaired prognosis compared to the general population in terms of recurrent ischaemia and mortality.<sup>11,12</sup>

In general, cardiovascular mortality rates have declined during the past 50 years and this has also translated into better survival in RA. Presumably due to marked advances in RA pharmacotherapy and improved control of disease activity, recent studies suggest that the gap in cardiovascular mortality between RA patients diagnosed in the 2000s and the general population may be narrowing or even closing.<sup>13,14</sup>

Risk of cerebrovascular accidents is increased among RA patients by 40% compared to the general population.<sup>6</sup> Alongside ASCVD, RA patients are at increased risk of other types of CVD, such as AF and heart failure.<sup>2</sup> Heart failure in RA may partly result from increased prevalence of CHD, but the risk of non-ischaemic heart disease and diastolic dysfunction is also aggravated.<sup>15</sup>

### Traditional Risk Factors do not Fully Explain Excess CVD Risk in RA

RA patients are more likely to suffer from insulin resistance, abnormal fat distribution, be physically inactive and smoke cigarettes compared to non-RA controls.<sup>16</sup> Tobacco smoking is the strongest environmental risk factor for RA. A large international RA cohort study demonstrated that about a quarter of cardiovascular events were attributable to smoking (population attributable risk being 37.2% for men and 18.1% for women).<sup>17</sup> Diabetes may be more prevalent in RA than in controls, although significant heterogeneity exists between studies.<sup>16</sup> Data regarding overrepresentation of hypertension among RA patients is controversial, but most reports indicate no difference.<sup>16</sup>

Traditional CVD risk factors, however, do not fully explain the observed excess cardiovascular morbidity and mortality in RA. RA-related factors, such as disease activity and burden, extra-articular disease manifestations, elevated inflammatory markers and RF/ACPA positivity attribute to about 30% of the 10-year risk of cardiovascular events.<sup>17</sup> A well-powered US-based registry study examined the relationship between longitudinally measured RA disease activity and the risk of CVD and demonstrated that treating RA from high disease activity to remission was associated with an approximate 50% reduction in risk of cardiovascular events.<sup>18</sup> Both

exposure to each RA flare and cumulative disease activity seem to lead to an increased risk of CVD.<sup>19</sup> In some studies, excess cardiovascular mortality risk has been confined only to seropositive RA patients.<sup>20</sup>

### Inflammation Drives CVD Risk in RA and the General Population

Development of an atherosclerotic plaque is not passive deposition of cholesterol in the vascular wall but an active inflammatory process involving the innate and adaptive immune system. Next, we will highlight the role of selected inflammatory mediators, which are central to pathogenesis of RA, on the development of atherosclerosis, such as nucleotide-binding oligomerisation domain, leucine-rich repeat and pyrin domain-containing protein 3 (NLRP3) inflammasome, and proinflammatory cytokines interleukin 1 (IL-1), tumour necrosis factor alpha (TNF- $\alpha$ ) and interleukin 6 (IL-6). These inflammatory mediators are all interconnected in signalling pathways and they are targets for drugs that are used to treat rheumatic diseases and have gained interest as potential drugs for secondary prevention of ASCVD (Figure 1).

Similar to high-sensitivity C-reactive protein (hsCRP), the role of IL-6 in prediction of cardiovascular events in the general population has been confirmed in primary and secondary prevention cohorts.<sup>5</sup> To highlight a few studies, increased levels of IL-6 were associated with the risk of future MI in a prospective study with almost 15,000 apparently healthy men.<sup>21</sup> In a secondary analysis of the CIRT study, IL-6 and hsCRP predicted major recurrent cardiovascular events in patients with prior MI or multivessel CHD who also had diabetes or metabolic syndrome, despite modern CVD preventive polypharmacotherapy and high rates of coronary revascularisation.<sup>22</sup> IL-6 is also a marker of poor prognosis in acute coronary syndrome (ACS).<sup>23</sup>


Based on two Mendelian randomisation analyses, IL-6 signalling may have a causal role in the development of ASCVD.<sup>24,25</sup> In MI patients, IL-6 production occurs at the site of plaque rupture, as indicated by higher concentration of IL-6 locally compared to levels in systemic circulation.<sup>26</sup> IL-6 may also be involved in ischaemia-reperfusion myocardial injury.<sup>27</sup> In contrast, CRP is produced in the liver in response to IL-6, and in fact, hsCRP seems to be an easy-to-use downstream marker of IL-6 activation rather than a causal agent in atherogenesis.

NLRP3 inflammasome is a macromolecular protein complex of the innate immune system, which can be activated by many stimuli, such as urate crystals in gout or cholesterol crystals within an atherosclerotic plaque.<sup>28</sup> IL-1 $\beta$  and NLRP3 inflammasome are important IL-6 upstream activators. Preclinical evidence links IL-1 to impaired vasodilatation, oxidative stress, plaque formation, growth and rupture.<sup>29</sup> IL-1 $\alpha$  may contribute to infarct size, whereas IL-1 $\beta$  is associated to adverse cardiac remodelling.<sup>29</sup> Based on an ex vivo rodent model of myocardial ischaemia reperfusion, NLRP3 inflammasome may contribute to infarct size.<sup>30</sup> Levels of TNF- $\alpha$  have also been demonstrated to predict future cardiovascular events.<sup>31</sup> TNF- $\alpha$  is an important regulator of vascular homeostasis and contributes to the development of endothelial dysfunction and a prothrombotic state.<sup>32</sup> Among RA patients, increased concentrations of both TNF- $\alpha$  and IL-6 are associated with increased prevalence of coronary artery calcification, independently of Framingham risk score or the presence of diabetes.<sup>33</sup>

### Systemic Inflammation in RA Modifies Traditional CVD Risk Factors

The associations of lipids with CVD risk in RA are complex and essentially linked with inflammation. The levels of total cholesterol (TC), HDL

**Figure 2: The Inverse Relationship Between Changes in Inflammatory and Lipid Parameters**



Usually, before treatment of RA is initiated and when RA disease activity is high, inflammatory markers may increase and lipid levels decrease. Antirheumatic treatment suppresses synovial and systemic inflammation effectively and lipid levels increase. Similar outbursts of inflammatory markers and concomitant reductions in lipid levels can occur during flares of rheumatoid arthritis. CRP = C-reactive protein; RA: rheumatoid arthritis. Source: Choy et al. 2014.<sup>36</sup> Reproduced with permission from Oxford University Press.


cholesterol (HDL-C) and LDL cholesterol (LDL-C) during active RA and ongoing inflammation typically decrease (Figure 2), a trend that is seen also in septicaemia and other inflammatory states.<sup>34</sup> Low lipid levels during inflammation are, however, not associated with lower CVD risk, but rather the opposite.<sup>34</sup> This phenomenon is referred to as the lipid paradox.<sup>35</sup> In 2011, a retrospective cohort study among 651 RA patients reported a non-linear association of CVD risk and TC levels, with increasing CVD risk at TC levels below 4 mmol/L.<sup>35</sup> Recently, this finding was confirmed in a study pooling four of the largest North American RA cohorts with information on coronary arterial calcium (CAC) scores: in contrast to controls, the association of LDL-C with CAC score in RA patients was U-shaped and the largest relative difference in coronary atherosclerosis between the RA and control groups was found among those with an LDL-C lower than 1.8 mmol/L.<sup>36</sup> In addition, it was shown that CAC scores were higher in RA patients than in controls across LDL-C levels.<sup>36</sup> A possible mechanism for the increased CVD risk despite low LDL-C is that inflammation promotes the oxidation of LDL-C, which is proatherogenic.<sup>37</sup> HDL-C particles of RA patients may be unable to prevent LDL oxidation more often than those in controls.<sup>38</sup>

In addition to lipid levels, inflammation modifies other CVD risk factors such as insulin resistance, body composition and blood pressure (Figure 3). A 2020 study exploring the effects of CRP levels on systolic blood pressure in both an RA cohort and a large non-RA outpatient cohort revealed a biphasic association of systolic blood pressure with CRP (with a positive association at CRP <6 mg/l and inverse at CRP >6 mg/l).<sup>39</sup> Body mass index (BMI) may also have a paradoxical effect on mortality in RA: patients with the lowest BMI seem to have the highest mortality rates.<sup>40</sup> This might be due to the presence of rheumatoid cachexia, sarcopenia and frailty among those with prolonged high disease activity and hence higher CVD risk. Proinflammatory cytokines TNF- $\alpha$  and IL-6 induce insulin resistance, and many antirheumatic agents, including hydroxychloroquine, TNF- $\alpha$ , IL-1- and IL-6-inhibitors, seem to improve markers of glucose metabolism in RA patients.<sup>41</sup>

### Prevention of CVD in RA Estimation of Cardiovascular Risk

The accurate estimation of CVD risk in RA and other IJDs is challenging and remains an area of active research. Patients tend to underestimate

**Figure 3: The Interplay Between Synovial and Systemic Inflammation and Cardiovascular Risk Factors in Rheumatoid Arthritis**



IL = interleukin; TNF = tumour necrosis factor.

their CVD risk, warranting more patient education on this topic.<sup>42</sup> In 2017, the European League Against Rheumatism (EULAR) published an update on recommendations for CVD risk management in RA and other IJDs.<sup>43</sup> These recommendations emphasise that the rheumatologist is responsible for making sure that CVD risk management is addressed in patients with RA and suggest CVD risk assessment for all RA patients at least once every 5 years and reconsidered after major changes in antirheumatic therapy. Lipids should ideally be measured when RA is in remission due to the impact of inflammation on lipoprotein levels.<sup>43</sup>

CVD risk management among RA patients is performed either by rheumatologists during outpatient visits, GPs collaborating with rheumatologists or, less often, cardiologists cooperating with rheumatologists in a cardio-rheumatology clinic.<sup>44</sup> A simple suggestion for CVD risk assessment in rheumatology clinics is that if blood pressure and lipids are recorded along with the already standardised recording of gender, age and smoking status, CVD risk can be calculated based on the Systematic Coronary Risk Evaluation (SCORE) or other CVD risk prediction algorithms. If the recommended treatment threshold is exceeded, patients could be referred to their primary care physician or a cardiologist for CVD prevention measures.<sup>45</sup>

Most established CVD risk calculators, such as SCORE, and the Reynolds and Framingham risk scores, incorporate only traditional risk factors and underestimate CVD risk among RA patients.<sup>46</sup> QRISK2 and QRISK3 include RA as an independent CVD risk factor with a weight of 1.4 and have been shown to inaccurately predict CVD risk in RA.<sup>46</sup> The 2018 American College of Cardiology and American Heart Association (ACC/AHA) guidelines on the management of blood cholesterol suggest that RA should be considered as a 'risk enhancer', promoting statin therapy in patients with intermediate or borderline risk.<sup>47</sup> The 2015/2016 EULAR recommendations state that CVD risk scores should be adapted for patients with RA by a 1.5 multiplication factor if RA is not already included in the algorithm in use.<sup>43</sup>

Substantial efforts have been made to develop an RA-specific CVD risk prediction model, but with only modest results. An international consortium – A Trans-Atlantic Cardiovascular Consortium for Rheumatoid Arthritis (ATACC-RA) – including 13 RA cohorts from 10 countries developed two risk prediction models that included either a measure of disease activity or functional status but disappointingly, have been found to not improve discrimination when compared with established CVD prediction models.<sup>48</sup> A similar well-powered registry study from North America presented a model incorporating measures of disease activity, disability, daily use of prednisolone and disease duration.<sup>49</sup> It performed well in

internal validation, but failed to improve discrimination compared to established risk calculators in external validation.<sup>49,50</sup>

Ultrasonography of carotid arteries is a promising non-invasive method to improve CVD risk prediction in RA. It is already incorporated into clinical practice in a few rheumatology clinics. Multiple studies have concluded that RA patients have higher prevalence of asymptomatic carotid atherosclerosis than non-RA controls.<sup>51</sup> Carotid ultrasound may markedly improve CVD risk evaluation in RA as was shown in a study of 355 patients with IJDs where 30–60% of patients with an estimated low-to-moderate CVD risk according to established CVD risk calculators had carotid plaques and it was therefore reclassified to very high-risk category with an indication for lipid-lowering therapy.<sup>52</sup>

### Management of Traditional Cardiovascular Risk Factors

Management of traditional CVD risk factors among RA patients is fundamental and includes advice about maintaining a healthy diet, regular exercise, weight control, smoking cessation and treatment with antihypertensive and lipid-lowering agents. Compelling evidence comparing different primary prevention strategies in RA are lacking and risk factor management is recommended to be carried out according to national guidelines.<sup>2,43</sup> This means that treatment strategies, thresholds and targets for antihypertensive, anti-diabetic and lipid-lowering therapies are the same in RA and the general population.

It may be asked whether primary prevention of CVD with statins could be beneficial to RA patients irrespective of their estimated CVD risk. A multicentre randomised controlled trial (RCT) TRACE-RA compared the effect of 40 mg daily atorvastatin versus placebo on hard CVD endpoints in RA patients.<sup>53</sup> Due to unexpectedly low cardiovascular event rates, the study was terminated prematurely after recruiting only 3,002 RA patients with a median follow-up of 2.5 years. The reduction in cardiovascular events in the atorvastatin versus the placebo arm was estimated to be 34% but was not statistically significant probably due to lack of power caused by short observation period and low number of patients. Although, most importantly, this large risk reduction of 34% has been found in other large statin studies with hard CVD endpoints.<sup>54</sup>

Statin seem to be equally safe and effective among RA patients and the general population. A post hoc analysis of two large secondary prevention trials demonstrated that the effect of statins on reduction of lipids and cardiovascular event rates among patients with and without IJDs was comparable, despite lower baseline lipid levels in the IJD group.<sup>55</sup> Statins may also ameliorate RA disease activity and reduce CRP and erythrocyte sedimentation rate (ESR) levels through their pleiotropic effects.<sup>56</sup>

Smoking cessation in patients with RA is of utmost importance. An international multicenter study showed that smoking cessation among RA patients is a predictor of reduced rate of cardiovascular events (for former versus current smokers HR 0.70; 95% CI [0.51–0.95]; p=0.02, and for never versus current smokers HR 0.48 ; 95% CI [0.34–0.69]; p<0.001).<sup>57</sup> Smoking cessation was also associated with lower disease activity.<sup>57</sup> A trial of intensive smoking cessation intervention on RA disease activity (NCT02901886) is currently under way. Every RA patient who smokes should receive advice for smoking cessation and, in our opinion, smoking cessation programmes would ideally be implemented in rheumatology clinics.

Patient education is the key to lifestyle modifications. All patients should receive sufficient information on CVD risk associated with RA in oral and

written form. Heart friendly diet recommendations are the same for RA patients as for the general population. Even brief standardised advice on a heart friendly diet may result in a change in nutritional habits, as implied by a small pilot study among IJD patients.<sup>58</sup> Barriers to physical activity in RA include pain, fatigue and stiffness, but on the other hand, exercise may aid symptom management, pain relief and joint function.<sup>59</sup>

### Antirheumatic Treatment and Cardiovascular Risk in RA and Beyond

The antirheumatic medications used in the treatment of RA comprise conventional synthetic disease-modifying antirheumatic drugs (DMARDs) such as methotrexate, biological DMARDs such as TNF- $\alpha$ , IL-6- or IL-1-inhibitors, and targeted synthetic DMARDs, such as Janus kinase (JAK) inhibitors. The pharmacodynamics for conventional synthetic DMARDs is not fully elucidated, whereas biological and targeted synthetic DMARDs affect specific stages of inflammatory signalling.

Suppression of disease activity with antirheumatic treatment is fundamental in lowering CVD risk. Although the mechanisms remain poorly understood, observational studies demonstrate an association of many DMARDs and lowered CVD risk in RA. An increasing number of clinical trials have also shown positive effects of several DMARDs on surrogate markers of CVD among RA patients, but RCTs with hard CVD endpoints are lacking.<sup>60–63</sup>

In the following sections, we will discuss the cardiovascular effects of DMARDs among RA patients and summarise the published RCTs that explore anti-inflammatory therapies in secondary prevention of ASCVD in patients without IJD. Potential anti-inflammatory therapies for ASCVD, which are not used in the treatment of RA and do not target the central IL-1, TNF- $\alpha$  and IL-6 pathways are not included in this review.

### Methotrexate

Methotrexate is the anchor drug of RA treatment and the most important part of the initial antirheumatic treatment strategy. It can be combined with other synthetic or biological DMARDs. Its effects may be mediated at least partly via increased cellular adenosine release.<sup>64</sup> Methotrexate may also have direct atheroprotective effects: in a model of cholesterol-fed rabbits, methotrexate reduced the size of atherosclerotic lesions by 75% and also diminished macrophage migration and presence of apoptotic cells in the vessel wall.<sup>65</sup>

From the beginning of the 2000s, mounting evidence from observational studies among RA cohorts point to an association of methotrexate use and reduced cardiovascular morbidity and mortality. In one prospective study of 1,240 RA patients with a mean follow-up of 6 years, after adjustment for relevant prognostic factors, cardiovascular mortality was reduced by as much as 70% and overall mortality by 60% in patients on methotrexate versus those never treated with methotrexate.<sup>66</sup>

According to a 2015 meta-analysis, methotrexate use for IJDs was associated with a 28% reduction in fatal and non-fatal cardiovascular events, based on eight observational studies.<sup>67</sup> A recently published, well-powered retrospective study using Medicare claims data on RA patients who were initiating biological DMARDs showed that concomitant methotrexate use was associated with a decreased risk of cardiovascular events also among biological DMARD users.<sup>68</sup> Unmeasured confounding is naturally a threat to the validity of these findings.

Ridker et al. conducted a large RCT of low-dose methotrexate compared with placebo in 4,786 patients with previous MI or multivessel CHD and

either type 2 diabetes or metabolic syndrome (CIRT).<sup>69</sup> The study was terminated after a median follow-up of 2.3 years and methotrexate did not affect the rate of a composite CVD endpoint or the levels of IL-1 $\beta$ , IL-6 and CRP. It may be speculated whether an enrichment strategy with hsCRP to guide enrollment of patients into the study would have affected the outcome of the trial (median hsCRP at randomisation was only 1.6 mg/l). In contrast to ASCVD patients with low-grade inflammation, high RA disease activity often results in high-grade inflammation, which in most patients is well-controlled with methotrexate. Thus, it seems logical that the effect of methotrexate on CVD endpoints in RA is different from patients without IJD.

### Other Conventional Synthetic DMARDs

Hydroxychloroquine, originally an antimalarial drug, may have cardioprotective effects through favourable changes in lipid and glucose metabolism.<sup>70</sup> A safety pilot study of hydroxychloroquine compared with placebo for patients with MI to prevent recurrent cardiovascular events (NCT02648464) has now recruited 125 patients and if no important safety signals arise, the study will most likely be expanded to involve 2,500 patients. In one cross-sectional cohort of 4,363 patients with RA from 15 countries, use of not only methotrexate and biological DMARDs but also use of leflunomide and sulfasalazine was associated with a reduction in the risk of cardiovascular events.<sup>71</sup> Leflunomide causes hypertension, and blood pressure must be monitored during treatment.<sup>72</sup>

### TNF- $\alpha$ Inhibitors

If treatment with conventional synthetic DMARDs fails and RA disease activity is high, there is an indication for treatment with biological DMARDs. The most frequently prescribed biological DMARDs are TNF- $\alpha$  inhibitors, which have been in clinical use for two decades. In a 2015 meta-analysis of observational cohort studies among patients with IJD, anti-TNF- $\alpha$  therapy was associated with a 30% reduction in all cardiovascular events, 41% reduction in MIs, 43% reduction in strokes and up to 70% reduction in major adverse cardiac events (MACEs) when compared to other antirheumatic treatments.<sup>67</sup> However, the decrease in ACS risk may not be similar in all patients treated with TNF- $\alpha$  inhibitors but conditional to treatment response. In a Swedish registry study among TNF- $\alpha$  inhibitor initiators with RA, 1-year risk of ACS was reduced by 50% among those with good treatment response compared to non-responders.<sup>73</sup> In addition, compared with the general population, non-responders had a doubled risk of ACS but those with a good response had equal risk.

TNF- $\alpha$  inhibitors may improve endothelial function and carotid intima-media thickness among RA patients.<sup>61,63</sup> However, in contrast to RA patients with high disease activity, patients with low disease activity or remission present no difference in carotid ultrasound findings or arterial stiffness compared with non-RA controls, suggesting that deceleration of these pre-atherosclerotic changes may be related to effective disease control rather than a certain DMARD.<sup>74</sup>

In the 1990s, preclinical data suggested that TNF- $\alpha$  played a role in the evolution and progression of heart failure. However, RCTs on TNF- $\alpha$  inhibitors etanercept and infliximab for moderate-to-severe heart failure did not show a beneficial effect on hospitalisation or death due to chronic heart failure.<sup>75,76</sup> In fact, in the group with the highest infliximab dose of 10 mg/kg, the combined risk of death or hospitalisation for heart failure through 28 weeks compared to placebo was almost tripled.<sup>76</sup> As a consequence, New York Heart Association III–IV heart failure is a contraindication for TNF- $\alpha$  inhibitors.

### IL-6 Inhibitors

Tocilizumab and sarilumab are anti-IL-6 receptor monoclonal antibodies that are used in the treatment of RA. They are highly effective in suppressing systemic inflammation and normalising the production of acute phase reactants, especially CRP. IL-6 inhibitors increase lipid levels, which has raised concerns about their cardiovascular safety. A Phase IV non-inferiority trial addressed this issue by comparing tocilizumab to etanercept among 3,080 patients with seropositive RA and at least one CVD risk factor. During a mean follow-up of 3.2 years, the occurrence of MACE did not differ between the groups.<sup>77</sup>

Recently, signs of an atheroprotective effect of IL-6 receptor blockade in RA have been published. A meta-analysis of observational studies suggested that exposure to tocilizumab was associated with a lower risk of MACE but not stroke compared to exposure to TNF- $\alpha$  inhibitors.<sup>78</sup> Despite elevations in TC, HDL-C and LDL-C, tocilizumab treatment incurs no change in TC/HDL-C ratio, and it may modify HDL particles towards an anti-inflammatory composition and also reduce lipoprotein (a) levels, which altogether may modify the risk towards a more favourable CVD outcome.<sup>60,79</sup> Concomitant statin treatment with tocilizumab seems to be effective for lipid reduction.<sup>80</sup> Of note, in patients using IL-6 inhibitors treatment with fluvastatin, pravastatin or rosuvastatin should be chosen, since IL-6 inhibitors may affect cytochrome P450 3A4 (CYP3A4) activity and thus the efficacy of other statins.<sup>2</sup>

The effect of IL-6 receptor blockade has also been explored for ACS. In an RCT including 121 patients with non-ST-elevation MI (STEMI) scheduled for coronary angiography, a single dose of tocilizumab not only lowered hsCRP levels as expected, but also attenuated the increase in troponin T during hospitalisation, suggesting a smaller infarct size.<sup>81</sup> These findings are being further explored in an ongoing study (NCT03004703), in which 200 patients with first-time STEMI will be randomised to receive either tocilizumab or placebo prior to percutaneous coronary intervention with a primary endpoint of myocardial salvage index measured by cardiac MRI 3–7 days after the intervention.

### IL-1 inhibitors

There are currently three IL-1 inhibitors available on the market: the IL-1 receptor antagonist anakinra, the soluble decoy receptor rilonacept and the monoclonal IL-1 $\beta$  antibody canakinumab. Of these, only anakinra is approved for the treatment of RA and its efficacy compared to other biological DMARDs is limited. In contrast, IL-1 inhibitors are highly effective in the treatment of certain autoinflammatory diseases. Evidence from rather small clinical trials imply that anakinra may reduce glycated haemoglobin levels among RA patients with diabetes and improve cardiac function, i.e. left ventricular longitudinal strain and coronary flow reserve.<sup>82,83</sup>

Cardiovascular effects of anakinra have been explored outside rheumatic diseases. In an RCT among 182 non-STEMI patients, 14-day treatment with anakinra attenuated inflammatory response following MI.<sup>84</sup> In two pilot studies with a total of 70 patients hospitalised for STEMI, anakinra not only blunted inflammatory response but also improved left ventricle systolic volume index in cardiac MRI and seemed to reduce the incidence of new-onset heart failure compared to placebo, suggesting a favourable effect on left ventricle remodelling and function.<sup>85</sup>

A pivotal proof-of-concept trial for the inflammatory hypothesis of atherothrombosis, CANTOS, recruited 10,061 patients with prior MI and persistent elevation of hsCRP >2 mg/l. Canakinumab 150 mg, but not 50 mg, reduced the incidence of a combined endpoint of stroke, MI or CVD

death by 15% (HR 0.85; 95% CI [0.74–0.98]; p=0.021) compared with placebo over 48 months of follow-up.<sup>86</sup> A similar reduction in the primary endpoint was observed for canakinumab 300 mg, but this did not meet the prespecified threshold for statistical significance.

### Other Biological DMARDs and JAK Inhibitors

Abatacept, an inhibitor of co-stimulation of T-cells, was found to be associated with a decreased risk of a composite CVD endpoint compared to conventional DMARDs in a large prospective observational study from the US.<sup>87</sup> A similar effect was not observed for rituximab, a monoclonal anti-CD20 antibody.

JAK inhibitors are targeted synthetic DMARDs that modulate the cytokine receptor-mediated intracellular signalling cascades. In phase II and III RCTs, a safety signal of a dose-dependent increase in the risk of venous thromboembolism (VTE) was identified with JAK inhibitor tofacitinib.<sup>88</sup> However, a recent meta-analysis of RCTs reporting JAK inhibitor safety data revealed no increased risk of all CVD, MACE or VTE, with the exception that baricitinib 2 mg daily was safer than 4 mg with regard to all CVD.<sup>89</sup> Caution should be taken when prescribing JAK inhibitors to patients with risk factors for VTE. Similar to IL-6 inhibitors, JAK inhibitors increase cholesterol levels, a side-effect that can be well-controlled with statins.<sup>90</sup>

### Non-steroidal Anti-inflammatory Drugs, Glucocorticoids and Colchicine

Non-steroidal anti-inflammatory drugs (NSAIDs) are often used for rheumatic pain. Long-term use of NSAIDs, especially cyclo-oxygenase-2 selective inhibitors, has been associated with increased risk of CVD. This also holds true in patients with IJD.<sup>67</sup> NSAIDs should be used with caution, especially in the presence of CVD risk factors or CVD.<sup>43,91</sup>

Glucocorticoids have well-known side-effects that amplify CVD risk, including weight gain, hypertension and insulin resistance. Observational studies have demonstrated that low-dose glucocorticoid use in IJD is associated with an approximate 50% increase in CVD risk.<sup>67</sup> This association is dependent on glucocorticoid dosage and cumulative exposure.<sup>92</sup> It is unclear, however, to which extent these results are affected by confounding by indication. This question may be answered by the GLORIA study which is assessing the harm, benefit and costs of 5 mg prednisolone daily versus placebo added to standard care in RA patients >65 years (NCT02585258). Glucocorticoids should be used for RA with the minimum effective dosage and tapered in case of remission or low disease activity.

Although not used in the treatment of RA but in gout, colchicine, an anti-inflammatory drug that inhibits tubulin polymerisation, microtubule generation and putatively also NLRP3 inflammasome activation, deserves attention (*Figure 1*).<sup>93</sup> Two large RCTs have shown that colchicine reduces the risk of cardiovascular events compared to placebo after acute MI and in chronic CHD.<sup>94,95</sup>

### Conclusion

RA patients have an increased ASCVD burden that is linked to disease activity, systemic inflammation, traditional CVD risk factors and a paradoxical decrease in lipid levels. Rheumatoid synovitis and unstable atherosclerotic plaque share common inflammatory pathophysiological mechanisms, such as expression of proinflammatory cytokines IL-1, TNF- $\alpha$  and IL-6. Increased CVD risk in RA may be reduced by addressing CVD risk factors and suppression of inflammation with effective antirheumatic

therapy. However, awareness of this phenomenon is still low in RA patients and healthcare professionals and CVD risk factors often remain undertreated in this high-risk population.

Several DMARDs, especially methotrexate, TNF- $\alpha$  inhibitors and IL-6 inhibitors, seem to improve various biomarkers of CVD and reduce cardiovascular events in observational studies among RA cohorts. It is uncertain, however, whether these observations are due to reduced disease activity in general or to specific atheroprotective effects. The former theory is supported by the notion that CVD risk and its surrogate markers seem to correlate with treatment response to antirheumatic medication. More research is warranted on tailoring of DMARD treatment

to RA patients with regard to their individual CVD risk. Furthermore, RA patients would most probably benefit from more structured CVD prevention programmes, such as in diabetes care.

Anti-inflammatory therapies – especially IL-1 inhibitors and colchicine – are promising agents to reduce inflammation-mediated residual risk of recurrent cardiovascular events among ASCVD patients. Further evaluation of IL-6 and IL-1 inhibitors as adjunct therapy for MI is warranted. Lessons from RA, a natural model of high-grade inflammation, can be used to understand the inflammatory mechanisms of atherosclerosis also in the general population and to identify novel targets for ASCVD treatment.  $\square$

- WHO. *Global Health Estimates 2016: Deaths by Cause, Age, Sex, by Country and by Region, 2000–2016*. Geneva: WHO, 2018.
- Semb AG, Ikdahl E, Wibetoe G et al. Atherosclerotic cardiovascular disease prevention in rheumatoid arthritis. *Nat Rev Rheumatol* 2020;16:361–79. <https://doi.org/10.1038/s41584-020-0428-y>; PMID: 32494054.
- Ikdahl E, Wibetoe G, Rollefstad S, et al. Guideline recommended treatment to targets of cardiovascular risk is inadequate in patients with inflammatory joint diseases. *Int J Cardiol* 2019;274:311–8. <https://doi.org/10.1016/j.ijcard.2018.06.111>; PMID: 30007486.
- Lindhardtsen J, Ahlehoff O, Gislason GH, et al. Initiation and adherence to secondary prevention pharmacotherapy after myocardial infarction in patients with rheumatoid arthritis: a nationwide cohort study. *Ann Rheum Dis* 2012;71:1496–501. <https://doi.org/10.1136/annrheumdis-2011-200806>; PMID: 22402144.
- Lawler PR, Bhatt DL, Godoy LC, et al. Targeting cardiovascular inflammation: next steps in clinical translation. *Eur Heart J* 2021;42:113–31. <https://doi.org/10.1093/eurheartj/ehaa099>; PMID: 32176778.
- Avina-Zubieta JA, Thomas J, Sadatsafavi M, et al. Risk of incident cardiovascular events in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Ann Rheum Dis* 2012;71:1524–9. <https://doi.org/10.1136/annrheumdis-2011-200726>; PMID: 22425941.
- Avina-Zubieta JA, Choi HK, Sadatsafavi M, et al. Risk of cardiovascular mortality in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Arthritis Rheum* 2008;59:1690–7. <https://doi.org/10.1002/art.24092>; PMID: 19035419.
- Lindhardtsen J, Ahlehoff O, Gislason GH, et al. The risk of myocardial infarction in rheumatoid arthritis and diabetes mellitus: a Danish nationwide cohort study. *Ann Rheum Dis* 2011;70:929–34. <https://doi.org/10.1136/ard.2010.143396>; PMID: 21389043.
- Kerola AM, Kerola T, Kauppi MJ, et al. Cardiovascular comorbidities antedating the diagnosis of rheumatoid arthritis. *Ann Rheum Dis* 2013;72:1826–9. <https://doi.org/10.1136/annrheumdis-2012-202398>; PMID: 23178207.
- Maradit-Kremers H, Crowson CS, Nicola PJ, et al. Increased unrecognized coronary heart disease and sudden deaths in rheumatoid arthritis: a population-based cohort study. *Arthritis Rheum* 2005;52:402–11. <https://doi.org/10.1002/art.20853>; PMID: 15693010.
- McCoy SS, Crowson CS, Maradit-Kremers H, et al. Long-term outcomes and treatment after myocardial infarction in patients with rheumatoid arthritis. *J Rheumatol* 2013;40:605–10. <https://doi.org/10.3899/jrheum.120941>; PMID: 23418388.
- Palomäki A, Kerola AM, Malmberg M, Rautava P, Kytö V. Patients with rheumatoid arthritis have impaired long-term outcomes after myocardial infarction – a nationwide case-control registry study. *Rheumatology (Oxford)* 2021 <https://doi.org/10.1093/rheumatology/keab204>; PMID: 33667301; epub ahead of press.
- Kerola AM, Nieminen TVM, Virta LJ, et al. No increased cardiovascular mortality among early rheumatoid arthritis patients: a nationwide register study in 2000–2008. *Clin Exp Rheumatol* 2015;33:391–8. PMID: 25936374.
- Lacaille D, Avina-Zubieta JA, Sayre EC, et al. Improvement in 5-year mortality in incident rheumatoid arthritis compared with the general population – closing the mortality gap. *Ann Rheum Dis* 2017;76:1057–63. <https://doi.org/10.1136/annrheumdis-2016-209562>; PMID: 28031164.
- mantel A, Holmqvist M, Andersson DC, et al. Association between rheumatoid arthritis and risk of ischemic and nonischemic heart failure. *J Am Coll Cardiol* 2017;69:1275–85. <https://doi.org/10.1016/j.jacc.2016.12.033>; PMID: 28279294.
- Liao KP, Solomon DH. Traditional cardiovascular risk factors, inflammation and cardiovascular risk in rheumatoid arthritis. *Rheumatology (Oxford)* 2013;52:45–52. <https://doi.org/10.1093/rheumatology/kes243>; PMID: 22986289.
- Crowson CS, Rollefstad S, Ikdahl E, et al. Impact of risk factors associated with cardiovascular outcomes in patients with rheumatoid arthritis. *Ann Rheum Dis* 2018;77:48–54. <https://doi.org/10.1136/annrheumdis-2017-211735>; PMID: 28877868.
- Solomon DH, Reed GW, Kremer JM, et al. Disease activity in rheumatoid arthritis and the risk of cardiovascular events. *Arthritis Rheumatol* 2015;67:1449–55. <https://doi.org/10.1002/art.39098>; PMID: 25776112.
- Myasoedova E, Chandran A, Ilhan B, et al. The role of rheumatoid arthritis (RA) flare and cumulative burden of RA severity in the risk of cardiovascular disease. *Ann Rheum Dis* 2016;75:560–5. <https://doi.org/10.1136/annrheumdis-2014-206411>; PMID: 25637001.
- Gonzalez A, Icen M, Kremers HM, et al. Mortality trends in rheumatoid arthritis: the role of rheumatoid factor. *J Rheumatol* 2008;35:1009–14. PMID: 18412312.
- Ridker PM, Rifai N, Stampfer MJ, et al. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation* 2000;101:1767–72. <https://doi.org/10.1161/01.cir.101.15.1767>; PMID: 10769275.
- Ridker PM, MacFadyen JG, Glynn RJ, et al. Comparison of interleukin-6, C-reactive protein, and low-density lipoprotein cholesterol as biomarkers of residual risk in contemporary practice: secondary analyses from the Cardiovascular Inflammation Reduction Trial. *Eur Heart J* 2020;41:2952–61. <https://doi.org/10.1093/eurheartj/ehaa160>; PMID: 32221587.
- Su D, Li Z, Li X, et al. Association between serum interleukin-6 concentration and mortality in patients with coronary artery disease. *Mediators Inflamm* 2013;2013:726178. <https://doi.org/10.1155/2013/726178>; PMID: 23818744.
- Interleukin-6 Receptor Mendelian Randomisation Analysis (IL6R MR) Consortium. The interleukin-6 receptor as a target for prevention of coronary heart disease: a Mendelian randomisation analysis. *Lancet* 2012;379:1214–24. [https://doi.org/10.1016/S0140-6736\(12\)60110-X](https://doi.org/10.1016/S0140-6736(12)60110-X); PMID: 22421340.
- Georgakis MK, Malik R, Gill D, et al. Interleukin-6 signaling effects on ischemic stroke and other cardiovascular outcomes: a Mendelian randomization study. *Circ Genom Precis Med* 2020;13:e002872. <https://doi.org/10.1161/CIRCGEN.119.002872>; PMID: 32397738.
- Maier W, Altwegg LA, Corti R, et al. Inflammatory markers at the site of ruptured plaque in acute myocardial infarction: locally increased interleukin-6 and serum amyloid A but decreased C-reactive protein. *Circulation* 2005;111:1355–61. <https://doi.org/10.1161/01.CIR.0000158479.58589.0A>; PMID: 15753219.
- Sawa Y, Ichikawa H, Kagisaki K, et al. Interleukin-6 derived from hypoxic myocytes promotes neutrophil-mediated reperfusion injury in myocardium. *J Thorac Cardiovasc Surg* 1998;116:511–7. [https://doi.org/10.1016/s0022-5223\(98\)70018-2](https://doi.org/10.1016/s0022-5223(98)70018-2); PMID: 9731794.
- Duewell P, Kono H, Rayner KJ, et al. NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature* 2010;464:1357–61. <https://doi.org/10.1038/nature08938>; PMID: 20428172.
- Buckley LF, Abbate A. Interleukin-1 blockade in cardiovascular diseases: a clinical update. *Eur Heart J* 2018;39:2063–9. <https://doi.org/10.1093/eurheartj/ehy128>; PMID: 29584915.
- Sandanger O, Ranheim T, Vinge LE, et al. The NLRP3 inflammasome is up-regulated in cardiac fibroblasts and mediates myocardial ischaemia-reperfusion injury. *Cardiovasc Res* 2013;99:164–74. <https://doi.org/10.1093/cvr/cvt091>; PMID: 23580606.
- Kaptoge S, Seshasai SR, Gao P, et al. Inflammatory cytokines and risk of coronary heart disease: new prospective study and updated meta-analysis. *Eur Heart J* 2014;35:578–89. <https://doi.org/10.1093/eurheartj/ehi367>; PMID: 24026779.
- Dixon WG, Symmons DP. What effects might anti-TNF $\alpha$  treatment be expected to have on cardiovascular morbidity and mortality in rheumatoid arthritis? A review of the role of TNF $\alpha$  in cardiovascular pathophysiology. *Ann Rheum Dis* 2007;66:1132–6. <https://doi.org/10.1136/ard.2006.063867>; PMID: 17251223.
- Rho YH, Chung CP, Oeser A, et al. Inflammatory mediators and premature coronary atherosclerosis in rheumatoid arthritis. *Arthritis Rheum* 2009;61:1580–5. <https://doi.org/10.1002/art.25009>; PMID: 19877084.
- Robertson J, Peters MJ, McInnes IB, et al. Changes in lipid levels with inflammation and therapy in RA: a maturing paradigm. *Nat Rev Rheumatol* 2013;9:513–23. <https://doi.org/10.1038/nrrheum.2013.91>; PMID: 23774906.
- Myasoedova E, Crowson CS, Kremers HM, et al. Lipid paradox in rheumatoid arthritis: the impact of serum lipid measures and systemic inflammation on the risk of cardiovascular disease. *Ann Rheum Dis* 2011;70:482–7. <https://doi.org/10.1136/ard.2010.135871>; PMID: 21216812.
- Giles JT, Wasko MCM, Chung CP, et al. Exploring the lipid paradox theory in rheumatoid arthritis: associations of low circulating low-density lipoprotein concentration with subclinical coronary atherosclerosis. *Arthritis Rheumatol* 2019;71:1426–36. <https://doi.org/10.1002/art.40889>; PMID: 30883031.
- Fernandez-Ortiz AM, Ortiz AM, Perez S, et al. Effects of disease activity on lipoprotein levels in patients with early arthritis: can oxidized LDL cholesterol explain the lipid paradox theory? *Arthritis Res Ther* 2020;22:213–8. <https://doi.org/10.1186/s13075-020-02307-8>; PMID: 32917272.
- McMahon M, Grossman J, FitzGerald J, et al. Proinflammatory high-density lipoprotein as a biomarker for atherosclerosis in patients with systemic lupus erythematosus and rheumatoid arthritis. *Arthritis Rheum* 2006;54:2541–9. <https://doi.org/10.1002/art.21976>; PMID: 16868975.
- Yu Z, Kim SC, Vanni K, et al. Association between inflammation and systolic blood pressure in RA compared to patients without RA. *Arthritis Res Ther* 2018;20:107–9. <https://doi.org/10.1186/s13075-018-1597-9>; PMID: 29855349.
- Escalante A, Haas RW, del Rincon I. Paradoxical effect of body mass index on survival in rheumatoid arthritis: role of comorbidity and systemic inflammation. *Arch Intern Med* 2005;165:1624–9. <https://doi.org/10.1001/archinte.165.14.1624>; PMID: 16043681.
- Nicolau J, Lequerre T, Bacquet H, et al. Rheumatoid arthritis, insulin resistance, and diabetes. *Joint Bone Spine* 2017;84:411–6. <https://doi.org/10.1016/j.jbspin.2016.09.001>; PMID: 27777170.
- Alonso-Molero J, Prieto-Pena D, Mendoza G, et al. Misperception of the cardiovascular risk in patients with rheumatoid arthritis. *Int J Environ Res Public Health* 2020;17:5954. <https://doi.org/10.3390/ijerph17165954>; PMID: 32824469.
- Agca R, Heslinga SC, Rollefstad S, et al. EULAR recommendations for cardiovascular disease risk management in patients with rheumatoid arthritis and other forms of inflammatory joint disorders: 2015/2016 update. *Ann Rheum Dis* 2017;76:17–28. <https://doi.org/10.1136/annrheumdis-2016-209775>; PMID: 27697765.
- Weijers JM, Semb, AG, Rollefstad S, et al. Strategies for implementation of guideline recommended cardiovascular risk management for patients with rheumatoid arthritis: results from a questionnaire survey of expert rheumatology centers. *Rheumatol Int* 2020;40:523–7. <https://doi.org/10.1007/s00271-020-00777-7>; PMID: 32824469.

- org/10.1007/s00296-020-04533-4; PMID: 32088752.
45. Semb AG, Rollefstad S, van Riel P, et al. Cardiovascular disease assessment in rheumatoid arthritis: a guide to translating knowledge of cardiovascular risk into clinical practice. *Ann Rheum Dis* 2014;73:1284–8. <https://doi.org/10.1136/annrheumdis-2013-204792>; PMID: 24608403.
  46. Arts EE, Popa C, Den Broeder AA, et al. Performance of four current risk algorithms in predicting cardiovascular events in patients with early rheumatoid arthritis. *Ann Rheum Dis* 2015;74:668–74. <https://doi.org/10.1136/annrheumdis-2013-204024>; PMID: 24389293.
  47. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019;139:e1046–81. <https://doi.org/10.1161/CIR.0000000000000624>; PMID: 30423391.
  48. Crowson CS, Rollefstad S, Kitas GD, et al. Challenges of developing a cardiovascular risk calculator for patients with rheumatoid arthritis. *PLoS One* 2017;12:e0174656. <https://doi.org/10.1371/journal.pone.0174656>; PMID: 28334012.
  49. Solomon DH, Greenberg J, Curtis JR, et al. Derivation and internal validation of an expanded cardiovascular risk prediction score for rheumatoid arthritis: a Consortium of Rheumatology Researchers of North America registry study. *Arthritis Rheumatol* 2015;67:1995–2003. <https://doi.org/10.1002/art.39195>; PMID: 25989470.
  50. Crowson CS, Gabriel SE, Semb AG, et al. Rheumatoid arthritis-specific cardiovascular risk scores are not superior to general risk scores: a validation analysis of patients from seven countries. *Rheumatology (Oxford)* 2017;56:1102–10. <https://doi.org/10.1093/rheumatology/kex038>; PMID: 28339992.
  51. Jamthikar AD, Gupta D, Puuvula A, et al. Cardiovascular risk assessment in patients with rheumatoid arthritis using carotid ultrasound B-mode imaging. *Rheumatol Int* 2020;40:1921–39. <https://doi.org/10.1007/s00296-020-04691-5>; PMID: 32857281.
  52. Semb AG, Ikdhall E, Hisdal J, et al. Exploring cardiovascular disease risk evaluation in patients with inflammatory joint diseases. *Int J Cardiol* 2016;223:331–6. <https://doi.org/10.1016/j.ijcard.2016.08.129>; PMID: 27543704.
  53. Kitas GD, Nightingale P, Armitage J, et al. A multicenter, randomized, placebo-controlled trial of atorvastatin for the primary prevention of cardiovascular events in patients with rheumatoid arthritis. *Arthritis Rheumatol* 2019;71:1437–49. <https://doi.org/10.1002/art.40892>; PMID: 30983166.
  54. Pedersen TR, Kjekshus J, Berg K, et al. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994;344:1383–9. [https://doi.org/10.1016/S0140-6736\(94\)90566-5](https://doi.org/10.1016/S0140-6736(94)90566-5); PMID: 7968073.
  55. Semb AG, Kvien TK, DeMicco DA, et al. Effect of intensive lipid-lowering therapy on cardiovascular outcome in patients with and those without inflammatory joint disease. *Arthritis Rheum* 2012;64:2836–46. <https://doi.org/10.1002/art.34524>; PMID: 22576673.
  56. Lv S, Liu Y, Zou Z, et al. The impact of statins therapy on disease activity and inflammatory factor in patients with rheumatoid arthritis: a meta-analysis. *Clin Exp Rheumatol* 2015;33:69–76. PMID: 25327393.
  57. Roelsgaard IK, Ikdhall E, Rollefstad S, et al. Smoking cessation is associated with lower disease activity and predicts cardiovascular risk reduction in rheumatoid arthritis patients. *Rheumatology (Oxford)* 2020;59:1997–2004. <https://doi.org/10.1093/rheumatology/kez557>; PMID: 31782789.
  58. Fagerhøj MG, Rollefstad S, Olsen SJ, Semb AG. The effect of brief versus individually tailored dietary advice on change in diet, lipids and blood pressure in patients with inflammatory joint disease. *Food Nutr Res* 2018;62. <https://doi.org/10.29219/fnr.v62.1512>; PMID: 30202399.
  59. Veldhuijzen van Zanten JJ, Rouse PC, Hale ED, et al. Perceived barriers, facilitators and benefits for regular physical activity and exercise in patients with rheumatoid arthritis: a review of the literature. *Sports Med* 2015;45:1401–12. <https://doi.org/10.1007/s40279-015-0363-2>; PMID: 262619268.
  60. McInnes IB, Thompson L, Giles JT, et al. Effect of interleukin-6 receptor blockade on surrogates of vascular risk in rheumatoid arthritis: MEASURE, a randomised, placebo-controlled study. *Ann Rheum Dis* 2015;74:694–702. <https://doi.org/10.1136/annrheumdis-2013-204345>; PMID: 24368514.
  61. Del Porto F, Lagana B, Lai S, et al. Response to anti-tumour necrosis factor alpha blockade is associated with reduction of carotid intima-media thickness in patients with active rheumatoid arthritis. *Rheumatology (Oxford)* 2007;46:1111–5. <https://doi.org/10.1093/rheumatology/kem089>; PMID: 17449484.
  62. Li HZ, Xu XH, Lin N, Lu HD. Metabolic and cardiovascular benefits of hydroxychloroquine in patients with rheumatoid arthritis: a systematic review and meta-analysis. *Ann Rheum Dis* 2019;78:e21. <https://doi.org/10.1136/annrheumdis-2018-213157>; PMID: 29453218.
  63. Hurlimann D, Forster A, Noll G, et al. Anti-tumor necrosis factor-alpha treatment improves endothelial function in patients with rheumatoid arthritis. *Circulation* 2002;106:2184–7. <https://doi.org/10.1161/01.cir.0000037521.71373.44>; PMID: 12390945.
  64. Chan ES, Cronstein BN. Methotrexate – how does it really work? *Nat Rev Rheumatol* 2010;6:175–8. <https://doi.org/10.1038/nrrheum.2010.5>; PMID: 20197777.
  65. Bulgarelli A, Martins Dias AA, Caramelli B, et al. Treatment with methotrexate inhibits atherogenesis in cholesterol-fed rabbits. *J Cardiovasc Pharmacol* 2012;59:308–14. <https://doi.org/10.1097/FJC.0b013e318241c385>; PMID: 22113347.
  66. Choi HK, Hernan MA, Seeger JD, et al. Methotrexate and mortality in patients with rheumatoid arthritis: a prospective study. *Lancet* 2002;359:1173–7. [https://doi.org/10.1016/S0140-6736\(02\)08213-2](https://doi.org/10.1016/S0140-6736(02)08213-2); PMID: 11955534.
  67. Roubille C, Richer V, Starnino T, et al. The effects of tumour necrosis factor inhibitors, methotrexate, non-steroidal anti-inflammatory drugs and corticosteroids on cardiovascular events in rheumatoid arthritis, psoriasis and psoriatic arthritis: a systematic review and meta-analysis. *Ann Rheum Dis* 2015;74:480–9. <https://doi.org/10.1136/annrheumdis-2014-206624>; PMID: 25561362.
  68. Xie F, Chen L, Yun H, et al. Benefits of methotrexate use on cardiovascular disease risk among rheumatoid arthritis patients initiating biologic disease-modifying antirheumatic drugs. *J Rheumatol* 2020. <https://doi.org/10.3899/jrheum.191326>; PMID: 33060309; epub ahead of press.
  69. Ridker PM, Everett BM, Pradhan A, et al. Low-dose methotrexate for the prevention of atherosclerotic events. *N Engl J Med* 2019;380:752–62. <https://doi.org/10.1056/NEJMoa1809798>; PMID: 30415610.
  70. Rempenault C, Combe B, Barnette T, et al. Metabolic and cardiovascular benefits of hydroxychloroquine in patients with rheumatoid arthritis: a systematic review and meta-analysis. *Ann Rheum Dis* 2018;77:98–103. <https://doi.org/10.1136/annrheumdis-2017-211836>; PMID: 28970215.
  71. Naranjo A, Sokka T, Descalzo MA, et al. Cardiovascular disease in patients with rheumatoid arthritis: results from the QUEST-RA study. *Arthritis Res Ther* 2008;10:R30. <https://doi.org/10.1186/ar2383>; PMID: 18325087.
  72. Rozman B, Praprotnik S, Logar D, et al. Leflunomide and hypertension. *Ann Rheum Dis* 2002;61:567–9. <https://doi.org/10.1136/ard.61.6.567>; PMID: 12006342.
  73. Ljung L, Rantapaa-Dahlqvist S, Jacobsson LT, et al. Response to biological treatment and subsequent risk of coronary events in rheumatoid arthritis. *Ann Rheum Dis* 2016;75:2087–94. <https://doi.org/10.1136/annrheumdis-2015-208995>; PMID: 26984007.
  74. Arida A, Proterogerou AD, Konstantonis G, et al. Atherosclerosis is not accelerated in rheumatoid arthritis of low activity or remission, regardless of antirheumatic treatment modalities. *Rheumatology (Oxford)* 2017;56:934–9. <https://doi.org/10.1093/rheumatology/kew506>; PMID: 28160488.
  75. Mann DL, McMurray JJ, Packer M, et al. Targeted anticytokine therapy in patients with chronic heart failure: results of the Randomized Etorcept Worldwide Evaluation (RENEWAL). *Circulation* 2004;109:1594–602. <https://doi.org/10.1161/01.CIR.0000124490.27666.B2>; PMID: 15023878.
  76. Chung ES, Packer M, Lo KH, et al. Randomized, double-blind, placebo-controlled, pilot trial of infliximab, a chimeric monoclonal antibody to tumor necrosis factor- $\alpha$ , in patients with moderate-to-severe heart failure: results of the anti-TNF Therapy Against Congestive Heart Failure (ATTACH) trial. *Circulation* 2003;107:3133–40. <https://doi.org/10.1161/01.CIR.0000077913.60364.D2>; PMID: 12796126.
  77. Giles JT, Sattar N, Gabriel S, et al. Cardiovascular safety of tocilizumab versus etanercept in rheumatoid arthritis: a randomized controlled trial. *Arthritis Rheumatol* 2020;72:31–40. <https://doi.org/10.1002/art.41095>; PMID: 31469238.
  78. Singh S, Fumery M, Singh AG, et al. Comparative risk of cardiovascular events with biologic and synthetic disease-modifying antirheumatic drugs in patients with rheumatoid arthritis: a systematic review and meta-analysis. *Arthritis Care Res (Hoboken)* 2020;72:561–76. <https://doi.org/10.1002/acr.23875>; PMID: 30875456.
  79. Cacciapaglia F, Anelli MG, Rinaldi A, et al. Lipids and atherogenic indices fluctuation in rheumatoid arthritis patients on long-term tocilizumab treatment. *Mediators Inflamm* 2018;2018:2453265. <https://doi.org/10.1155/2018/2453265>; PMID: 30405318.
  80. Soubrier M, Pei J, Durand F, et al. Concomitant use of statins in tocilizumab-treated patients with rheumatoid arthritis: a post hoc analysis. *Rheumatol Ther* 2017;4:133–49. <https://doi.org/10.1007/s40744-016-0049-8>; PMID: 27900570.
  81. Kleaveland O, Kunszt G, Brattlie M, et al. Effect of a single dose of the interleukin-6 receptor antagonist tocilizumab on inflammation and troponin T release in patients with non-ST-elevation myocardial infarction: a double-blind, randomized, placebo-controlled phase 2 trial. *Eur Heart J* 2016;37:2406–13. <https://doi.org/10.1093/eurheartj/ehw171>; PMID: 27161611.
  82. Ruscitti P, Masedu F, Alvaro S, et al. Anti-interleukin-1 treatment in patients with rheumatoid arthritis and type 2 diabetes (TRACK): a multicentre, open-label, randomised controlled trial. *PLoS Med* 2019;16:e1002901. <https://doi.org/10.1371/journal.pmed.1002901>; PMID: 31513665.
  83. Ikonomidis I, Pavlidis G, Katsimbri P, et al. Differential effects of inhibition of interleukin 1 and 6 on myocardial, coronary and vascular function. *Clin Res Cardiol* 2019;108:1093–101. <https://doi.org/10.1007/s00392-019-01443-9>; PMID: 30859382.
  84. Morton AC, Rothman AM, Greenwood JP, et al. The effect of interleukin-1 receptor antagonist therapy on markers of inflammation in non-ST elevation acute coronary syndromes: the MRC-ILA Heart Study. *Eur Heart J* 2015;36:377–84. <https://doi.org/10.1093/eurheartj/ehu272>; PMID: 25079365.
  85. Abbate A, Van Tassel BW, Biondi-Zoccai G, et al. Effects of interleukin-1 blockade with anakinra on adverse cardiac remodeling and heart failure after acute myocardial infarction [from the Virginia Commonwealth University-Anakinra Remodeling Trial (2) (VCU-ART2) pilot study]. *Am J Cardiol* 2013;111:1394–400. <https://doi.org/10.1016/j.amjcard.2013.01.287>; PMID: 23453459.
  86. Ridker PM, Everett BM, Thuren T, et al. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med* 2017;377:1119–31. <https://doi.org/10.1056/NEJMoa1707914>; PMID: 28845751.
  87. Ozen G, Pedro S, Michaud K. The risk of cardiovascular events associated with disease-modifying antirheumatic drugs in rheumatoid arthritis. *J Rheumatol* 2020. <https://doi.org/10.3899/jrheum.200265>; PMID: 32801134; epub ahead of press.
  88. Sandborn WJ, Panes J, Sands BE, et al. Venous thromboembolic events in the tofacitinib ulcerative colitis clinical development programme. *Aliment Pharmacol Ther* 2019;50:1068–76. <https://doi.org/10.1111/apt.15514>; PMID: 31599001.
  89. Xie W, Huang Y, Xiao S, et al. Impact of Janus kinase inhibitors on risk of cardiovascular events in patients with rheumatoid arthritis: systematic review and meta-analysis of randomised controlled trials. *Ann Rheum Dis* 2019;78:1048–54. <https://doi.org/10.1136/annrheumdis-2018-214846>; PMID: 31088790.
  90. Taylor PC, Kremer JM, Emery P, et al. Lipid profile and effect of statin treatment in pooled phase II and phase III baricitinib studies. *Ann Rheum Dis* 2018;77:988–95. <https://doi.org/10.1136/annrheumdis-2017-212461>; PMID: 29463520.
  91. Lindhardtsen J, Gislason GH, Jacobsen S, et al. Non-steroidal anti-inflammatory drugs and risk of cardiovascular disease in patients with rheumatoid arthritis: a nationwide cohort study. *Ann Rheum Dis* 2014;73:1515–21. <https://doi.org/10.1136/annrheumdis-2012-203137>; PMID: 23749610.
  92. Avina-Zubieta JA, Abrahamowicz M, De Vera MA, et al. Immediate and past cumulative effects of oral glucocorticoids on the risk of acute myocardial infarction in rheumatoid arthritis: a population-based study. *Rheumatology (Oxford)* 2013;52:68–75. <https://doi.org/10.1093/rheumatology/kes353>; PMID: 23192907.
  93. Demidowich AP, Davis AI, Dedhia N, et al. Colchicine to decrease NLRP3-activated inflammation and improve obesity-related metabolic dysregulation. *Med Hypotheses* 2016;92:67–73. <https://doi.org/10.1016/j.mehy.2016.04.039>; PMID: 27241260.
  94. Tardif JC, Kouz S, Waters DD, et al. Efficacy and safety of low-dose colchicine after myocardial infarction. *N Engl J Med* 2019;381:2497–505. <https://doi.org/10.1056/NEJMoa1921388>; PMID: 31733140.
  95. Nidorf SM, Fiolet ATL, Mosterd A, et al. Colchicine in patients with chronic coronary disease. *N Engl J Med* 2020;383:1838–47. <https://doi.org/10.1056/NEJMoa2021372>; PMID: 32865380.
  96. Choy E, Ganeshalingam K, Semb AG, et al. Cardiovascular risk in rheumatoid arthritis: recent advances in the understanding of the pivotal role of inflammation, risk predictors and the impact of treatment. *Rheumatology (Oxford)* 2014;53:2143–54. <https://doi.org/10.1093/rheumatology/keu224>; PMID: 24907149.