

Forbidden links, trait matching and modularity in plant-hummingbird networks: Are specialized modules characterized by higher phenotypic floral integration?

Jaume Izquierdo-Palma¹, Maria del Coro Arizmendi¹, Carlos Lara² and Juan Francisco Ornelas³

¹Laboratorio de Ecología, UBIPRO, Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México, Tlalnepantla de Baz, Estado de México, Mexico

²Centro de Investigación en Ciencias Biológicas, Universidad Autónoma de Tlaxcala, San Felipe Ixtacuixtla, Tlaxcala, Mexico

³Departamento de Biología Evolutiva, Instituto de Ecología A.C., Xalapa, Veracruz, Mexico

ABSTRACT

Background: Plant-pollinator mutualistic networks show non-random structural properties that promote species coexistence. However, these networks show high variability in the interacting species and their connections. Mismatch between plant and pollinator attributes can prevent interactions, while trait matching can enable exclusive access, promoting pollinators' niche partitioning and, ultimately, modularity. Thus, plants belonging to specialized modules should integrate their floral traits to optimize the pollination function. Herein, we aimed to analyze the biological processes involved in the structuring of plant-hummingbird networks by linking network morphological constraints, specialization, modularity and phenotypic floral integration.

Methods: We investigated the understory plant-hummingbird network of two adjacent habitats in the Lacandona rainforest of Mexico, one characterized by lowland rainforest and the other by savanna-like vegetation. We performed monthly censuses to record plant-hummingbird interactions for 2 years (2018–2020). We also took hummingbird bill measurements and floral and nectar measurements. We summarized the interactions in a bipartite matrix and estimated three network descriptors: connectance, complementary specialization (H_2'), and nestedness. We also analyzed the modularity and average phenotypic floral integration index of each module.

Results: Both habitats showed strong differences in the plant assemblage and network dynamics but were interconnected by the same four hummingbird species, two Hermits and two Emeralds, forming a single network of interaction. The whole network showed low levels of connectance (0.35) and high specialization ($H_2' = 0.87$). Flower morphologies ranged from generalized to specialized, but trait matching was an important network structurer. Modularity was associated with morphological specialization. The Hermits *Phaethornis longirostris* and *P. striigularis* each formed a module by themselves, and a third module was formed by the less-specialized Emeralds: *Chlorestes candida* and *Amazilia tzacatl*. The floral integration values were

Submitted 15 July 2020
Accepted 29 January 2021
Published 10 March 2021

Corresponding author
Maria del Coro Arizmendi,
coro@unam.mx

Academic editor
Stuart Pimm

Additional Information and
Declarations can be found on
page 20

DOI 10.7717/peerj.10974

© Copyright
2021 Izquierdo-Palma et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

higher in specialized modules but not significantly higher than that formed by generalist species.

Conclusions: Our findings suggest that biological processes derived from both trait matching and “forbidden” links, or nonmatched morphological attributes, might be important network drivers in tropical plant-hummingbird systems while morphological specialization plays a minor role in the phenotypic floral integration. The broad variety of corolla and bill shapes promoted niche partitioning, resulting in the modular organization of the assemblage according to morphological specialization. However, more research adding larger datasets of both the number of modules and pollination networks for a wider region is needed to conclude whether phenotypic floral integration increases with morphological specialization in plant-hummingbird systems.

Subjects Animal Behavior, Biodiversity, Ecology, Plant Science, Zoology

Keywords Interaction networks, Hummingbirds, Biotic interactions, Pollination, Mexico, Forbidden links, Biotic specialization, Modularity, Phenotypic floral integration, Neotropical rainforest

INTRODUCTION

Plant and animal species are integrated in complex networks of interdependencies forming ecological communities playing an important role in the generation of Earth’s biodiversity (*Bascompte & Jordano, 2007*). Plant-pollinator mutualistic networks are linked by trophic interactions where plants act as primary producers and animals as a special subset of primary consumers that feed on nectar and/or pollen (*Bascompte & Jordano, 2007; Waser et al., 1996*). The strength of their interactions varies, resulting in large, complex networks in which interacting species impose reciprocal selective pressures as they interrelate over ecological and evolutionary time (*Bascompte & Jordano, 2007; Thomson & Wilson, 2008; Waser et al., 1996*). Over the last decade, the study of mutualistic networks has changed radically, along with the theory (*Bascompte & Jordano, 2007; Blüthgen et al., 2007; Olesen et al., 2007; Ings et al., 2009; Vázquez et al., 2009; Dalsgaard et al., 2011*), and new powerful analytical tools have been proposed (*Dormann, Gruber & Fründ, 2008; Dormann & Strauss, 2014*). These advances have shown that non-random structural properties can be characterized by certain network metrics. For example, most interaction networks show a nested structure (i.e., specialists interact with subsets of species with which generalists also interact) and varying levels of connectivity among species. Both properties facilitate species coexistence by minimizing competition relative to facilitation, supporting greater biodiversity (*Bascompte et al., 2003; Olesen et al., 2006; Verdú & Valiente-Banuet, 2008; Bastolla et al., 2009; Sugihara & Ye, 2009*). Furthermore, most ecological networks are strongly asymmetric (i.e., a plant species might heavily depend on a pollen-vector species that, in turn, is only weakly dependent on that plant species). Thus, the community is structured around a central core of generalists, offering robustness and resilience to the random loss of species (*Vázquez & Aizen, 2003*;

Bascompte, Jordano & Olesen, 2006; Guimarães et al., 2006; Blüthgen et al., 2007; Bascompte & Jordano, 2013).

Although the discussion on the architecture of mutualistic networks is quite settled, the underlying structuring mechanisms are still being debated (*Maruyama et al., 2014; Vizentin-Bugoni, Maruyama & Sazima, 2014; Araujo et al., 2018*). Mutualistic networks tend to be very heterogeneous in the number of interacting species at each level and the distribution of their connections, such as, for example, between animal-dispersed fruits and their dispersers or animal-pollinated angiosperms and their pollinators (*Bascompte & Jordano, 2007*). Two major biological mechanisms that influence the structure of networks are the “complementary traits” and “barrier traits” (*Santamaría & Rodríguez-Gironés, 2007*). For complementary traits, the interaction is determined by the similarity between the reward that the plant has to offer and the resource that the pollinator seeks. This mechanism can progressively generate co-specialization. For example, flowers that are mainly pollinated by birds are red in color, matching the perceptual system of their avian pollinators who show a preference for this color when looking for nectar resources (*Niovi Jones & Reithel, 2001*). For the mechanism of barrier traits, the interaction is related to the ability of the pollinator to reach the reward offered by the flower. Only those pollinators whose traits allow them to overcome the floral barriers are able to access the reward. For example, Hermit hummingbirds with typically long and curved bills have access to flowers with long and curved corolla tubes that hummingbird species with short and straight bills cannot access (*Maglianesi et al., 2014*). Consequently, competition for shared floral resources is reduced (*Feinsinger, 1976; Ings et al., 2009; McGuire et al., 2009; Abrahamczyk & Kessler, 2010; Maglianesi et al., 2014*). These exploitation barriers imposed by the mismatch of biological attributes that lead to a decrease in the connectivity and/or strength of the interactions in ecological networks can be called “forbidden links” (*Santamaría & Rodríguez-Gironés, 2007; Olesen et al., 2011*).

Species abundances can be as important or even more important as species traits in structuring the ecological interaction networks of local communities (e.g., common in insect-pollination networks; *Vázquez, Chacoff & Cagnolo, 2009*). However, plant-hummingbird mutualistic networks are considered a specialized system, particularly those located near the Equator due to higher productivity and the relatively stable and predictable availability of resources throughout the year (*Feinsinger & Colwell, 1978; Dalsgaard et al., 2011; Belmaker, Sekercioglu & Jetz, 2012; Zanata et al., 2017*). Hence, several studies have shown that mismatches in species morphology and phenology play a major role in structuring interactions in plant-hummingbird systems (*Stiles, 1975, 1978; Lara, 2006; Maglianesi et al., 2014; Maruyama et al., 2014; Vizentin-Bugoni, Maruyama & Sazima, 2014; Sonne et al., 2019*). Hummingbird clades have characteristic morphologies that influence resource use, flight capabilities, competitive skills and environmental filtering, important mechanisms structuring hummingbird communities (*Rodríguez-Flores et al., 2019*). Furthermore, hummingbird morphological traits, such as bill length and curvature and body mass, have also been hypothesized to play a role in the specialization of hummingbird interactions. Meanwhile, corolla length and curvature

and nectar volume are floral traits associated with the specialization of hummingbird-pollinated plants (Maruyama *et al.*, 2014, 2018; Maglianesi *et al.*, 2014, 2015; Dehling *et al.*, 2016). Thus, specialization in trophic resources' use by hummingbird assemblages, which vary in its type and/or its strength, can arise in niche segregation of the community due to the differential access to flowers (Feinsinger & Colwell, 1978).

The extent to which connectivity is limited by both morphological mismatches and spatio-temporal constraints often results in resource segregation, leading to niche partitioning and, consequently, a modular structure in ecological networks. Network modularity reflects the tendency of a set of species to interact predominantly with species within the set and less frequently with species in other sets. Modularity implies that species can be grouped (i.e., modules) in such a way that weakly interlinked subsets of species are strongly connected internally (Olesen *et al.*, 2007). Modules can provide information on the dynamics of ecological communities by identifying specialized functional groups of pollinators and floral traits (Newman, 2006; Olesen *et al.*, 2007; Danieli-Silva *et al.*, 2012; Dormann & Strauss, 2014; Maruyama *et al.*, 2014). Some studies support the idea that modularity increases with the increased specialization of the interacting species (Lewinsohn *et al.*, 2006; Guimarães *et al.*, 2006) and is positively related with network size (Olesen *et al.*, 2007). Thus, in little-connected and highly nested networks, which are common in the tropics (Olesen & Jordano, 2002), there is a higher probability of modularity (Fortuna *et al.*, 2010).

Specialized modules drive strong plant-pollinator relationships, leading to the diffuse co-evolution of complementary morphological traits (Sazatornil *et al.*, 2016). Furthermore, selection imposed by pollinators can be an important mechanism shaping the patterns of variation and covariation of floral traits, particularly in plant species with specialized pollination. Such is the case in the *Phaethornis-Heliconia* pollination system (Armbruster, 1991; Fenster, 1991; Herrera *et al.*, 2002; Pérez, Arroyo & Medel, 2007; Ordano *et al.*, 2008). In this context, flowers can be seen as suites or units of traits that require a precise configuration and arrangement of their sexual organs for proper pollination (Bissell & Diggle, 2008). From this multi-trait view, some authors have argued that selection imposed by specialized pollinators reduces phenotypic variability and favors the integration of subsets of floral traits (*Pleiades*), that is, correlations among traits within functional units usually involved in one ecological function (Berg, 1960; Stebbins, 1970; Conner & Via, 1993; Rosas-Guerrero *et al.*, 2011). However, other factors may be linked with phenotypic floral integration, such as the breeding system (Anderson & Busch, 2006; Rosas-Guerrero *et al.*, 2011) and developmental-genetic factors (Conner, 2002; Smith & Rausher, 2008). Nonetheless, the relationship between specialization and niche partitioning in hummingbird assemblages and its possible effects on the phenotypic floral integration of interacting plant species are poorly understood (Berg, 1960; Stebbins, 1970; Conner & Via, 1993; Rosas-Guerrero *et al.*, 2011).

In this study, we assessed the association among biological processes involved in the structuring of plant-hummingbird networks, niche segregation by ecological specialization, and phenotypic floral integration of the plant assemblages. We used the phenotypic floral integration index as a measurable estimate of the magnitude and pattern

of covariation among sets of functionally related floral traits to obtain new insights into the link between modularity and specialization in plant-hummingbird mutualistic networks (Ordano *et al.*, 2008; Rosas-Guerrero *et al.*, 2011; Dormann & Strauss, 2014). To achieve this, we investigated the network architecture by descriptors commonly used in similar studies and identified the possible underlying biological processes, that is, trait matching and forbidden links. Then, we analyzed the modularity and floral integration index of each module to identify patterns of covariation. In particular, we examined the understory plant-hummingbird mutualistic networks in two distinct adjacent habitats in Mexico, addressing the following questions: (1) Are there differences between the two habitats in the composition of interacting species and their network metrics? (2) What is the main network structuring each habitat? And, (3) is there a relationship between module specialization and phenotypic floral integration? To our knowledge, the linking of network morphological constraints, specialization, modularity and phenotypic floral integration is a new approach in the study of plant-hummingbird interaction networks.

MATERIALS & METHODS

Study area

Fieldwork was carried out at the Chajul Biological Station located in the Montes Azules Biosphere Reserve (16°06' N; 90°56' E) within the Lacandon region in southern Mexico a few kilometers from the Guatemalan border. The study area covers an extension of ~331,200 ha and is situated from 150 to 1,500 m above sea level (m.a.s.l.). Mean rainfall in Chajul is around 3,000 mm, of which ca. 70% is concentrated from June to October. The short dry season occurs between February and April. The mean annual temperature is 22.5 °C (Siebe *et al.*, 1996; Carabias, De la Maza & Cadena, 2015). The dominant vegetation around the field station is lowland evergreen tropical rainforest (hereafter “rainforest” habitat) with some variability influenced by the soil properties and proximity to water bodies. Near streams and rivers, there is riparian vegetation and sections of flooded plains. In areas surrounding the field station where the anthropic impact has been more intense in recent times, the vegetation mainly consists of secondary forest and abandoned fields in different stages of ecological succession. There is also some hilly terrain (highest elevation: 230 m.a.s.l) with thin and poor soils, and the vegetation here is savanna-like with low and scattered trees and an understory characterized by abundant grass (*Scleria melaleuca*; hereafter “savanna” habitat) (Miranda & Hernández, 1963; Rzedowski & Huerta, 1994; Siebe *et al.*, 1996). We collected data from January 2018 to January 2020 along trails 6,700 m long in both study habitats.

Phenology of hummingbirds and their plants

At monthly intervals, we recorded the hummingbird species and their numbers along six study trails in both habitats. Walking censuses began around 7:00 AM and ended at 1:30 PM. All hummingbird-pollinated plant species (individuals or floral patches) flowering within 2.5 m on each side of the trails were counted at a maximum height of 5 m. We focused on the understory plant community for logistical reasons. Flowering plants in the canopy are difficult to see from the ground, which may result in an underestimation of

the number of plant individuals and interactions with their pollinators. Binoculars (Nikon 10 × 42) and a field guide were used to identify hummingbirds (Arizmendi & Berlanga, 2014), and plant specimens were identified at the Chajul Biological Station herbarium. The scientific names of plants were validated using “The Plant List” online database (<http://www.theplantlist.org/>) and hummingbird scientific names using the IOC World Bird List (www.worldbirdnames.org).

Floral measures and bill morphology

Because hummingbirds commonly visited the flowers of the plant species we observed during the censuses, we quantified several floral traits presumed to be associated with hummingbird pollinator attraction and pollen transfer efficiency (Wolf, Stiles & Hainsworth, 1976; Stiles, 1995). The flower morphology was characterized by measuring the corolla length and curvature, as these are the primary constraints determining the ability of hummingbirds to reach the nectar. We measured the effective corolla length (i.e., distance from the nectary to the distal portion of the flower, which determines how far the bill of the feeding bird fits into the flower) (Wolf, Stiles & Hainsworth, 1976). In addition, we calculated the flower curvature following similar methodology used in hummingbird bills (Stiles, 1995; Rico-Guevara & Araya-Salas, 2015). We calculated a corolla curvature index as the arc:chord ratio of the corolla. Arc length was measured following the dorsal profile of the corolla from the calyx to the corolla tip, and the chord was measured as the straight-line distance from the calyx to the corolla tip. These measures were taken from lateral photographs using the ImageJ software (Schneider, Rasband & Eliceiri, 2012). Because the placement of pollen on the vector and its subsequent reception on the stigma is crucial to plant fitness, we measured the average stamen length and style length. For each plant species recorded in both habitats, we measured from 2 to 180 flowers collected from at least two individuals. Accumulated nectar was also quantified to determine reward availability for hummingbirds. From each plant species, 2–30 buds about to open were selected and placed in mesh bags (1-mm bridal tulle) to exclude hummingbird visitors and allow nectar to accumulate. After the flowers opened, the accumulated nectar was extracted, and the nectar was removed and measured after 24 h of nectar accumulation using calibrated micropipettes (5 µL) and a digital caliper (error: 0.1 mm). The sugar concentration (percentage sucrose) was measured by a hand-held pocket refractometer (range concentration 0–32° Brix units (°Bx); Atago, Tokyo, Japan). To characterize the main floral types of the whole plant assemblage, we performed a principal components analysis (PCA) of the measured floral traits (morphology and nectar) after discarding highly correlated variables through a correlation analysis.

The hummingbirds’ bill morphology was measured as the length of the exposed culmen and its curvature from voucher specimens housed at the collection of the Museo de Zoología, Facultad de Ciencias, Universidad Nacional Autónoma de México (MZFC, UNAM) (*Phaethornis longirostris* ($n = 30$), *P. striigularis* ($n = 20$), *Amazilia tzacatl* ($n = 30$), and *Chlorestes candida* ($n = 28$)). As hummingbirds can project their tongues to drink nectar (Paton & Collins, 1989), bill measurements that ignore tongue extension can underestimate birds’ capacity to access nectar. Because precise measurements of tongue

length are unavailable for different hummingbird species, we added one-third to the bill length for each species (*Vizentin-Bugoni, Maruyama & Sazima, 2014*). To examine differences in bill shape, we calculated a bill curvature index as the arc:chord ratio of the exposed culmen (maxillary curvature) (*Stiles, 1995; Rico-Guevara & Araya-Salas, 2015*). Arc length was measured following the dorsal profile of the bill from the feathered base to the tip, and the chord was measured as the straight-line distance from the feathered base to the tip (*Phaethornis longirostris* ($n = 18$), *P. striigularis* ($n = 14$), *Amazilia tzacatl* ($n = 13$), and *Chlorestes candida* ($n = 20$)). These measures were taken from lateral photographs of the complete bill using the ImageJ software (*Schneider, Rasband & Eliceiri, 2012*). Furthermore, we obtained the average measures of total body weight to relate them with agonistic behavior and nutritional requirements (*Phaethornis longirostris* ($n = 15$), *P. striigularis* ($n = 11$), *Amazilia tzacatl* ($n = 10$) and *Chlorestes candida* ($n = 12$)). To obtain a closer relationship between the trait match of plant species and their hummingbird visitors, we calculated the average of each floral trait of the flowering plants visited by the hummingbird species. Then, comparisons were made with the average bill length and curvature.

Plant-hummingbird interactions

We built the plant-hummingbird mutualistic networks from a plant-centered approach (*Jordano, 1987; Bosch et al., 2009*). We did not record other pollinator interactions despite have been reported that many plants visited by hummingbirds are also visited by insects (*Dalsgaard et al., 2009*). Nevertheless, this fact should not affect to test our hypotheses due to the high specialization of our study system. Plant-insect pollination studies require another methodology with closer observations and adapted to the daily foraging activity of each invertebrate species. Because our study was based on mutualistic relationships, we only considered hummingbirds as potential pollinators. We recorded legitimate hummingbird visits, that is, when hummingbirds contacted the reproductive structures of the flowers. Each visit was defined as the moment a hummingbird probed one flower until it left the flowering plant/patch. We conducted from 8 to 50 h focal observations of each plant species (*Vizentin-Bugoni et al., 2016*). Most observations were conducted by video recording (GoPro Hero5), but in some cases (i.e., large floral patches or epiphytes with difficult access), we used binoculars (Nikon 10 × 42) to prevent underestimating the interactions. The observations were conducted from 07:00 AM to 11:30 AM, the time period of maximum foraging activity based on preliminary observations. Whenever possible, we conducted the observations at different plant individuals and locations to capture maximum variability. We observed a total of 657 h of plant-hummingbird interactions.

Analysis of interaction networks

We summarized the plant-hummingbird interactions in a bipartite matrix with each cell indicating the frequency of interactions. Because the two habitats are adjacent, they can form a single network. For this reason, we built a single interaction network for both habitats. However, to have a glimpse of the possible underlying biological processes

modeling interactions in each habitat, we also built two separate interaction networks corresponding with the rainforest and savanna communities. Using these matrices, we estimated several network metrics of structure and specialization, which are detailed at following: (1) Connectance was calculated as the proportion of possible links in the network that are actually realized. If nonmatching species traits can prevent the occurrence of certain interactions (forbidden links), connectance is an estimate of how interactions are constrained in the communities. (2) Complementary specialization (H_2') estimates the exclusiveness of interactions considering the ecological specialization of a species (i.e., how connected a species is) and how these interactions differ among species. The H_2' index is useful for comparing ecological networks, as it is less affected by community size or sampling intensity (Blüthgen et al., 2007). (3) Nestedness was calculated using the ANINHADO software (Guimarães & Guimarães, 2006). We used two estimators, the NODF index, which uses qualitative presence/absence data and wNODF, which considers quantitative interaction data (Almeida-Neto et al., 2008; Almeida-Neto & Ulrich, 2011). (4) Modularity (Q), as defined above, was estimated for both quantitative and qualitative matrices. For the quantitative matrices, we used the QuaBiMo optimization algorithm (Dormann & Strauss, 2014). As the QuaBiMo algorithm has an iterative searching algorithm (values can slightly differ between runs), we chose the highest values from 10 independent runs. The modularity of the qualitative matrix was estimated in MODULAR (Marquitti et al., 2014), a stochastic algorithm, using Barber's metric for bipartite networks (Barber, 2007) and following the recommended program settings (Marquitti et al., 2014; Appendix 3). We estimated the significance of each run against 100 null matrices obtained with two null models: the Erdős-Rényi (ER) model (Marquitti et al., 2014) and one proposed by Bascompte et al. (2003). We also ran a modularity analysis considering both habitats together. If the habitats functioned as separated units, separate modules corresponding with each community would be generated.

To evaluate the statistical significance of the estimated network metrics, we compared the observed values to 1,000 random values calculated from the null matrices. These matrices were generated using a randomization algorithm that conserves the total number of interactions per row and column in the matrix (Patefield's r2dtable algorithm). Such a null model is not prone to type I errors (Dormann, Gruber & Fründ, 2008). The network indices (connectance, H_2' , NODF, wNODF and Q) were expressed as z-scores (observed – mean(null)/sd(null)), and the statistical significance was assessed by Z-tests. The interaction networks and networks metrics were built and estimated using the *bipartite* package version 2.11 (Dormann, Gruber & Fründ, 2008) in R software (R Development Core Team, 2018).

Analysis of phenotypic floral integration

To obtain a measurable estimate of the magnitude (i.e., degree to which the traits are tied) and pattern (i.e., arrangement of the relationships among traits) of covariation among sets of functionally related floral traits, we estimated the phenotypic integration index (PINT). We also expressed the PINT as a percentage depending on the maximum possible integration levels (RelPINT). PINT and RelPINT were estimated using the package




Figure 1 Networks of hummingbirds and their nectar plants with identified modules indicated by colors. (A) Ecological network comprised by the plant and the hummingbird assemblages from the two habitats, identifying modules for plants and pollinators. (B) Ecological network obtained from the rainforest habitat. (C) Ecological network from the savanna habitat. The thickness of the lines is proportional to the strength of the interactions. Circles from A to Z represent the plant species: (A) *Aechmea tillandsioides*, (B) *Aechmea bracteata*, (C) *Androlepis skinneri*, (D) *Billbergia viridiflora*, (E) *Bromelia pinguin*, (F) *Calathea lutea*, (G) *Catopsis berteroniana*, (H) *Costus pictus*, (I) *Costus scaber*, (J) *Erythrina folkersii*, (K) *Heliconia aurantiaca*, (L) *Heliconia collinsiana*, (M) *Heliconia latispatha*, (N) *Heliconia librata*, (O) *Heliconia wagneriana*, (P) *Justicia aurea*, (Q) *Malvaviscus arboreus*, (R) *Odontonema callistachyum*, (S) *Odontonema tubaeforme*, (T) *Palicourea triphylla*, (U) *Psychotria poeppigiana*, (V) *Stromanthe macrochlamys*, (W) *Tillandsia bulbosa*, (X) *Tillandsia pruinosa*, (Y) *Tillandsia streptophylla*, (Z) *Vriesea heliconioides*. Circles H1–2 represent the hummingbird Hermits clade: *Phaethornis longirostris* (H1) and *Phaethornis striigularis* (H2). Circles E1–2 represent the hummingbird Emeralds clade: *Chlorestes candida* (E1) and *Amazilia tzacatl* (E2). [Full-size !\[\]\(43e4e85f3e07c993849f98a1fa87e486_img.jpg\) DOI: 10.7717/peerj.10974/fig-1](https://doi.org/10.7717/peerj.10974/fig-1)

PHENIX (Torices & Muñoz-Pajares, 2015) in R software (R Development Core Team, 2018); both are based on a correlation matrix following Wagner (1984). We calculated the PINT for each plant species (except those lacking sufficient data) and the average PINT of plants in both communities. Since flowers with floral traits of a similar size, i.e., corolla, stamen, and style length, produce high PINT values simply by correlation, we included nectar metrics as floral traits. The reward traits were added to mitigate high PINT values unrelated with floral specialization. We obtained the average PINT across the species of each module of the overall interaction network to link phenotypic floral integration patterns and ecological specialization and assessed differences across these species with one-way ANOVAs. We also compared the average PINT across habitats following the same procedure.

RESULTS

Hummingbirds and their floral resources

The plant-hummingbird data set comprised a total of 3,403 interactions between 26 plant species belonging to eight families and four hummingbird species. In the rainforest habitat, we recorded 1,069 interactions with 18 plant species belonging to eight families (Acanthaceae, Bromeliaceae, Costaceae, Fabaceae, Heliconiaceae, Malvaceae, Marantaceae, Rubiaceae) (Figs. 1 and 2). In the savanna habitat, we recorded 953 interactions with eight plant species belonging to two families (Bromeliaceae and Rubiaceae) (Figs. 1 and 2). The hummingbird assemblage was the same in both communities and composed of year-round species, including two species in the Emeralds clade (*Chlorestes candida*




Figure 2 (A) Rainforest and (B) savanna habitats with some examples of understory plant species visited by hummingbirds (C–J) photographed in the study site. Plant species correspond to: (C) *Jus-ticia aurea* (Acanthaceae), (D) *Heliconia wagneriana* (Heliconiaceae), (E) *Bromelia pinguin* (Bromeliaceae) and (F) *Costus scaber* (Costaceae) from the rainforest assemblage and (G) *Palicourea triphylla* (Rubiaceae), (H) *Tillandsia pruinosa* (Bromeliaceae), (I) *Psychotria poeppigiana* (Rubiaceae) and (J) *Androlepis skinneri* (Bromeliaceae) from the savanna assemblage. Photo credit: Jaume Izquierdo-Palma.

Full-size DOI: [10.7717/peerj.10974/fig-2](https://doi.org/10.7717/peerj.10974/fig-2)

and *Amazilia tzacatl*) and two in the Hermits clade (*Phaethornis longirostris* and *Phaethornis striigularis*). In the rainforest habitat, we recorded three additional species: *Anthracothonax prevostii*, *Heliothryx barroti* and *Phaeochroa cuvierii*. However, they only made illegitimate visits, acting as nectar robbers. For this reason, these species were not included in the mutualistic network described below. The plant assemblages were distinct in each community, with no shared species, yet the hummingbird species were the same. Thus, both habitats are considered as a single interaction network interconnected by the hummingbirds.

Despite we did not sample other plant-pollinator interactions, insects as floral visitors were rare in general and acted mainly as illegitimate visitors based on our observations. Most of the plant assemblage exhibited a clear ornithophilic syndrome (acting by itself as an insect deterrent) but some plant species showed less strict floral barriers against generalized pollinators, such as *Catopsis berteroniana*. From the video recordings we identified legitimate visits by insects in *Calathea lutea* and *Stromanthe macrochlamys*, both from the Marantaceae family. Despite *Phaethornis striigularis* was the only hummingbird visitor registered in both of these plant species, we observed insects (butterflies, bees and hoverflies) visiting and even opening the flowers using their complex explosive pollination mechanism (Ley & Claßen-Bockhoff, 2009). We also noticed abundant legitimate visits by different butterfly species in *Psychotria poeppigiana*. Thus, hummingbirds acted as occasional visitors to those species. In addition, we observed some visits in *Costus pictus* by euglossine bees, but we could not determine whether they were legitimate or illegitimate. In some plant species, such as *Heliconia librata* and *Heliconia aurantiaca*, we identified small butterflies, ants and small bees (mainly stingless bees of the genus *Trigona*) obtaining nectar from the bracts, acting as nectar robbers, but none of these observations were considered in this study.

Plant-hummingbird interaction networks

The complete network had low levels of connectance (0.35, z-score = -2.82, $p = 0.005$) and high levels of H_2 compared to the null matrices (0.87, z-score = 15.97, $p < 0.0001$), showing ecological specialization between hummingbirds and plants. The values of NODF (38.33, z-score = -3.960, $p = 0.002$) and wNODF (11.41, z-score = -3.89, $p < 0.001$) were statistically significant, showing lower levels of nestedness than expected. The Q value (0.51, z-score = 16.43, $p < 0.001$) indicated significant modularity that was higher than expected. We obtained three modules: one formed by *Phaethornis longirostris*, another by *Phaethornis striigularis*, and a final formed by the two Emerald species, *Chlorestes candida* and *Amazilia tzacatl*. The modules did not separate the habitats, but the habitats were related with the ecological specialization of species. The module formed by *P. longirostris* only included plants species from the rainforest assemblage (Fig. 3; Table S1). As we obtained the same modules using the QuaBiMo and MODULAR software, we only used the results from QuaBiMo because our network was quantitative.

The Hermits were the main clade of floral visitors. *Phaethornis longirostris* visited 16 plant species and was the only hummingbird species recorded in 11 of these, all belonging to the rainforest assemblage (Figs. 1 and 2). The strength of the interaction (represented by the number of visits/h) between *P. longirostris* and *Heliconia wagneriana* is remarkable, with a mean of 47.44 visits/h, far above any other interaction. This visitation rate can be explained by the fact that *H. wagneriana* grows in large patches and *P. longirostris* is the only hummingbird capable of obtaining nectar from their long, curved flowers (Fig. 4). Thus, they remained near the *H. wagneriana* patches during the flowering period, taking advantage of their abundance despite being considered trapezoiders. *Phaethornis striigularis* visited the flowers of 15 plant species and was the only visitor recorded to eight of them, five of which were in the rainforest and two in the savanna




Figure 3 Average floral integration index (PINT) found in each module. Colors indicate the modules found in the complete network including the rainforest and savanna assemblages. One module is formed by *Amazilia tzacatl* (*A. tzacatl*) and *Chlorestes candida* (*Ch. candida*), one by *Phaethornis striigularis* (*P. striigularis*) and one by *Phaethornis longirostris* (*P. longirostris*). Hummingbird illustration credit: Marco Antonio Pineda Maldonado/Banco de Imágenes CONABIO.

Full-size DOI: 10.7717/peerj.10974/fig-3



Figure 4 Trait matching between corresponding pairs of morphological traits in three plant species and their exclusive hummingbird visitor, *Phaethornis longirostris*, in the study area. Plant species correspond to: (A) *Billbergia viridiflora* (Bromeliaceae), (B) *Heliconia aurantiaca* (Heliconiaceae), and (C) *Heliconia wagneriana* (Heliconiaceae). Photo credit: Jaume Izquierdo-Palma.

Full-size DOI: 10.7717/peerj.10974/fig-4

assemblage (Figs. 1 and 2). *Amazilia tzacatl* and *Chlorestes candida* visited seven and six plant species, respectively. These Emeralds always acted as generalist foragers of generalist plants species in both habitats. This is probably due to the trait mismatch between their bills and the specialized corollas, so they were not the only visitors recorded to any of the flowering plants.

When dividing the complete network by habitat, each community differed considerably in its network topography. The plant-hummingbird interaction network in the rainforest habitat had a low level of connectance (0.35, z -score = -2.88 , $p = 0.004$) and

Table 1 Contribution of morphological and nectar variables in the PCA analysis. Variables contribution in the PCA analysis related to floral types according to the floral measurements (corolla length and curvature) and nectar metrics (volume and concentration) from plant species visited legitimately by hummingbirds in the study area. Total variance explained: PC1 (50.5%), PC1 (19.5%) and PC3 (18.18%).

	PC1	PC2	PC3
Corolla length (mm)	-0.724813	-0.554099	0.104947
Curvature (degrees)	-0.732850	0.057970	0.572058
Nectar volume (μ l)	-0.710645	-0.131267	-0.617961
Nectar concentration ($^{\circ}$ Bx)	-0.672742	0.672500	-0.083464

high level of H_2' (0.83, z-score = 4.95, $p < 0.0001$), similar to the complete network. The NODF (25.77, z-score = -0.18, $p = 0.86$) was not statistically significant, yet the wNODF (10.15, z-score = -2.32, $p = 0.02$) was lower than expected. In the rainforest habitat, *Phaethornis longirostris* was the main floral visitor to 13 plant species followed by *Phaethornis striigularis*, which visited seven species (Fig. 1). On the other hand, *Amazilia tzacatl* and *Chlorestes candida* only visited three or two plant species, respectively. The plant-hummingbird interaction network in the savanna habitat showed higher levels of connectance compared to the rainforest community, although these were lower than expected (0.59, z-score = -2.64, $p = 0.009$). The H_2' value (0.47, z-score = 3, $p = 0.003$) was intermediate, suggesting less niche specialization than in the rainforest community (Fig. 1). The NODF (70.59, z-score = 0.71, $p = 0.47$) and wNODF (41.91, z-score = 1.37, $p = 0.17$) were higher but not significantly higher as compared to the rainforest network. In the savanna habitat, *P. striigularis* was the main visitor to eight plant species. On the other hand, we only recorded a few visits of *P. longirostris* to three plant species, with a visitation rate of 0.02 to 0.10 visits/h. For this reason, the latter species can be considered a rare visitor to the savanna habitat. Unlike the rainforest interaction network, the two Emerald species had a greater role as floral visitors and behaved as territorial in the savanna habitat, as indicated by the strength of some of their floral interactions (Fig. 1).

Plant-hummingbird trait matching

From the PCA analysis, we obtained three principal components that accumulated 88.18% of the total variance (Tables 1 and 2). Floral traits selected after the correlation analysis were corolla length, corolla curvature, nectar volume, and nectar concentration (Table S2). The first component was related with straight, small-sized flowers with dilute nectar in small quantities (PC1: 50.5% of total variance). The main plant families matching with this category were Bromeliaceae (4 species), Rubiaceae (2), Acanthaceae (1), and Marantaceae (1). Fifty-three percent were from the rainforest and the remaining 47% from the savanna. Eighty percent of the plant species were visited by *Phaethornis striigularis*, 20% by *P. longirostris*, 40% by *Amazilia tzacatl*, and 33% by *Chlorestes candida*. The second factor was related with small flowers with high nectar concentration (PC2: 19.5% of total variance). Only two species belonged to this factor, *Calathea lutea* and

Table 2 Plant species contribution in the three principal components according to their floral and nectar measures. Plant species contribution in the PCA analysis related to floral types according to their floral measures and nectar metrics. Family is indicated for each plant species.

	PC1	PC2	PC3	Family
<i>Justicia aurea</i>	-0.20912	-1.38380	0.63834	Acanthaceae
<i>Odontonema callistachyum</i>	1.54261	-0.33942	0.78373	Acanthaceae
<i>Odontonema tubaeforme</i>	0.26520	0.01332	1.74687	Acanthaceae
<i>Aechmea tillandsioides</i>	1.05441	0.41421	-0.49576	Bromeliaceae
<i>Billbergia viridiflora</i>	-1.32573	0.53509	0.88704	Bromeliaceae
<i>Vriesea heliconioides</i>	0.35424	0.19862	0.72610	Bromeliaceae
<i>Costus pictus</i>	-2.96626	0.26148	-0.55565	Costaceae
<i>Costus scaber</i>	-1.85085	-0.13146	0.73183	Costaceae
<i>Erythrina folkersii</i>	0.36308	-2.31230	-0.23929	Fabaceae
<i>Heliconia aurantiaca</i>	-1.53691	-0.74214	0.29206	Heliconiaceae
<i>Heliconia collinsiana</i>	-2.58638	-0.00323	-0.54788	Heliconiaceae
<i>Heliconia latispatha</i>	-1.44918	-0.29543	-2.40521	Heliconiaceae
<i>Heliconia librata</i>	0.41088	0.55341	-0.39214	Heliconiaceae
<i>Heliconia wagneriana</i>	-1.02959	-0.66353	0.50188	Heliconiaceae
<i>Malvaviscus arboreus</i>	0.80604	-0.78644	-0.57486	Malvaceae
<i>Calathea lutea</i>	-1.40318	2.44192	0.32448	Marantaceae
<i>Stromanthe macrochlamys</i>	1.56472	1.00354	-0.29824	Marantaceae
<i>Aechmea bracteata</i>	1.89721	0.36199	-0.16328	Bromeliaceae
<i>Androlepis skinneri</i>	1.07203	0.24615	-1.24655	Bromeliaceae
<i>Catopsis berteroniana</i>	1.68092	-0.31029	-0.72595	Bromeliaceae
<i>Tillandsia streptophylla</i>	0.34628	-0.13030	0.57020	Bromeliaceae
<i>Tillandsia bulbosa</i>	0.43713	-0.06154	0.69774	Bromeliaceae
<i>Palicourea triphylla</i>	1.27467	0.39677	0.13276	Rubiaceae
<i>Psychotria poeppigiana</i>	1.28778	0.73338	-0.38821	Rubiaceae

Stromanthe macrochlamys, both found in the rainforest habitat. As explained above, both of these plant species were visited mainly by insects apart from *Phaethornis striigularis*. Lastly, the third factor was related with flowers with moderate corolla curvature and low nectar volume (PC3: 18.18% of the total variance). The associated plant species mainly belonged to the Bromeliaceae (5 species) and Acanthaceae (3) families. Seventy-five percent were from the rainforest and 25% from the savanna. In this case, the two Hermit species were the only visitors, and they visited the same number of plant species.

We observed trait matching between plants (floral traits) and hummingbirds (bill morphology) mainly in species with specialized interactions (Fig. 4; Table 3). Plant species exclusively visited by *Phaethornis longirostris* were differentiated by their long and curved corollas (i.e., *Heliconia* species). The average bill length of *P. longirostris* was 53.3 ± 2.88 mm ($n = 30$), practically identical to the average corolla length of the flowers they exclusively visited, allowing them access the nectar. This Hermit species had the most curved bill in the study area ($31.82^\circ \pm 4.33$, $n = 18$), and the flowers it visited also had

Table 3 Hummingbird species (or groups) associated with the average floral traits across plant species they visited legitimately. Average floral measures and nectar metrics across plant species visited exclusively by *Phaethornis longirostris*, *Phaethornis striigularis* and visited by multiple species (*Amazilia tzacatl*, *Chlorestes candida*, *Phaethornis longirostris* and/or *Phaethornis striigularis*).

Species	Corolla length (mm)	Stamen length (mm)	Style length (mm)	Curvature (degrees)	Nectar volume (μl)	Nectar concentration (°Bx)
<i>Phaethornis longirostris</i>	53.45 ± 15.96 (n = 11)	57.46 ± 7.15 (n = 10)	56.09 ± 6.92 (n = 10)	13.45 ± 6.75 (n = 9)	26.75 ± 16.74 (n = 10)	24.19 ± 4.73 (n = 10)
<i>Phaethornis striigularis</i>	22.79 ± 11.49 (n = 8)	26.48 ± 14.92 (n = 8)	26.42 ± 16.35 (n = 7)	9.24 ± 5.93 (n = 8)	5.98 ± 5.16 (n = 7)	23.54 ± 6.30 (n = 7)
<i>Amazilia tzacatl</i> , <i>Chlorestes candida</i> , <i>Phaethornis longirostris</i> , <i>Phaethornis striigularis</i>	18.30 ± 9.28 (n = 7)	17.56 ± 11.13 (n = 7)	20.04 ± 10.98 (n = 6)	1.62 ± 2.77 (n = 7)	24.06 ± 24.68 (n = 7)	22.55 ± 2.47 (n = 7)

higher curvature in their corollas. However, its bill was approximately three times more curved than the corolla of its visited flowers. In addition, this Hermit species was the largest hummingbird of the assemblage, with an average body weight of 5.50 g (± 0.83 , $n = 15$), which seems related with the highest average nectar volume and sugar concentration of its visited plant species. We also obtained trait matching between *Phaethornis striigularis* and the flowers they exclusively visited, mainly small- to medium-sized flowers with some degree of curvature (higher than 5°). The average bill length of *P. striigularis* was 27.39 mm (± 1.23 , $n = 20$), close to the average corolla length of its visited flowers. The average bill curvature was 25.21° (± 3.40 , $n = 14$) although, as observed with the other Hermit species, the bill curvature was higher than the average corolla curvature. *Phaethornis striigularis* was the smallest hummingbird of the assemblage, with an average body weight of 2.61 g (± 1.32 , $n = 11$). The average nectar volume of the flowers they exclusively visited was approximately 4.5 times lower than those visited by *P. longirostris*. However, the sugar concentration remained similar. Finally, small- to medium-sized flowers with less than 5° of corolla curvature were visited by several hummingbird species, mainly by the Emerald species and *P. striigularis*.

Seven out of 26 plant species received visits by two or more hummingbird species. In these seven species the average corolla length was shorter than the bill length of the two Emerald species, which was 23.52 mm (± 1.53 , $n = 28$) in *Chlorestes candida* and 27.9 mm (± 1.97 , $n = 30$) in *Amazilia tzacatl*, similar to the average bill length of *P. striigularis*. The main difference of the Emerald species was related to the bill curvature, with these species having the straighter bills of the assemblage, or an average bill curvature of 17.84° (± 3.13 , $n = 20$) for *Ch. candida* and 16.75° (± 2.93 , $n = 13$) for *A. tzacatl*. Correspondingly, the flowers they visited were straight or had little curvature in their corollas. Regarding body weight, the two Emerald species had intermediate values between the two *Phaethornis* species, or 3.35 g (± 0.45 g, $n = 12$) for *Ch. candida* and 4.93 g (± 0.96 g, $n = 10$) for *A. tzacatl*. The average nectar volume of the flowers they visited was similar to that of the flowers visited by *P. longirostris*, although the average sugar concentration was lower, corresponding with 22.55°Bx (± 2.47 , $n = 123$) (Table 3).

Phenotypic floral integration

We obtained phenotypic integration values for 22 out of the 26 plant species (Fig. 3; Table S1). The average floral integration of the plant assemblage in our study site (with nectar variables) was 19.38%, around the average (21.5%) for the angiosperms examined by *Ordano et al. (2008)*. Sufficient data were not available for the following plant species: *Aechmea tillandsioides*, *Billbergia viridiflora*, *Bromelia pinguin* and *Tillandsia pruinosa*. In comparing the average PINT of the plant assemblages of each habitat, the savanna community had a slightly higher value (rainforest = 0.80, $n = 15$; savanna = 0.92, $n = 7$), but it was not statistically significant ($F = 0.35$, $df = 1$, $p = 0.56$). The results of the PINT analysis across modules suggest that specialized modules had higher values, even though they were statistically similar. The plant species integrated to the *Phaethornis longirostris* module had higher values (PINT = 0.92, RelPINT = 22.79%, $n = 9$), followed by those integrated to the *Phaethornis striigularis* module (PINT = 0.82, RelPINT = 20.50%, $n = 8$). Meanwhile, the module integrated by the two Emerald species had lower values (PINT = 0.74, RelPINT = 20.36%, $n = 5$), although the results from the ANOVA test showed that these differences were not significant ($F = 0.25$, $df = 2$, $p = 0.79$). Therefore, the relationship between the ecological specialization of modules and their phenotypic floral integration index was unclear according to these data (Fig. 3).

DISCUSSION

As expected, we found that forbidden links and trait matching promote modularity in the plant-hummingbird system of the Lacandona rainforest. However, the low number of modules and small pollination networks likely affected the non-significant relationship between ecological specialization and phenotypic floral integration.

Our results suggest that the adjacent habitats, interconnected by the same hummingbird species, did not function as separate units but instead form a single plant-hummingbird interaction network. Thus, it is possible that two plants from the rainforest and savanna are more intimately linked through their shared hummingbird species than two plants from the same habitat with different hummingbird pollinators. According to *Bergamo et al. (2017)*, the overlap of pollinators can influence the visitation patterns and potentially lead to indirect interactions (e.g., facilitation or competition), especially with plants with a similar floral phenotype. Nevertheless, we found that strong habitat differences in plant composition might impact some of the structural parameters when analyzed separately. The rainforest habitat was characterized by plant species with long and curved corollas, whereas small- and medium-sized flowers with straight corollas or with a little curvature (even non-ornithophilous) characterized the savanna habitat (*Arizmendi & Ornelas, 1990; Araújo, Sazima & Oliveira, 2013; Maruyama et al., 2013*). The lack of flowers with long corollas is probably the cause of the almost complete absence of *Phaethornis longirostris* in the savanna habitat. Floral morphology has been shown to play an important role in tropical hummingbird-pollination systems, influencing not only the visitors but also the strength of their interactions. For example, in plant-hummingbird interactions on the West Indies, most specialized hummingbird-pollinated plants were found in highlands and were mainly pollinated by large, long-billed

hummingbirds, whereas highly generalist plants were found in dry and warm lowlands and were pollinated by small, short-billed hummingbirds in addition to insect species (Dalsgaard *et al.*, 2009). In another case study in Brazil, Maruyama *et al.* (2014) highlighted the importance of traits as determinants of interaction frequencies and associated them with morphological specialization and habitat occupancy, the main network structurers, in a Neotropical savanna/forest network.

In relation to network metrics, our results showed low levels of connectance and high complementary specialization in accordance with other mutualistic networks in tropical forests (Vizentin-Bugoni, Maruyama & Sazima, 2014; Maglianesi *et al.*, 2015; Araujo *et al.*, 2018). The relationship between plants and hummingbirds resulted highly asymmetric: Many plants only received visits from a single hummingbird species, whereas some hummingbirds visited more than ten plant species. However, reciprocally specialized interactions are rare in nature, even in networks considered specialized (Joppa *et al.*, 2009). Despite the low number of hummingbird species in the habitats sampled, they showed high variation in their morphological traits such as body size, bill length and foraging behavior. Morphological and behavioral differences among species enabled them to be classified into three roles in the organization of the community: *Phaethornis longirostris* is a high-reward trapliner and *P. striigularis* is a low-reward trapliner (frequently acting as a nectar robber when it is unable to access the nectar reward). And, depending on the patch quality, *Chlorestes candida* and *Amazilia tzacatl* act as territorial and generalist species (Feinsinger & Colwell, 1978). Therefore, in our network, both trait matching and forbidden links could be playing a major role in niche partitioning, shaping the network structure (Dalsgaard *et al.*, 2011). Morphological resemblance has been found to allow the exclusive access of some species (e.g., *Phaethornis* species) to the most specialized flowers (Bergamo *et al.*, 2018; Sonne *et al.*, 2019; Sonne *et al.*, 2020). Moreover, forbidden links regulated the interactions of the two Emerald species with less specialized bill morphologies that were unable to access flowers with long and curved corollas. Thus, variation in feeding strategies and degrees of specialization with respect to specific floral resources might be crucial for the coexistence of hummingbird species (Abrahamczyk & Kessler, 2015; Maglianesi *et al.*, 2015; Rodríguez-Flores *et al.*, 2019; Sonne *et al.*, 2020).

We found that modularity was not significantly related with habitat occupancy but rather with morphological specialization (Maruyama *et al.*, 2014). Interestingly, both Hermit species formed modules integrated by only one species. Differences in bill length and curvature may promote specialization in specific floral morphologies, as reported by Rodríguez-Flores & Stiles (2005) for the Colombian Amazon. The Hermits clade is considered the most specialized hummingbird group in regard to food resources and is highly diverse in the rainforests of South America (Rodríguez-Flores & Stiles, 2005). Given that several plant species were visited exclusively by *P. longirostris* and *P. striigularis*, these hummingbirds could be acting as “key” species for the maintenance of the plant community (De Araújo, Hoffmann & Sazima, 2018). Hermits have been previously reported to play this role in other studies and to interact with more plants than other

hummingbird species, for example, *Phaethornis eurynome* in the Atlantic rainforest (Vizentin-Bugoni, Maruyama & Sazima, 2014) and *P. petrei* in the Neotropical savanna of Brazil (Maruyama et al., 2014; De Araújo, Hoffmann & Sazima, 2018). The two Emerald species, unlike the Hermits, were also observed feeding in canopy trees, always on non-specialized flowers where nectar is easily accessible (e.g., *Inga vera* in Fabaceae and *Quararibea funebris* in Bombacaceae). Additionally, in the understory, the role of the two Emerald species was more important for plant species with less restrictive morphological floral barriers, where they usually behave as territorial. Thus, in the absence of morphological specialization, the dominance hierarchy, which is correlated with body size, might play an important role in the Emeralds' niche partitioning (Rodríguez-Flores & Arizmendi, 2016, López-Segoviano, Bribiesca & Arizmendi, 2018; Márquez-Luna et al., 2019).

Contrary to expectations, we did not find higher floral integration in specialized modules or differences between habitats. Some studies have reported the absence of evidence for pollinator-mediated selection on correlated traits (e.g., Conner, 2002; Herrera et al., 2002; Meng et al., 2008). In both studied habitats, *Phaethornis longirostris* and *P. striigularis* were the only visitors to many plant species, which was consequently reflected in the floral integration. Plant species with specialized pollination systems should experience stronger or more consistent stabilizing or directional selection on floral traits than species with generalized pollination (Rosas-Guerrero et al., 2011). However, high covariation among floral and vegetative traits could be the default situation (Armbruster et al., 1999).

In this context, pollination by various functional groups would decrease the homogeneity of the pollination function and, as a result, the correlational selection on relevant floral characters and nectar properties (Berg, 1960; Ordano et al., 2008). For this reason, the fact of not having considered other invertebrate visitors should not alter our hypothesis, since specialized modules are less likely to be visited by other pollinator taxa. Some studies such as Rosas-Guerrero et al. (2011) on *Ipomoea* and Pérez, Arroyo & Medel (2007) on *Schizanthus* support the idea that floral integration in pollinator-dependent species is shaped by pollinator-mediated selection and is stronger in specialized relationships. However, these studies tested differences between different functional groups of pollinators or morphospecies, for example, between bird pollination and insect pollination systems. Thus, differences can be higher across plant species with different pollination systems when comparisons are conducted among species from the same family. Nevertheless, all plants included in both assemblages received legitimate visits from hummingbirds despite differences in their floral specialization.

Studies on plant-pollinator mutualistic networks have provided important information for understanding the underlying processes that structure communities. However, the impact of pollinators on their nutritional plants, especially those with a greater degree of specificity, has received little attention. Herein, we used a new approach with the aim of linking the underlying network structuring processes with the consequent modularity related to ecological specialization and the consequences for phenotypic floral integration in two adjacent habitats in the Lacandona rainforest of Mexico. However, we noted a

main limitation in our study, which is the low number of modules and pollination networks that can limited the assessed relationships between ecological specialization of modules and their phenotypic floral integration. To assess whether specialized modules are characterized by higher phenotypic floral integration one would need larger datasets of both the number of modules and pollination networks. Thus, generalizations about the processes involved in the modular organization of plant-hummingbird networks in terms of morphological specialization (i.e., phenotypic floral integration could be made). Therefore, future work reviewing plant-hummingbird interactions and levels of flower integration should include datasets from a wider region to obtain both a higher number of specialized or generalist modules and networks of various sizes. Although the number of modules and network size were the main limitations in our study to properly assess the relationships between ecological specialization of modules and their phenotypic floral integration, it is also possible that the establishment of modules failed because the interaction network was built based on the legitimate visitation rates of their floral visitors (potential pollinators) instead of the pollinating efficiency of each hummingbird species. Therefore, we also suggest pollination experiments to determine the effectiveness and relative role of each visitor as pollinators and the breeding system of the plants species in the mutualistic networks as alternatives to determine the number and specialization levels of their modules. Clearly, this is a critical question for future research to address.

CONCLUSIONS

Mutualistic networks vary in their number of connections and the strength of interactions among species with distinct ecological specializations. Herein, we found that the plant composition of two adjacent habitats in the Lacandona rainforest may impact some of the structural parameters of the studied hummingbird-plant networks. Although the plant assemblages were distinct, the two habitats were highly interconnected by the hummingbirds, meaning that they formed a single interaction network. Forbidden links and trait matching were important mechanisms shaping the network topology, and they varied their relative importance according to the specialization of the species involved and habitat sampled. Modularity was associated with morphological specialization and, indirectly, with the habitat affinity of species. However, the low number of modules and pollination networks in our study limited our assessment of how ecological specialization affects the phenotypic floral integration among modules. Future research should seek to include datasets from a wider region to obtain a higher number of specialized or generalist modules and networks of various sizes in order to delineate further the relationship between plant-hummingbird interactions and their levels of floral integration.

ACKNOWLEDGEMENTS

We thank Natura Mexicana A.C., especially Javier de la Maza and Julia Carabias for the logistical support and allowing access to their facilities. We thank the staff of the Estación Biológica Chajul for trail maintenance, security, and accommodation. We thank Stuart Pimm, Pietro Kiyoshi Maruyama, and one anonymous reviewer who provided

constructive comments that improved the manuscript. JI-P thanks Sergio Izquierdo and Fulvio Eccardi for being a constant source of inspiration as well as Posgrado en Ciencias Biológicas (Universidad Nacional Autónoma de México) and Universidad Autónoma de Tlaxcala for the logistical support, fieldwork equipment, and supplies. This work constitutes the partial fulfillment of JI-P's doctorate at the Universidad Nacional Autónoma de México. This paper is dedicated to the memory of Ricardo Frias, who suddenly passed in April 2020. He was a tireless defender of the Lacandona rainforest and considerably improved this manuscript with his invaluable comments and ideas.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by Universidad Nacional Autónoma de México (UNAM) PAPIIT research funds IN216617 and Consejo Nacional de Ciencia y Tecnología (CONACyT) with the JI-P doctoral scholarship (No. 855974). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Universidad Nacional Autónoma de México (UNAM) PAPIIT Research Funds: IN216617. Consejo Nacional de Ciencia y Tecnología (CONACyT): 855974.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Jaume Izquierdo-Palma conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Maria del Coro Arizmendi conceived and designed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Carlos Lara conceived and designed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Juan Francisco Ornelas conceived and designed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

Raw data and R scripts are available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.10974#supplemental-information>.

REFERENCES

- Abrahamczyk S, Kessler M. 2010.** Hummingbird diversity, food niche characters, and assemblage composition along a latitudinal precipitation gradient in the Bolivian lowlands. *Journal of Ornithology* **151**(3):615–625 DOI [10.1007/s10336-010-0496-x](https://doi.org/10.1007/s10336-010-0496-x).
- Abrahamczyk S, Kessler M. 2015.** Morphological and behavioural adaptations to feed on nectar: how feeding ecology determines the diversity and composition of hummingbird assemblages. *Journal of Ornithology* **156**(2):333–347 DOI [10.1007/s10336-014-1146-5](https://doi.org/10.1007/s10336-014-1146-5).
- Almeida-Neto M, Guimarães P, Guimarães PR, Loyola RD, Ulrich W. 2008.** A consistent metric for nestedness analysis in ecological systems: reconciling concept and measurement. *Oikos* **117**(8):1227–1239 DOI [10.1111/j.0030-1299.2008.16644.x](https://doi.org/10.1111/j.0030-1299.2008.16644.x).
- Almeida-Neto M, Ulrich W. 2011.** A straightforward computational approach for measuring nestedness using quantitative matrices. *Environmental Modelling & Software* **26**(2):173–178 DOI [10.1016/j.envsoft.2010.08.003](https://doi.org/10.1016/j.envsoft.2010.08.003).
- Anderson IA, Busch JW. 2006.** Relaxed pollinator-mediated selection weakens floral integration in self-compatible taxa of *Leavenworthia* (Brassicaceae). *American Journal of Botany* **93**(6):860–867 DOI [10.3732/ajb.93.6.860](https://doi.org/10.3732/ajb.93.6.860).
- Araujo AC, Martín González AM, Sandel B, Maruyama PK, Fischer E, Vizontin-Bugoni J, Araújo FP, Coelho AG, Faria RR, Kohler G, Las-Casas FMG, Lopes AV, Machado AO, Machado CG, Machado IC, McGuire JA, Moura AC, Oliveira GM, Oliveira PE, Rocca MA, Rodrigues LC, Rodrigues M, Rui AM, Sazima I, Sazima M, Varassin IG, Wang Z, Dalsgaard B, Svenning JC. 2018.** Spatial distance and climate determine modularity in a cross-biomes plant-hummingbird interaction network in Brazil. *Journal of Biogeography* **45**(8):1846–1858 DOI [10.1111/jbi.13367](https://doi.org/10.1111/jbi.13367).
- Araújo FP, Sazima M, Oliveira PE. 2013.** The assembly of plants used as nectar sources by hummingbirds in a Cerrado area of Central Brazil. *Plant Systematics and Evolution* **299**(6):Sazima M, Oliveira–Sazima M, Oliveira1133 DOI [10.1007/s00606-013-0783-0](https://doi.org/10.1007/s00606-013-0783-0).
- Arizmendi MC, Berlanga H. 2014.** *Colibríes de México y Norteamérica*. México: CONABIO, 160.
- Arizmendi MC, Ornelas JF. 1990.** Hummingbirds and their floral resources in a tropical dry forest in Mexico. *Biotropica* **22**(2):172–180 DOI [10.2307/2388410](https://doi.org/10.2307/2388410).
- Armbruster WS. 1991.** Multilevel analysis of morphometric data from natural plant populations: insights into ontogenetic, genetic, and selective correlations in *Dalechampia scandens*. *Evolution* **45**(5):1229–1244 DOI [10.1111/j.1558-5646.1991.tb04389.x](https://doi.org/10.1111/j.1558-5646.1991.tb04389.x).
- Armbruster WS, Di Stilio VNS, Tuxill JD, Flores TC, Velásquez Runk JL. 1999.** Covariance and decoupling of floral and vegetative traits in nine Neotropical plants: a re-evaluation of Berg's correlation-pleiades concept. *American Journal of Botany* **86**(1):39–55 DOI [10.2307/2656953](https://doi.org/10.2307/2656953).
- Barber MJ. 2007.** Modularity and community detection in bipartite networks. *Physical Review E* **76**(6):066102 DOI [10.1103/PhysRevE.76.066102](https://doi.org/10.1103/PhysRevE.76.066102).
- Bascompte J, Jordano P. 2007.** Plant-animal mutualistic networks: the architecture of biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **38**(1):567–593 DOI [10.1146/annurev.ecolsys.38.091206.095818](https://doi.org/10.1146/annurev.ecolsys.38.091206.095818).
- Bascompte J, Jordano P. 2013.** *Mutualistic networks*. Princeton: Princeton University Press.
- Bascompte J, Jordano P, Melián CJ, Olesen JM. 2003.** The nested assembly of plant animal mutualistic networks. *Proceedings of the National Academy of Sciences* **100**:9383–9387 DOI [10.1073/pnas.1633576100](https://doi.org/10.1073/pnas.1633576100).
- Bascompte J, Jordano P, Olesen JM. 2006.** Asymmetric coevolutionary networks facilitate biodiversity maintenance. *Science* **312**(5772):431–433 DOI [10.1126/science.1123412](https://doi.org/10.1126/science.1123412).

- Bastolla U, Fortuna MA, Pascual-García A, Ferrera A, Luque B, Bascompte J. 2009.** The architecture of mutualistic networks minimizes competition and increases biodiversity. *Nature* **458(7241)**:1018–1020 DOI [10.1038/nature07950](https://doi.org/10.1038/nature07950).
- Belmaker J, Sekercioglu CH, Jetz W. 2012.** Global patterns of specialization and coexistence in bird assemblages. *Journal of Biogeography* **39(1)**:193–203 DOI [10.1111/j.1365-2699.2011.02591.x](https://doi.org/10.1111/j.1365-2699.2011.02591.x).
- Berg RL. 1960.** The ecological significance of correlation pleiades. *Evolution* **14(2)**:171–180 DOI [10.1111/j.1558-5646.1960.tb03076.x](https://doi.org/10.1111/j.1558-5646.1960.tb03076.x).
- Bergamo PJ, Wolowski M, Maruyama PK, Vizentin-Bugoni J, Carneiro LG, Sazima M. 2017.** The potential indirect effects among plants via shared hummingbird pollinators are structured by phenotypic similarity. *Ecology* **98(7)**:1849–1858 DOI [10.1002/ecy.1859](https://doi.org/10.1002/ecy.1859).
- Bergamo PJ, Wolowski M, Maruyama PK, Vizentin-Bugoni J, Sazima M. 2018.** Trait patterns across space and time suggest an interplay of facilitation and competition acting on Neotropical hummingbird-pollinated plant communities. *Oikos* **127(11)**:1690–1700 DOI [10.1111/oik.05571](https://doi.org/10.1111/oik.05571).
- Bissell EK, Diggle PK. 2008.** Floral morphology in *Nicotiana*: architectural and temporal effects on phenotypic integration. *International Journal of Plant Sciences* **169(2)**:225–240 DOI [10.1086/523875](https://doi.org/10.1086/523875).
- Blüthgen N, Menzel F, Hovestadt T, Fiala B, Blüthgen N. 2007.** Specialization, constraints, and conflicting interests in mutualistic networks. *Current Biology* **17(4)**:341–346 DOI [10.1016/j.cub.2006.12.039](https://doi.org/10.1016/j.cub.2006.12.039).
- Bosch J, Martín González AM, Rodrigo A, Navarro D. 2009.** Plant-pollinator networks: adding the pollinator's perspective. *Ecology Letters* **12(5)**:409–419 DOI [10.1111/j.1461-0248.2009.01296.x](https://doi.org/10.1111/j.1461-0248.2009.01296.x).
- Carabias J, De la Maza J, Cadena R. 2015.** *Conservación y desarrollo sustentable en la Selva Lacandona. 25 años de actividades y experiencias*. Ciudad de México: Natura y Ecosistemas Mexicanos.
- Conner JK. 2002.** Genetic mechanisms of floral trait correlations in a natural population. *Nature* **420(6914)**:407–410 DOI [10.1038/nature01105](https://doi.org/10.1038/nature01105).
- Conner J, Via S. 1993.** Patterns of phenotypic and genetic correlations among morphological and life-history traits in wild radish, *Raphanus raphanistrum*. *Evolution* **47(2)**:704–711 DOI [10.1111/j.1558-5646.1993.tb02128.x](https://doi.org/10.1111/j.1558-5646.1993.tb02128.x).
- Dalsgaard B, González AMM, Olesen JM, Ollerton J, Timmermann A, Andersen LH, Tossas AG. 2009.** Plant-hummingbird interactions in the West Indies: floral specialisation gradients associated with environment and hummingbird size. *Oecologia* **159(4)**:757–766 DOI [10.1007/s00442-008-1255-z](https://doi.org/10.1007/s00442-008-1255-z).
- Dalsgaard B, Magård E, Fjeldså J, Martín González AM, Rahbek C, Olesen JM, Ollerton J, Alarcón R, Araujo AC, Cotton PA, Lara C, Machado CG, Sazima I, Sazima M, Timmermann A, Watts S, Sandel B, Sutherland WJ, Svenning JC. 2011.** Specialization in plant-hummingbird networks is associated with species richness, contemporary precipitation and quaternary climate-change velocity. *PLOS ONE* **6(10)**:e25891 DOI [10.1371/journal.pone.0025891](https://doi.org/10.1371/journal.pone.0025891).
- Danieli-Silva A, de Souza JMT, Donatti AJ, Campos RP, Vicente-Silva J, Freitas L, Varassin IG. 2012.** Do pollination syndromes cause modularity and predict interactions in a pollination network in tropical high-altitude grasslands? *Oikos* **121(1)**:35–43 DOI [10.1111/j.1600-0706.2011.19089.x](https://doi.org/10.1111/j.1600-0706.2011.19089.x).

- De Araújo FP, Hoffmann D, Sazima M. 2018.** The planalto hermit, *Phaethornis pretrei*-a key species in a Neotropical savanna community in Central Brazil. *Journal of Natural History* 52(37–38):2385–2396 DOI 10.1080/00222933.2018.1536767.
- Dehling DM, Jordano P, Schaefer HM, Böhning-Gaese K, Schleuning M. 2016.** Morphology predicts species' functional roles and their degree of specialization in plant-frugivore interactions. *Proceedings of the Royal Society B: Biological Sciences* 283(1823):20152444 DOI 10.1098/rspb.2015.2444.
- Dormann CF, Gruber B, Fründ J. 2008.** Introducing the bipartite package: analyzing ecological networks. *R News* 8:8–12.
- Dormann CF, Strauss R. 2014.** A method for detecting modules in quantitative bipartite networks. *Methods in Ecology and Evolution* 5(1):90–98 DOI 10.1111/2041-210X.12139.
- Feinsinger P. 1976.** Organization of a tropical guild of nectarivorous birds. *Ecological Monographs* 46(3):257–291 DOI 10.2307/1942255.
- Feinsinger P, Colwell RK. 1978.** Community organization among Neotropical nectar-feeding birds. *American Zoologist* 18(4):779–795 DOI 10.1093/icb/18.4.779.
- Fenster CB. 1991.** Selection on floral morphology by hummingbirds. *Biotropica* 23(1):98–101 DOI 10.2307/2388696.
- Fortuna MA, Stouffer DB, Olesen JM, Jordano P, Mouillot D, Krasnov BR, Poulin R, Bascompte J. 2010.** Nestedness versus modularity in ecological networks: two sides of the same coin? *Journal of Animal Ecology* 79:811–817 DOI 10.1111/j.1365-2656.2010.01688.x.
- Guimarães JPR, Guimarães P. 2006.** Improving the analyses of nestedness for large sets of matrices. *Environmental Modelling & Software* 21(10):1512–1513 DOI 10.1016/j.envsoft.2006.04.002.
- Guimarães PR, Rico-Gray V, Furtado dos Reis S, Thompson JN. 2006.** Asymmetries in specialization in ant-plant mutualistic networks. *Proceedings of the Royal Society B: Biological Sciences* 273(1597):2041–2047 DOI 10.1098/rspb.2006.3548.
- Herrera CM, Cerdá X, Garcia MB, Guitián J, Medrano M, Rey PJ, Sánchez-Lafuente AM. 2002.** Floral integration, phenotypic covariance structure and pollinator variation in bumblebee-pollinated *Helleborus foetidus*. *Journal of Evolutionary Biology* 15(1):108–121 DOI 10.1046/j.1420-9101.2002.00365.x.
- Ings TC, Montoya JM, Bascompte J, Blüthgen N, Brown L, Dormann CF, Edwards F, Figueroa D, Ute J, Jones JI, Lauridsen RB, Ledger ME, Lewis HM, Olesen JM, Van Veen FJF, Warren PH, Woodward G. 2009.** Ecological networks-beyond food webs. *Journal of Animal Ecology* 78(1):253–269 DOI 10.1111/j.1365-2656.2008.01460.x.
- Joppa LN, Bascompte J, Montoya JM, Solé RV, Sanderson J, Pimm SL. 2009.** Reciprocal specialization in ecological networks. *Ecology Letters* 12(9):961–969 DOI 10.1111/j.1461-0248.2009.01341.x.
- Jordano P. 1987.** Patterns of mutualistic interactions in pollination and seed dispersal: connectance, dependence asymmetries, and coevolution. *American Naturalist* 129(5):657–677 DOI 10.1086/284665.
- Lara C. 2006.** Temporal dynamics of flower use by hummingbirds in a highland temperate forest in Mexico. *Ecoscience* 13(1):23–29 DOI 10.2980/1195-6860(2006)13[23:TDOFUB]2.0.CO;2.
- Lewinsohn TM, Inácio Prado P, Jordano P, Bascompte J, Olesen JM. 2006.** Structure in plant–animal interaction assemblages. *Oikos* 113(1):174–184 DOI 10.1111/j.0030-1299.2006.14583.x.
- Ley AC, Claßen-Bockhoff R. 2009.** Pollination syndromes in African Marantaceae. *Annals of Botany* 104(1):41–56 DOI 10.1093/aob/mcp106.

- López-Segoviano G, Bribiesca R, Arizmendi MC. 2018. The role of size and dominance in the feeding behaviour of coexisting hummingbirds. *Ibis* **160**(2):283–292 DOI [10.1111/ibi.12543](https://doi.org/10.1111/ibi.12543).
- Maglianesi MA, Blüthgen N, Böhning-Gaese K, Schleuning M. 2014. Morphological traits determine specialization and resource use in plant-hummingbird networks in the Neotropics. *Ecology* **95**(12):3325–3334 DOI [10.1890/13-2261.1](https://doi.org/10.1890/13-2261.1).
- Maglianesi MA, Blüthgen N, Böhning-Gaese K, Schleuning M. 2015. Functional structure and specialization in three tropical plant-hummingbird interaction networks across an elevational gradient in Costa Rica. *Ecography* **38**(11):1119–1128 DOI [10.1111/ecog.01538](https://doi.org/10.1111/ecog.01538).
- Marquitti FMD, Guimarães JPR, Pires MM, Bittencourt LF. 2014. MODULAR: software for the autonomous computation of modularity in large network sets. *Ecography* **37**(3):221–224 DOI [10.1111/j.1600-0587.2013.00506.x](https://doi.org/10.1111/j.1600-0587.2013.00506.x).
- Maruyama PK, Oliveira GM, Ferreira C, Dalsgaard B, Oliveira PE. 2013. Pollination syndromes ignored: importance of non-ornithophilous flowers to Neotropical savanna hummingbirds. *Naturwissenschaften* **100**(11):1061–1068 DOI [10.1007/s00114-013-1111-9](https://doi.org/10.1007/s00114-013-1111-9).
- Maruyama PK, Sonne J, Vizentin-Bugoni J, Martin González AM, Zanata TB, Abrahamczyk S, Alarcón R, Araujo AC, Araújo FP, Baquero AC, Chávez-González E, Coelho AG, Cotton PA, Dehling DM, Fischer E, Kohler G, Lara C, Las-Casas FMG, Machado AO, Machado CG, Maglianesi MA, Malucelli TS, Marín-Gómez OH, Oliveira PE, Ornelas JF, Ortiz-Pulido R, Ramírez-Burbano MB, Rocca MA, Rodrigues LC, Rosero-Lasprilla L, Rui AM, Sandel B, Svenning JCH, Tinoco BA, Varassin IG, Watts S, Rahbek C, Sazima M, Schleuning M, Dalsgaard B. 2018. Functional diversity mediates macroecological variation in plant-hummingbird interaction networks. *Global Ecology and Biogeography* **27**(10):1186–1199 DOI [10.1111/geb.12776](https://doi.org/10.1111/geb.12776).
- Maruyama PK, Vizentin-Bugoni J, Oliveira GM, Oliveira PE, Dalsgaard B. 2014. Morphological and spatio-temporal mismatches shape a Neotropical savanna plant-hummingbird network. *Biotropica* **46**(6):740–747 DOI [10.1111/btp.12170](https://doi.org/10.1111/btp.12170).
- McGuire JA, Witt CC, Remsen JV, Dudley R, Altshuler DL. 2009. A higher-level taxonomy for hummingbirds. *Journal of Ornithology* **150**(1):155–165 DOI [10.1007/s10336-008-0330-x](https://doi.org/10.1007/s10336-008-0330-x).
- Meng JL, Zhou XH, Zhao ZG, Du GZ. 2008. Covariance of floral and vegetative traits in four species of Ranunculaceae: a comparison between specialized and generalized pollination systems. *Journal of Integrative Plant Biology* **50**(9):1161–1170 DOI [10.1111/j.1744-7909.2008.00722.x](https://doi.org/10.1111/j.1744-7909.2008.00722.x).
- Miranda F, Hernández E. 1963. Los tipos de vegetación de México y su clasificación. *Botanical Sciences* **28**(28):29–179 DOI [10.17129/botsci.1084](https://doi.org/10.17129/botsci.1084).
- Márquez-Luna U, Lara C, Corcuera P, Valverde PL. 2019. Factors affecting the dominance hierarchy dynamics in a hummingbird assemblage. *Current Zoology* **65**(3):261–268 DOI [10.1093/cz/zoy057](https://doi.org/10.1093/cz/zoy057).
- Newman ME. 2006. Modularity and community structure in networks. *Proceedings of the National Academy of Sciences* **103**:8577–8582 DOI [10.1073/pnas.0601602103](https://doi.org/10.1073/pnas.0601602103).
- Niovi Jones K, Reithel JS. 2001. Pollinator-mediated selection on a flower color polymorphism in experimental populations of *Antirrhinum* (Scrophulariaceae). *American Journal of Botany* **88**(3):447–454 DOI [10.2307/2657109](https://doi.org/10.2307/2657109).
- Olesen JM, Bascompte J, Dupont YL, Elberling H, Rasmussen C, Jordano P. 2011. Missing and forbidden links in mutualistic networks. *Proceedings of the Royal Society B: Biological Sciences* **278**(1706):725–732 DOI [10.1098/rspb.2010.1371](https://doi.org/10.1098/rspb.2010.1371).
- Olesen JM, Bascompte J, Dupont YL, Jordano P. 2006. The smallest of all worlds: pollination networks. *Journal of Theoretical Biology* **240**(2):270–276 DOI [10.1016/j.jtbi.2005.09.014](https://doi.org/10.1016/j.jtbi.2005.09.014).

- Olesen JM, Bascompte J, Dupont YL, Jordano P. 2007. The modularity of pollination networks. *Proceedings of the National Academy of Sciences* **104**:19891–19896 DOI [10.1073/pnas.0706375104](https://doi.org/10.1073/pnas.0706375104).
- Olesen JM, Jordano P. 2002. Geographic patterns in plant-pollinator mutualistic networks. *Ecology* **83**:2416–2424 DOI [10.1890/0012-9658\(2002\)083\[2416:GPIPPM\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[2416:GPIPPM]2.0.CO;2).
- Ordano M, Fornoni J, Boege K, Domínguez CA. 2008. The adaptive value of phenotypic floral integration. *New Phytologist* **179**(4):1183–1192 DOI [10.1111/j.1469-8137.2008.02523.x](https://doi.org/10.1111/j.1469-8137.2008.02523.x).
- Paton DC, Collins BG. 1989. Bills and tongues of nectar-feeding birds: a review of morphology, function and performance, with intercontinental comparisons. *Australian Journal of Ecology* **14**(4):473–506 DOI [10.1111/j.1442-9993.1989.tb01457.x](https://doi.org/10.1111/j.1442-9993.1989.tb01457.x).
- Pérez F, Arroyo MTK, Medel R. 2007. Phylogenetic analysis of floral integration in *Schizanthus* (Solanaceae): does pollination truly integrate corolla traits? *Journal of Evolutionary Biology* **20**(5):1730–1738 DOI [10.1111/j.1420-9101.2007.01393.x](https://doi.org/10.1111/j.1420-9101.2007.01393.x).
- R Development Core Team. 2018. *R: a language and environment for statistical computing*. Vienna: The R Foundation for Statistical Computing. Available at <http://www.R-project.org>.
- Rico-Guevara A, Araya-Salas M. 2015. Bills as daggers? A test for sexually dimorphic weapons in a lekking hummingbird. *Behavioral Ecology* **26**(1):21–29 DOI [10.1093/beheco/aru182](https://doi.org/10.1093/beheco/aru182).
- Rodríguez-Flores CI, Arizmendi MC. 2016. The dynamics of hummingbird dominance and foraging strategies during the winter season in a highland community in Western Mexico. *Journal of Zoology* **299**(4):262–274 DOI [10.1111/jzo.12360](https://doi.org/10.1111/jzo.12360).
- Rodríguez-Flores CI, Ornelas JF, Wethington S, Arizmendi MC. 2019. Are hummingbirds generalists or specialists? Using network analysis to explore the mechanisms influencing their interaction with nectar resources. *PLOS ONE* **14**(2):e021855 DOI [10.1371/journal.pone.0211855](https://doi.org/10.1371/journal.pone.0211855).
- Rodríguez-Flores CI, Stiles FG. 2005. Análisis ecomorfológico de una comunidad de colibríes ermitaños (Trochilidae, Phaethorninae) y sus flores en la amazonia colombiana. *Ornitología Colombiana* **3**:3–27.
- Rosas-Guerrero V, Quesada M, Armbruster WS, Pérez-Barrales R, Smith SD. 2011. Influence of pollination specialization and breeding system on floral integration and phenotypic variation in *Ipomoea*. *Evolution: International Journal of Organic Evolution* **65**(2):350–364 DOI [10.1111/j.1558-5646.2010.01140.x](https://doi.org/10.1111/j.1558-5646.2010.01140.x).
- Rzedowski J, Huerta L. 1994. *Vegetación de México*. México: Limusa, Noriega Editores.
- Santamaría L, Rodríguez-Gironés MA. 2007. Linkage rules for plant-pollinator networks: trait complementarity or exploitation barriers? *PLoS Biology* **5**(2):e31 DOI [10.1371/journal.pbio.0050031](https://doi.org/10.1371/journal.pbio.0050031).
- Sazatornil FD, More M, Benitez-Vieyra S, Cocucci AA, Kitching IJ, Schlumpberger BO, Oliveira PE, Sazima M, Amorim FW. 2016. Beyond neutral and forbidden links: morphological matches and the assembly of mutualistic hawkmoth-plant networks. *Journal of Animal Ecology* **85**(6):1586–1594 DOI [10.1111/1365-2656.12509](https://doi.org/10.1111/1365-2656.12509).
- Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH image to imageJ: 25 years of image analysis. *Nature Methods* **9**(7):671–675 DOI [10.1038/nmeth.2089](https://doi.org/10.1038/nmeth.2089).
- Siebe C, Martínez-Ramos M, Segura-Warnholtz G, Rodríguez-Velázquez J, Sánchez-Beltrán S. 1996. Soil and vegetation patterns in the tropical rainforest at Chajul, Southeast Mexico. In: *Proceedings of the International Congress on Soils of Tropical Forest Ecosystems*. 40–58.
- Smith RA, Rausher MD. 2008. Selection for character displacement is constrained by the genetic architecture of floral traits in the ivyleaf morning glory. *Evolution* **62**(11):2829–2841 DOI [10.1111/j.1558-5646.2008.00494.x](https://doi.org/10.1111/j.1558-5646.2008.00494.x).

- Sonne J, Vizentin-Bugoni J, Maruyama PK, Araujo AC, Chávez-González E, Coelho AG, Cotton PA, Marín-Gómez OH, Lara C, Lasprilla LR, Machado CG, Maglianesi MA, Malucelli TS, Martín González AM, Oliveira GM, Oliveira PE, Ortiz-Pulido R, Rocca MA, Rodrigues LC, Sazima I, Simmons BI, Tinoco B, Varassin IG, Vasconcelos MF, O'Hara B, Schleuning M, Rahbek C, Sazima M, Dalsgaard B. 2020. Ecological mechanisms explaining interactions within plant-hummingbird networks: morphological matching increases towards lower latitudes. *Proceedings of the Royal Society B* **287**(1922):20192873 DOI [10.1098/rspb.2019.2873](https://doi.org/10.1098/rspb.2019.2873).
- Sonne J, Zanata TB, Martín González AM, Torres NLC, Fjeldså J, Colwell RK, Tinoco BA, Rahbek C, Dalsgaard B. 2019. The distributions of morphologically specialized hummingbirds coincide with floral trait matching across an Andean elevational gradient. *Biotropica* **51**(2):205–218 DOI [10.1111/btp.12637](https://doi.org/10.1111/btp.12637).
- Stebbins GL. 1970. Adaptive radiation of reproductive characteristics in angiosperms, I: pollination mechanisms. *Annual Review of Ecology and Systematics* **1**(1):307–326 DOI [10.1146/annurev.es.01.110170.001515](https://doi.org/10.1146/annurev.es.01.110170.001515).
- Stiles FG. 1975. Ecology, flowering phenology, and hummingbird pollination of some Costa Rican *Heliconia* species. *Ecology* **56**(2):285–301 DOI [10.2307/1934961](https://doi.org/10.2307/1934961).
- Stiles FG. 1978. Temporal organization of flowering among the hummingbird foodplants of a tropical wet forest. *Biotropica* **10**(3):194–210 DOI [10.2307/2387905](https://doi.org/10.2307/2387905).
- Stiles FG. 1995. Behavioral, ecological and morphological correlates of foraging for arthropods by the hummingbirds of a tropical wet forest. *Condor* **97**(4):853–878 DOI [10.2307/1369527](https://doi.org/10.2307/1369527).
- Sugihara G, Ye H. 2009. Cooperative network dynamics. *Nature* **458**(7241):979–980 DOI [10.1038/458979a](https://doi.org/10.1038/458979a).
- Thomson JD, Wilson P. 2008. Explaining evolutionary shifts between bee and hummingbird pollination: convergence, divergence, and directionality. *International Journal of Plant Sciences* **169**(1):23–38 DOI [10.1086/523361](https://doi.org/10.1086/523361).
- Torices R, Muñoz-Pajares AJ. 2015. PHENIX: an R package to estimate a size-controlled phenotypic integration index. *Applications in Plant Sciences* **3**(5):1400104 DOI [10.3732/apps.1400104](https://doi.org/10.3732/apps.1400104).
- Verdú M, Valiente-Banuet A. 2008. The nested assembly of plant facilitation networks prevents species extinctions. *American Naturalist* **172**(6):751–760 DOI [10.1086/593003](https://doi.org/10.1086/593003).
- Vizentin-Bugoni J, Maruyama PK, Debastiani VJ, Duarte LDS, Dalsgaard B, Sazima M. 2016. Influences of sampling effort on detected patterns and structuring processes of a Neotropical plant-hummingbird network. *Journal of Animal Ecology* **85**(1):262–272 DOI [10.1111/1365-2656.12459](https://doi.org/10.1111/1365-2656.12459).
- Vizentin-Bugoni J, Maruyama PK, Sazima M. 2014. Processes entangling interactions in communities: forbidden links are more important than abundance in a hummingbird-plant network. *Proceedings of the Royal Society B: Biological Sciences* **281**(1780):20132397 DOI [10.1098/rspb.2013.2397](https://doi.org/10.1098/rspb.2013.2397).
- Vázquez DP, Aizen MA. 2003. Null model analyses of specialization in plant-pollinator interactions. *Ecology* **84**(9):2493–2501 DOI [10.1890/02-0587](https://doi.org/10.1890/02-0587).
- Vázquez DP, Blüthgen N, Cagnolo L, Chacoff NP. 2009. Uniting pattern and process in plant-animal mutualistic networks: a review. *Annals of Botany* **103**(9):1445–1457 DOI [10.1093/aob/mcp057](https://doi.org/10.1093/aob/mcp057).
- Vázquez DP, Chacoff NP, Cagnolo L. 2009. Evaluating multiple determinants of the structure of mutualistic networks. *Ecology* **90**(8):2039–2046 DOI [10.1890/08-1837.1](https://doi.org/10.1890/08-1837.1).

- Wagner GP. 1984.** On the eigenvalue distribution of genetic and phenotypic dispersion matrices: evidence for a nonrandom organization of quantitative character variation. *Journal of Mathematical Biology* **21**(1):77–95 DOI [10.1007/BF00275224](https://doi.org/10.1007/BF00275224).
- Waser NM, Chittka L, Price MV, Williams NM, Ollerton J. 1996.** Generalization in pollination systems, and why it matters. *Ecology* **77**(4):1043–1060 DOI [10.2307/2265575](https://doi.org/10.2307/2265575).
- Wolf LL, Stiles FG, Hainsworth FR. 1976.** Ecological organization of a tropical, highland hummingbird community. *Journal of Animal Ecology* **45**(2):349–379 DOI [10.2307/3879](https://doi.org/10.2307/3879).
- Zanata TB, Dalsgaard B, Passos FC, Cotton PA, Roper JJ, Maruyama PK, Fischer E, Schleuning M, Martín González AM, Vizentin-Bugoni J, Franklin DC, Abrahamczyk S, Alárcon R, Araujo Aécia C, Araújo FP, Azevedo-Junior SMde, Baquero AC, Böhning-Gaese K, Carstensen DW, Chupil H, Coelho AG, Faria Rério R, Hořák D, Ingversen TT, Janeček Štěpán, Kohler G, Lara C, Las-Casas FMG, Lopes AV, Machado AO, Machado CG, Machado IC, Maglianesi Mía A, Malucelli TS, Mohd-Azlan J, Moura AC, Oliveira GM, Oliveira PE, Ornelas JF, Riegert J, Rodrigues Léia C, Rosero-Lasprilla L, Rui AM, Sazima M, Schmid B, Sedláček Ořej, Timmermann A, Vollstädt MGR, Wang Z, Watts S, Rahbek C, Varassin IG. 2017.** Global patterns of interaction specialization in bird-flower networks. *Journal of Biogeography* **44**(8):1891–1910 DOI [10.1111/jbi.13045](https://doi.org/10.1111/jbi.13045).