

Analysis of Clinical Isolation Characteristics of Nontuberculous Mycobacteria and Drug Sensitivity of Rapidly Growing Mycobacteria in the General Hospital of Guangzhou, China

Xiaoyi Liu^{1,*}, Zhiwei Lin^{2,*}, Yiwen Li³, Zhiwei Zhong¹, Aiwu Wu¹, Yueting Jiang³

¹KingMed School of Laboratory Medicine, Guangzhou Medical University, Guangzhou, People's Republic of China; ²Department of Clinical Laboratory of the First Affiliated Hospital of Guangzhou Medical University, State Key Laboratory of Respiratory Disease, National Center for Respiratory Medicine, National Clinical Research Center for Respiratory Disease, Guangzhou Institute of Respiratory Health, Guangzhou, People's Republic of China; ³Department of Clinical Laboratory, the Key Laboratory of Advanced Interdisciplinary Studies Center, the First Affiliated Hospital of Guangzhou Medical University, National Center for Respiratory Medicine, Guangzhou, People's Republic of China

*These authors contributed equally to this work

Correspondence: Aiwu Wu; Yueting Jiang, Email aiwwu66@163.com; jyting8899@126.com

Purpose: The clinical distribution characteristics of nontuberculous mycobacteria (NTM) in general hospital were explored to guide the clinical diagnosis and treatment of NTM infection.

Methods: Samples with positive mycobacterium culture in the First Affiliated Hospital of Guangzhou Medical University were collected and identified through PCR. Phenotypic drug sensitivity experiments were conducted on 44 *Mycobacteroides abscessus* isolated from clinical departments with broth microdilution method, and *rrl*, *rrs* and *erm (41)* genes associated with drug resistance were detected.

Results: From September 2020 to July 2023, 314 mycobacterium-positive isolates were separated from patients in the First Affiliated Hospital of Guangzhou Medical University, with 147 (46.8%) NTM isolates were included in our study. The samples were respiratory tract specimens mainly, with 64% bronchoalveolar lavage fluid. Of 144 cases identified, samples were from 133 patients (60 males and 73 females; gender ratio of 0.82:1). NTM was mainly isolated from the people aged 40 and above, especially females ($\chi^2 = 10.688$, $P = 0.014$). *M. abscessus* (61, 42.36%), *M. intracellulare* (35, 24.31%) were the two most NTMs in this hospital. Clinical strains of *M. abscessus* exhibited high resistance to antibiotics, except for cefoxitin (31.8%), linezolid (25.0%), amikacin (0%), and clarithromycin (18.2%). Among 8 strains of *M. abscessus* with clarithromycin acquired resistance, just 4 strains (50.0%) showed mutations (A2270G, A2271G) in *rrl* gene, but a new mutation (C2750T) was detected in 1 strain. Among 14 strains of *M. abscessus* with clarithromycin-induced resistance, 13 (93.0%) strains had T28 *erm (41)* gene and 1 (7.0%) strain had C28 *erm (41)* gene.

Conclusion: *M. avium-intracellulare* complex was gradually becoming predominant strain in Guangzhou area. The resistant situation of *M. abscessus* in general hospital had shown severe. Potential mutation in *rrl* gene associated with clarithromycin acquired resistance of *M. abscessus* were found, but drug-resistant mechanism remained unclear.

Keywords: nontuberculous mycobacterium, drug resistance, clarithromycin, *Mycobacteroides abscessus*

Introduction

Nontuberculous mycobacteria (NTMs) belong to the genus mycobacterium except, the *Mycobacterium tuberculosis* complex (MTBC, including *Mycobacterium bovis*, *Mycobacterium africanum*, and *Mycobacterium tuberculosis*) and *Mycobacterium leprae*, which can exist in natural environments, such as soil, rivers, and human settlements.¹ More than 190 types of NTMs have been discovered (<http://www.bacterio.net/mycobacterium.html>), most of which are parasitic

bacteria and few are conditional pathogens. NTMs can be divided into two categories by growth rate: rapidly growing mycobacteria (RGMs) and slow-growing mycobacteria (SGMs). Common conditionally pathogenic NTMs in clinical settings include *Mycobacterium avium*, *Mycobacterium kansasii*, *Mycobacterium intracellulare*, *Mycobacteroides abscessus*, *Mycobacteroides chelonae*, *Mycobacterium marinum*, and *Mycolcibacterium fortuitum*, which cause lung diseases and are related to infections in the central nervous system, skin, and bone. Patients with acquired immune deficiency syndrome (AIDS) were the high incidence population of NTM disease, and most of them were disseminated NTM disease. *M. chelonae* and *M. fortuitum* were associated with skin, lung, bone, and central nervous system infections. *M. avium* and *M. intracellulare* were the most commonly pathogenic SGM in NTM. And *M. kansasii* was the second most common pathogen causing lung diseases in SGM.^{2–4}

Proportion of NTM clinical isolates varies among locations in the Chinese mainland. In South China, the highest clinical isolation rate of NTM is *M. abscessus*.^{5,6} The top five NTM strains isolated in Guangzhou area from 2018 to 2021 were *Mycobacterium avium–intracellulare* complex, *M. chelonae–abscessus* complex, *M. kansasii*, *M. fortuitum*, and *Mycobacterium gordonae*,⁷ and the highest isolation rate in RGM was obtained in the *M. abscessus* culture. *M. intracellulare* was the most common clinical isolates of NTM in Beijing and Sichuan, China, while the clinical isolation rate of *M. kansasii* in Shanghai, China, was the highest.^{8–10} A statistical study on the distribution of isolate strains in NTM lung disease patients in Europe also revealed that *Mycobacterium avium–intracellulare* complex and *M. kansasii* were the two most common pathogens in the European region, with *Mycobacterium avium–intracellulare* complex being the main pathogen in northern Europe and *M. kansasii* being the main pathogen in southern Europe.¹¹ The isolate quantity of *M. avium* and *M. kansasii* ranked first and second in the NTM of general hospitals in the San Francisco and Virginia, United States.^{12,13} NTM infection is often misdiagnosis as *Mycobacterium tuberculosis* infection due to the biological similarities between NTM and MTBC and inadequate experience in the diagnosis and treatment of NTM infections.

Traditional clinical laboratory tests for NTM, including acid-fast staining smear microscopy and Roche solid medium isolation and culture, are still used in nontuberculous-designated medical institutions. Distinguishing among Mycobacteria species is difficult even in mycobacteria culture. Designated medical institutions for tuberculosis prevention and control are difficult to carry out because of their high prices and rigorous operations; nevertheless, advanced molecular diagnostic technologies can directly identify mycobacterium species.^{14–16} Based on the different situations of NTM isolation across the country and the fact that clinical laboratory testing methods of NTM in general hospitals were neglected, this study combined with the isolation of NTM at the First Affiliated Hospital of Guangzhou Medical University to improve the data and explore the clinical distribution characteristics of NTM in the general hospital and aimed to guide the clinical diagnosis and drug treatment of NTM infection.

Materials and Methods

Study Design and Setting

This is a survey on the epidemiology and drug-sensitive test of NTM in general hospital in Guangzhou, China. The study was set at the First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China, which was one of the general hospitals in Guangzhou area.

The study sample included 147 patients and their clinical specimens between September 2020 and July 2023. The clinical data of the subjects were collected and analyzed retrospectively, including age, gender, medical history, and clinical diagnosis. Based on gender as the independent variable, this study analyzed statistical differences in age and medical history between different genders. Drug sensitive test. And broth microdilution method and polymerase-chain reaction (PCR) were used in drug-sensitive analysis.

Inclusive Criteria

The inclusive criteria were as follows: (1) Positive results within the reporting period (42 days) for bottles cultured on a mycobacterium liquid culture apparatus (BACTEC MGIT 960, BD, America); (2) smear obtained from the culture solution is positive for acid-fast bacteria; (3) Colloidal gold test for detecting the MPT64 antigen showed negative result.

Identification

The NTM positive liquid culture of BACTEC MGIT 960 was inactivated by using 70% ethanol and ultraviolet ray for 2 hours. Then, nucleic acid from the inactivated bacterial solution was extracted and stored at -20°C .

Mycobacterium 16S-23S ribosomal deoxyribonucleic acid (rDNA) internal transcribed spacer sequence (ITS) and ribonucleic acid (RNA) polymerase β subunit coding gene (*rpoB*) were used as the target genes for polymerase-chain reaction (PCR). Two pairs of forward and reverse primers were used (Table 1), and the ITS PCR reaction system was 50 μL , which contained a template (2 μL , 20.4–36.3 $\text{ng}/\mu\text{L}$), forward primer and reverse primer (1.8 μL each), 2 \times Accurate Taq DNA polymerase (25 μL ; Accurate Biology), and ddH₂O (20.4 μL). The amplification program was as follows: 5 min at 94 $^{\circ}\text{C}$; 45 cycles of 45s at 94 $^{\circ}\text{C}$, 1 min at 55 $^{\circ}\text{C}$, and 1 min at 72 $^{\circ}\text{C}$; 10 min at 72 $^{\circ}\text{C}$. The *rpoB* gene PCR reaction system was 50 μL and contained a template (4 μL), forward primer and reverse primer (2.5 μL each), 2 \times Accurate Taq DNA polymerase (25 μL), and ddH₂O (16 μL). The amplification program was as follows: 5 min at 94 $^{\circ}\text{C}$; 35 cycles of 45s at 94 $^{\circ}\text{C}$, 1 min at 52 $^{\circ}\text{C}$, and 1 min at 72 $^{\circ}\text{C}$; 10 min at 72 $^{\circ}\text{C}$. The amplified product was subjected to 130 V and 20 mins of electrophoresis with 2% agarose gel as the carrier.

Primer synthesis and DNA bidirectional sequencing were completed by Shenggong Biotechnology Co., Ltd. (Shanghai, China), and the sequencing primers were the same as the amplification primers. The sequencing results were compared with BLASTN sequences in the NCBI GenBank standard database for the identification of specific bacterial species (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Antimicrobial Drug Sensitivity Testing of RGM

The *Mycobacterium*-positive liquid culture bottles were transferred to Columbia blood agar plates (Autobio Diagnostics, China) and cultured for 5 days in 35 $^{\circ}\text{C}$ in 5% CO₂. Commercial susceptibility test kits (Sensititre RAPMYCOI, ThermoFisher Scientific, America) were used for drug sensitivity tests. The minimum inhibitory concentration (MIC) of 15 antibiotic drugs against the clinical RGM strains were determined using the broth microdilution method according to the Clinical and Laboratory Standards Institute (CLSI) M24 E3 file (2018),¹⁷ and the breakpoints of drugs were identified using the CLSI M62 ED1 file (2018).¹⁸ The procedures in the operating manual were strictly followed.

We designed primers for the *rrl*, *erm* (41), and *rrs* genes were as follows (Table 1), and PCR was performed. The PCR products were then sent to Shenggong Biotechnology Co., Ltd. for sequencing, and BLAST analysis was performed using the NCBI standard database and the standard strain ATCC19977.

Statistical Analysis

Clinical and experimental data were statistically analyzed with IBM SPSS v21.0 software. The comparative analysis of age constituent ratio between age groups used chi-square test or Fisher's exact test. And Wilcoxon rank sum test was used in the comparison of average age between male and female. $P < 0.05$ indicated the statistically significant difference.

Table 1 Target Sequences and Primers for PCR Amplification

Primers	Sequence	Length(bp)
<i>rpoB</i>	F 5'-CACGAGGTGCTGGAAGGA-3' R 5'-CGGGCGATAGCGGAGT-3'	250–750
16S-23S rDNA ITS	F 5'- GAAGTCGTAACAAGG –3' R 5'- CACGAGGTGCTGGAAGGA –3'	250–750
<i>rrl</i>	F 5'- GAATTGAAACGCTGGCACACT –3' R 5'- GGTGTCCTACTCTTCGGTTCC –3'	3464
<i>rrs</i>	F 5'-ACCAACGATGGTGTGTCCAT-3' R 5'-CTTGTCGAACCGCATAACCCT-3'	1727
<i>erm</i> (41)	F 5'- GAAGCTGGCAGGCAACG –3' R 5'- CCGTCGGCACGCAGTA –3'	872

Results

Sample and Clinical Information Analysis

From September 2020 to July 2023, a total of 314 mycobacterium-positive isolates were separated from patients in the First Affiliated Hospital of Guangzhou Medical University, with 147 (46.8%) NTM isolates were included in our study. A total of 144 cases of Mycobacterium infections (68 cases of RGM and 76 cases of SGM) were identified by Sanger sequencing and real-time quantitative PCR (Xiamen Zeesan Biotech, China). The specimens were collected from 85 bronchoalveolar lavage fluids (57.8%), 21 bronchoscopy sputum (14.3%), 12 lung tissues (8.2%), 6 sputum samples (4.1%), 4 pus samples (2.7%), 3 deep sputum samples (2.0%), 3 secretions (2.1%), 2 urine samples (1.3%), and 11 other specimens (7.5%; [Figure 1](#)).

A total of 144 NTM samples were collected from 133 patients (60 males, average age of 62.0 ± 18.8 years; 73 females, average age of 55.5 ± 12.7 years) with a gender ratio of 0.82:1 ([Table 2](#)). A total of 70 patients were diagnosed with NTM infection, and the proportion of NTM positive patients diagnosed clinically with NTM infection reached 51.9%. There was a statistical difference in the NTM separation rate among different age groups and between genders ($\chi^2 = 10.688$, $P = 0.014$). Male patients were mainly over 60 years old, whereas female patients were mainly 40–60 years old. However, there was no statistical difference in average age between males and females in the Wilcoxon rank sum test ($Z = -1.172$, $P = 0.241$). Patients previously diagnosed with NTM or tuberculosis were more likely to be diagnosed with NTM infections again or clinically isolated with NTM (39/135, 28.9%) ([Table 3](#)).

Identification of NTM

A total of 144 cases were identified ([Table 4](#)). A total of 61 *M. abscessus* (42.36%), 35 *M. intracellulare* (24.31%), 16 *M. avium* (11.11%), 5 *Mycobacterium colombiense* (3.47%), 10 *M. kansasii* (6.94%), 3 *M. fortuitum* (2.08%), and *M. scrofulaceum* (2/1.39%), and *M. phocaicum* (2/1.39%) specimens were identified.




Figure 1 Clinical isolation of Mycobacterium from September 2020 to July 2023 at the First Affiliated Hospital of Guangzhou Medical University (a). Sample types of 147 NTM isolates (b). NTM, nontuberculous mycobacteria.
Abbreviation: BALF, bronchoalveolar lavage fluid.

Table 2 NTM Positive Cases Between Different Age Groups

Age (Years Old)	Male	Female	Total
≤20	3 (2.26%)	0 (0.00%)	3 (2.26%)
21–40	6 (4.51%)	10 (7.52%)	16 (12.03%)
41–60	16 (12.03%)	35 (26.32%)	51 (38.35%)
>60	35 (26.32%)	28 (21.05%)	63 (47.37%)
Total	60 (45.11%)	73 (54.89%)	133 (100.00%)

Abbreviation: NTM, nontuberculous mycobacteria.

Table 3 Medical History Statistics of NTM Positive Cases

Medical History	Male	Female	Total
Previously diagnosed with NTM infection or tuberculosis	15	24	39
Lung cancer	5	7	12
Lung transplantation	9	1	10
COPD	7	1	8
Smoke	20	0	20
Other tumors/cancers	2	0	2
Accident injury	2	2	4

Abbreviations: NTM, nontuberculous mycobacteria; COPD, chronic obstructive pulmonary disease.

Table 4 Identification of NTM in the First Affiliated Hospital of Guangzhou Medical University

Strains	Number	Percentage (%)	Classification
<i>M. abscessus</i>	61	42.36	RGM
<i>M. intracellulare</i>	35	24.31	SGM
<i>M. avium</i>	16	11.11	SGM
<i>M. colombiense</i>	5	3.47	SGM
<i>M. kansasii</i>	10	6.94	SGM
<i>M. fortuitum</i>	3	2.08	RGM
<i>M. phocaicum</i>	2	1.39	RGM
<i>M. scrofulaceum</i>	2	1.39	SGM
<i>M. senegalense</i>	1	0.69	RGM
<i>M. arupense</i>	1	0.69	SGM
<i>M. szulgai</i>	1	0.69	SGM
<i>M. asiaticum</i>	1	0.69	SGM
<i>M. simiae</i>	1	0.69	SGM
others	5	3.47	/
Total	144	100	

Abbreviations: NTM, nontuberculous mycobacteria; RGM, rapidly growing mycobacteria; SGM, slowly growing mycobacteria.

Results of Antimicrobial Drug Sensitivity Testing

As the NTM with the highest proportion in our study, we completed phenotypic drug sensitivity tests on 44 clinical strains of *M. abscessus*, and detected a total of 7 categories and 15 clinical drugs used for against RGM infection: β -lactam, aminoglycosides, macrolides, quinolones, sulfonamides, tetracyclines, and oxazolidinone (Tables 5 and 6). The drug resistance rates of ciprofloxacin, moxifloxacin, minocycline and doxycycline had reached over 80%; resistance rates of imipenem and tobramycin exceeded 60%; *M. abscessus* shown good sensitivities to ceftazidime, linezolid, amikacin and clarithromycin. However, after extending the cultivation time to 14 days of clarithromycin, rates of susceptible and intermediate were lowered and the increase in drug resistance rate from 18.2% to 50.0%.

Table 5 Results of Drug Sensitivity Tests on 44 Strains of *Mycobacteroides abscessus*

Drugs	Numbers of Isolates		
	Susceptible	Intermediate	Resistance
CIP	1 (2.3%)	4 (9.1%)	39 (88.6%)
IMI	1 (2.3%)	12 (27.3%)	31 (70.4%)
MXF	2 (4.5%)	4 (9.1%)	38 (86.4%)
TOB	1 (2.3%)	14 (31.8%)	29 (65.9%)
MIN	2 (4.5%)	3 (6.8%)	39 (88.7%)
DOX	3 (6.8%)	4 (9.1%)	37 (84.1%)
SXT	9 (20.5%)	/	35 (79.5%)
FOX	1 (2.3%)	29 (65.9%)	14 (31.8%)
LZD	27 (61.4%)	6 (13.6%)	11 (25.0%)
AMI (IV)	44 (100%)	0	0
CLA (5 days)	35 (79.5%)	1 (2.3%)	8 (18.2%)
CLA (14 days)	20 (45.5%)	2 (4.5%)	22 (50.0%)
FEP	–	–	–
AXO	–	–	–
TGC	–	–	–
AUG2	–	–	–

Abbreviations: CIP, ciprofloxacin; IMI, imipenem; MXF, moxifloxacin; TOB, tobramycin; MIN, minocycline; DOX, doxycycline; SXT, trimethoprim-sulfamethoxazole; FOX, cefoxitin; LZD, linezolid; AMI (IV), amikacin (intravenous); CLA, clarithromycin; FEP, cefepime; AXO, ceftriaxone; TGC, tigecycline; AUG2, amoxicillin/potassium clavulanate 2:1. Cefepime, ceftriaxone, tigecycline, amoxicillin/potassium clavulanate 2:1 had no breakpoints in CLSI M62 EDI file (2018).¹⁸ Minocycline's breakpoints referred about *Nocardia* spp. in CLSI M62 EDI file (2018).¹⁸

Table 6 Number and Distribution of *Mycobacteroides abscessus* Drugs Concentration

Drugs	Range	MIC50	MIC90	No. of Strains at Different Drug Concentrations											
				0.12	0.25	0.5	1	2	4	8	16	32	64	128	
CIP	0.12–4	>4	>4	0	0	0	1	4	39	/	/	/	/	/	
IMI	2–64	64	>64	/	/	/	/	0	1	4	8	4	27	/	
MXF	0.25–8	>8	>8	/	0	0	2	4	8	30	/	/	/	/	
TOB	1–16	8	16	0	0	0	0	1	14	21	8	0	0	0	
MIN	1–8	>8	>8	/	/	/	2	1	2	39	/	/	/	/	
DOX	0.12–16	>16	>16	0	0	1	2	2	2	1	36	/	/	/	
SXT	0.25/4.75–8/152	8/152	>8/152	/	1	1	3	4	12	23	/	/	/	/	
FOX	4–128	64	128	/	/	/	/	/	0	0	1	13	16	14	
LZD	1–32	8	32	/	/	/	0	0	11	16	6	11	0	0	
AMI	1–64	4	8	0	0	0	0	9	20	13	2	0	/	/	
CLA ^a	0.06–16	0.5	16	8	10	6	5	3	1	1	7	/	/	/	
CLA ^b	0.06–16	>16	>16	6	4	6	2	1	2	0	22	/	/	/	
FEP	1–32	>32	>32	/	/	/	0	0	0	0	2	42	/	/	
AXO	4–64	>64	>64	/	/	/	/	/	0	0	0	0	44	/	
TGC	0.015–4	0.5	2	1	8	13	13	8	0	/	/	/	/	/	
AUG2	2/1–64/32	>64/32	>64/32	/	/	/	/	0	0	0	0	0	44	/	

Notes: ^aCLA for 5 days. ^bCLA for 14 days. The drug concentration unit was µg/mL. There were also 3 strains of clarithromycin (5 days) and 1 strain of clarithromycin (14 days) with MIC <0.06 µg/mL.

Abbreviations: CIP, ciprofloxacin; IMI, imipenem; MXF, moxifloxacin; TOB, tobramycin; MIN, minocycline; DOX, doxycycline; SXT, trimethoprim-sulfamethoxazole; FOX, cefoxitin; LZD, linezolid; AMI (IV), amikacin (intravenous); CLA, clarithromycin; FEP, cefepime; AXO, ceftriaxone; TGC, tigecycline; AUG2, amoxicillin/potassium clavulanate 2:1. MIC₅₀, the minimum drug concentration for inhibiting 50% of the experimental strain; MIC₉₀, the minimum drug concentration for inhibiting 90% of the experimental strain.

Clarithromycin Resistance Gene Detection in *Mycobacteroides abscessus*

Two mechanisms underlie the resistance of *M. abscessus* to clarithromycin. One was a point mutation in the *rml* gene encoding peptidyltransferase at sites 2058 or 2059 (*Escherichia coli* numbering). Drug sensitivity results showed resistance within 3–5 days.¹⁹ We sequenced the *rml* gene related to the clarithromycin-acquired resistance of *M. abscessus* in 8 clinical strains with MIC for clarithromycin of ≥ 8 $\mu\text{g/mL}$ in 3–5 days. Three strains (No.237–1, 237–2 and 262) showed a mutation in the *rml* gene at the A2271G site, and 1 strain (No.295) showed the same mutation at the A2270C site. The mutation sites of the *rml* gene in a drug-resistant strain (No.206) were 837T deletion and C2750T, and the mutation sites of the same gene in 1 strain (No.188) were G17A and insertion of G at 1288 and 1303 sites (Figure 2). 2 drug-resistant strains (Nos.142 and 284) had no mutations in the *rml* gene. Nucleotides 2016–2625 (*E. coli* numbering) were aligned in the domain V regions in the *rml* gene, and thus position 2750, which was near the peptidyl transferase center in the No.206 strain, was likely associated with clarithromycin resistance.

The other underlying mechanism was the adenine methylation at site 2058 of the *rml* gene. This process was attributed to erythromycin methylase encoded by functional *erm* (41) genes.²¹ Drug sensitivity results required long-term cultivation to show resistance. The No. 28 base pairs of the *erm* (41) gene exhibit polymorphism with two sequence types: T28 and C28. We also detected the *erm* (41) gene that can induce the drug resistance of *M. abscessus* on 44 clinical isolates in our study. Except for 1 clinical strain that failed to detect the *erm* (41) gene, among the remaining 43 strains, a total of 17 strains (39.5%) had the T28 *erm* (41) gene, consistent with the drug-sensitive phenotype. The MIC of clarithromycin was ≥ 16 $\mu\text{g/mL}$ when cultivation was extended to 14 days, and 23 sensitive strains (53.5%) of the T28 *erm* (41) gene showed sequence deletions at 61–62 and 156–429 nt sites. Three sensitive strains (7.0%) had the C28 *erm* (41) gene and exhibited sensitive phenotype after 14 days of cultivation.

Moreover, the sensitivity rate of amikacin was 100%, and T1373A, A1375G, C1376T, and G1458T mutations were not found in the *rrs* gene.

Discussion

The clinical isolation rate of NTM from the First Affiliated Hospital of Guangzhou Medical University in 2020–2023 was 46.8%, increasing from 39.13% in 2021 to 45.65% in 2022 (Figure 1). The rate was higher than that in Guangzhou Chest Hospital in 2018–2021 (35.45%).⁷ We isolated 68 strains (47.2%) of RGM and 76 strains (52.8%) of SGM, which were consistent with the number in previous research performed in Guangzhou area.^{7,20} In Guangzhou, the clinical isolation rate of *M. abscessus* (61/42.36%) was the highest, followed by the clinical isolation




Figure 2 Base deletion, substitution, and insertion in the *rml* gene of No. 206 (a and b). Alignment with *rml* gene of ATCC19977 and No. 206 (c and d).

rates of *M. intracellulare* (35/24.31%), *M. avium* (16/11.11%), and *M. colombiense* (5/3.47%). These rates are inconsistent with those reported in Nanjing, Chengdu, and Shanghai.^{10,22,23} Compared with the clinical isolates of NTM in Guangzhou 5 years ago,²⁴ the *M. avium* complex including *M. avium*, *M. intracellulare*, and *M. colombiense* isolates was higher in number, and the proportion of *M. intracellulare* and *M. colombiense* has considerably increased. Therefore, the epidemic status and clinical distribution characteristics of the *M. avium* complex in Guangzhou in the past 5 years should be examined.

In Guangzhou, NTMs were common in middle-aged and elderly people, especially in women, unlike in Sichuan and Chongqing.^{25–27} Patients who had previously been diagnosed with tuberculosis or NTM infections were more likely to possess NTMs because of multidrug resistance of NTMs, especially *M. abscessus* complex, to first-line antituberculosis drugs.²⁸ NTMs had high resistance to most first-line antituberculosis drugs, and common pathogenic strains of *M. intracellulare*, *M. avium*, and *M. chelonae* exhibit multidrug-resistant characteristics and are resistant to at least eight antibiotics.^{29,30} Thus, drugs were usually ineffective against NTM infections, which eventually recur. Drug sensitivity tests should be conducted timely on patients with multidrug-resistance NTMs.

We identified 44 clinical RGM strains through a phenotypic drug sensitivity test. *M. abscessus* was less sensitive to conventional anti-NTM infections drugs, and the sensitivity of *M. abscessus* isolate strains from Guangzhou to linezolid was only 61.4%, and the resistance rate was up to 25.0%, which was much higher than that reported in the Beijing general hospital from 2012 to 2015 (9.7%).³¹ This difference may be attributed to two reasons. First, linezolid was commonly used to treat bacterial lung infections, especially those caused by methicillin-resistant *Staphylococcus aureus*. Patients often come into contact with this drug and are prone to develop drug-resistant strains.³² Second, linezolid could be used in the treatment of drug-resistant pulmonary tuberculosis disease and extrapulmonary tuberculosis disease,³³ and most strains in this study were isolated from patients who had been infected with *M. tuberculosis* or NTMs, indicating that previous antibiotic treatment was ineffective and may have increased linezolid resistance. Fortunately, first-line drugs for treating NTM infections, such as amikacin and clarithromycin, showed low resistance rates. However, the resistance rate of *M. abscessus* for clarithromycin reached 50.0% at prolonged cultivation time. Identifying clarithromycin resistance targets in *M. abscessus* is more important than identifying amikacin resistance targets, and strains that produce acquired and inducible resistance are needed for the identification of resistance genes and genotypes. In this study, we discovered a new mutation in the *rrl* gene. This mutation was related to the clarithromycin resistance of *M. abscessus* in one clinical strain,³⁴ and the proportion of common mutation sites was 50.0% (4/8). Other strains without resistance mutation sites exhibited different resistance mechanisms that need to be further explored. The MIC_{50s} and MIC_{90s} of β -lactam drugs were higher than those of other antibiotics in *M. abscessus* possibly because of the highly active β -lactamase (Bla_{Mab}) encoded by the MAB_2875 gene and the thick cell wall of *M. abscessus* with a large amount of lipids.^{35,36} Furthermore, although tigecycline had no drug breakpoints in CLSI guidelines, we determined the MICs for *M. abscessus*, which can further improve the experimental data for reference.

However, samples were collected only from the First Affiliated Hospital of Guangzhou Medical University and the samples from other general hospitals were not included in the study, which cannot fully reflect the NTM isolation in other general hospitals in Guangzhou. Moreover, the MPT64 antigen test results of the specimens included in this study were all negative, but the MPT64 antigen of *Mycobacterium tuberculosis* complex was positive. So, the possibility of mixed infection of *Mycobacterium tuberculosis* complex and NTM was not considered. For this study, it was still necessary to expand the sample size, improve the research data required for the analysis of clinical isolation characteristics of NTM in Guangzhou general hospitals, and systematically explain the prevalence of NTM in the Guangzhou area.

Conclusion

The clinical isolation rates of NTMs in the First Affiliated Hospital of Guangzhou Medical University were higher than the Guangzhou Chest Hospital. NTMs might be isolated more easily in female patients and people who had been diagnosed with tuberculosis or NTM infections. The isolation rate of *M. avium-intracellulare* complex had shown a significant upward trend, gradually becoming a dominant strain in Guangzhou. The drug resistance rates of

M. abscessus from the general hospital in Guangzhou had increased, especially to linezolid and clarithromycin. Common drug-resistant mutation sites in the *rrl* gene accounted for a small proportion of the resistance.

Ethics

This study had been reviewed by the Medical Ethics Committee of the First Affiliated Hospital of Guangzhou Medical University and had obtained ethical permission, with ethics number 2022121.

Acknowledgments

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Halstrom S, Price P, Thomson R. Review: environmental mycobacteria as a cause of human infection. *Int J Mycobacteriol*. 2015;4(2):81–91. doi:10.1016/j.ijmyco.2015.03.002
2. Griffith DE, Akshamit T, Brown-Elliott BA, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med*. 2007;175(4):367–416. doi:10.1164/rccm.200604-571ST
3. Lai CC, Hsueh PR. Diseases caused by nontuberculous mycobacteria in Asia. *Fut Microbiol*. 2014;9(1):93–106. doi:10.2217/fmb.13.138
4. Oatway WH. Nontuberculous mycobacteria and associated diseases. *Am Rev Respir Dis*. 1979;120(6):1389–1390. doi:10.1164/arrd.1979.120.6.1389
5. Zhou L, Xu D, Liu H, et al. Trends in the prevalence and antibiotic resistance of non-tuberculous mycobacteria in Mainland China, 2000–2019: systematic review and meta-analysis. *Front Public Health*. 2020;8:295. doi:10.3389/fpubh.2020.00295
6. Liu CF, Song YM, He WC, et al. Nontuberculous mycobacteria in China: incidence and antimicrobial resistance spectrum from a nationwide survey. *Infect Diseases Poverty*. 2021;10(1):59. doi:10.1186/s40249-021-00844-1
7. Chen H, Lai X, Chen P, et al. Analysis of the identification results of non-tuberculous clinical isolates of mycobacterium tuberculosis in Guangzhou from 2018 to 2021. *J Pra Med*. 2022;38(17):2191–2197.
8. Huang JJ, Li YX, Zhao Y, et al. Prevalence of nontuberculous mycobacteria in a tertiary hospital in Beijing, China, January 2013 to December 2018. *BMC Microbiol*. 2020;20(1):158. doi:10.1186/s12866-020-01840-5
9. Wu J, Zhang Y, Li J, et al. Increase in nontuberculous mycobacteria isolated in Shanghai, China: results from a population-based study. *PLoS One*. 2014;9(10):e109736. doi:10.1371/journal.pone.0109736
10. Wang DM, Liu H, Zheng YL, Xu YH, Liao Y. Epidemiology of nontuberculous mycobacteria in tuberculosis suspects, Southwest of China, 2017–2022. *Front Cell Infect Microbiol*. 2023;13:1282902.
11. Dahl VN, Laursen LL, He Y, Zhang YA, Wang MS. Species distribution among patients with nontuberculous mycobacteria pulmonary disease in Europe. *J Infect*. 2023;87(5):469–472. doi:10.1016/j.jinf.2023.03.010
12. Sheu LC, Tran TM, Jarlsberg LG, Marras TK, Daley CL, Nahid P. Non-tuberculous mycobacterial infections at San Francisco General Hospital. *Clin Respir J*. 2015;9(4):436–442. doi:10.1111/crj.12159
13. Mullen B, Houghton ER, Colston J, et al. Geographic variation and environmental predictors of nontuberculous mycobacteria in laboratory surveillance. *Emerg Infect Dis*. 2024;30(3):548–554. doi:10.3201/eid3003.231162
14. Baliga S, Murphy C, Sharon L, et al. Rapid method for detecting and differentiating Mycobacterium tuberculosis complex and non-tuberculous mycobacteria in sputum by fluorescence in situ hybridization with DNA probes. *Int J Infect Dis*. 2018;75:1–7. doi:10.1016/j.ijid.2018.07.011
15. Karami-Zarandi M, Bahador A, Gizaw Feysia S, et al. Identification of non-tuberculosis mycobacteria by line probe assay and determination of drug resistance patterns of isolates in Iranian patients. *Arch Razi Inst*. 2019;74(4):375–384. doi:10.22092/ari.2019.127144.1372
16. Yao L, Gui X, Wu X, et al. Rapid identification of nontuberculous mycobacterium species from respiratory specimens using nucleotide MALDI-TOF MS. *Microorganisms*. 2023;11(8):1975. doi:10.3390/microorganisms11081975
17. CLSI. Susceptibility Testing of Mycobacteria, Nocardia spp, and Other Aerobic Actinomycetes. 3rd ed. *CLSI Standard M24*. Clinical and Laboratory Standards Institute; 2018.
18. CLSI. Performance Standards for Susceptibility Testing of Mycobacteria, Nocardia spp, and Other Aerobic Actinomycetes. 1st ed. *CLSI Supplement M62*. Clinical and Laboratory Standards Institute; 2018.
19. Wallace RJ, Meier A, Brown BA, et al. Genetic basis for clarithromycin resistance among isolates of Mycobacterium chelonae and Mycobacterium abscessus. *Antimicrob Agents Chemother*. 1996;40(7):1676–1681. doi:10.1128/AAC.40.7.1676
20. Liu D, Zheng H, He W, et al. Analysis of the identification of non-tuberculous mycobacteria clinical isolates and drug sensitivity test results in Guangzhou area. *Chin J Antitubercul*. 2019;41(05):534–540.
21. Bastian S, Veziris N, Roux AL, et al. Assessment of clarithromycin susceptibility in strains belonging to the Mycobacterium abscessus group by erm(41) and rrl sequencing. *Antimicrob Agents Chemother*. 2011;55(2):775–781. doi:10.1128/AAC.00861-10
22. Zhang R, Luo S, Wang N, Zhang H, Wu X. Epidemiology of nontuberculous mycobacteria in Nanjing and MAB_0540 mutations associated with clofazimine resistance in mycobacterium abscessus. *Infect Drug Resist*. 2023;16:2751–2764. doi:10.2147/IDR.S408986
23. Ying R, Yang J, Wu X, Yu F, Sha W. Antimicrobial susceptibility testing using the MYCO test system and mic distribution of 8 drugs against clinical isolates of nontuberculous mycobacteria from Shanghai. *Microbiology Spectrum*. 2023;11(2):e0254922. doi:10.1128/spectrum.02549-22

24. Chen H, Hu J, Chen P, et al. Epidemic status and distribution characteristics of non-tuberculous mycobacteria in Guangzhou, 2018–2019. *J Molec Imag.* 2021;44(02):378–382.
25. Luo J, Yu QJ, Lin YL, Yi FL, Xiong L, Xie Y. Clinical characteristics of nontuberculous mycobacterium infection Cases in Sichuan, China in 2016–2021: a Retrospective Study. *Sichuan Da Xue Xue Bao Yi Xue Ban.* 2022;53(5):890–895. doi:10.12182/20220960503
26. Zhang H, Luo M, Zhang K, et al. Species identification and antimicrobial susceptibility testing of non-tuberculous mycobacteria isolated in Chongqing, Southwest China. *Epidemiol Infect.* 2020;149:e7.
27. Dedrick RM, Abad L, Storey N, et al. The problem of Mycobacterium abscessus complex: multi-drug resistance, bacteriophage susceptibility and potential healthcare transmission. *Clin Microbiol Infect.* 2023;29(10):1335.e9–1335.e16. doi:10.1016/j.cmi.2023.06.026
28. Fujiwara K, Uesugi F, Furuuchi K, et al. Minimum inhibitory concentrations before and after antibacterial treatment in patients with mycobacterium abscessus pulmonary disease. *Microbiology Spectrum.* 2021;9(3):e0192821. doi:10.1128/Spectrum.01928-21
29. Li F, Li GL, Pang H, et al. Preliminary study on drug susceptibility profile and resistance mechanisms to macrolides of clinical isolates of non-tuberculous mycobacteria from China. *Biomed Environ Sci.* 2018;31(4):290–299.
30. Li G, Pang H, Guo Q, et al. Antimicrobial susceptibility and MIC distribution of 41 drugs against clinical isolates from China and reference strains of nontuberculous mycobacteria. *Internat J Antimicrob Age.* 2017;49(3):364–374. doi:10.1016/j.ijantimicag.2016.10.024
31. Li YM, Tong XL, Xu HT, Ju Y, Cai M, Wang C. Prevalence and antimicrobial susceptibility of mycobacterium abscessus in a General Hospital, China. *Biomed Environ Sci.* 2016;29(2):85–90. doi:10.3967/bes2016.009
32. Kato H, Hagihara M, Asai N, et al. Meta-analysis of vancomycin versus linezolid in pneumonia with proven methicillin-resistant *Staphylococcus aureus*. *J Global Antimicrob Resist.* 2021;24:98–105. doi:10.1016/j.jgar.2020.12.009
33. Chinese Medical Association. [Consensus on linezolid in the treatment of tuberculosis(2022 update)]. *Zhonghua jie he he hu xi za zhi.* 2022;45(10):988–995. doi:10.3760/cma.j.cn112147-20220320-00220
34. Li B, Yang S, Chu H, et al. Relationship between antibiotic susceptibility and genotype in mycobacterium abscessus clinical isolates. *Front Microbiol.* 2017;8:1739. doi:10.3389/fmicb.2017.01739
35. Story-Roller E, Maggioncalda EC, Cohen KA, Lamichhane G. Mycobacterium abscessus and β -Lactams: emerging Insights and Potential Opportunities. *Front Microbiol.* 2018;9:2273. doi:10.3389/fmicb.2018.02273
36. Luthra S, Rominski A, Sander P. The role of antibiotic-target-modifying and antibiotic-modifying enzymes in mycobacterium abscessus drug resistance. *Front Microbiol.* 2018;9:2179. doi:10.3389/fmicb.2018.02179

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>