

RESEARCH ARTICLE

Open Access

Improved mycobacterial protein production using a *Mycobacterium smegmatis* *groEL1*ΔC expression strain

Elke E Noens^{1*}, Chris Williams¹, Madhankumar Anandhakrishnan¹, Christian Poulsen², Matthias T Ehebauer¹ and Matthias Wilmanns¹

Abstract

Background: The non-pathogenic bacterium *Mycobacterium smegmatis* is widely used as a near-native expression host for the purification of *Mycobacterium tuberculosis* proteins. Unfortunately, the Hsp60 chaperone GroEL1, which is relatively highly expressed, is often co-purified with polyhistidine-tagged recombinant proteins as a major contaminant when using this expression system. This is likely due to a histidine-rich C-terminus in GroEL1.

Results: In order to improve purification efficiency and yield of polyhistidine-tagged mycobacterial target proteins, we created a mutant version of GroEL1 by removing the coding sequence for the histidine-rich C-terminus, termed GroEL1ΔC. GroEL1ΔC, which is a functional protein, is no longer able to bind nickel affinity beads. Using a selection of challenging test proteins, we show that GroEL1ΔC is no longer present in protein samples purified from the *groEL1*ΔC expression strain and demonstrate the feasibility and advantages of purifying and characterising proteins produced using this strain.

Conclusions: This novel *Mycobacterium smegmatis* expression strain allows efficient expression and purification of mycobacterial proteins while concomitantly removing the troublesome contaminant GroEL1 and consequently increasing the speed and efficiency of protein purification.

Background

Heterologous expression of recombinant proteins in *Escherichia coli* can result in the production of insoluble inclusion bodies. Recent statistics show that less than half of the *M. tuberculosis* (Mtb) proteins expressed in *E. coli* are soluble [1]. Therefore, the non-pathogenic bacterium *Mycobacterium smegmatis* is often used as an alternative, more closely related host for the expression of mycobacterial proteins. Furthermore, *M. smegmatis* may also provide mycobacterium-specific chaperones, which can help correct folding of Mtb proteins [1].

During nickel affinity purification, it has been observed that a protein of 56 kDa is co-purified with polyhistidine-tagged recombinant proteins while using *M. smegmatis* as an expression system. This contaminant was previously identified as the Hsp60 chaperone

GroEL1 of *M. smegmatis* [1-3]. The protein sequence of GroEL1 shows a histidine-rich C-terminus (7 out of 11 amino acids are histidines), which is likely to be the reason for the observed nickel sepharose binding [1,2].

Unlike most other bacteria, mycobacteria possess two Hsp60 chaperone *groEL* genes, one of which is arranged in the bicistronic *groESL* operon [4]. *M. smegmatis* also encodes a third Hsp60 protein (Msmeg1978), which is more distantly related to GroEL1 (Msmeg1583) and GroEL2 (Msmeg0880) [3]. Although *groEL1* of *M. smegmatis* can be found in the same operon as *groES*, an arrangement indispensable for the chaperone function in bacteria, its histidine-rich tail is distinct from the more typical glycine-methionine-rich C-terminal region found in GroEL2 [3]. Furthermore, *groEL2* is an essential gene and exists in all actinobacteria, in contrast to *groEL1* [3,5]. Recently, it has been shown that *groEL2* and *groES* are expressed more strongly than *groEL1*, which might have arisen from a difference in stability of the predicted post-transcriptionally cleaved mRNAs for

* Correspondence: e.noens@embl-hamburg.de

¹European Molecular Biology Laboratory (EMBL), Hamburg Outstation, c/o DESY, Building 25a, Notkestraße 85, 22603 Hamburg, Germany
Full list of author information is available at the end of the article

groES and *groEL1* [5]. Consistent with the current chaperone model in mycobacteria, one chaperone, here GroEL2, would act as the main house keeping chaperone in *M. smegmatis*, with the other chaperones (GroEL1 and Msmeg1978) adopting more specialised functions. Indeed, GroEL1 of *M. tuberculosis* was recently identified as being associated with nucleotides, suggesting a role as a DNA chaperone, while GroEL1 of *M. smegmatis* was found to have a role in mycolic acid biosynthesis during biofilm formation [5,6,3].

The co-purification of GroEL1 with histidine-tagged recombinant proteins can be particularly problematic since native GroEL1 is expressed at relatively high levels, meaning that in the case of a low yield of recombinant protein, GroEL1 may well compete with the protein of interest for binding sites on nickel affinity beads. Minimal sample manipulation is recommended during protein purification to improve efficiency. Therefore, additional steps required to remove GroEL1 can result in a significant loss of the protein of interest.

In this article, we describe an *M. smegmatis* expression strain containing a mutant version of GroEL1, termed GroEL1ΔC, which consists of a *groEL1* gene without a coding sequence for the histidine-rich C-terminal tail. We show that GroEL1ΔC is a functional protein, which no longer co-purifies when using nickel affinity purification and we provide evidence that proteins purified from this strain are correctly folded, active and that they behave identically to those purified from the original expression strain. Taken together, our data demonstrate that *M. smegmatis groEL1ΔC* is a competent protein expression strain, which allows the efficient removal of the troublesome contaminant GroEL1 without the requirement of additional purification steps.

Methods

Bacterial strains and media

The *E. coli* strains DH5α (Invitrogen) and HB101 (Pro-mega) were used for cloning of expression constructs and the target substrate to generate the mutant version of *groEL1* using standard procedures [7]. Transformants were selected in Luria Broth containing the appropriate antibiotics.

M. smegmatis mc²155 was used as the parent (wild type) strain for the *groEL1ΔC* strain. Both *M. smegmatis* strains were maintained in Middlebrook 7H9 or 7H10 medium supplemented with 0.2% (v/v) glycerol, 10% ADC, 0.05% (v/v) tween-80 and the appropriate antibiotics.

For biofilm formation, 10 ml of biofilm media was inoculated with 10 μl of saturated culture and incubated at 30°C without disturbance [3,8].

For the expression of the recombination proteins in *M. smegmatis* in order to create the mutant form of

groEL1, 0.2% succinate (w/v) was added as a carbon source to 7H9 medium supplemented with 0.2% (v/v) glycerol, 0.05% (v/v) tween and the appropriate antibiotics. Expression of his-tagged recombinant proteins in *M. smegmatis* was performed in 7H9 medium supplemented with 0.2% (w/v) glucose as carbon source. Acetamide was added to a final concentration of 0.2% (w/v) at 0.5 OD₆₀₀ and at 2.5 OD₆₀₀ for the expression of the recombination proteins and his-tagged recombinant proteins, respectively.

Plasmids, constructs and oligonucleotides

All plasmids and constructs are summarised in Table 1 and oligonucleotides are listed in Table 2. pJV53 was used to express the recombination proteins [9]. pYUB854 was used for the preparation of the target substrate to create the *groEL1ΔC* strain [10]. pGH542, harbouring a δγ resolvase, was used to generate an unmarked deletion [11]. Using the primer pairs Msmeg1583-F1.2 & Msmeg1583-R1 and Msmeg1583-F2 & Msmeg1583-R2.1, two 500 bp fragments, homologous to the fragments +1067/+1587 and +1621/+2176 relative to the translational start of Msmeg1583, were amplified

Table 1 Plasmids and constructs used in this study

Plasmid/ construct	Description	Reference
pJV53	Che9c recombination proteins under control of the acetamidase promoter in pLAM12	[9]
pYUB854	Hyg ^R cassette flanked by γδ-res sites and 2 MCSs	[10]
pGH542	Expressing an γδ resolvase and tetracycline resistant	[11]
pEN15	pYUB854 with a 520 bp fragment harbouring <i>groEL1</i> (+1067/+1587, relative to <i>groEL1</i>) inserted upstream of the Hyg ^R cassette and a 555 bp fragment downstream of <i>groEL1</i> including the STOP codon of <i>groEL1</i> , inserted downstream of the Hyg ^R cassette	This paper
pMyNT	Mycobacterial overexpression vector	Geerlof et al., unpublished data
pMyNT/ PrcA-B	Rv2109-2110 in pMYNT, Rv2110 is N-terminally his-tagged	[12]
pMyNT/ AccD5E5	Rv3280-3281 in pMYNT. Only his-tagged Rv3280 seems to express using this construct.	This paper
pMyNT/ AccA3	Rv3285 in pMyNT	This paper
pMyNT/ CFP10- ESAT6	Rv3874-3875 in pMYNT, Rv3874 is N-terminally his-tagged	[12]
pMyNT/ ACPS	Rv2523 in pMYNT	This paper

Table 2 Primers used in this study

Primer	Sequence (5'-3')	Location 5'	Relative to
Msmeg1583-F1.2	GCGC CTTAAG CGACTGGGATCGCGAGAAGCTGC	+1067	Msmeg1583
Msmeg1583-R1	GCGC TCTAGA CTCGTCTCGTCGGCCGGCTTG	+1587	Msmeg1583
Msmeg1583-F2	GCGC AAGCTT GATCCATTTACGCGACACCCCC	+1620	Msmeg1583
Msmeg1583-R2.1	GCGC ACTAGT GGTGTTCGATCGTCTGGCCGATG	+2176	Msmeg1583
accD5E5-F	GATC TCATGA GTATGACAAGCGTTACC G	+1	Rv3280
accD5E5-R	GTCA AAGCTT TTATCGGCGCATGTGCG	+2161	Rv3280
accA3-F	GATC CCATGG GTATGGCTAGTCACGCC	+2	Rv3285
accA3-R	GTCA AAGCTT TTACTTGATCTCGGCGAGC	+1803	Rv3285
Rv2523-F	CATG CCATGG GCATCGTCCGGTGTGGGG	+1	Rv2523
Rv2523-R	CCC AAGCTT ACGGGGCCTCCAGGATGGC	+391	Rv2523

Restriction sites are presented in bold face. CTTAAG = *EcoRI*, TCTAGA = *XbaI*, CCATGG = *NcoI*, TCATGA = *BspHI*. AAGCTT = *HindIII*, ACTAGT = *SpeI*.

and subsequently ligated *AflIII-XbaI* (F1.2-R1) and *HindIII-SpeI* (F2-R2.1) into pYUB854, creating pEN15.

For the expression of *M. tuberculosis* proteins in *M. smegmatis*, the pMyNT expression vector was used [Geerlof *et al.*, unpublished data]. pMyNT/ACPS, pMyNT/AccA3 and pMyNT/AccD5 were made as follows: PCR was performed with primer pair Rv2523-F & Rv2523-R for ACPS, accA3-F & accA3-R for AccA3 and accD5E5-F & accD5E5-R for AccD5 and the resulting fragments were digested with *NcoI-HindIII* and inserted into *NcoI-HindIII* digested pMyNT.

Creation of the *groEL1ΔC* mutant

The *groEL1ΔC* mutant was created using the mycobacterial recombineering method [9]. pEN15 was digested with *AflIII* and *SpeI* to create the linear target substrate, which was introduced into mc²155 electrocompetent cells, expressing the recombinase genes on pJV53 and in this way creating hygromycin-resistant transformants. The hygromycin-resistance cassette was removed using $\delta\gamma$ resolvase, expressed on pGH542, generating an unmarked deletion [11].

Southern blot analysis

Genomic DNA (5ug) was isolated as described [9], digested with the appropriate enzymes, separated on a 0.9% agarose gel and transferred to a positively charged nylon membrane (Roche). For DNA probe labelling, hybridisation and detection, the DIG high prime DNA labelling and detection starter kit 1 (Roche) was used.

Growth curves

Bacterial growth was followed by measuring the optical densities at a wavelength of 600 nm as a function of time. Cultures were prepared with 7H9 expression medium (0.2% (w/v) glucose as carbon source) in identical triplicates for each strain. Duplicate samples were taken

every 4 hours for 40 hours. When the optical density at 600 nm exceeded 1.5, samples were diluted in order to remain within the linear range of the detector.

Protein expression and purification

All methods related to protein expression in *M. smegmatis* were carried out as described [12,13]. Protein-protein complexes from operon-encoded proteins were expressed using the native operon structure [9]. In brief, pellets from 500 ml cultures were dissolved in 30 ml lysis buffer containing 50 mM Tris-HCl pH 8.0, 300 mM NaCl, 0.5 M urea with protease inhibitor cocktail (Sigma) and 1 mg/ml DNase I (Serva). Resuspended cells were sonicated four times, each for 5 min (with a 0.3 s pulse and 0.7 s rest) at 5 min intervals to prevent overheating, using a Bandelin VW3200 probe at 45% amplitude. The supernatant was collected after centrifugation (30,000 × g) for 1 h at 4°C, filtered through a 0.44 μm filter and loaded onto a nickel affinity sepharose (NiAC) column. After washing with 10 column volumes of 50 mM Tris-HCl pH 8.0, 300 mM NaCl and 20 mM imidazole, proteins were eluted in 50 mM Tris-HCl, 100-150 mM NaCl and 250-500 mM imidazole and subjected to size exclusion chromatography using either a Superdex 75 (16/60) column (GE Healthcare) or, for large protein complexes, a Superose 6 (10/300) (GE Healthcare) with 25 mM Tris-HCl pH 8.0, 150 mM NaCl and 1 mM DTT as buffer. The collected protein samples were analysed by SDS-PAGE and concentrated accordingly.

Circular Dichroism (CD) spectrum analysis

CD measurements were performed on a Jasco J-810 spectropolarimeter. Prior to measurement, samples were dialysed into 10 mM potassium phosphate, 150 mM NaCl, pH 7.4. Spectra were recorded between 182 and 260 nm in a 2 mm cuvette with machine settings as

follows: 1 nm bandwidth, 1 sec response, 1 nm data pitch, 100 nm/min scan speed, cell length of 0.1 cm. Each curve presented is the average of three separate measurements.

Coupled enzyme assay

Enzymatic activity of the AccD5-AccA3 complex was estimated by a coupled enzyme assay that follows the rate of ATP hydrolysis spectrophotometrically [14]. The production of ADP during the reaction was coupled to pyruvate kinase and lactate dehydrogenase, and the oxidation of NADH was probed at 340 nm. The assay mixture contained 7 units of pyruvate kinase, 10 units of lactate dehydrogenase, 50 mM NaHCO₃, 3 mM ATP, 0.5 mM phosphoenol pyruvate, 0.2 mM NADH, 0.3 mg/ml BSA, 100 mM K₂HPO₄ pH 7.6 and 5 mM MgCl₂ and varying concentrations of propionyl-coenzyme A. Reactions were initiated by the addition of enzyme to the assay mixture and were maintained at 30°C. Data were acquired using a Tecan infinite M1000 microplate reader. The kinetic parameters K_m and V_{max} were determined by fitting the mean velocities *versus* the substrate concentration to the Michaelis-Menten equation of enzyme kinetics using nonlinear regression analysis, executed by the program Prism 5 (GraphPad Software™).

Results and Discussion

Creation of the *groEL1ΔC* strain


Currently, the role of GroEL1 in protein folding is uncertain. A closer look at the structure of *E. coli* GroEL [15] indicates that, although the C-terminal region of the protein is not easily accessible, pointing towards the central cavity of the wheel-like structure adopted by oligomeric GroEL, the extreme C-terminal 20 amino acids are absent from the model. Similarly, the GroEL structure of *Paracoccus denitrificans* also lacks these residues [16]. These observations suggest that the C-terminal region of GroEL is highly flexible and could reach out of the central cavity, allowing in this way *M. smegmatis* GroEL1 to bind nickel affinity beads. Additionally, as native GroEL1 from *M. tuberculosis* is oligomeric [17], nickel binding would require only one accessible histidine-rich region. Therefore, we decided to change only the last eleven amino acids of the protein, rather than to make a full knock out strain, in order to minimise changes to the expression strain. A precise chromosomal deletion of fragment 1588-1620, relative to the translational start of *groEL1* (Msmeg1583), was created using the mycobacterial recombineering technique [9] (Figure 1A). Southern hybridisation (Figure 1B) was used to verify that correct homologous recombination had taken place in the hygromycin-resistant and the unmarked deletion strain (Figure 1B). The latter strain, in which the hygromycin resistance cassette has been

removed, has the C-terminal eleven residues of GroEL1, containing seven histidines, replaced by six non-histidine residues, which are part of the “scar” sequence left behind after removal of the resistance cassette (Figure 1A). The stop codon of this recombinant version of GroEL1 is TAA, which although rare, is recognised in high G+C *mycobacteria* [18]. This unmarked deletion strain, referred to as *M. smegmatis groEL1ΔC*, is used in all further experiments.

Ojha *et al.* reported that the last 18 amino acids of GroEL1 are essential for the formation of mature biofilms [3]. Therefore, to test the functionality of the GroEL1ΔC protein, we compared biofilm formation in both the wild type and *groEL1ΔC* strains. Both strains were able to form mature biofilms after an incubation time of 7 days at 30°C, indicating that GroEL1ΔC is indeed fully functional (Figure 2A). Taking into account the data from Ojha *et al.*, our results could suggest that either the amino acids important for biofilm formation are upstream of those removed in the GroEL1ΔC protein, or that removal of the last 18 residues may affect the folding of at least a part of GroEL1. Additionally, as this newly created strain was constructed for the overexpression of mycobacterial proteins, its growth in 7H9 expression medium was compared to the original expression strain *M. smegmatis mc²155* (Figure 2B). We observed no significant differences in growth between the two strains, with both reaching an OD₆₀₀ of between 2.5-3.0 after approximately 18 hours, at which time expression is usually induced.

GroEL1ΔC is absent during nickel affinity purification of proteins expressed in *M. smegmatis groEL1ΔC*

To demonstrate the absence of GroEL1ΔC as a contaminant when using the *M. smegmatis groEL1ΔC* expression strain, we determined the expression and purification efficiency of our strain in comparison to the wild type strain using five different constructs, representing a variety of different protein molecules, including the mycobacterial proteasome, the CFP10-ESAT6 complex, the AccD5-AccA3 dodecameric acyl-CoA carboxylase complex and the holo-acyl-carrier protein synthase (for details, see Table 3). Additionally, we also used the empty pMyNT vector, to check for GroEL1 binding in the absence of a his-tagged protein. All constructs were transformed into both *M. smegmatis mc²155* and *groEL1ΔC* and the resulting transformants were cultured in 7H9 expression medium and induced by the addition of acetamide to a final concentration of 35 mM. Eighteen hours after induction, the cells were collected by centrifugation, lysed and the soluble protein fraction was passed over a nickel affinity column, with the elution fraction being analysed by SDS-PAGE (Figure 3). While GroEL1 was visible in samples purified



from *M. smegmatis* mc²155 (Figure 3, lanes a), the protein was noticeably absent in five out of six protein samples isolated from the *groEL1ΔC* strain (Figure 3, lanes b). Due to the fact that AccD5 has a similar size to GroEL1, we were unable to determine its presence or absence in samples of the purified acyl-CoA carboxylase complex by SDS-PAGE. Therefore, samples isolated from gel (Figure 3) were analyzed by mass spectrometry (Additional file 1). While numerous peptides from both GroEL1 and AccD5 could be identified from gel slices deriving from the mc²155 strain, only AccD5 peptides could be detected in the sample obtained

from the *groEL1ΔC* strain (Additional file 1). Likewise, MALDI-TOF mass spectrometry was performed on the other protein samples, verifying the absence of GroEL1 peptides in the protein samples derived from *M. smegmatis groEL1ΔC* (data not shown).

Proteins purified from *M. smegmatis groEL1ΔC* behave identically to those purified from the wild type strain
M. smegmatis encodes three forms of the Hsp60 chaperone GroEL: Msmeg1583 (GroEL1), Msmeg0880 (GroEL2) and Msmeg1978. However, the precise molecular function of each protein remains unclear.




Figure 2 Biofilm formation and growth rates of *M. smegmatis* mc² 155 and *M. smegmatis* *groEL1ΔC* are comparable. (A) Both *M. smegmatis* mc² 155 (WT) and *M. smegmatis* *groEL1ΔC* strains are able to form biofilms after an incubation time of 7 days at 30°C. (B) Growth curve of *M. smegmatis* mc²155 (WT = black) and *M. smegmatis* *groEL1ΔC* strains (grey) in 7H9 expression medium. The arrow represents the typical time of induction in *M. smegmatis*.

Changing the last 18 amino acids of GroEL1 does not alter growth but does result in a strong defect in biofilm formation [3]. To confirm that the newly created recombinant version of GroEL1 has no effect on the correct folding and, ultimately, the function of the proteins expressed in *M. smegmatis* *groEL1ΔC*, a number of different proteins and protein complexes have been expressed and analysed.

In the previous section, we have shown that it is possible to express and purify potentially challenging protein complexes, such as the proteasome complex PrcA-B and the CFP10-ESAT6 complex, from the recombinant *groEL1ΔC* strain. These data imply that the proteins isolated from the *groEL1ΔC* strain are correctly folded, since we were able to observe all components after purification. In both examples, complex formation

Table 3 List of test proteins used to validate the *groEL1ΔC* expression strain

ORF	Annotation	Description	Expressed ...	Mol. Mass (kDa)
Rv2109c	PrcA	α - and β -subunit of the mycobacterial proteasome ($\alpha_7\beta_7\beta_7\alpha_7$ subunit organisation)	Using native operon content, producing a 730 kDa multimeric complex	26.8
Rv2110c	PrcB			30.3
Rv3285	AccA3	α - and β -subunit from acyl-CoA carboxylase AccD5-AccA3 complex ($\alpha_3\beta_3\beta_3\alpha_3$ subunit organisation)	As monomeric proteins, mixed to form a acyl-CoA carboxylase complex of 740 kDa	63.8
Rv3280	AccD5			59.4
Rv3874	CFP10	Potential virulence factor CFP10-ESAT6 complex	Using native operon content, producing a heterodimeric (1:1) complex	10.8
Rv3875	ESAT6			9.9
Rv2523c	ACPS	Holo-acyl-carrier protein synthase	As monomeric protein	14




Figure 3 GroEL1ΔC is absent from protein samples purified from *M. smegmatis* groEL1ΔC. SDS-PAGE analysis of protein samples isolated from cells expressing either CFP10-ESAT6 heterodimer (1), a holo-acyl-carrier protein synthase (ACPS) (2), the β- and α-subunit D5 (3) and A3 (4) of the acyl-CoA carboxylase AccD5-AccA3 complex, the PrcA-B complex (5) or the empty expression vector (empty pMyNT) (6). **a**, proteins expressed in *M. smegmatis* mc²155; **b**, proteins expressed in *M. smegmatis* groEL1ΔC. Purified proteins of interest are labelled with #. Visible presence of GroEL1 is depicted with *. Squares identify the bands excised for peptide mass fingerprinting. GroEL1 = 56 kDa.

requires direct protein-protein interactions between subunits of the complex as only one subunit is his-tagged.

Taking our analysis one step further, we directly tested the structural and functional properties of proteins isolated from the *groEL1ΔC* strain. We used the five expression constructs described above and transformed them into both *M. smegmatis* mc²155 and *groEL1ΔC*. Proteins were expressed and purified using a nickel affinity column as described above. AccD5 and AccA3 protein samples were mixed in a 1:1 stoichiometry to form the high-molecular-weight AccD5-AccA3 complex. Size exclusion chromatography was performed on all samples as a final purification step.

Circular dichroism (CD) spectroscopy is a powerful tool used to visualise the secondary structure properties of protein samples. We observed that the four protein samples isolated from *groEL1ΔC* gave virtually identical CD spectra to those purified from the wild type strain (Figure 4), implying that they are correctly folded. Furthermore, the CD spectra of the CFP10-ESAT6 complexes, showing a protein with high helical content, are comparable to those collected previously [12] and are in line with the X-ray structure, which consists of a four-helical bundle complex (PDB ID: 3FAV) [12].

Additionally, we have demonstrated carboxylase activity of the acyl-CoA carboxylase AccD5-AccA3 complex,

isolated from *groEL1ΔC*, using an enzyme-coupled reaction (Figure 5). Using propionyl-CoA as a substrate, AccD5-AccA3 showed carboxylase activity with a $K_m = 0.1301 \pm 0.0198$ mM and a $V_{max} = 1.333 \pm 0.049$ mM $\text{min}^{-1} \text{mg}^{-1}$, data which are similar to the parameters determined using the AccD5-AccA3 complex isolated from *E. coli* [19], indicating that the AccD5-AccA3 complex isolated from *groEL1ΔC* is a functional carboxylase. Carboxylase activity requires the α-subunit of the carboxylase to be post-translationally biotinylated [19], implying that the subunits of this large megasynthase are folded correctly and, in the case of the α-subunit, correctly post-translationally modified, when isolated from *groEL1ΔC*.

Conclusions

We have developed an *M. smegmatis* expression strain that allows efficient expression and purification of mycobacterial proteins, multi-subunit protein complexes and post-translationally modified proteins while concomitantly removing the troublesome contaminant GroEL1 and consequently increasing the speed and efficiency of protein purification. The *M. smegmatis* *groEL1ΔC* strain is particularly suitable for laboratories performing *in vitro* activity assays and structural studies on mycobacterial proteins and protein complexes.




Figure 4 Proteins isolated from both strains give virtually identical CD spectra. CD spectra of the multimeric proteasome complex PrcA-B (A), the dodecameric acyl-CoA carboxylase AccD5-AccA3 complex (B), CPF10-ESAT6 heterodimer (C), and monomeric protein ACPS (D) expressed in *M. smegmatis* mc²155 (WT = black) and *M. smegmatis* *groEL1ΔC* (grey) are virtually identical. For A and B, a concentration between 170 and 200 nM was used while for C and D, concentrations were between 5 and 10 μM.




Figure 5 Kinetics of AccD5-AccA3 isolated from *M. smegmatis* *groEL1ΔC*. Carboxylation activity of the acyl-CoA carboxylase AccD5-AccA3 complex isolated from *groEL1ΔC* was measured using an enzyme-coupled reaction with propionyl-CoA as substrate, providing $K_m = 0.1469 \text{ mM}$ and $V_{\text{max}} = 28.5 \text{ mOD/min}$.

Additional material

Additional file 1: GroEL1 is absent from an AccD5 protein sample derived from *M. smegmatis* *groEL1ΔC*. Results of peptide mass fingerprinting analysis of samples excised from SDS-PAGE gel (Figure 3, boxes). Shown in red are the peptides that could be identified. (a) Sample derived from *M. smegmatis* mc²155. (b) Sample derived from *M. smegmatis* *groEL1ΔC*.

Abbreviations

PCR: Polymerase chain reaction; kDa: kilo Dalton; Hsp60: Heat shock protein 60; ADC: Albumine-dextrose-catalase; DMSO: dymethylsulfoxide; NiAc: Nickel affinity sepharose column; SDS-PAGE: sodium dodecyl sulfate polyacrylamide gel electroporesis; MALDI-TOF: matrix-assisted laser desorption/ionization reflection time-of-flight.

Acknowledgements and Funding

We thank Arie Geerlof for the pMyNT expression vector, Young-Hwa Song for her contribution in the early stages of the work, the Proteomics Core Facility of EMBL Heidelberg for performing peptide mass fingerprinting, the

Mandelkow Lab (Max Planck Institute for Structural Molecular Biology, Hamburg, Germany) for access to their circular dichroism spectroscope. CW is funded by a Rubicon post-doctoral fellowship (825.08.023) from the Netherlands organization for scientific research (NWO). ME is funded by an EMBO long-term fellowship (ALTF-7272008). The project has been supported by grants awarded to MW from BMBF (Pathogenomik Plus PTJ-BIO 0313801L), from the European Commission Framework VII (NATT, 222965 and SystemTB, 241587) and from the DFG (SPP1170, WI 1058/6-3).

Author details

¹European Molecular Biology Laboratory (EMBL), Hamburg Outstation, c/o DESY, Building 25a, Notkestraße 85, 22603 Hamburg, Germany. ²The Scripps Research Institute, La Jolla, 92037, USA.

Authors' contributions

EN and CP designed the study. EN made the *groEL1ΔC* strain, tested its functionality and growth, expressed and purified all proteins described except the AccD5-AccA3 complex and wrote the manuscript. CW carried out the CD measurements, provided technical assistance and participated in writing the manuscript. MA carried out all experiments concerning AccD5-AccA3. CP provided expression constructs. ME participated in testing the strain's growth and feasibility. MW organized the funding, supervised the work and helped revising the manuscript. All authors read and approved the final manuscript.

Received: 17 December 2010 Accepted: 25 March 2011

Published: 25 March 2011

References

1. Goldstone RM, Moreland NJ, Bashiri G, Baker EN, Lott JS: **A new Gateway® vector and expression protocol for fast and efficient recombinant protein expression in *Mycobacterium smegmatis***. *Protein Expr Purif* 2008, **57**:81-87.
2. Poulsen C, Ahkter Y, Jeon AH, Schmitt-Ulms G, Meyer HE, Stühler K, Wilmanns M, Song YH: **Proteome-wide identification of mycobacterial pupylation targets**. *Mol Syst Biol* 2010, **6**:386-394.
3. Ojha A, Anand M, Bhatt A, Kremer L, Jacobs WR Jr, Hatfull GF: **GroEL1: a dedicated chaperone involved in mycolic acid biosynthesis during biofilm formation in *Mycobacteria***. *Cell* 2005, **123**:861-873.
4. Rinke de Wit TF, Bekelie S, Osland A, Miko TL, Hermans PW, van Soelingen D, Drijfhout JW, Schönningh R, Janson AA, Thole JE: ***Mycobacteria* contain two *groEL* genes: the second *Mycobacterium leprae* *groEL* gene is arranged in an operon with *groES***. *Mol Microbiol* 1992, **6**(14):1995-2007.
5. Rao T, Lund PA: **Differential expression of the multiple chaperonins of *Mycobacterium smegmatis***. *FEMS Microbiol Lett* 2010, **310**:24-31.
6. Basu D, Khare G, Singh S, Tyagi A, Khosla S, Mande SC: **A novel nucleoid-associated protein of *Mycobacterium tuberculosis* is a sequence homolog of GroEL**. *Nucleic Acids Res* 2009, **37**(15):4944-4954.
7. Sambrook J, Fritsch EF, Maniatis T: **Molecular cloning: a laboratory manual**. Cold Spring harbor: Cold Spring harbor laboratory press; 1989.
8. Recht J, Kolter R: **Glycopeptidolipid acetylation affects sliding motility and biofilm formation in *Mycobacterium smegmatis***. *J Bacteriol* 2001, **183**:5718-5724.
9. van Kessel JC, Hatfull GF: **Recombineering in *Mycobacterium tuberculosis***. *Nat Methods* 2007, **4**(2):147-152.
10. Bardarov S, Bardarov S Jr, Pavelka Ms Jr, Sambandamurthy V, Larsen M, Tufariello J, Chan J, Hatfull G, Jacobs WR Jr: **Specialized transduction: an efficient method for generating marked and unmarked targeted gene disruptions in *Mycobacterium tuberculosis*, *M. bovis* BCG and *M. smegmatis***. *Microbiology* 2002, **148**:3007-3017.
11. Piuiri M, Hatfull GF: **A peptidoglycan hydrolase motif within the mycobacteriophage TM4 tape measure protein promotes efficient infection of stationary phase cells**. *Mol Microbiology* 2006, **62**:1569-1585.
12. Poulsen C, Holton S, Geerlof A, Wilmanns M, Song YH: **Stoichiometric protein complex formation and over-expression using the prokaryotic native operon structure**. *FEBS Lett* 2010, **584**:669-674.
13. Daugelat S, Kowall J, Matthow J, Bumann D, Winter R, Hurwitz R, Kaufmann SH: **The RD1 proteins of *Mycobacterium tuberculosis*: expression in *Mycobacterium smegmatis* and biochemical characterization**. *Microbes Infect* 2003, **5**:1082-1095.

14. Diacovich L, Peiru S, Kurth D, Rodriguez E, Podesta F, Khosla C, Gramajo H: **Kinetic and Structural Analysis of a New Group of Acyl-CoA Carboxylases Found in *Streptomyces coelicolor* A3(2)**. *J Biol Chem* 2002, **277**:31228-31236.
15. Xu Z, Horwich AL, Sigler PB: **The crystal structure of the asymmetric GroEL-GroES-(ADP)7 chaperonin complex**. *Nature* 1997, **388**(6644):741-50.
16. Fukami TA, Yohda M, Taguchi H, Yoshida M, Miki K: **Crystal structure of chaperonin-60 from *Paracoccus denitrificans***. *J Mol Biol* 2001, **312**(3):501-9.
17. Kumar CM, Khare G, Srikanth CV, Tyagi AK, Sardesai AA, Mande SC: **Facilitated oligomerization of mycobacterial GroEL: Evidence for phosphorylation-mediated oligomerization**. *J Bacteriol* 2009, **191**:6525-6538.
18. Andersson SGE, Sharp PM: **Codon usage in the *Mycobacterium tuberculosis* complex**. *Microbiology* 1996, **142**:915-925.
19. Gago G, Kurth D, Diacovich L, Tsai SC, Gramajo H: **Biochemical and structural characterization of an essential acyl Coenzyme A carboxylase from *Mycobacterium tuberculosis***. *J Bacteriol* 2006, **188**:477-486.

doi:10.1186/1472-6750-11-27

Cite this article as: Noens et al.: Improved mycobacterial protein production using a *Mycobacterium smegmatis* *groEL1ΔC* expression strain. *BMC Biotechnology* 2011 **11**:27.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

