



Elsevier has created a [Monkeypox Information Center](#) in response to the declared public health emergency of international concern, with free information in English on the monkeypox virus. The Monkeypox Information Center is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its monkeypox related research that is available on the Monkeypox Information Center - including this research content - immediately available in publicly funded repositories, with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the Monkeypox Information Center remains active.

## Letter to the Editor

### Monkeypox is a global public health emergency: The role of repurposing cholesterol lowering drugs not to be forgotten

On July 23, 2022, World Health Organization (WHO) declared the current monkeypox outbreak a Public Health Emergency of International Concern (PHEIC).<sup>1</sup> Less than a week later New York and San Francisco declared a state of emergency due to an alarming rise in the number of monkeypox cases.<sup>2, 3</sup> By July 29, 2022; 22, 141 cases of monkeypox had been reported globally in the countries that have not been reported this disease historically.<sup>4</sup> Of these cases 4,906 were reported from the USA.

As a preventive measure effective smallpox vaccination is currently offered in many countries as Pre-Exposure Prophylaxis for risk groups to prevent monkeypox but it is not widely available and vaccination programs have only just been initiated.<sup>5</sup> Thus, the number of monkeypox patient cases will continue to rise. There are several antiviral medications including cidofovir, brincidofovir, and tecovirimat to treat monkeypox.<sup>6</sup> Of these, the most effective medication against monkeypox is tecovirimat which inhibits the function of a major envelope protein required for the production of extracellular virus. Tecovirimat thus prevents the mature virus from leaving an infected cell. The recommended duration of tecovirimat treatment is 14 days.<sup>6</sup> Currently there is very little data of the effectiveness of tecovirimat treatment against monkeypox although several clinical trials are ongoing.<sup>6, 7</sup>

It can be expected that clinical trials regarding effectiveness of tecovirimat will last very long. So, at the current situation repurposing of cholesterol lowering drugs as adjuvants to tecovirimat, including statins, PCSK9 inhibitors and fenofibrates is an attractive strategy and has already been applied in the COVID-19 pandemic.<sup>8-10</sup>

The interest to repurposing cholesterol lowering drugs, particularly statins, is best known among COVID-19 patients.<sup>11</sup> Moreover, there are several examples of this repurposing strategy regarding other severe infections. Statins have been repurposed in the treatment of influenza and Ebola.<sup>12</sup> With Ebola infection, statins may result in the production of fusion-inefficient Ebola virus particles.<sup>13</sup> PCSK9 inhibitors have been used to treat Dengue virus infection because PCSK9 inhibitors enhance secretion of type I inter-

ferons.<sup>14</sup> Fenofibrate has been shown to decrease mortality and morbidity among Japanese encephalitis patients.<sup>15</sup> Compared to Japanese encephalitis vaccination Sehkal and coauthors mention the potential usefulness of fenofibrate especially when treating endemic infections.<sup>15</sup>

While currently there is clearly a lack of studies repurposing cholesterol lowering drugs for the treatment of Monkeypox more attention needs to be given how such viral infections effect the re-distribution of cellular cholesterol.<sup>16</sup> For example, the efficient endosomal re-distribution of cholesterol is essential for viral life cycles. If this cholesterol re-distribution can be disrupted the viral life cycles may be inhibited or even collapse.<sup>16</sup> Fenofibrate also seems to have this ability, at least *in vitro*.<sup>17</sup> Taken together, monkeypox appears to become yet another global public health problem, and, as such, it is a reminiscent of the current COVID-19 pandemic and its predecessors. Therefore, the potential beneficial role of repurposing cholesterol lowering drugs should not be forgotten in this newly emerging viral outbreak. Patients with severe hypercholesterolemia should not stop taking cholesterol-lowering medication after contracting Monkeypox. If tecovirimat medication is started for Monkeypox, simvastatin should be changed to another statin, analogously when treating SARS-CoV-2 infection with Paxlovid.<sup>18</sup>

## Declaration of Competing Interest

AV has received consultancy fees Amgen and Novartis. PTK has received consultancy fees, lecture honoraria, and/or travel fees from Amgen, Novartis, Raisio Group, and Sanofi. FR has received research grants, honoraria, or consulting fees for professional input and/or lectures from Sanofi, Regeneron, Amgen, and Novartis.

Alpo Vuorio, MD\*

*Mehiläinen Airport Health Centre, Vantaa, Finland  
University of Helsinki, Department of Forensic Medicine,  
Helsinki, Finland*

Frederick Raal, PhD

*Faculty of Health Sciences, University of Witwatersrand,  
Johannesburg, South Africa*

Petri T. Kovanen, MD  
 Wihuri Research Institute, Helsinki, Finland

<https://doi.org/10.1016/j.jaACL.2022.08.003>

\*Correspondence author.  
 E-mail address: [alpo.vuorio@gmail.com](mailto:alpo.vuorio@gmail.com) (A. Vuorio)

Received August 4, 2022  
 Accepted August 16, 2022

## References

1. Nuzzo JB, Borio LL. The WHO declaration of monkeypox as a global public health emergency. *JAMA*. 2022 Published online July 27. doi:10.1001/jama.2022.12513.
2. New York State. 8-29-2022. Declaring a disaster in the State of New York. [governor.ny.gov/executive-order/no-20-declaring-disaster-state-new-york](http://governor.ny.gov/executive-order/no-20-declaring-disaster-state-new-york)
3. City and County of San Francisco. 8-28-2022. Office of the Mayor. San Francisco to declare local public health emergency for monkeypox. [sfmayor.org/article/san-francisco-declare-local-public-health-emergency-monkeypox](http://sfmayor.org/article/san-francisco-declare-local-public-health-emergency-monkeypox)
4. Centers for Disease Control and Prevention. 8-29-2022. Monkeypox outbreak global map. [cdc.gov/poxvirus/monkeypox/response/2022/world-map.html](https://cdc.gov/poxvirus/monkeypox/response/2022/world-map.html)
5. Centers for Disease Control and Prevention. 8-28-2022. Considerations for monkeypox vaccination. [cdc.gov/poxvirus/monkeypox/considerations-for-monkeypox-vaccination.html](https://cdc.gov/poxvirus/monkeypox/considerations-for-monkeypox-vaccination.html)
6. Siegrist EA, Sassine J. Antivirals with activity against monkeypox: a clinically oriented review. *Clin Infect Dis*. 2022;ciac622. doi:10.1093/cid/ciac622.
7. Adler H, Gould S, Hine P, et al. Clinical features and management of human monkeypox: a retrospective observational study in the UK. *Lancet Infect Dis*. 2022;22:1153–1162 Erratum in: *Lancet Infect Dis*. 2022 Jul;22(7):e177. Erratum in: *Lancet Infect Dis*. 2022 Jul;22(7):e177. doi:10.1016/S1473-3099(22)00228-6.
8. Vuorio A, Kovanen PT. Statins as adjuvant therapy for COVID-19 to calm the stormy immunothrombosis and beyond. *Front Pharmacol*. 2021;11:579548. doi:10.3389/fphar.2020.579548.
9. Vuorio A, Kovanen PT. PCSK9 inhibitors for COVID-19: an opportunity to enhance the antiviral action of interferon in patients with hypercholesterolaemia. *J Intern Med*. 2021;289:749–751. doi:10.1111/joim.13210.
10. Vuorio A, Brinck J, Kovanen PT. Continuation of fibrate therapy in patients with metabolic syndrome and COVID-19: a beneficial regime worth pursuing. *Ann Med*. 2022;54:1952–1955. doi:10.1080/07853890.2022.2095667.
11. Talasaz AH, Sadeghipour P, Aghakouchakzadeh M, et al. Investigating lipid-modulating agents for prevention or treatment of COVID-19: JACC state-of-the-art review. *J Am Coll Cardiol*. 2021;78:1635–1654. doi:10.1016/j.jacc.2021.08.021.
12. Fedson DS. Treating the host response to emerging virus diseases: lessons learned from sepsis, pneumonia, influenza and Ebola. *Ann Transl Med*. 2016;4:421. doi:10.21037/atm.2016.11.03.
13. Lee J, Kreutzberger AJB, Odongo L, et al. Ebola virus glycoprotein interacts with cholesterol to enhance membrane fusion and cell entry. *Nat Struct Mol Biol*. 2021;28:181–189. doi:10.1038/s41594-020-00548-4.
14. Gan ES, Tan HC, Le DHT, et al. Dengue virus induces PCSK9 expression to alter antiviral responses and diseases outcomes. *J Clin Invest*. 2020;130:5223–5234. doi:10.1172/JCI1137536.
15. Sehgal N, Kumawat KL, Basu A, Ravindranath V. Fenofibrate reduces mortality and precludes neurological deficits in survivors in murine model of Japanese encephalitis viral infection. *PLoS One*. 2012;7:e35427. doi:10.1371/journal.pone.0035427.
16. Glitscher M, Hildt E. Endosomal cholesterol in viral infections - a common denominator? *Front Physiol*. 2021;12:750544. doi:10.3389/fphys.2021.750544.
17. Glitscher M, Martiñ DH, Woytinek K, et al. Targeting cholesterol metabolism as efficient antiviral strategy against the hepatitis E virus. *Cell Mol Gastroenterol Hepatol*. 2021;12:159–180. doi:10.1016/j.jcmgh.2021.02.002.
18. Vuorio A, Kovanen PT, Raal F. Cholesterol-lowering drugs for high-risk hypercholesterolemia patients with COVID-19 while on Paxlovid™ therapy. *Future Virol*. 2022 Ahead of print 2. doi:10.2217/fvl-2022-0060.