

# Awareness of varicella-zoster virus among undergraduate students at the University of Namibia

Nestor Tomas, Festus Kampanza

Department of General Nursing Science, School of Nursing, Faculty of Health Sciences, University of Namibia (UNAM), Rundu, Namibia

## Abstract

Although varicella-zoster virus (VZV) is preventable and curable, the disease remains a public health problem in Namibia. Access to vaccination in the country remains a challenge for many citizens including students. No previous study has been conducted to assess awareness on the prevention and control of VZV in Namibia. The overall purpose of the study was to investigate the awareness of hostel students on prevention and control measures for VZV in Namibia. The objective of the study was to assess and describe the awareness of hostel students on the prevention and control measures for VZV in Namibia. A quantitative descriptive study using an online questionnaire was used. Simple random sampling was used to select 165 respondents. Descriptive statistics and a non-parametric Kruskal Wallis test were performed to test the mean differences between variables and the significance of the data. A great proportion of the respondents, 70.7% (n=118), were aware that VZV is caused by a virus (0.71±0.45) while 76% (n=127) (0.76±0.42) were aware that VZV can be vaccinated against. The study found a strong association between aetiology (p=0.03), available information (p=0.00), mode of transmission (p=0.02), and knowledge on contagious diseases (p=0.03). The results were inconclusive as to whether the level of awareness is high or low, owing to the differences in scores for each variable. The study recommends that a mobile clinic be established on campus to offer health education information.

## Introduction

The varicella-zoster virus (VZV) causes chickenpox, which is an acute and highly contagious disease that results in blister-like rashes, itching, fatigue and fever.<sup>1</sup> Varicella occurs worldwide and, in the absence of a vaccination programme, affects nearly every person by mid-adulthood. Prior to the

introduction of a vaccine, VZV afflicted about 4 million people a year, led to more than 10 500 hospitalisations, and about 100 to 150 deaths.<sup>2</sup> Although the death rate is relatively low (1 in 60.000 cases) it may lead to serious complications, namely, viral and bacterial pneumonia, meningitis, encephalitis, seizures, skin infections and coma.<sup>3</sup> The epidemiology of the disease differs between temperate and tropical climates but is common in developing countries where poor sanitation and vaccination are challenges.<sup>4</sup>

In developing countries generally, the status of varicella infection and vaccination is unclear. Some Middle East countries have reported on varicella burden, seroprevalence, complications and the high cost of medical care and hospitalisation. To develop a vaccination protocol and appropriate preventive health care measures against diseases in different countries, it is very important to know the seroprevalence of any disease for an individual country.<sup>5</sup>

The African continent has several risk factors that could result in an increased burden of VZV disease; these include the high prevalence of HIV/AIDS, primary infections among the elderly mainly found in tropical countries, and diabetes prevalence. These factors may overstretch the already challenged healthcare systems in Africa, preventing the management of the complications of many diseases, including VZV. Furthermore, VZV causes varicella and herpes zoster. Most available data on VZV epidemiology are from temperate, industrialised countries and cannot be used to guide decisions on an immunisation policy against VZV in Africa, given Africa's limited resources and healthcare system challenges.<sup>6</sup> Every decade since the 1940s, health policymakers, professionals and providers have launched new global and national initiatives in an attempt to address the health challenges and needs of populations, particularly those living in sub-Saharan Africa. However, only a few reforms have been successful.<sup>4</sup>

A systematic review of twenty articles from 13 African countries revealed that VZV seroprevalence is high among adults and co-morbidity with VZV is common. Furthermore, there is a lack of quality data to develop VZV control programmes, including vaccination, in Africa.<sup>6</sup> In comparison to developed countries where vaccination is available, in developing countries such as Namibia, VZV vaccination is not provided at public health facilities, while the private sector charges exorbitant fees, thus excluding 85% of the population who depend on public health services. Hence, many VZV victims present with severe con-

Correspondence: Nestor Tomas, Department of General Nursing Science, School of Nursing, Faculty of Health Sciences, University of Namibia (UNAM), Rundu, Namibia. Tel. 264-66-268-6009. E-mail: ntomas@unam.na

Key words: Awareness; students; prevention; control; varicella-zoster virus.

Contributions: FK and NT were both involved in the protocol development and writing up of the manuscript. NT performed data analysis and edited the manuscript. Both authors edited and approved the final version for publication.

Conflict of interest: The authors declare no conflict of interest.

Funding: None.

Availability of data and materials: The data sets for this study are available from the authors upon reasonable requests.

Ethical approval: Ethical approval for the study was obtained from the University of Namibia Research Committee, reference no. SoNEC 67/2020. To ensure privacy and anonymity the researcher gathered data from respondents using an online questionnaire which they could respond to in their comfort of their homes and no personal details were required. This was in line with Covid-19 guidelines of social distancing. Confidentiality was maintained by keeping the electronic data password protected and accessed only by the researcher and supervisor.

Informed consent: The researcher explained the study objectives and purpose to the respondents and that participation in the study was voluntary. Willing respondents signed the informed consent but still were free to withdraw from the study at any time without any penalties or loss of privileges.

Received for publication: 7 May 2021.  
Accepted for publication: 20 April 2022.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright: the Author(s), 2022

Licensee PAGEPress, Italy  
Journal of Public Health in Africa 2022; 13:1923  
doi:10.4081/jphia.2022.1923

*Publisher's note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

ditions, resulting in serious complications or death.<sup>2</sup> Although VZV is preventable and curable, the disease still remains a public health problem in Namibia, and access to vaccination remains a challenge for many citizens, including students. Apart from similar studies that have been conducted in other settings, no previous study has been conducted to assess the prevention and control of VZV in students' dormitories in Namibia. The researchers who were affiliated to the University of Namibia (UNAM), Rundu Campus, and observed nine cases of VZV in 2018, with those students being allowed to freely come into contact with other students and some even attending classes during their illness. This study aims to determine the awareness of hostel students on the prevention of and control measures for VZV at UNAM.

The goal of the study was to investigate the awareness of hostel students on the prevention of and control measures for VZV at UNAM. Its objectives were to assess and describe the awareness of hostel students on the prevention of and control measures for VZV at UNAM Rundu Campus.

## Materials and Methods

In this study, the researchers used a quantitative research design, with each step being standardised to reduce bias when collecting and analysing the data. The results for this approach are more valid, reliable and generalisable to a larger population.<sup>7</sup> The researcher conducted the study with students who board at a satellite campus of UNAM in the Eastern part of the country. In

2020, the campus had a total of 280 boarding students from three faculties: the Faculty of Health Sciences, Faculty of Education and Faculty of Economic and Management Sciences. The researcher chose this campus because it is a fast-growing campus with an undergraduate student population of 4000.

The study population comprised of 280 boarding students from the three faculties. The eligibility criteria for the study included boarding students who could communicate well in English, were willing and who gave consent to participate in the study.

Solvin's formula for estimating sample size

$$\frac{N}{(1+N \times a^2)}$$

was used to estimate the population sample with "N" being the total population and "a" being the total confidence limit at 95%, representing a margin of error of 0.05. A total of 165 boarding students were accordingly selected for this study.

Respondents were randomly selected from a population of 280 boarding students. This sampling technique ensures that all available people or objects for a study have an equal chance of being part of the study.<sup>8</sup> The inclusion criteria included all boarding undergraduate students who were 18 years

**Table 1. Demographic characteristics of the respondents (n=165).**

	Frequency	Percent (%)
<b>Gender</b>		
Female	93	56.0
Male	72	44.0
Total	165	100.0
<b>Age</b>		
18–25 Years	145	88.0
26–35 Years	20	12.0
Total	165	100.0
<b>Course of study</b>		
1 <sup>st</sup> year	11	6.6
2 <sup>nd</sup> year	37	22.4
3 <sup>rd</sup> year	46	27.8
4 <sup>th</sup> year	71	43.0
Total	165	100.0
Bachelor of Economics	12	7.2
Bachelor of Education	97	58.7
Bachelor of Nursing	56	33.9
Total	165	100.0

**Table 2. Respondents' awareness of VZV.**

Variables	Options	Frequency	Percentage (%)	Mean (SD)
Aetiology	Bacteria	49	29.3	0.71 (0.45)
	Virus	118	70.7	
Information is available on campus	No	50	29.9	0.70 (0.45)
	Yes	117	70.1	
Transmission	Sex	38	22.8	0.77 (0.42)
	Contact	129	77.2	
VZV can be vaccinated against	No	40	24.0	0.76 (0.42)
	Yes	127	76.0	
Chicken pox increases absenteeism	Strongly disagree	6	3.6	4.23 (0.93)
	Disagree	3	1.8	
	Neutral	13	7.8	
	Agree	70	41.9	
	Strongly agree	75	44.9	
Chicken pox is contagious	Strongly disagree	2	1.2	4.16 (0.87)
	Disagree	6	3.6	
	Neutral	22	13.2	
	Agree	71	42.5	
	Strongly agree	66	39.5	

and above, from the three faculties on the campus.

In this study, data was collected from 165 respondents using an online questionnaire. This data collection tool was used for the study, as it is considered relevant for use when the researcher seeks to collect data from a larger sample.<sup>8</sup> Furthermore, an online questionnaire was considered suitable for collecting data during the Covid-19 pandemic, in line with social distance guidelines. The researcher developed the questionnaire in English as all the respondents were conversant with the English language. The researcher approached the respondents via the boarding students' WhatsApp group, giving a link to the online questionnaire. Subsequently, a total of 165 respondents completed the online questionnaire.

Data was analysed with SPSS version 26.0. Descriptive statistics and inferential statistics (frequency, mean, standard deviation and percentage) and a non-parametric Kruskal Wallis test were performed to test the mean differences between variables for the significance of unequally distributed data. The acceptable level of significance was  $p=0.05$ .

## Results

A total of 165 online questionnaires were distributed among hostel students via the institution's WhatsApp groups. The study attracted a 100% ( $n=165$ ) response rate.

Table 1 outlines the characteristics of the respondents. Accordingly, this study reveals that more females, 56% ( $n=93$ ), than males, 44% ( $n=72$ ), took part in the study. The majority of the respondents were in their fourth year, 43% ( $n=71$ ) and between the age of 18 and 25 years, 88% ( $n=145$ ). Most respondents, 58.7% ( $n=97$ ), were enrolled for the Bachelor of Education.

The majority of respondents, 70.7% ( $n=118$ ), were aware that VZV is caused by a virus, 29.3% ( $n=49$ ) ( $0.71\pm 0.45$ ) believed that VZV is caused by a bacteria, while 22.8% ( $n=38$ ) ( $0.77\pm 0.42$ ) believed VZV is sexually transmitted. Regarding vaccination, 76% ( $n=127$ ) ( $0.76\pm 0.42$ ) indicated that VZV can be vaccinated against, whereas 42.5% ( $n=71$ ) ( $4.16\pm 0.87$ ) agreed that chicken pox is contagious (Table 2).

Table 3 highlights respondents' aware-

ness of VZV among the different courses of study. The study found no statistical significance between VZV-related absenteeism ( $p=0.89$ ) and the course of study. The study did, however, find a strong association between aetiology ( $p=0.003$ ), available information ( $p=0.00$ ), mode of transmission ( $p=0.02$ ), knowledge on contagious diseases ( $p=0.03$ ) and course of study.

## Discussion

The majority of respondents, 70.7% ( $n=118$ ), knew that VZV is a viral disease; this was supported by a great proportion, 70.1% ( $n=117$ ), who were in agreement that information on VZV exists on campus. These findings are supported by a survey carried out among biology students which revealed that about 98% of the students believed VZV to be a viral disease.<sup>9,10</sup>

Awareness on how VZV is transmitted is important in the prevention and control of VZV spread. The study found 77.2% ( $n=129$ ) to have the awareness that VZV is spread through close contact but is not a sexually transmitted disease. These findings correspond to Shiel's,<sup>1</sup> whose study

**Table 3. Respondents' awareness of VZV among different courses of studies.**

		Course of study			P*
		Education	Nursing	Economic	
Aetiology	Bacteria	39	9	1	0.003
	Virus	59	47	10	
	Total	56	11	165	
Information is available on campus	No	40	9	1	0.002
	Yes	58	47	10	
	Total	56	11	165	
Transmission	Sex	28	6	4	0.023
	Contact	70	50	7	
	Total	56	11	165	
Need for vaccination	No	11	26	2	0.000
	Yes	87	30	9	
	Total	56	11	165	
VZV increases absenteeism	Strongly disagree	4	1	1	0.896
	Disagree	2	1	0	
	Neutral	8	4	1	
	Agree	41	25	3	
	Strongly agree	43	25	6	
	Total	56	11	165	
VZV is contagious	Strongly disagree	1	1	0	0.031
	Disagree	6	0	0	
	Neutral	15	6	1	
	Agree	45	20	5	
	Strongly agree	31	29	5	
	Total	56	11	165	
The most affected organ is?	Respiratory system	7	10	1	0.050
	Integumentary system	26	18	2	
	Reproductive system	22	14	4	
	Cardiovascular system	30	8	3	
	Digestive system	13	6	1	
	Total	56	11	165	

\*Kruskal Wallis Test

reaffirmed that VZV spreads from one person to another by means of direct contact and through the air.

Regarding vaccination, 76% (n=127) of respondents agreed that vaccination is generally warranted to prevent VZV. These findings are in line with those of Leung *et al.*<sup>11</sup> that vaccination should be offered to all susceptible individuals to prevent hospital outbreaks, as well as academic and economic loss for students. A study conducted in Sri Lanka on the susceptibility of new entrant university students to VZV revealed that of the 458 eligible students those who were unvaccinated had low levels of immunoglobulin G (IgG) antibodies compared to those who were vaccinated.<sup>12</sup>

A significant number of respondents, 44.9% (n=75), felt that contracting VZV led to the disruption of classes and increased absenteeism. This concurs with a similar study by Kurukulasooriya *et al.*<sup>13</sup> which found that VZV outbreaks can lead to the disruption of school attendance and activities; students with varicella may miss classes and activities for seven days or more.

## Conclusions

Based on the study results it was evident that students had an awareness of VZV. Although it may be concluded that the findings were statistically significant with regard to the course of study ( $p < 0.05$ ), no statistical significance was found between absenteeism, course of study and awareness of VZV. Greater efforts to disseminate information about VZV disease, its preven-

tion and control need to be implemented at UNAM Rundu campus. Existing health programs on school health can be particular vital in creating awareness among university boarding students.

## References

1. William C, Shiel JR. Medical Definition of Chickenpox. Calif: Medicine Net. Available from: <https://www.medicinenet.com/chickenpox/definition.htm>. Accessed: August 8, 2019.
2. The Namibian. Chickenpox can be deadly. Available from: <https://www.namibian.com.na/157382/archive-read/Chickenpox-can-be-deadly>. Accessed June 25, 2019.
3. CDC. Prevention of varicella: Recommendations of the advisory committee on immunization practices (ACIP). Centers for disease control and prevention website. Available from: <https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/varicella.html>. Accessed May 27, 2019.
4. WHO. Varicella and herpes zoster vaccines. Available from: <http://www.who.int/wer/2014/wer8925.pdf?ua=1>. Accessed June 26, 2019.
5. Turab AM, Chehadeh W. Varicella infection in the Middle East: Prevalence, complications, and vaccination. *J Res Med Sci.* 2018 Apr 26;23:19.
6. Hussey H, Abdullahi L, Collins J, et al. Varicella zoster virus-associated morbidity and mortality in Africa - a systematic review. *BMC Infect Dis* 2017;17:717.
7. Dowd M. Advantages & disadvantages of qualitative & quantitative research. 2018. Available from: <https://www.the-classroom.com/advantages-disadvantages-of-qualitative-quantitative-research-12082716.html>. Accessed March 27, 2020.
8. Brink H, Van der Walt C, Van Rensburg G. Fundamentals of research methodology for healthcare professionalism. 4th ed. Cape Town: Juta & Co, 2018.
9. Qadir M. Survey about awareness of VZV among biology students. *J Human Virol Retrovirol* 2019;7.
10. Ho EW, Ereno IL, Ibrahim M, Yeo CL. Knowledge, attitudes, and practices regarding chickenpox disease and its prevention in Singapore: Comparison between parents and medical students. *Proc Singapore Healthcare* 2012;21:257-64.
11. Meyers J, Narasimhan P, MacIntyre CRM, et al. Epidemic varicella zoster virus among university students, India. *Emerg Infect Dis* 2018;24:366-9.
12. Leung J, Marin M, Leino V, et al. Varicella immunization requirements for US colleges: 2014–2015 academic year. *J Am Coll Health* 2016;64:490-5..
13. Kurukulasooriya GMPCP, Thevanesam V, Agampodi S, et al. Susceptibility of new entrant university students in Sri Lanka to varicella zoster infection. *Asia-Pacific J Public Health* 2010:1941-2479.