

None-endoscopic Screening for Esophageal Squamous Cell Carcinoma- A Review

Gholamreza Roshandel^{1,2}, Shahryar Semnani^{2*}, Reza Malekzadeh¹

1. Digestive Diseases Research Center (DDRC), Tehran University of Medical Sciences (TUMS), Tehran, Iran.
2. Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran

ABSTRACT

Esophageal cancer (EC) is the eighth most common cancer and sixth most frequent cause of cancer mortality worldwide. Esophageal squamous cell carcinoma (ESCC) is the most common type of EC. ESCC develops by progression from premalignant lesions, which are called esophageal squamous dysplasia (ESD). Prevention is the most effective strategy for controlling this disease. Generally, two methods may be defined for ESCC prevention. The aim of the first preventive method is to prevent the initiation of ESD by avoiding the known risk factors, or primary prevention. Secondary prevention focuses on detection of the disease in its early curable stage, thus preventing its progression into advanced stages. Endoscopy with iodine staining and biopsy is the diagnostic choice for ESD. However it is invasive and expensive, and not accepted by asymptomatic ESD cases. Therefore, it is necessary to find a non-endoscopic screening method. Despite the large number of studies conducted worldwide, no approved method has been developed for ESCC screening. Regarding the multi-factorial nature of ESCC, it is proposed that the use of a combination of various criteria, such as cytological examination, risk factors, genetic alteration, and molecular markers may result in the development of a comprehensive and effective ESCC screening program.

KEYWORDS

Esophageal squamous cell carcinoma; Screening; Non-endoscopic; Review.

Please cite this paper as:

Roshandel Gh, Semnani Sh, Malekzadeh R. None-endoscopic screening for Esophageal Squamous Cell Carcinoma- A review. *Middle East J Dig Dis* 2012;4:111-24.

Epidemiology of ESCC

Esophageal cancer (EC) is the eighth most common cancer and the sixth most frequent cause of cancer mortality worldwide.¹ The highest rates for EC are reported from a geographical area that extends from northern Iran to north-central China, so-called the "Asian belt of EC".²⁻⁴ Despite its declining trend, EC remains the most frequent malignancy with a high incidence rate in northern Iran.^{5,6} Southeast Africa (Kenya, Zimbabwe), parts of South America (Brazil, Uruguay) and western Europe (France) are known as intermediate risk areas for EC.³ Other parts of the world, including the US, are considered to have low incidence rates of EC.¹ Adenocarcinoma and squamous cell carcinoma (SCC) are the main morphologies of EC.^{2,7} In the 1960s, esophageal squamous cell carcinoma (ESCC) comprised about 90% of EC cases.³ Recent studies have suggested a declining trend in the rate of ESCC as well

* Corresponding Author:

Dr Shahryar Semnani, MD
Golestan Research Center of Gastroenterology and Hepatology, Shahid Nabavi clinic, 4th Azae alley, 5th Azar street, Gorgan, Iran.
Tel: +98 171 2340835
Fax: +98 171 2369210
Email: sh_semnani@yahoo.com
Received: 11 Nov. 2011
Accepted: 28 Jan. 2012

as an increase in the rate of esophageal adenocarcinoma in the US and some European countries.⁸⁻¹⁰ Currently, it is estimated that the morphology in about 70% of worldwide EC cases are ESCC.¹¹ But, the situation seems to be different in the developing world. Reports from developing countries have shown that the shifting pattern is not the same as western countries. For example, ESCC comprises more than 90% of EC cases in northeastern Iran.^{12,13} Thus, ESCC is the most important type of EC in the developing world and a major health problem in high risk areas.

Pathogenesis of ESCC

ESCC develops by progression from dysplastic lesions within the squamous epithelium of the esophagus.³ In other words, esophageal squamous dysplasia (ESD) is the premalignant precursor lesion for ESCC.^{14,15} Various factors may cause dysplastic changes in normal epithelial cells of the esophagus, including Tobacco smoking, opium consumption, nass chewing, hot tea consumption, an excessive alcohol consumption, drinking mate, low intake of fresh fruits and vegetables, environmental factors, low socioeconomic status and viral infections.¹⁶⁻²¹ So, ESCC is a complex and multi-factorial disease. Over months to years ESDs grow into tumor mass (ESCC).²² The absence of true serosal layer in the esophageal wall makes ESCC a progressive cancer with relatively rapid invasion into neighboring structures.²³ The clinical manifestations in most of ESCC patients do not present in early stages and this results in delayed diagnosis of the disease. Distant metastasis and bone marrow invasion may be detected in 30 and 40 percent of ESCC cases, respectively.²⁴ So, ESCC cases usually present in late stages and the prognosis in most of them is poor. The overall five-year survival of ESCC patients is as low as 9%.²²

Controlling ESCC

When a patient is diagnosed with ESCC, the appropriate treatment modality is selected according to the tumor stage.³ In the early stages, curative therapeutic methods are considered. The thera-

peutic option in patients with submucosal ESCC or carcinoma in situ is endoscopic mucosal resection (EMR). Surgical resection, radiotherapy, and chemotherapy are selected for localized ESCC,³ whereas endoscopic palliative therapies are recommended for patients with advanced stage ESCC. These include laser therapy, argon plasma coagulation, esophageal dilation, and esophageal stent replacement.³ Although the above mentioned therapeutic modalities have been known to be useful in ESCC patients, prevention, however, is the most effective strategy for disease control. Generally, two methods may be defined for ESCC prevention. The aim of the first preventive method is to prevent the initiation of the ESD, or primary prevention. However, ESD is a multi-factorial condition whose risk factors have not been completely identified. Therefore primary prevention may not warrant complete elimination of ESD and consequently ESCC.

The second method focuses on detection of the disease in an early curable stage, preventing its progression into advanced stages (secondary prevention). Although endoscopy with iodine staining and biopsy is the diagnostic choice for ESD,^{3,25} it is an invasive and expensive method not particularly accepted by asymptomatic ESD cases.²⁶ Therefore, it is necessary to find a non-endoscopic screening method which takes into consideration a combination of various criteria such as cytological examination, risk factors, and molecular markers.²⁷

Screening Methods for ESCC

By definition, a screening program might have potential benefit for a condition if the following assumptions are true. At first, the disease in all or most cases should start from a detectable preclinical phase. Secondly, in the absence of intervention, most or all cases in the preclinical phase progress into the clinical phase.²⁸ Both assumptions are true for ESCC.²² Thus, a simple, minimally invasive, accurate, and cost effective screening program will be helpful for controlling ESCC. Recently, some studies have been conducted to find a screening method for ESCC. Various criteria were used in these studies, including patients' risk factors, cytopathological

examination of esophageal secretions, serum markers, and genetic profiling. A detailed discussion about these criteria is given in the following sections.

1- Risk Factors

The risk factors for ESD are reported to be similar to those of ESCC,²⁹ and both may be used as possible criteria for ESCC screening. Xibin et al. have reported a negative relationship between household income, residential space, education, and EC. They also noted that consumption of beans, vegetables, and vinegar were protective against EC.³⁰ By measuring urine 1-hydroxypyrene glucuronide (1-OHPG), a stable polycyclic aromatic hydrocarbon (PAHs) metabolite, Kamangar et al. have suggested that high exposure to PAH may be a possible risk factor for ESCC in northeastern Iran.³¹ Islami et al. have reported that dimensions of socio-economic status (SES), which included a higher level of education, wealth, and being married were inversely related to ESCC.³² Drinking high-temperature beverages was shown to be an important risk factor for ESCC.³³ The results of a study from Iran suggested human papillomavirus (HPV) infection as a potential risk factor for ESCC in high risk areas.³⁴

In addition, Nasrollahzadeh et al. showed that the risk of ESCC in individuals who smoked both cigarettes and opium substances (OR=2.35) was higher than those who smoked only cigarettes (OR=1.98) or opium (OR=2.12).³⁵ Wu et al. reported that the combined usage of three substances (cigarette smoking, betel chewing, and alcohol consumption) increased the risk of ESCC (OR=39.2, 95%CI: 13.2-116.1). They found that patients who used only one of these substances had a 1.5-fold (95%CI: 0.6-3.8) risk for ESCC.³⁶

Wei et al. attempted to use a questionnaire and physical examination data to develop a risk model for triaging subjects for endoscopies.²⁹ The aim of their study was to define a screening program using previously known risk factors for ESCC. The researchers considered ten risk factors of age, sex, smoking status, ethanol patch test flushing response, number of persons in the subject's household, household income, family history of cancer,

systolic blood pressure, heating stove type, and quintile of tooth loss. The sensitivity and specificity of their final model for predicting dysplasia was 57% and 54%, respectively. As such, they concluded that risk factors, alone, were not appropriate and successful measures with which to develop a screening program for ESCC. They have proposed that the addition of cytological or molecular data from subjects would result in the development of an efficient screening program.

Generally, researchers need to consider some important criteria for using a risk factor as a screening test for ESCC. The risk factor should have a strong association with ESCC. Variations in exposure to the risk factor within the population should be mentioned. The risk of disease should be considered in all quintiles of the distribution of the risk factor within the population. In other words, not only should those at both ends of the distribution of exposure (high and low) be included, but also those in the middle of the distribution.³⁷ If appropriately used, risk factors could play an important role in ESCC screening. Table 1 shows the characteristics of major risk factors for ESCC and ESD.

2- Cyto-Pathological Examination of Esophageal Secretion

Non-endoscopic cytological methods have been used for early diagnosis of ESCC since the 1970s.⁵⁶ Lazarus et al. have used abrasive brush cytology for early detection of EC, with a high sensitivity (90%) and specificity (99.9%) for their method.⁵⁷ Roth et al., in 1997, assessed the validity of balloon and sponge samples for detecting ESD and ESCC.²⁶ The sensitivity and specificity of balloon sampling for the detection of ESD was 47%, whereas it was 88% for ESCC. The sensitivity and specificity of the sponge sampler for identifying ESD or ESCC was 24% and 92%, respectively. The methods were not adequately sensitive to be used as screening programs within the community. The researchers suggested that improving samplers would increase the sensitivity of the test. In a complementary study to improve the validity of their screening test, Pan et al. have used new mechanical and inflatable balloons for identifying ESD or ESCC.⁵⁸ The mechanical balloons had a sensitivity of 39% whereas,

Table 1: Risk factors for esophageal squamous carcinoma or esophageal dysplasia

Risk factors	Author	Location	Statistic		
			Type	Values	
Positive family history of cancer	Yes vs. no	Wei et al. ²⁹	China	OR (CI 95%)	1.57 (1.13-2.18)
	Yes vs. no	Wang et al. ³⁸	China	OR (CI 95%)	3.83 (1.13-12.97)
	Yes vs. no	Akbari et al. ³⁹	Iran	OR (CI 95%)	3.6 (2.3-5.7)
Relationship between parents	Related vs. no relationship	Akbari et al. ³⁹	Iran	OR (<i>p</i> -value)	4.1 (0.006)
Systolic blood pressure	per 10 mm Hg increase	Wei et al. ²⁹	China	OR (CI 95%)	1.11(1.03-1.19)
Heating stove without chimney	Yes vs. no	Wei et al. ²⁹	China	OR (CI 95%)	2.22 (1.27-3.86)
Oral health	12–31 vs. 0-4 teeth lost	Wei et al. ²⁹	China	OR (CI 95%)	1.91 (1.17-3.15)
	6-15 vs. 0-5 teeth lost	Guha et al. ⁴⁰	Central Europe	OR (CI 95%)	2.84 (1.26-6.41)
	6-15 vs. 0-5 teeth lost	Guha et al. ⁴⁰	Latin America	OR (CI 95%)	2.18 (1.04-4.59)
	Extremely poor vs. good	Sepehr et al. ⁴¹	Iran	OR (CI 95%)	4.76 (1.48-15.31)
	Decayed, missing, or filled teeth =32 vs. ≤15	Abnet et al. ⁴²	Iran	OR (CI 95%)	2.1 (1.19-3.7)
	No regular oral hygiene vs. daily tooth brushing	Abnet et al. ⁴²	Iran	OR (CI 95%)	2.37 (1.42-3.97)
Source of drinking water	Other than tap water	Xibin et al. ³⁰	China	OR (CI 95%)	5.49 (1.43-21.1)
Cigarette smoking	>30 pack-years vs. none	Wu et al. ³⁶	Taiwan	OR (CI 95%)	3.7 (1.6-8.7)
	≤ 3.5 packs/wk vs. none	Tai et al. ⁴³	Taiwan	OR (CI 95%)	6.08 (1.43-25.94)
	≥ 15 cigarettes/day vs. none	Vizcaino et al. ⁴⁴	Zimbabwe	OR (CI 95%)	4.3 (2.8-6.7)
	≥ 80 packs-years vs. none	Vaughan et al. ⁴⁵	USA	OR (CI 95%)	16.9 (4.1-69.1)
	> 11 cigarettes/day vs. none	Nasrollahzadeh et al. ³⁵	Iran	OR (CI 95%)	1.98 (1.2-3.25)
	≥15 cigarettes/day vs. none	Castelletto et al. ⁴⁶	Argentina	OR (CI 95%)	3 (1.5-5.7)
Areca (betel nut) chewing	> 495 betel/year vs. none	Wu et al. ³⁶	Taiwan	OR (CI 95%)	9.4 (1.8-48.3)
Alcohol consumption	> 1220 g-year vs. none	Wu et al. ³⁶	Taiwan	OR (CI 95%)	9.8 (4.2-22.6)
	> 158 g/wk vs. none	Tai et al. ⁴³	Taiwan	OR (CI 95%)	20.58 (1.72-245.62)
	≥20 vs. 0-6 drinks/week	Vaughan et al. ⁴⁵	USA	OR (CI 95%)	9.5 (4-22.3)
	≥200ml/day vs. none	Castelletto et al. ⁴⁶	Argentina	OR (CI 95%)	5.7 (2.2-15.2)
Esophageal lesions (esophagitis,...)	Yes vs. no	Wang et al. ³⁸	China	OR (CI 95%)	11.63 (1.13-119.33)
<i>Helicobacter pylori</i> infection	Yes vs. no	Wang et al. ³⁸	China	OR (CI 95%)	3.19 (1.11-9.15)
Eating breakfast	Yea vs. no	Sharp et al. ⁴⁷	UK	OR (CI 95%)	0.18 (0.07-0.48)
Aspirin consumption	Daily use vs. none	Sharp et al. ⁴⁷	UK	OR (CI 95%)	0.08 (0.01-0.56)
Fresh fruit consumption	Weekly vs. less often	Sepehr et al. ⁴¹	Iran	OR (CI 95%)	3.18 (1.14-8.9)
Opium consumption	Yes vs. no	Nasrollahzadeh et al. ³⁵	Iran	OR (CI 95%)	2.12 (1.21–3.74)

Risk factors	Author	Location	Statistic		
			Type	Values	
Drinking mate	Heavy drinkers of very hot vs. light drinkers of cold/warm/hot	Castellsague et al. ⁴⁸	South America	OR (CI 95%)	4.14 (2.24–7.67)
	Very hot vs. warm	De Stefani et al. ⁴⁹	Uruguay	OR (CI 95%)	5.76 (2.92-11.35)
Tea temperature	Very hot vs. warm	Islami et al. ⁵⁰	Iran	OR (CI 95%)	8.16 (3.93-16.91)
	Hot vs. others	Cook-Mozaffari et al. ⁵¹	Iran	OR (<i>p</i> -value)	Men=1.72; Women=2.17 (<i><</i> 0.01)
	Hot vs. not hot	Onuk et al. ⁵²	Turkey	OR (CI 95%)	8.7 (2.5-30.2)
	Very hot vs. cold/warm	Castellsague et al. ⁴⁸	South America	OR (CI 95%)	3.73(1.41-9.89)
	High temperature vs. never drinking	Wu et al. ⁵³	China	OR (CI 95%)	4.2 (2.3-7.6)
Drinking coffee	very hot vs. cold/warm	Castellsague et al. ⁴⁸	South America	OR (CI 95%)	2.29 (1.37-3.81)
	Burning hot vs. others	De Jong et al. ⁵⁴	Singapore	OR (<i>p</i> -value)	Men=4.22; Women=4.09 (<i><</i> 0.01)
Interval between tea being poured and drunk (minutes)	<i><</i> 2 vs. <i>≥</i> 4	Islami et al. ⁵⁰	Iran	OR (CI 95%)	5.41 (2.63-11.14)
Formal education	Middle school or higher vs. no school	Islami et al. ³²	Iran	OR (CI 95%)	0.2 (0.06-0.65)
Eating barbecued meat	<i>≥</i> 1 vs. <i><</i> 1 per week	Castelletto et al. ⁴⁶	Argentina	OR (CI 95%)	2.4 (1.2-4.8)
PAH content (8E11 antibody of the oesophageal epithelium)	Fifth quintile vs. first quintile	Abedi-Ardekani et al. ⁵⁵	Iran	OR (CI 95%)	26.6 (5.21-135)

the inflatable balloons was 46%. Unfortunately, the sensitivity of the new balloons was still inadequate to be used as a population-based screening program. The result of this study emphasized the need to consider molecular markers to increase the validity of a program for detecting ESD.

Lao-Sirieix et al. used a capsule sponge (cytosponge) for cytological examination of the esophagus. They suggested that this device had good validity for early detection of Barrett's esophagus.⁵⁹ Recently, Kadri et al. have shown that coupling a Barrett's specific immunomarker with a cytosponge would improve its sensitivity and specificity for detecting Barrett's lesions; the combination would be a suitable screening program to be used in the primary care setting.⁶⁰ Therefore, coupling a cytosponge with an ESD specific marker may be considered as a promising method for developing

an applicable ESCC screening program.

3- Serum Markers

In a study from Japan, the peripheral blood samples of EC patients were assessed for Δ Np63 gene expression by Δ Np63-specific RT-PCR.⁶¹ Δ Np63 mRNA was detected in blood samples of 52% of primary ESCC and in 60% of recurrent ESCC cases. No Δ Np63 expression was observed in controls. Therefore, the researchers have concluded that Δ Np63 is a good and highly specific blood marker for early detection of ESCC.⁶¹ Hibi et al. found a high rate of promoter methylation of the p16 gene in the serum of ESCC patients, which suggested that p16 gene promoter methylation may be used as a serum marker for identifying precursor lesions of ESCC.⁶²

In a study from India, Kannan et al. have con-

cluded that the Thomson–Friedenreich (TF) antigen was an appropriate marker for the early diagnosis of ESCC.⁶³ Yang et al. studied the expression of squamous cell carcinoma antigen 2 (SCCA2) in peripheral blood of patients with ESCC and ESD, as well as normal individuals.⁶⁴ Their results showed significantly higher SCCA2 mRNA expression in ESCC and ESD cases than normal subjects. They also compared the SCCA2 levels measured by two methods, including enzyme-linked immunosorbent assay (ELISA) and SCCA2 mRNA expression. The results showed a significant correlation between ELISA SCCA2 levels in the serum and SCCA2 mRNA expression levels in the peripheral blood. They have suggested that SCCA2 is a good biomarker for the detection of premalignant esophageal lesions.⁶⁴

Chung et al. used multiplex tissue immunoblotting to quantify the expression of some proteins in esophageal carcinogenesis.⁶⁵ Their results have shown overexpression of secreted protein, acidic and rich in cysteine (SPARC). The possibility of measuring SPARC in serum makes it a potential biomarker for early detection of ESCC.⁶⁵ Munck-Wikland reported elevated serum levels of tumor markers carcinoembryonic antigen (CEA; 39%), CA 50 (41%), and CA 19-9 (13%) in ESCC patients. The sensitivity of considering these markers together for detecting ESCC was 59%.⁶⁶

In a study from Japan, a proteomics-based approach was used to identify tumor antigen in an ESCC cell line (TE2) and related autoantibodies in serum of ESCC patients.⁶⁷ They found an autoantibody against peroxiredoxin VI, and suggested that it was a potential biomarker for early detection of ESCC.⁶⁷

4- Genetic Alterations

Alterations in oncogenes, tumor suppressor genes as well as alterations in microRNA expression have been known to involve in the pathogenesis of ESCC.⁶⁸

4-1- Oncogenes

4-1-1- Cyclin D1: Jiang et al. suggested changes

in the expression of cyclin D1 gene in ESCC patients.⁶⁹

4-1-2- *Fart1*: Saitoh et al. reported the overexpression of *Frat1* mRNA in ESCC patients.⁷⁰

4-1-3- *HoxD9* and *Pbx1*: The results of a study from China showed over-expression of *HoxD9* and *Pbx1* genes in ESCC tissues.⁷¹

4-1-4- Vascular endothelial growth factor (VEGF) gene: A significant over-expression of VEGF gene was found in ESCC tissues.⁷²

4-1-5- *Akr1c2* gene: The up-regulation of *Akr1c2* gene suggested it to play a possible role in the development of ESCC.⁷³

4-2- Tumor suppressor genes

Mutations in *p53*^{74,75} and *MTS1*⁷⁶ genes have been reported in ESCC cases. The *p53* gene mutation was found in the esophagitis area as well as in dysplastic areas of the esophagus.⁷⁷ Gao et al.⁷⁸ and Wang et al.⁷⁹ have reported that *p53* protein accumulation occurs quite frequently in ESD cases, therefore concluding that the *p53* mutation is an early event in the pathogenesis of ESCC.

Xing et al. noted a loss of heterozygosity (LOH) of the retinoblastoma (*Rb*) gene in 55% of ESCC cases.⁸⁰ According to these researchers, *Rb* LOH was significantly more frequent in tumors with *p53* mutations. They have concluded that concurrent alteration of *Rb* and *p53* genes is an important mechanism for the development of ESCC.⁸⁰

The results of a study from Japan have shown that alteration of the deleted in lung cancer 1 (*DLC1*) gene may be involved in development of ESCC.⁸¹ According to other studies, changes in the *p16INK4a* and *p15INK4b* genes have been found in 68% and 50% of ESCC patients, which suggests these changes are important in the pathogenesis of ESCC.⁸² Guo et al. have also found frequent methylation of the *p16INK4a* gene in ESCC patients.⁸³

In a study from the US, loss of heterozygosity in adenomatous polyposis coli (*APC*) or *MCC* genes was found in 80% of ESCC patients. Therefore, it was proposed that alterations in the *APC* and *MCC* genes have important roles in the pathogenesis and progression of ESCC.⁸⁴ Zare et al. found *APC* pro-

moter hypermethylation in ESCC patients, thus indicating that it could be an appropriate candidate molecular marker in ESCC cases.⁸⁵

Inactivation of the WW domain containing oxidoreductase (WWOX) gene, as reported by Kuroki et al. may be a possible mechanism in the pathogenesis of ESCC.⁸⁶

Other genes, such as the cysteine-rich protein with Kazal motifs (RECK) gene showed significant reduction in ESCC tissues when compared with normal tissues.⁷²

Mal gene: Kazemi-Noureini et al. reported down-regulation of mal gene in ESCC patients.⁷³
CDKN2A gene: Frequent methylation of CDKN2A was reported in ESCC patients.⁸³

S100A2 gene: Cao et al. found a significant down-regulation of S100A2 gene in ESCC tissues.⁸⁷

P63 gene: The expression of P63 gene was significantly decreased in ESCC tissues when compared with normal esophageal epithelium.⁸⁷

FHIT gene: Loss of FHIT tumor suppressor gene was reported in precursor lesions of ESCC, indicating its possible role in developing ESCC.⁸⁸

4-3- Other Genetic Alterations

According to Ishii et al., DNA methylation may play an important role in the progression of ESD into ESCC.⁸⁹ Guo et al. have noted an increasing trend in the number of methylated genes along with increasing cellular atypia in EC cases.⁸³ In a research by Li et al, an association between increasing telomerase activity and ESCC precursor lesions was noted. Their results showed telomerase activity in 60% of esophageal metaplasia and in about 90% of esophageal dysplasia.⁹⁰

Expression of matrix metalloproteinase-7 (MMP-7) increases in some cancers, including EC.⁶⁸ In addition, MMP-26 could be a potential biomarker for the diagnosis of EC.⁹¹ Increasing matrix MMP-3 and MMP-10 occurs due to upregulation of their genes in ESCC patients, which suggests they may also be potential diagnostic markers.⁹²

In a study from China, proliferating cell nuclear antigen (PCNA) was expressed in 55% of normal esophageal tissue, in 75% of ESD and 93% of

ESCC tissues. This study also indicated a positive correlation between overexpression of p53 and PCNA during different stages in the development of ESCC from normal esophageal epithelium.⁹³

Ishii et al. have proposed that inactivation of the FEZ1 gene may play a role in the development of ESCC.⁹⁴ A member of the human frizzled gene (FzE3) was expressed in 86% of poorly differentiated ESCCs.⁹⁵ By decreasing expression of the APC gene and increasing the B-catenin mediated signals, FzE3 may play a possible role in the pathogenesis of ESCC.⁹⁵ Overexpression of ornithine decarboxylase (ODC) mRNA was seen in about 91% of ESCC cases, therefore it may be important in developing ESCC.⁹⁶ Liu et al has suggested a role for translocation of annexin I protein in the pathogenesis of ESCC.⁹⁷ Qi et al. suggested that annexin II protein was a suitable biomarker for screening and detecting precursor lesions in ESCC patients.⁹⁸ Overexpression of the gene amplified in squamous cell carcinoma 1 (GASC1) was also shown to be related to the development of ESCC.⁹⁹

Yue et al. suggested the inactivation of esophageal cancer related gene 4 (ECRG4) due to hypermethylation as an important mechanism in the development of ESCC.¹⁰⁰ Gholamin et al. found a frequent over-expression of Interleukin-10 (IL-10) gene in ESCC patients.¹⁰¹ Over-expression of transforming growth factor β (TGF- β) gene was frequent in ESCC cases.¹⁰¹ Akbari et al. reported a significant association between ADH1B gene and the risk of ESCC in northeast of Iran.¹⁰² Tsuda et al. reported high rates of amplification of hst1 and int2 genes in ESCC patients.¹⁰³ Allelic losses on chromosomes 5q¹⁰⁴ and 17q¹⁰⁵ were found in ESCC patients. Kashyap et al. analyzed the mRNA expression profiles of 20 ESCC patients by whole genome DNA microarrays and Significant downregulations were identified in the pathway of several genes in arachidonic acid metabolic including ALOX15B, GPX3 and PTGDS in ESCC.¹⁰⁶ Adams et al. used quantitative methylation-specific PCR techniques to evaluate the accuracy of methylation in selected genes in esophageal balloon cytology specimens for identifying ESD.¹⁰⁷ The results showed that the

maximum sensitivity and specificity for individual genes were 34% and 99%. To increase the validity, they used panels of multiple genes and found a sensitivity of 50% and a specificity of 65% for a panel of four genes (AHRR, p16INK4a, CLDN3, and MTIG). They concluded that further studies on gene methylation in balloon cytology specimens will result in designing more accurate and suitable screening methods for ESCC.¹⁰⁷ Akbari et al. found mutations in Fanconi anemia genes (including FANCD2, FANCE, FANCL and FANCA)¹⁰⁸ as well as in BRCA2 (FANCD1) gene¹⁰⁹ in ESCC cases from northeast of Iran. Eisenberger et al. used a Microsatellite DNA analysis using a panel of 12 microsatellite markers on chromosome 9p (p16), chromosome 17p (p53), chromosome 18q (DPC4-gene), chromosome 18p and chromosomes 5 p and q (APC gene) in ESCC patients.¹¹⁰ Their results showed that 93% and 96% of cases had at least one microsatellite DNA alterations in their tumor and serum, respectively.¹¹⁰ Sepehr et al. reported a higher frequency of polymorphisms of CYP1A1 m1, CYP1A1 m2, CYP2A6*9, and ADH2*1 genes in population with high ESCC rates in northeast of Iran.¹¹¹ The risk of ESCC may increase in rare combination of GSTT1 null genotype and GSTP1 Val/Val variant.¹¹² Zhang et al. used whole-genome microarray method to analyze genome-wide mRNA expression profiles in ESCC patients.¹¹³ They found that 263 genes were significantly downregulated, including MFAP4, LYVE1, ANGPTL1, DPT, PPP1R1A, RERGL, NKX31 and SCARA5.¹¹³ They also reported significant upregulation in 104 genes, including AURKA, CDC20, CDCA3, TTK, RHPN2, DEPDC1, RASEF, IGF2BP3, TRIP13, CTHRC1 and COL5A2.¹¹³ In a study from India, genomewide mRNA profiling showed that fibroblast activation protein (FAP) as well as oral cancer overexpressed 2 (ORAOV2) were overexpressed in 98% and 68% of cases, respectively. In addition, Overexpression of Osteopontin (OPN) was found in 97% of the ESCC cases. So, genomewide mRNA profiling was a good approach to identify new biomarkers for ESCC.¹⁰⁶ Fu et al. used proteomics approach to assess overexpression of alpha-actinin

4 (ACTN4) and 67 kD a laminin receptor (67LR) in ESCC cases. ACTN4 and 67LR were overexpressed in 58.9% and 48.8% of samples suggesting them as suitable biomarkers for ESCC.¹¹⁴ Galectin-7 was also showed a significant upregulation in ESCC cases using a proteomic analysis, indicating its potential role in development of ESCC. So, galectin-7 may potentially be served as a marker for ESCC.¹¹⁵ In a study from China, CDC25B mRNA expression showed an increase in ESD and ESCC cases comparing to normal individuals, suggesting CDC25B protein as a potential biomarker for ESCC screening.¹¹⁶ The expression of plasminogen activator inhibitor-1 (PAI-1) was reported to be higher in advanced than early stages of ESCC.¹¹⁷ Abnet et al. reported high prevalence of mitochondrial DNA (mtDNA) mutation (the common deletion) in ESCC tissues, concluding that common deletion in mtDNA may be useful in developing ESCC screening programs.¹¹⁸ Marjani et al. found a significant higher levels of polycyclic aromatic hydrocarbons (PAHs)-DNA adducts in ESCC tissues than normal ones, suggesting PAH-DNA adducts as possible marker for ESCC.¹¹⁹

4-4- Alterations in the Expression of micro-RNA (miRNA)

Feber et al. reported decreased expression of miRNA-203 and miRNA-205 and increased expression of miRNA-21 in ESCC patients.¹²⁰ miRNA-21 expression was elevated, whereas expression of miRNA-375 was lower in ESCC cases than noncancerous ones.¹²¹ Ogawa et al. have suggested that miRNA-129 may play an important role in the pathogenesis of ESCC.¹²² Lee et al. have shown that miRNA-373 may suppress an oncogene and play a role in developing ESCC.¹²³ Significant elevation of miRNA-21 expression was found in ESCC tissues.¹²⁴ Guo et al. used advanced microRNA microarray techniques to assess the expression of miRNAs in ESCC patients. Their results showed over-expression of three miRNAs (hsa-miR-25, hsa-miR-424, and hsa-miR-151) as well as down-regulation of four miRNAs (hsa-miR-100, hsa-miR-99a, hsa-miR-29c, and mmu-miR-140).¹²⁵ In a study from Japan, the expression of miR-205 and

miR-10a was significantly altered, which suggested their possible role in pathogenesis of ESCC.¹²⁶

Designing a Comprehensive Screening Program for ESCC

Regarding the multi-factorial nature of ESCC, some researchers propose that the use of a combination of the above mentioned criteria may result in the development of an effective screening program for ESCC. Nicolas Perez et al. have reported that the use of cytology or Lugol chromo-endoscopy in high ESCC risk populations or individuals will result in early detection of ESCC cases, suggesting this combination strategy to be a good screening program for ESCC.¹²⁷ In a study from Iran, individuals with GSTP1 Ile/Ile variants have more susceptibility to ESCC in smokers, while non-smokers with this genotype seem to be protected. Therefore, assessment of GSTP1 genotype together with smoking habits may be important in determining the risk of ESCC.¹¹² According to Montesano et al., a strong relationship between p53 mutation and tobacco smoking in ESCC patients exists.¹²⁸ In a study from Japan, Yokoyama et al. have developed a health risk appraisal (HRA) model to determine high risk individuals for ESCC. They used a genetic alteration (inactivation of aldehyde dehydrogenase-2; ALDH2) and some risk factors (alcohol drinking, tobacco smoking, vegetable and fruit consumption). The results showed that the higher sensitivity and specificity when both genetic changes and risk factors were considered in the model.¹²⁹

Despite the large number of studies conducted worldwide, to date, no approved method has been developed for ESCC screening.³ A possible explanation for this situation is that ESCC is a multi-factorial disease, where most research has focused on one or some limited aspects of the disease. Therefore, we need to consider more complex and multi-aspect designs for developing a comprehensive and effective ESCC screening program.

CONFLICT OF INTEREST

The authors declare no conflict of interest related to this work.

REFERENCES

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. GLOBOCAN 2008, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10 Lyon: International Agency for Research on Cancer, 2010.
2. Bird-Lieberman EL, Fitzgerald RC. Early diagnosis of oesophageal cancer. *Br J Cancer* 2009;**101**:1-6.
3. Das A. Tumors of the Esophagus. In: Feldman M, Friedman L, Brandt LJ. Sleisenger and Fordtran's gastrointestinal and liver disease: pathophysiology, diagnosis, management, 9th ed, vol. 1. Philadelphia: Saunders Elsevier, 2010:745-70.
4. Mahboubi E, Kmet J, Cook PJ, Day NE, Ghadirian P, Salmasizadeh S. Oesophageal cancer studies in the Caspian Littoral of Iran: the Caspian cancer registry. *Br J Cancer* 1973;**28**:197-214.
5. Semnani S, Sadjadi A, Fahimi S, Nouraei M, Naeimi M, Kabir J, et al. Declining incidence of esophageal cancer in the Turkmen Plain, eastern part of the Caspian Littoral of Iran: a retrospective cancer surveillance. *Cancer Detect Prev* 2006;**30**:14-9.
6. Islami F, Kamangar F, Aghcheli K, Fahimi S, Semnani S, Taghavi N, et al. Epidemiologic features of upper gastrointestinal tract cancers in Northeastern Iran. *Br J Cancer* 2004;**90**:1402-6.
7. Parkin DM, Whelan SL, Ferlay J, Teppo L, Thomas DB. Cancer Incidence in Five Continents, Vol. VIII Lyon: IARC, 2002.
8. Brown LM, Devesa SS, Chow WH. Incidence of adenocarcinoma of the esophagus among white Americans by sex, stage, and age. *J Natl Cancer Inst* 2008;**100**:1184-7.
9. Pera M, Manterola C, Vidal O, Grande L. Epidemiology of esophageal adenocarcinoma. *J Surg Oncol* 2005;**92**:151-9.
10. Bollschweiler E, Wolfgarten E, Gutschow C, Holscher AH. Demographic variations in the rising incidence of esophageal adenocarcinoma in white males. *Cancer* 2001;**92**:549-55.
11. Parkin DM, Bray F, Ferlay J, Pisani P. Estimating the world cancer burden: Globocan 2000. *Int J Cancer* 2001;**94**:153-6.
12. Kamangar F, Malekzadeh R, Dawsey SM, Saidi F. Esophageal cancer in Northeastern Iran: a review. *Arch Iran Med* 2007;**10**:70-82.
13. Islami F, Kamangar F, Nasrollahzadeh D, Moller H, Bof-feta P, malekzadeh R. Oesophageal cancer in Golestan Province, a high-incidence area in northern Iran - A review. *Eur J Cancer* 2009;**45**:3156-65.
14. Wang GQ, Abnet CC, Shen Q, Lewin KJ, Sun XD, Roth MJ, et al. Histological precursors of oesophageal squamous cell carcinoma: results from a 13 year prospective follow up study in a high risk population. *Gut* 2005;**54**:187-92.
15. Dawsey SM, Lewin KJ, Wang GQ, Liu FS, Nieberg RK, Yu Y, et al. Squamous esophageal histology and subsequent risk of squamous cell carcinoma of the esophagus. A prospective follow-up study from Linxian, China. *Cancer* 1994;**74**:1686-92.

16. Lavergne D, de Villiers EM. Papillomavirus in esophageal papillomas and carcinomas. *Int J Cancer* 1999;**80**:681-4.
17. Kamangar F, Chow W-H, C. Abnet C, M. Dawsey S. Environmental Causes of Esophageal Cancer. *Gastroenterol Clin North Am* 2009;**38**:27-57.
18. Islami F, Pourshams A, Semnani S. Prevalence of Esophageal Cancer Risk Factors among Turkmen and Non-Turkmen Ethnic Groups in a High Incidence Area in Iran. *Arch Iran Med* 2010;**13**:111-5.
19. Ghadirian P. Thermal irritation and esophageal cancer in northern Iran. *Cancer* 1987;**60**:1909-14.
20. Nouarie M, Pourshams A, Kamangar F, Sotoudeh M, Derakhshan MH, Akbari MR, et al. Ecologic study of serum selenium and upper gastrointestinal cancers in Iran. *World J Gastroenterol* 2004;**10**:2544-6.
21. Semnani S, Roshandel G, Zendeabad A, Keshtkar A, Rahimzadeh H, Abdolahi N, et al. Soils selenium level and esophageal cancer: An ecological study in a high risk area for esophageal cancer. *J Trace Elem Med Biol* 2010;**24**:174-7.
22. Kumar V, Abbas AK, Fausto N, Aster JC. Robbins and Cotran pathologic basis of disease, ed. 8th, vol. 2 Philadelphia: Saunders Elsevier 2010.
23. 23. Ludeman L, Shepherd NA. Serosal involvement in gastrointestinal cancer: Its assessment and significance. *Histopathology* 2005;**47**:123-31.
24. Thorban S, Roder JD, Nekarda H, Funk A, Siewert JR, Pantel K. Immunocytochemical detection of disseminated tumor cells in the bone marrow of patients with esophageal carcinoma. *J Natl Cancer Inst* 1996;**88**:1222-7.
25. Dawsey SM, Fleischer DE, Wang GQ, Zhou B, Kidwell JA, Lu N, et al. Mucosal iodine staining improves endoscopic visualization of squamous dysplasia and squamous cell carcinoma of the esophagus in Linxian, China. *Cancer* 1998;**83**:220-31.
26. Roth MJ, Liu SF, Dawsey SM, Zhou B, Copeland C, Wang GQ, et al. Cytologic detection of esophageal squamous cell carcinoma and precursor lesions using balloon and sponge samplers in asymptomatic adults in Linxian, China. *Cancer* 1997;**80**:2047-59.
27. Chung CS, Lee YC, Wang CP, Ko JY, Wang WL, Wu MS, et al. Secondary Prevention of Esophageal Squamous Cell Carcinoma in Areas Where Smoking, Alcohol, and Betel Quid Chewing are Prevalent. *J Formos Med Assoc* 2010;**109**:408-21.
28. Gordis L. Epidemiology, ed. 4th Philadelphia: Saunders Elsevier, 2009.
29. Wei WQ, Abnet CC, Lu N, Roth MJ, Wang GQ, Dye BA, et al. Risk factors for oesophageal squamous dysplasia in adult inhabitants of a high risk region of China. *Gut* 2005;**54**:759-63.
30. Xibin S, Meilan H, Moller H, Evans HS, Dixin D, Wenjie D, et al. Risk factors for oesophageal cancer in Linzhou, China: a case-control study. *Asian Pac J Cancer Prev* 2003;**4**:119-24.
31. Kamangar F, Strickland PT, Pourshams A, Malekzadeh R, Boffetta P, Roth MJ, et al. High Exposure to Polycyclic Aromatic Hydrocarbons May Contribute to High Risk of Esophageal Cancer in Northeastern Iran. *Anticancer Res* 2005;**25**:425-8.
32. Islami F, Kamangar F, Nasrollahzadeh D, Aghcheli K, Sotoudeh M, Abedi-Ardekani B, et al. Socio-economic status and oesophageal cancer: results from a population-based case-control study in a high-risk area. *Int J Epidemiol* 2009;**38**:978-88.
33. Islami F, Boffetta P, Ren JS, Pedoeim L, Khatib D, Kamangar F. High temperature beverages and foods and esophageal cancer risk—A systematic review. *Int J Cancer* 2009;**125**:491-524.
34. Farhadi M, Tahmasebi Z, Merat S, Kamangar F, Nasrollahzadeh D, Malekzadeh R. Human papillomavirus in squamous cell carcinoma of esophagus in a high-risk population. *World J Gastroenterol* 2005;**11**:1200-3.
35. Nasrollahzadeh D, Kamangar F, Aghcheli K, Sotoudeh M, Islami F, Abnet CC, et al. Opium, tobacco, and alcohol use in relation to oesophageal squamous cell carcinoma in a high-risk area of Iran. *Br J Cancer* 2008;**98**:1857-63.
36. Wu MT, Lee YC, Chen CJ, Yang PW, Lee CJ, Wu DC, et al. Risk of betel chewing for oesophageal cancer in Taiwan. *Br J Cancer* 2001;**85**:658-60.
37. Wald NJ, Hackshaw AK, Frost CD. When can a risk factor be used as a worthwhile screening test? *BMJ* 1999;**319**:1562-5.
38. Wang Z, Tang L, Sun G, Tang Y, Xie Y, Wang S, et al. Etiological study of esophageal squamous cell carcinoma in an endemic region: a population-based case control study in Huaian, China. *BMC cancer* 2006;**6**:287.
39. Akbari MR, Malekzadeh R, Nasrollahzadeh D, Amanian D, Sun P, Islami F, et al. Familial risks of esophageal cancer among the Turkmen population of the Caspian littoral of Iran. *Int J Cancer* 2006;**119**:1047-51.
40. Guha N, Boffetta P, Wunsch Filho V, Eluf Neto J, Shangina O, Zaridze D, et al. Oral Health and Risk of Squamous Cell Carcinoma of the Head and Neck and Esophagus: Results of Two Multicentric Case-Control Studies. *Am J Epidemiol* 2007;**166**:1159-73.
41. Sepehr A, Kamangar F, Fahimi S, Saidi F. Poor oral health as a risk factor for esophageal squamous dysplasia in northeastern Iran. *Anticancer Res* 2005;**25**:543-6.
42. Abnet CC, Kamangar F, Islami F, Nasrollahzadeh D, Brennan P, Aghcheli K, et al. Tooth Loss and Lack of Regular Oral Hygiene Are Associated with Higher Risk of Esophageal Squamous Cell Carcinoma. *Cancer Epidemiol Biomarkers Prev* 2008;**17**:3062-8.
43. Tai SY, Wu IC, Wu DC, Su HJ, Lu CY, Huang JL, et al. Cigarette smoking and alcohol drinking and esophageal cancer risk in Taiwanese women. *World J Gastroenterol* 2010;**16**:1518-21.
44. Vízcaíno AP, Parkin DM, Skinner ME. Risk factors associated with oesophageal cancer in Bulawayo, Zimbabwe. *Br J Cancer* 1995;**72**:769-73.
45. Vaughan TL, Davis S, Kristal A, Thomas DB. Obesity, alcohol, and tobacco as risk factors for cancers of the esophagus and gastric cardia: adenocarcinoma versus

- squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 1995;**4**:85-92.
46. Castelletto R, Castellsague X, Munoz N, Iscovich J, Chopta N, Jmelnitsky A. Alcohol, tobacco, diet, mate drinking, and esophageal cancer in Argentina. *Cancer Epidemiol Biomarkers Prev* 1994;**3**:557-64.
 47. Sharp L, Chilvers CED, Cheng KK, McKinney PA, Logan RFA, Cook-Mozaffari P, et al. Risk factors for squamous cell carcinoma of the oesophagus in women: a case-control study. *Br J Cancer* 2001;**85**:1667-70.
 48. Castellsague X, Munoz N, De Stefani E, Victora CG, Castelletto R, Rolon PA. Influence of mate drinking, hot beverages and diet on esophageal cancer risk in South America. *Int J Cancer* 2000;**88**:658-64.
 49. De Stefani E, Boffetta P, Fagundes RB, Deneo-Pellegrini H, Ronco AL, Acosta G, et al. Nutrient patterns and risk of squamous cell carcinoma of the esophagus: a factor analysis in Uruguay. *Anticancer Res* 2008;**26**:2499-506.
 50. Islami F, Pourshams A, Nasrollahzadeh D, Kamangar F, Fahimi S, Shakeri R, et al. Tea drinking habits and esophageal cancer in a high risk area in northern Iran: population based case-control study. *BMJ* 2009;**338**:b929.
 51. Cook-Mozaffari PJ, Azordegan F, Day NE, Ressicaud A, Sabai C, Aramesh B. Oesophageal cancer studies in the Caspian Littoral of Iran: results of a case-control study. *Br J Cancer* 1979;**39**:293-309.
 52. Onuk MD, Oztopuz A, Memik F. Risk factors for esophageal cancer in Eastern Anatolia. *Hepatogastroenterology* 2002;**49**:1290-2.
 53. Wu M, Liu AM, Kampman E, Zhang ZF, Van't Veer P, Wu DL, et al. Green tea drinking, high tea temperature and esophageal cancer in high- and low-risk areas of Jiangsu Province, China: a population-based case-control study. *Int J Cancer* 2009;**124**:1907-13.
 54. De Jong UW, Breslow N, Hong JG, Sridharan M, Shanmugaratnam K. Aetiological factors in esophageal cancer in Singapore Chinese. *Int J Cancer* 1974;**13**:291-303.
 55. 55. Abedi-Ardekani B, Kamangar F, Hewitt SM, Hainaut P, Sotoudeh M, Abnet CC, et al. Polycyclic aromatic hydrocarbon exposure in esophageal tissue and risk of esophageal squamous cell carcinoma in north-eastern Iran. *Gut* 2010;**59**:1178-83.
 56. 56. Nabeya K, Onozawa K, Ri S. Brushing cytology with capsule for esophageal cancer. *Chir Gastroenterol* 1979;**13**:101-7.
 57. Lazarus C, Jaskiewicz K, Sumeruk RA, Nainkin J. Brush cytology technique in the detection of esophageal carcinoma in the asymptomatic, high risk subject; a pilot survey. *Cytopathology* 1992;**3**:291-6.
 58. Pan QJ, Roth MJ, Guo HQ, Kochman ML, Wang GQ, Henry M, et al. Cytologic Detection of Esophageal Squamous Cell Carcinoma and Its Precursor Lesions Using Ballon Samplers and Liquid-Based Cytology in Asymptomatic Adults in Linxian, China. *Acta Cytol* 2008;**52**:14-23.
 59. Lao-Sirieix P, Rous B, O'Donovan M, Hardwick RH, DeBiram I, Fitzgerald RC. Non-endoscopic immunocytological screening test for Barrett's oesophagus. *BMJ* 2007;**56**:1033-34.
 60. Kadri SR, Lao-Sirieix P, O'Donovan M, DeBiram I, Das M, Blazeby JM, et al. Acceptability and accuracy of a non-endoscopic screening test for Barrett's oesophagus in primary care: cohort study. *BMJ* 2010;**341**:c4372.
 61. Koike M, Hibi K, Kasai Y, Ito K, Akiyama S, Nakao A. Molecular Detection of Circulating Esophageal Squamous Cell Cancer Cells in the Peripheral Blood. *Clin Cancer Res* 2002;**8**:2879-82.
 62. Hibi K, Taguchi M, Nakayama H, Takase T, Kasai Y, Ito K, et al. Molecular Detection of p16 Promoter Methylation in the Serum of Patients with Esophageal Squamous Cell Carcinoma. *Clin Cancer Res* 2001;**7**:3135-8.
 63. Kannan S, Lakku RA, Niranjali D, Jayakumar K, Steven AH, Taralakhmi VV, et al. Expression of peanut agglutinin-binding mucin-type glycoprotein in human esophageal squamous cell carcinoma as a marker. *Mol Cancer* 2003;**5**:38.
 64. Yang YF, Li H, Xu XQ, Diao YT, Fang XQ, Wang Y, et al. An expression of squamous cell carcinoma antigen 2 in peripheral blood within the different stages of esophageal carcinogenesis. *Dis Esophagus* 2008;**21**:395-401.
 65. Chung J-Y, Braunschweig T, Hu N, Roth M, Traicoff JL, Wang Q-H, et al. A Multiplex Tissue Immunoblotting Assay for Proteomic Profiling: A Pilot Study of the Normal to Tumor Transition of Esophageal Squamous Cell Carcinoma. *Cancer Epidemiol Biomarkers Prev* 2006;**15**:1403-8.
 66. Munck-Wikland E, Kuylensstierna R, Wahren B, Lindholm J, Haglund S. Tumor markers carcinoembryonic antigen, CA 50, and CA 19-9 and squamous cell carcinoma of the esophagus pretreatment screening. *Cancer* 1988;**62**:2281-6.
 67. Fujita Y, Nakanishi T, Hiramatsu M, Mabuchi H, Miyamoto Y, Miyamoto A, et al. Proteomics-Based Approach Identifying Autoantibody against Peroxiredoxin VI as a Novel Serum Marker in Esophageal Squamous Cell Carcinoma. *Clin Cancer Res* 2006;**12**:6415-20.
 68. McCabe ML, Dlamini Z. The molecular mechanisms of esophageal cancer. *Int Immunopharmacol* 2005;**5**:1113-30.
 69. Jiang W, Zhang YJ, Kahn SM, Hollstein MC, Santella RM, Lu SH, et al. Altered expression of the cyclin D1 and retinoblastoma genes in human esophageal cancer. *Proc Natl Acad Sci U S A* 1993;**90**:9026-30.
 70. Saitoh T, Mine T, Katoh M. Molecular cloning and expression of proto-oncogene FRAT1 in human cancer. *Int J Oncol* 2002;**20**:785-9.
 71. Liu DB, Gu ZD, Cao XZ, Liu H, Li JY. Immunocytochemical detection of HoxD9 and Pbx1 homeodomain protein expression in Chinese esophageal squamous cell carcinomas. *World J Gastroenterol* 2005;**11**:1562-6.
 72. Li SL, Gao DL, Zhao ZH, Liu ZW, Zhao QM, Yu JX, et al. Correlation of matrix metalloproteinase suppressor genes RECK, VEGF, and CD105 with angiogenesis and biological behavior in esophageal squamous cell carcinoma. *World J Gastroenterol* 2007;**13**:6076-81.
 73. Kazemi-Noureini S, Colonna-Romano S, Ziaee AA, Mal-

- boobi MA, Yazdanbod M, Setayeshgar P, et al. Differential gene expression between squamous cell carcinoma of esophageus and its normal epithelium; altered pattern of mal, akr1c2, and rab11a expression. *World J Gastroenterol* 2004;**10**:1716-21.
74. Hollstein MC, Metcalf RA, Welsh JA, Montesano R, Harris CC. Frequent mutation of the p53 gene in human esophageal cancer. *Proc Natl Acad Sci U S A* 1990;**87**:9958-61.
 75. Hainaut P, Hernandez T, Robinson A, Rodriguez-Tome P, Flores T, Hollstein M, et al. IARC Database of p53 gene mutations in human tumors and cell lines: Updated compilation, revised formats and new visualisation tools. *Nucleic Acids Res* 1998;**26**:205-13.
 76. Zhou X, Tarmin L, Yin J, Jiang HY, Suzuki H, Rhyu MG, et al. The MTS1 gene is frequently mutated in primary human esophageal tumors. *Oncogene* 1994;**41**:3737-9.
 77. Mandard AM, Hainaut P, Hollstein M. Genetic steps in the development of squamous cell carcinoma of the esophagus. *Mutat Res* 2000;**462**:335-42.
 78. Gao H, Wang L-D, Zhou Q, Hong J-Y, Huang T-Y, Yang CS. p53 Tumor Suppressor Gene Mutation in Early Esophageal Precancerous Lesions and Carcinoma among High-Risk Populations in Henan, China. *Cancer Res* 1994;**54**:4342-6.
 79. Wang L-D, Hong J-Y, Qiu S-L, Gao H, Yang CS. Accumulation of p53 Protein in Human Esophageal Precancerous Lesions: A Possible Early Biomarker for Carcinogenesis. *Cancer Res* 1993;**53**:1783-7.
 80. Xing EP, Yang GY, Wang LD, Shi ST, Yang CS. Loss of heterozygosity of the Rb gene correlates with pRb protein expression and associates with p53 alteration in human esophageal cancer. *Clin Cancer Res* 1999;**5**:1231-40.
 81. Daigo Y, Nishiwaki T, Kawasoe T, Tamari M, Tsuchiya E, Nakamura Y. Molecular Cloning of a Candidate Tumor Suppressor Gene, DLC1, from Chromosome 3p21.3. *Cancer Res* 1999;**59**:1966-72.
 82. Xing EP, Nie Y, Wang L-D, Yang G-Y, Yang CS. Aberrant methylation of p16INK4a and deletion of p15INK4b are frequent events in human esophageal cancer in Linxian, China. *Carcinogenesis* 1999;**20**:77-84.
 83. Guo M, Ren J, House MG, Qi Y, Brock MV, Herman JG. Accumulation of Promoter Methylation Suggests Epigenetic Progression in Squamous Cell Carcinoma of the Esophagus. *Clin Cancer Res* 2006;**12**:4515-22.
 84. Boynton RF, Blount PL, Yin J, Brown VL, Huang Y, Tong Y, et al. Loss of heterozygosity involving the APC and MCC genetic loci occurs in the majority of human esophageal cancers. *Proc Natl Acad Sci U S A* 1992;**89**:3385-8.
 85. Zare M, Jazii F, Alivand M, Nasseri N, Malekzadeh R, Yazdanbod M. Qualitative analysis of Adenomatous Polypoid Coli promoter: Hypermethylation, engagement and effects on survival of patients with esophageal cancer in a high risk region of the world, a potential molecular marker. *BMC cancer* 2009;**9**:24.
 86. Kuroki T, Trapasso F, Shiraishi T, Alder H, Mimori K, Mori M, et al. Genetic alterations of the tumor suppressor gene WWOX in esophageal squamous cell carcinoma. *Cancer Res* 2002;**62**:2258-60.
 87. Cao LY, Yin Y, Li H, Jiang Y, Zhang HF. Expression and clinical significance of S100A2 and p63 in esophageal carcinoma. *World J Gastroenterol* 2009;**15**:8183-8.
 88. Mori M, Mimori K, Shiraishi T, Alder H, Inoue H, Tanaka Y, et al. Altered Expression of Fhit in Carcinoma and Precarcinomatous Lesions of the Esophagus. *Cancer Res* 2000;**60**:1177-82.
 89. Ishii T, Murakami J, Notohara K, Cullings HM, Sasamoto H, Kambara T, et al. Oesophageal squamous cell carcinoma may develop within a background of accumulating DNA methylation in normal and dysplastic mucosa. *Gut* 2007;**56**:13-9.
 90. Li C, Wu MY, Liang YR, Wu XY. Correlation between expression of human telomerase subunits and telomerase activity in esophageal squamous cell carcinoma. *World J Gastroenterol* 2003;**9**:2395-9.
 91. Zhao YG, Xiao AZ, Ni J, Man YG, Sang QX. Expression of matrix metalloproteinase-26 in multiple human cancer tissues and smooth muscle cells. *Ai Zheng* 2009;**28**:1168-75.
 92. Mukherjee S, Roth M, Dawsey S, Yan W, Rodriguez-Canales J, Erickson H, et al. Increased matrix metalloproteinase activation in esophageal squamous cell carcinoma. *J Transl Med* 2010;**8**:91.
 93. Chen H, Wang LD, Guo M, Gao SG, Guo HQ, Fan ZM, et al. Alterations of p53 and PCNA in cancer and adjacent tissues from concurrent carcinomas of the esophagus and gastric cardia in the same patient in Linzhou, a high incidence area for esophageal cancer in northern China. *World J Gastroenterol* 2003;**9**:16-21.
 94. Ishii H, Baffa R, Numata SI, Murakumo Y, Rattan S, Inoue H, et al. The FEZ1 gene at chromosome 8p22 encodes a leucine-zipper protein, and its expression is altered in multiple human tumors. *Proc Natl Acad Sci U S A* 1999;**96**:3928-33.
 95. Tanaka S, Akiyoshi T, Mori M, Wands JR, Sugimachi K. A novel Frizzled gene identified in human oesophageal carcinoma mediates APC/hcatenin signals. *Proc Natl Acad Sci U S A* 1998;**95**:10164-9.
 96. Mafune K, Tanaka Y, Mimori K, Mori M, Takubo K, Makuuchi M. Increased Expression of Ornithine Decarboxylase Messenger RNA in Human Esophageal Carcinoma. *Clin Cancer Res* 1999;**5**:4073-8.
 97. Liu Y, Wang HX, Lu N, Mao YS, Liu F, Wang Y, et al. Translocation of annexin I from cellular membrane to the nuclear membrane in human esophageal squamous cell carcinoma. *World J Gastroenterol* 2003;**9**:645-9.
 98. Qi YJ, Wang LD, Jiao XY, Feng XS, Fan ZM, Gao SS, et

- al. Dysregulation of Annexin II expression in esophageal squamous cell cancer and adjacent tissues from a high-incidence area for esophageal cancer in Henan province. *Ai Zheng* 2007;**26**:730-6.
99. Yang Z-Q, Imoto I, Fukuda Y, Pimkhaokham A, Shimada Y, Imamura M, et al. Identification of a Novel Gene, GASC1, within an Amplicon at 9p23-24 Frequently Detected in Esophageal Cancer Cell Lines. *Cancer Res* 2000;**60**:4735-9.
 100. Yue CM, Deng DJ, Bi MX, Guo LP, Lu SH. Expression of ECRG4, a novel esophageal cancer-related gene, downregulated by CpG island hypermethylation in human esophageal squamous cell carcinoma. *World J Gastroenterol* 2003;**9**:1174-8.
 101. Gholamin M, Moaven O, Memar B, Farshchian M, Naseh H, Malekzadeh R, et al. Overexpression and Interactions of Interleukin-10, Transforming Growth Factor β , and Vascular Endothelial Growth Factor in Esophageal Squamous Cell Carcinoma. *World J Surg* 2009;**33**:1439-45.
 102. Akbari MR, Malekzadeh R, Shakeri R, Nasrollahzadeh D, Foumani M, Sun Y, et al. Candidate Gene Association Study of Esophageal Squamous Cell Carcinoma in a High-Risk Region in Iran. *Cancer Res* 2009;**69**:7994-8000.
 103. Tsuda T, Tahara E, Kajiyama G, Sakamoto H, Terada M, Sugimura T. High Incidence of Coamplification of hst-1 and int-2 Genes in Human Esophageal Carcinomas. *Cancer Res* 1989;**49**:5505-8.
 104. Peralta RC, Casson AG, Wang R, Keshavjee S, Redston M, B. B. Distinct regions of frequent loss of heterozygosity of chromosome 5p and 5q in human esophageal cancer. *Int J Cancer* 1998;**78**:600-5.
 105. Iwaya T, Maesawa C, Ogasawara S, Tamura G. Tylosis esophageal cancer locus on chromosome 17q25.1 is commonly deleted in sporadic esophageal cancer. *Gastroenterology* 1998;**114**:1206-10.
 106. Kashyap MK, Marimuthu A, Kishore CJ, Peri S, Keerthikumar S, Prasad TS, et al. Genomewide mRNA profiling of esophageal squamous cell carcinoma for identification of cancer biomarkers. *Cancer Biol Ther* 2009;**8**:35-46.
 107. Adams L, Roth MJ, Abnet CC, Dawsey SP, Qiao Y-L, Wang G-Q, et al. Promoter Methylation in Cytology Specimens as an Early Detection Marker for Esophageal Squamous Dysplasia and Early Esophageal Squamous Cell Carcinoma. *Cancer Prev Res (Phila)* 2008;**1**:357-61.
 108. Akbari M, Malekzadeh R, Lepage P, Roquis D, Sadjadi A, Aghcheli K, et al. Mutations in Fanconi anemia genes and the risk of esophageal cancer. *Hum Genet* 2011;**129**:573-82.
 109. Akbari MR, Malekzadeh R, Nasrollahzadeh D, Amanian D, Islami F, Li S, et al. Germline BRCA2 mutations and the risk of esophageal squamous cell carcinoma. *Oncogene* 2007;**27**:1290-6.
 110. Eisenberger CF, Knoefel WT, Peiper M, Merkert P, Yekebas EF, Scheunemann P, et al. Squamous Cell Carcinoma of the Esophagus Can Be Detected by Microsatellite Analysis in Tumor and Serum. *Clin Cancer Res* 2003;**9**:4178-83.
 111. Sepehr A, Kamangar F, Abnet CC, Fahimi S, Pourshams A, Poustchi H, et al. Genetic polymorphisms in three Iranian populations with different risks of esophageal cancer, an ecologic comparison. *Cancer Lett* 2004;**213**:195-202.
 112. Moaven O, Raziee HR, Sima HR, Ganji A, Malekzadeh R, A'Rabi A, et al. Interactions between Glutathione-S-Transferase M1, T1 and P1 polymorphisms and smoking, and increased susceptibility to esophageal squamous cell carcinoma. *Cancer Epidemiol* 2010;**34**:285-90.
 113. Zhang X, Lin P, Zhu ZH, Long H, Wen J, Yang H, et al. Expression profiles of early esophageal squamous cell carcinoma by cDNA microarray. *Cancer Genet Cytogenet* 2009;**194**:23-9.
 114. Fu L, Qin YR, Xie D, Chow HY, Ngai SM, Kwong DLW, et al. Identification of alpha-actinin 4 and 67 kDa laminin receptor as stage-specific markers in esophageal cancer via proteomic approaches. *Cancer* 2007;**110**:2672-81.
 115. Zhu X, Ding M, Yu ML, Feng MX, Tan L-J, Zhao FK. Identification of galectin-7 as a potential biomarker for esophageal squamous cell carcinoma by proteomic analysis. *BMC Cancer* 2010;**10**:290.
 116. Shou JZ, Hu N, Takikita M, Roth MJ, Johnson LL, Giffen C, et al. Overexpression of CDC25B and LAMC2 mRNA and Protein in Esophageal Squamous Cell Carcinomas and Premalignant Lesions in Subjects from a High-Risk Population in China. *Cancer Epidemiol Biomarkers Prev* 2008;**17**:1424-35.
 117. Sakakibara T, Hibi K, Kodera Y, Ito K, Akiyama S, Nakao A. Plasminogen Activator Inhibitor-1 as a Potential Marker for the Malignancy of Esophageal Squamous Cell Carcinoma. *Clin Cancer Res* 2004;**10**:1375-8.
 118. Abnet C, Huppi K, Carrera A, Armistead D, McKenney K, Hu N, et al. Control region mutations and the 'common deletion' are frequent in the mitochondrial DNA of patients with esophageal squamous cell carcinoma. *BMC Cancer* 2004;**4**:30.
 119. Marjani HA, Biramijamal F, Rakhshani N, Hossein-Nezhad A, Malekzadeh R. Investigation of NQO1 genetic polymorphism, NQO1 gene expression and PAH-DNA adducts in ESCC. A case-control study from Iran. *Genet Mol Res* 2010;**9**:239-49.
 120. Feber A, Xi L, Luketich JD, Pennathur A, Landreneau RJ, Wu M, et al. MicroRNA expression profiles of esophageal cancer. *J Thorac Cardiovasc Surg* 2008;**135**:255-60.
 121. Mathe EA, Nguyen GH, Bowman ED, Zhao Y, Budhu A, Schetter AJ, et al. MicroRNA expression in squamous cell carcinoma and adenocarcinoma of the esophagus: associations with survival. *Clin Cancer Res* 2009;**15**:6192-200.
 122. Ogawa R, Ishiguro H, Kuwabara Y, Kimura M, Mitsui A, Katada T, et al. Expression profiling of micro-RNAs in human esophageal squamous cell carcinoma using RT-

PCR. *Med Mol Morphol* 2009;**42**:102-9.

123. Lee KH, Goan YG, Hsiao M, Lee CH, Jian SH, Lin JT, et al. MicroRNA-373 (miR-373) post-transcriptionally regulates large tumor suppressor, homolog 2 (LATS2) and stimulates proliferation in human esophageal cancer. *Exp Cell Res* 2009;**315**:2529-38.
124. Hiyoshi Y, Kamohara H, Karashima R, Sato N, Imamura Y, Nagai Y, et al. MicroRNA-21 regulates the proliferation and invasion in esophageal squamous cell carcinoma. *Clin Cancer Res* 2009;**15**:1915-22.
125. Guo Y, Chen Z, Zhang L, Zhou F, Shi S, Feng X, et al. Distinctive microRNA profiles relating to patient survival in esophageal squamous cell carcinoma. *Cancer Res* 2008;**68**:26-33.
126. Matsushima K, Isomoto H, Kohno S, Nakao K. MicroRNAs and Esophageal Squamous Cell Carcinoma. *Digestion* 2010;**82**:138-44.
127. Nicolas Perez D, Quintero E, Parra Blanco A. Screening the at-risk population for squamous cell carcinoma of the esophagus. *Gastroenterol Hepatol* 2005;**28**:337-46.
128. Montesano R, Holestein M, Hainui P. Genetic alterations in esophageal cancer and their relevance to etiology and pathogenesis: A review. *Int J Cancer* 1996;**69**:225-35.
129. Yokoyama T, Yokoyama A, Kumagai Y, Omori T, Kato H, Igaki H, et al. Health Risk Appraisal Models for Mass Screening of Esophageal Cancer in Japanese Men. *Cancer Epidemiol Biomarkers Prev* 2008;**17**:2846-54.