

Published in final edited form as:

Nat Ecol Evol. 2022 July ; 6(7): 1007–1023. doi:10.1038/s41559-022-01771-6.

A phylogenetic and proteomic reconstruction of eukaryotic chromatin evolution

Xavier Grau-Bové^{1,2}, Cristina Navarrete^{1,2}, Cristina Chiva², Thomas Pribasni³, Meritxell Antó⁴, Guifré Torruella⁵, Luis Javier Galindo⁵, Bernd Franz Lang⁶, David Moreira⁵, Purificación López-García⁵, Iñaki Ruiz-Trillo^{4,7}, Christa Schleper³, Eduard Sabidó^{1,2}, Arnau Sebé-Pedrós^{1,2,*}

¹Centre for Genomic Regulation (CRG), Barcelona Institute of Science and Technology (BIST), Barcelona 08003, Catalonia, Spain

²Universitat Pompeu Fabra (UPF), Barcelona 08003, Catalonia, Spain

³Department of Functional and Evolutionary Ecology, Archaea Biology Unit, University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria

⁴Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta, 37-49, Barcelona 08003, Catalonia, Spain

⁵Unité d'Ecologie Systématique et Evolution, CNRS, Université Paris-Saclay, AgroParisTech, Orsay, France

⁶Robert Cedergren Centre in Bioinformatics and Genomics, Department of Biochemistry, Université de Montréal, Montréal, Quebec H3C 3J7, Canada

⁷ICREA, Pg. Lluís Companys 23, Barcelona 08010, Catalonia, Spain

Abstract

Histones and associated chromatin proteins have essential functions in eukaryotic genome organization and regulation. Despite this fundamental role in eukaryotic cell biology, we lack a phylogenetically-comprehensive understanding of chromatin evolution. Here, we combine comparative proteomics and genomics analysis of chromatin in eukaryotes and archaea.

Proteomics uncovers the existence of histone post-translational modifications in Archaea.

However, archaeal histone modifications are scarce, in contrast with the highly conserved and abundant marks we identify across eukaryotes. Phylogenetic analysis reveals that chromatin-associated catalytic functions (e.g., methyltransferases) have pre-eukaryotic origins, whereas

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

*Corresponding author: arnau.sebe@crg.eu.

Author contributions

A.S.-P. conceived the project. X.G.-B., C.C., I.R.-T., C.S., E.S. and A.S.-P. designed experiments and analytical strategies. C.N., T.P., M.A. and A.S.-P. performed experiments. X.G.-B., C.C., and A.S.-P. analyzed the data. T.P., G.T., L.J.G., D.M., P.L.-G. and B.F.L. provided biological samples/cultures and genomic data. All authors contributed to data interpretation. X.G.-B. and A.S.-P. wrote the manuscript with input from all authors.

Declaration of interests

The authors declare no competing interests.

histone mark readers and chaperones are eukaryotic innovations. We show that further chromatin evolution is characterized by expansion of readers, including capture by transposable elements and viruses. Overall, our study infers detailed evolutionary history of eukaryotic chromatin: from its archaeal roots, through the emergence of nucleosome-based regulation in the eukaryotic ancestor, to the diversification of chromatin regulators and their hijacking by genomic parasites.

Introduction

The access to genetic information in eukaryotes is controlled by a manifold nucleoproteic interface called chromatin. This nucleosomal chromatin environment defines a repressive ground state for transcription and other DNA-templated processes in eukaryotic genomes^{1,2}. Multiple components associated with chromatin underlie elaborate eukaryotic genome regulation, allowing the differential access to genetic information in time/space and the maintenance of the resulting regulatory states^{3–6}. Moreover, chromatin-based regulation is essential in repressing parasitic genomic elements, like transposons and viruses^{7–11}.

The main protein components of eukaryotic chromatin are histones. All eukaryotes have four major types of histones (H2A, H2B, H3 and H4), which are combined as an octamer to form the basic repetitive unit of the chromatin: the nucleosome. Canonical histones are among the most highly conserved proteins across eukaryotes¹² and, in addition, unique histone variants (paralogs of one of the four major histone types) are found in many species, often associated with particular regulatory states^{13–17}. Histone chemical modifications, including acetylations and methylations play a central role in genome regulation and transgenerational epigenetic inheritance^{3,18–21}. These chemical moieties, known as histone post-translational modifications (hPTMs), are added and removed by specific enzymes ('writers', e.g., histone methyltransferases or acetylases; and 'erasers', e.g., histone demethylases and deacetylases). Some hPTMs (e.g., most acetylations) have a generic effect on nucleosome stability, while others are bound by specific proteins or protein complexes. These are often referred to as 'readers' and include proteins like HP1, which binds to H3K9me3, as well as a myriad of other proteins encoding Chromo, PHD, Tudor and Bromo structural domains, among others^{22–24}. Finally, nucleosome remodellers (like SNF2 proteins) and histone chaperones are additional important players in chromatin regulation, by mediating chromatin opening, nucleosomal assembly, and histone variant interchanges^{25–28}.

All eukaryotes studied to date possess histone-based chromatin organization, with the sole exception of dinoflagellates, which nonetheless encode for histone proteins in their genomes²⁹. Beyond eukaryotes, histones have also been identified in Archaea, where they have been shown to form nucleosomal structures^{30–33}. However, unlike eukaryotic histones, the few archaeal histones experimentally characterized so far (*i*) generally lack disordered *N*-terminal tails; (*ii*) do not have any known post-translational modifications³⁴; and (*iii*) do not seem to impose a widespread, genome-wide repressive transcriptional ground state^{33,35}. Thus, chromatin-based elaborate genome regulation is often considered a eukaryotic innovation^{36,37}.

From a phylogenetic perspective, our understanding of chromatin components and processes derives from a very small set of organisms, essentially animal, fungal and plant model species plus a few parasitic unicellular eukaryotes. Additional efforts have sampled specific aspects of chromatin regulation, such as histone modifications or their genome-wide distribution, in non-model animal species^{38,39}, fungi (*Neurospora crassa* and *Fusarium graminearum*)^{40,41}, and five other eukaryotes: the unicellular holozoan *Capsaspora owczarzakii*⁴², the dinoflagellate *Hematodinium* sp.²⁹, the brown alga *Ectocarpus siliculosus*⁴³, the amoebozoan *Dictyostelium discoideum*⁴⁴, and the ciliate *Tetrahymena thermophila*^{45,46}. However, these organisms represent a tiny fraction of eukaryotic diversity. Hence, we lack a systematic understanding of the evolution of eukaryotic chromatin modifications and components⁴⁷.

In order to infer the origin and evolutionary diversification of eukaryotic chromatin, we performed a joint comparative analysis of histone proteomics data from 30 different eukaryotic and archaeal taxa, including new data for 23 species. In parallel, we analyzed the complement of chromatin-associated gene families in an additional 172 eukaryotic genomes and transcriptomes. This comprehensive taxon sampling includes representatives of all major eukaryotic lineages, as well as multiple free-living members of enigmatic early-branching eukaryotes (e.g., jakobids, malawimonads, *Meteora* sp. and ancyromonads, as well as Collodictyonida, Rigifilida and Mantamonadida (CRuMS); Fig. 1a). In addition, in order to trace the pre-eukaryotic origins of these chromatin gene families, we systematically searched for orthologs in archaeal, bacterial and viral genomes. Specifically, we reconstructed the evolutionary history of enzymes involved in chromatin modification and remodelling; as well as the conservation of the hPTMs effected by these enzymes. Our comparative genomics and proteomics suggest a concurrent and early origin of canonical histones, a core of quasi-universal hPTMs, and their corresponding enzymatic effectors. We also identify independent expansions in hPTM reader gene families across eukaryotes and document evidence of the capture of these reader domains by parasitic genomic elements. Overall, this work provides a phylogenetically-informed framework to classify and compare chromatin components across the eukaryotic tree of life, and to further investigate the evolution of hPTM-mediated genome regulation.

Results

Comparative proteomics of eukaryotic histone modifications

We analyzed the phylogenetic distribution and evolutionary history of histone proteins. To this end, we surveyed the presence of histone-fold proteins across 172 eukaryotic and 4,226 archaeal taxa, using HMM searches (Fig. 1a,b and Supplementary Data 1). Histone proteins are found in all eukaryotic genomes. We clustered the identified 8,576 histone-encoding proteins using pairwise local alignments and then classified individual sequences in these clusters based on pairwise alignments to a reference database⁴⁸ (Fig. 1a and Extended Data Fig. 1a). This reveals four broad clusters corresponding to the four main eukaryotic histones (H2A, H2B, H3, and H4) and their variants (H2A.Z, macroH2A, and cenH3), as well as a fifth cluster composed of archaeal HMfB homologs. Finally, this classification also uncovers three large connected components composed of transcription factors with

histone-like DNA binding domains, which are widely distributed in eukaryotes (POLE3, POLE4, DR1) and/or archaea (NFYB). Further analysis of the genomic distribution of these histone genes shows a frequent occurrence of H3-H4 and H2A-H2B pairs in head-to-head orientation (5' to 5'), strongly indicating co-regulation across eukaryotes (Extended Data Fig. 1b,c and Supplementary Data 2).

Next, we investigated the distribution and conservation of hPTMs across major eukaryotic groups and Archaea, including methylations, acetylations, crotonylations, phosphorylations, and ubiquitylations. To this end, histones from 19 different eukaryotic species were extracted, chemically derivatized⁴⁹ and analyzed by mass-spectrometry (Fig. 1c and Supplementary Data 3), adding to previously available hPTM proteomics data for additional seven species. Our extensive taxon sampling covers all major eukaryotic groups, as well as hitherto unsampled early-diverging eukaryotic lineages—such as the malawimonad *Gefionella okellyi*, the discoban *Naegleria gruberi*, or the ancyromonad *Fabomonas tropica*—, thus providing a comprehensive comparative framework for evolutionary inference.

We focused first on hPTMs present in canonical histones, as defined by their highly conserved *N*-terminal regions, phylogenetic analyses, and sequence similarity to curated reference canonical histones (Fig. 1d; see Methods). hPTMs are detected in all canonical histones from all species. After correcting by sequence coverage, we observe that hPTMs are particularly abundant in H3 canonical histones (median = 23.5 hPTMs per species, mean = 24.3), compared with H2A, H2B and H4 (medians between 6.5 and 9, means between 9.5 and 13.4; Extended Data Fig. 2a). Holozoan canonical H2As (*Homo sapiens*, *Sycon ciliatum* and *Capsaspora owczarzaki*) represent an exception to this trend and contain similar number of modifications to H3s in these species. We also examined the reproducibility of hPTM detection across replicate samples, showing that the majority of hPTMs (87.5%) can be found in more than one sample (Extended Data Fig. 2b,c). Despite this, it is worth emphasizing that our data may contain false negatives, beyond the lack of coverage for particular residues that we systematically report. For example, some marks might be globally too scarce in the nucleosomes of a particular species, while other modifications like phosphorylations and ubiquitination are difficult to detect by mass-spectrometry without dedicated peptide-enrichment protocols.

Canonical H3 and H4 *N*-terminal tails contain the majority of phylogenetically-conserved hPTMs, in stark contrast with the relative paucity of conserved hPTMs in canonical H2A and H2B. A striking example of paneukaryotic conservation comes from the acetylation of the H4 K5, K8, K12 and K16 residues (Fig. 1d, second panel), all of which mark gene expression-permissive chromatin environments in multiple eukaryotic species²². A similar conservation pattern is observed in the acetylation of a group of *N*-terminal H3 lysines (K9, K14, K18, K23, K27) associated with similar functions, while other H3 acetylations are only found in a few species (e.g., residues K4, K56 and K79). While acetylations are highly conserved, only seven histone H3/H4 methylations are broadly conserved across eukaryotic lineages: H3K4me1/2/3, H3K9me1/2/3, H3K27me1/2/3, H3K36me1/2/3, H3K37me1/2/3 and, more sparsely, H3K79me1/2 and H4K20me1. Many of these broadly conserved marks have conserved roles in demarcating active (e.g., H3K4me) and repressive chromatin states (e.g., H3K9me and H3K27me)^{22,42,50}. The scarcity of conserved hPTMs in H2A and H2B

canonical histones can partially explained by their higher degree of sequence divergence (Fig. 1e), which is reflected in many non-homologous lysine residues (Fig. 1d). But even among homologous positions, we found little evidence of conservation, with the exception of H2A K5ac (associated to active promoters⁵¹) and, in fewer species, methylation of H2A K5 and H2B K5. Finally, we were also able to identify phosphorylations in serine and threonine residues and a few instances of ubiquitylation. In general, these marks show more restricted phylogenetic distributions than lysine acetylation or methylation, even in the tightly conserved H3 and H4 histones. We can identify conserved phosphorylations in H2A T120 and S122, which are shared by most opisthokonts, and the ubiquitylation of H2A K119 only in some holozoan species.

Mass-spectrometry analysis detected histone variants in all species included in our study, suggesting that they are relatively abundant in the chromatin of these eukaryotes (Fig. 1e). Most of these variants are lineage-specific, with the exception of the paneukaryotic variants H2A.Z, H3/cenH3 and H3.3; and the macroH2A variant found in holozoans and *Meteora* sp. (belonging to an orphan eukaryotic lineage). Interestingly, we find hPTMs in the vast majority of detected variants, both conserved and lineage-specific, particularly acetylations and methylations (Fig. 1e and Extended Data Fig. 2d). Overall, our comparative proteomic analysis suggests the existence of a highly conserved set of canonical hPTMs of ancestral eukaryotic origin in H3 and H4, which co-exists with less conserved hPTMs in H2A, H2B, and lineage-specific modifications in variant histones.

Archaeal histone post-translational modifications

In contrast with the paneukaryotic distribution of histones, sequence searches show that only a fraction of archaeal genomes encode for histones (28.1% of the taxa here examined; Fig. 2a). Archaeal histones exhibit a patchy phylogenetic distribution, similar to other gene families shared with eukaryotes⁵². Among others, histones are present in Euryarchaeota, the TACK superphylum and Asgard archaea^{12,53–56}. Asgard are generally considered to be the closest known archaeal relatives of eukaryotes^{57,58}, although this sister-group relationship has been challenged by some studies⁵⁹. Our extended sampling revealed that Asgard archaea histones, particularly in the Lokiarchaeota and Heimdallarchaeota clades⁵⁵, often have lysine-rich *N*-terminal tails in the manner of eukaryotic histones (Fig. 2a-c). These Asgard histones appear to be conserved across multiple taxa, albeit without direct sequence similarity compared to canonical eukaryotic histones (Extended Data Fig. 1d). When compared against eukaryotic sequences classified in HistoneDB⁴⁸, these archaeal histones clearly cluster in a separate group and are most similar to either eukaryotic H4 or, to a lesser degree, H3 canonical histones, in line with previous findings^{12,55,60}.

To identify potential archaeal hPTMs, we performed proteomics analysis of histones in three Euryarchaeota (the Methanobacteriota *Methanobrevibacter cuticularis* and the Halobacteriota *Methanospirillum stamsii* and *Methanosarcina spelaei*) and one Thaumarchaeota species (*Nitrososphaera viennensis*; Fig. 2b). Mass-spectrometry detects histone proteins in all of them: 2-4 in the euryarchaeotes (with 27-90% protein coverage) and one in the thaumarchaeote (80% protein coverage), including homologs with *N*-terminal tails encoded by each of the three euryarchaeotes in our survey (22-40 aa, 0.09-28 lysines

per residue; Fig. 2c). Moreover, this proteomics analysis finds evidence of hPTMs in archaeal histones. However, in comparison with eukaryotic histones, hPTMs are extremely scarce in archaeal histones. Specifically, we identify no hPTMs in *N. viennensis* and *M. spelaei* (one and two histones detected, respectively), three acetylations and one methylation in *M. stamsii* (in three out of four histones detected), and one acetylation and two methylations in *M. cuticularis* (in two out of four histones; Fig. 2b, top). Interestingly, we find conserved lysine residues with shared modifications in *M. stamsii* and *M. cuticularis* (methylation in K54 and acetylation in K57; Fig. 2b, bottom). This result indicates that highly-abundant hPTMs represent a eukaryotic innovation, likely linked to dynamic nucleosomal regulation in eukaryotes but not in Archaea.

Taxonomic distribution of chromatin-associated proteins

hPTMs are deposited and removed by specific modifying enzymes ('writers' and 'erasers'), while 'reader' protein domains found in diverse proteins bind and recognize specific hPTMs. For example, Bromo and Chromo domains bind acetylated and methylated lysine residues, respectively. In addition, the control of histone loading/eviction from specific genomic *loci* is mediated by chromatin remodellers, like SNF2 proteins²⁷, and histone chaperones²⁶. To date, the classification and evolutionary analysis of this chromatin machinery has been based on biased, partial taxonomic samplings and has not employed phylogenetic methods⁶¹ (with rare exceptions^{12,27}), often resulting in inaccurate orthologous relationships and confounded classification and naming schemes.

We sought to obtain a systematic, phylogenetics-based classification of histone remodellers, chaperones, readers, and modifiers in order to understand the evolutionary history of eukaryotic chromatin (Fig. 3a). To this end, we (*i*) compiled a taxa-rich dataset of 172 eukaryotic genomes and transcriptomes, covering all major eukaryotic supergroups and devoting particular attention to early-branching, non-parasitic lineages (Supplementary Data 1), as well as genomic data from 4,226 Archaea, 24,886 Bacteria and 185,579 viral taxa; (*ii*) defined a protein structural domain as a proxy for each gene family (Supplementary Data 4) and retrieved all genes in these genomes that contained these domains; and (*iii*) inferred accurate orthology groups from phylogenetic analyses of each gene class (next section).

We examined the taxonomic distribution and abundance of the major gene classes (Fig. 3b,c). Many domains with chromatin-associated functions in eukaryotes are also present in Archaea and Bacteria, albeit with scattered phylogenetic distributions (Fig. 3b and Extended Data Fig. 3a,b). Families with prokaryotic homologs include mostly catalytic gene classes (writer, eraser and remodeller enzymes), whereas readers and histone chaperones are virtually absent from prokaryotes (Fig. 3b). Histone fold-encoding genes constitute a case in point for this patchy distribution of chromatin proteins in prokaryotes: they are present in most archaeal phyla, but are absent in about half of the sampled genomes within each (Fig. 3b). Yet, there is a qualitative difference between the phylogenetic distribution of archaeal and bacterial chromatin-associated gene classes: whereas archaeal histones tend to co-occur with chromatin-associated gene classes, the bacterial complement of writers and erasers is much less conserved and is uncorrelated with the extremely rare presence of histone-like genes (Fig. 3d).

Within eukaryotes, most gene structural classes associated with chromatin functions are ubiquitously distributed across all lineages here surveyed, supporting an early eukaryotic origin for the core chromatin machinery (Fig. 3b and Extended Data Fig. 3d). In fact, the total number of chromatin writer, eraser and remodeller enzymes remains remarkably stable across eukaryotes (Fig. 3e). The only exception is the marked increase in genes encoding reader domains observed in lineages exhibiting complex multicellularity: animals, streptophyte plants, and, to a lesser degree, phaeophyte brown algae (Stramenopila). This occurs partially due to the addition of new gene classes (e.g., SAWADEE in the *Plantae s.l.* + Cryptista lineage, or ADD_DNMT3 in bilaterians and cnidarians), but also via the expansion of ancient, widely-distributed reader gene classes (e.g., Tudor, PHD, Chromo or Bromo domains). These taxonomic patterns indicate that chromatin modifying and remodelling catalytic activities originated in prokaryotes, while reader and chaperone structural domains are eukaryotic innovations.

Phylogenetics of chromatin modifiers and remodellers

To gain detailed insights into the origin and evolution of chromatin gene families, we used phylogenetic analysis to define orthology groups from paneukaryotic gene trees. We surveyed 172 eukaryotic species and defined a total of 1,713 gene families (orthogroups), 95% of which were conserved in two or more high-ranking taxonomic groups (as listed in Fig. 1a), and which included 51,426 genes in total (Supplementary Data 5). We annotated each gene family according to known members from eukaryotic model species. For simplicity, we use a human-based naming scheme throughout the present manuscript (unless otherwise stated), but we also provide a dictionary of orthologs in three additional model species (*Arabidopsis thaliana*, *Saccharomyces cerevisiae* and *Drosophila melanogaster*; see Supplementary Data 5). This phylogenetic classification scheme of eukaryotic chromatin gene families, as well as the sequences and associated phylogenetic trees, can be explored and retrieved in an interactive database: https://sebe-lab.shinyapps.io/chromatin_evolution

We first investigated the potential pre-eukaryotic origins of these gene families/orthogroups by comparing their phylogenetic distance to prokaryotic sequences and to other eukaryotic orthogroups (Fig. 4a). Most eukaryotic gene families are more closely related to other eukaryotes than to prokaryotic sequences, supporting the idea that writers, erasers, remodellers and readers diversified within the eukaryotic lineage, as previously noted for histones¹². This analysis also reveals a substantial fraction of eukaryotic gene families with close orthogroups in Archaea and Bacteria, which pinpoints components that were (i) inherited from a prokaryotic ancestor during eukaryogenesis; (ii) laterally transferred between eukaryotes and prokaryotes at later stages; or (iii) a combination of both phenomena. For example, we identified a well-supported sister-group relationship between the eukaryotic SIRT7 deacetylase and a clade of Asgard archaea Sirtuin enzymes (Heimdallarchaeota and Lokiarchaeota), a topology compatible with an archaeal origin or ancient transfers to/from Asgard and eukaryotes⁶²; whereas SIRT6 appears nested within other eukaryotic sequences (Fig. 4b, left). Likewise, the KAT14 acetylase is more closely related to bacterial enzymes than to other eukaryotic acetylases (Fig. 4b, right).

Next, we mapped the phylogenetic distribution of orthogroups in order to infer the origin and diversification of individual chromatin gene families (Fig. 4c and Extended Data Fig. 4a). Using probabilistic inference of ancestral gene content, we reconstruct a rich Last Eukaryotic Common Ancestor (LECA) complement of chromatin-associated gene families: 65 acetylases (amongst which 61 were conserved in at least two of the most deeply sampled eukaryotic early-branching lineages, namely Amorphea, Diaphoretickes, and Discoba); 20 deacetylases (19 in these early-branching eukaryotic lineages); 59 methyltransferases (55); 42 demethylases (38); 33 remodellers (33); and 25 chaperones (18) (Fig. 4c and Supplementary Data 5). The subsequent evolution of these families is characterized by relative stasis, with few new orthologous families emerging in later-branching eukaryotic lineages. Notable exceptions include the origin of KAT5 deacetylases and KMT5B/C SET methyltransferases in Opisthokonta; KAT8 and SIRT7 in Holozoa; and Viridiplantae-specific deacetylases (homologs of *A. thaliana* HDA7 and HDA14 deacetylases) and SETs (*A. thaliana* PTAC14); among others.

In spite of their broad distributions across eukaryotes, many chromatin modifier families exhibit variation in their protein domain architectures, likely conferring them functional properties such as distinct binding preferences (Extended Data Fig. 4b). For example, most CREBBP/EP300 acetylases consist of a catalytic HAT_KAT11 domain and two TAZ and ZZ zinc finger domains, but different lineages have acquired different reader domains: an acetylation-reading Bromo domain in holozoans and stramenopiles, PHD in plants and some stramenopiles, and no known reader domains in other lineages (e.g., in the fungal orthologs of the *S. cerevisiae* protein RTT109). A similar pattern is apparent in SET methyltransferase families sharing a core catalytic domain (SET) harboring variable DNA- and chromatin-interacting domains – animal SETDB1/2 homologs have MBD domains that bind CpG methylated DNA, while plants have SAD_SAR domains with the same function; and holozoan ASH1L homologs encode Bromo and BAH readers, whereas phaeophytes encode PHD domains (Extended Data Fig. 4b). Other architectures, however, are much more conserved, as exemplified by the presence of Tudor-knot and MYST zinc finger domains in most KAT5 deacetylases; or the ubiquitous co-occurrence of Helicase-C and SNF2_N domains in most remodellers (Extended Data Fig. 4b).

Specific examples of evolutionarily conserved chromatin gene families include the catalytic core and the subunits of well-studied chromatin complexes⁶³ like PRC1 (RING1/AB, PCGF), PRC2 (EZH1/2, SUZ12, EED, RBBP4/7) and Trithorax/MLL (MLL1/2/3/4, WRD5, ASH2L, RBBP5, DPY-30; Fig. 4d,e). However, when we compared the distribution of these complexes with the hPTMs they are related to, we found a generally poor co-occurrence (Fig. 4f-h). For example, organisms like *Dictyostelium discoideum* and *Creolimax fragrantissima* lack EZH1/2 orthologs, but we detected H3K27me3 in these species; while *Thecamonas trahens* and *Naegleria gruberi* lack Dot1 orthologs but have H3K79me marks. A poor correlation is also observed between the occurrence of H3K9me and that of SUV39H1 orthologs. An exception to this pattern is the ubiquitous distribution of H4K16ac and the acetylase family KAT5/8⁶⁴ (Fig. 4h). These patterns suggest that the specificity between hPTMs and their writers might not be completely conserved across eukaryotes, with distinct members of the same gene classes (e.g., methyltransferases) performing similar roles. In this context, reading domains present in writing/erasing

enzymes (directly in the same protein or as part of multi-protein complexes) are likely to play a major role in the re-purposing of chromatin catalytic activities.

Evolutionary expansion of chromatin readers

Multiple protein structural domains have been involved in the recognition of hPTMs, such as Bromo domains binding to acetylated lysines or Chromo, MBT and Tudor domains binding to methylated lysines^{23,24}. These are generally small domains and can be found both as stand-alone proteins as well as in combination with other domains, often catalytic activities such as hPTM writers, erasers and remodellers. Thus, they are central in the establishment of functional connections between chromatin states. To understand the contribution of these reading domains to the evolutionary diversification of chromatin networks, we studied in detail the phylogeny and protein architecture of reader domains across eukaryotes.

We quantified the co-occurrence frequency of reader and catalytic domains, finding (*i*) that most reader domains are present in genes without writer, eraser or remodeller domains (87%, Fig. 5a); and (*ii*) that most cases of reader-catalytic co-occurrence involve PHD, Chromo and Bromo domains (Extended Data Fig. 5a). For example, the conserved architecture of the paneukaryotic CHD3/4/5 re-modellers includes Chromo readers in most species and PHD domains specifically in animals and plants (Extended Data Fig. 4b). Likewise, PHD domains are often present in the KMT2A/B and KMT2C/D SET methyltransferase; and the ASH1L family has recruited Bromo and BAH domains in holozoans, and PHD in multicellular stramenopiles (Extended Data Fig. 4b). In spite of these redundancies, reader families typically have independent evolutionary histories, as illustrated by the fact that most reader domain-containing genes encode only one such domain (92%, Extended Data Fig. 5b).

We next performed phylogenetic analyses of individual reader domains and reconstructed the gains and losses of these reader gene families/orthogroups (Fig. 5a). Compared to the relative stasis of catalytic enzyme families, this reader-centric analysis revealed a strikingly different evolutionary pattern of lineage-specific bursts of innovation, particularly amongst PHD, Chromo and Bromo genes, as well as Tudor in animals (Fig. 5a and Extended Data Fig. 5c). PHD, Chromo and Bromo families also appeared as the most abundant in the reconstructed LECA reader domain repertoire, which amounted to 89 gene families (Fig. 5a, left). The distribution of gene family ages in extant species also corroborates that more readers have emerged at evolutionarily more recent nodes of the tree of life than catalytic gene families (Fig. 5b).

Co-option of chromatin machinery by transposable elements

Further examination of the domain co-occurrence networks of readers revealed that Chromo and PHD domains are often present together with protein domains found in transposable elements (TEs; Fig. 5c and Supplementary Data 6), including retrotransposons (e.g., retrotranscriptases and integrases; orange modules in Fig. 5c) and DNA transposons (e.g., DNA binding domains and transposases; red modules). It is known that some TEs show insertion-preferences associated to specific chromatin states⁶⁵, often mediated by direct chromatin tethering mechanisms⁶⁶. For example, the Chromo domain of the MAGGY gypsy

retrotransposon of the fungus *Magnaporthe grisea* targets H3K9me regions⁶⁷. Reciprocally, some protein domains of TE origin, often DNA-binding domains, have been co-opted into chromatin and transcriptional regulators⁶⁸. Thus, we decided to explore in detail the occurrence of chromatin-associated domain (readers, but also catalytic domains) linked to TEs in the 172 eukaryotic genomes in our dataset (Fig. 5d). Moreover, we used available RNA-seq datasets in many of these species to validate some of these TE fusions (Fig. 5d-e). A fully validated fusion gene would (i) come from a non-discontinuous gene model in the original assembly, and (ii) have evidence of expression, with reads mapping along the entire region between the TE-associated domain and the chromatin-associated domain (**Extended Data Fig. S6**).

We identified 823 predicted gene models containing both chromatin- and TE-associated domains (Fig. 5d). Whilst these TE fusions were not exclusive of reader domains, most such fusions involved PHD and Chromo-encoding genes; followed by SNF2_N remodellers, SET methyltransferases, and others. An homology search against a database of eukaryotic TEs revealed that most of these candidate TE fusions could be aligned to known retrotransposons or DNA transposons. For example, by way of validation, our analysis identifies the SETMAR human gene, a previously-described fusion between a SET methyltransferase and a Mariner-class DNA transposon⁶⁹. Overall, 31% of the candidate fusion genes were supported by valid gene models according to our stringent criteria (Fig. 5d). Interestingly, we find very few cases of hypothetical fusions between TEs and Bromo domains, which recognize K acetylations and are otherwise highly abundant across eukaryotes, and none of them is validated by RNA-seq data. This could be explained by the detrimental effect of targeting TE insertions to sites of active chromatin demarcated by histone acetylations, such as promoter and enhancer elements.

Some of these validated fusions have a broad phylogenetic distribution (Fig. 5e), such as a Gypsy-ERV retrotransposon with a C-terminal Chromo domain (Unk. Chromo 2.1 in Fig. 5e) that is widely distributed in animals and various microbial eukaryotes, and contains dozens of paralogs in vertebrate *Danio rerio* or the charophyte *Chara braunii*, many of which are expressed. Another widespread Gypsy-ERV retrotransposon with a Chromo domain is present in multiple expressed and highly similar copies in the fungus *Rhizopus delemar* (Fig. 5f,e), suggesting a successful colonization of this genome by this TE. By contrast, other TE fusions are taxonomically restricted to one or few related species, such as the fusion of hAT activator DNA transposons with Chromo CBX and CDY readers in the sponge *Ephydatia muelleri*; or multiple instances of fusions with Chromo and PHD readers in cnidarians. A common fusion in cnidarians involves different retrotransposon classes with PHD domains orthologous to the PYGO1/2 protein (Fig. 5e), which is known to recognize specifically H3K4me⁷⁰. Globally, this analysis reveals that recruitment of chromatin reading and even modifying domains by TE has occurred in many eukaryotic species, in a way that might facilitate the evasion from suppressing mechanisms in the host genomes as suggested by the expansion of Chromo-fused TEs in the genomes of *Chara braunii* (Viridiplantae), *Chromera velia* (Alveolata) and *Rhizopus delemar* (Fungi).

Chromatin components in viral genomes

In addition to TEs, chromatin is also involved in the suppression of another type of genomic parasites: viruses. Some chromatin-related genes, including histones, have been found in viral genomes, especially among the nucleocytoplasmic large DNA viruses – also known as giant viruses. Eukaryotic core histones have been even hypothesized to have evolved from giant virus homologs, after the discovery that certain *Marseilleviridae* genomes encoded deeply-diverging orthologs of the four canonical histones⁷¹. These viral histones have been recently shown to form nucleosome-like particles that package viral DNA^{72,73}.

We analyzed the distribution and abundance of chromatin-related protein domains among viruses, including data from 1,816 giant virus genomes. Based on structural domain searches, we identified 2,163 viral chromatin-related proteins (Fig. 5g and Supplementary Data 6). The majority of these proteins are encoded by giant viruses (55%), followed by *Caudovirales* (37%). Among these two groups, only giant virus genomes encode histones – specifically, the *Iridoviridae*, *Marseilleviridae*, *Mimiviridae*, *Pithoviridae*, and *Phycodnaviridae* families. Concordantly with previous studies⁷⁴, we also identify remodellers in all giant virus families; as well as less abundant components of the chromatin writer/eraser/reader toolkit (Fig. 5g).

We then investigated the phylogenetic affinities of these viral chromatin proteins, starting with histones (Fig. 5h). Our analysis recovers the phylogenetic affinity of *Marseilleviridae* histones with specific eukaryotic histone families⁷¹, and makes this pattern extensive to *Mimiviridae*, *Iridoviridae*, and *Pithoviridae* giant viruses (Fig. 5h), with the caveat of the ambiguous clustering of the H4-like viral histones with either H4 eukaryotic or archaeal HMF genes. In all these lineages, we identify genes encoding two histone-fold domains orthologous to H2B + H2A (inset table in Fig. 5h), whereas the H4 + H3 histone doublet genes appears to be exclusive to *Marseilleviridae*. By contrast, histone homologs in *Phycodnaviridae*, *Pandoraviridae* (also giant viruses), and *Polydnaviridae* (*incertae sedis*) are never found as either doublets or as early-branching homologs of eukaryotic histones, suggesting recent acquisition from eukaryotes.

Unlike histones, most of the viral chromatin-associated genes exhibited a mixture of prokaryotic and eukaryotic phylogenetic affinities and often lack affinity to any specific eukaryotic gene family (Fig. 5i and Extended Data Fig. 7). Viral readers, on the other hand, are often embedded within eukaryotic clades in gene trees and are similar to *bona fide* eukaryotic families, exhibiting topologies consistent with recent, secondary acquisitions. This is the case of BIRC2/3/XIAP readers widespread in the *Baculoviridae*, which encode BIR domains that are often hijacked from their hosts⁷⁵. We also find a number of viral Chromo-encoding genes, which fall in two main taxonomic categories: (i) giant virus homologs of the eukaryotic CBX1/3/5 family (present in *Mimiviridae*, *Iridoviridae* and *Phycodnaviridae*); and (ii) homologs from various *Adintoviridae*, which are closely related to animal Chromo genes encoding *rve* integrase domains⁷⁶ (Fig. 5i). Finally, we also identify a handful of eukaryotic-like viral genes with deep-branching positions relative to core eukaryotic gene families, as seen in histones (Fig. 5h). This includes *Mimiviridae* homologs of the eukaryotic methyltransferases SMYD1-5 and DOT1 (Extended Data Fig. 7d,e), as well as SNF remodeller families with homologs in distinct giant virus clades (HLTF/TTF2

in Phycodnaviridae, Mimiviridae and Iridoviridae). These results indicate that cases of horizontal transfer from eukaryotes to viruses are common in different chromatin-related gene families, including histones. Therefore, it is likely that basally-branching giant virus histones were similarly acquired from a stem eukaryotic lineage and this would explain the observed histone tree topology with extant eukaryotic species. In any case, most of the eukaryotic chromatin machinery appears to have cellular roots.

Discussion

Our comparative proteogenomics study reconstructs in detail the origin and evolutionary diversification of eukaryotic chromatin components, from post-translational modifications to gene family domain architectures. We looked first at the pre-eukaryotic roots of chromatin. Multiple aspects of archaeal chromatin have been studied in recent years, including nucleosomal patterns³¹ and the structure of the archaeal nucleosome³⁰. A recent taxonomic survey of archaeal nucleoid-associated proteins revealed multiple independent diversifications of DNA-wrapping proteins and a strong association between high levels of chromatinization and growth temperature, overall suggesting a structural, non-regulatory role for archaeal chromatin⁷⁷. Our proteomics data support this notion by showing the scarcity of hPTMs in four species belonging to two different archaeal lineages (Euryarchaeota and Thaumarchaeota). An earlier proteomics study reported the complete absence of hPTMs in the euryarchaeote *Methanococcus jannaschii*³⁴. Here we do identify a few instances of modified lysine residues in Euryarchaeota, which is in line with the recently reported acetylations in *Thermococcus gammatolerans* histones⁷⁸. It remains to be seen if hPTMs are frequently present in Asgard and other unsampled archaeal lineages, where other eukaryotic-like features have been found^{57,79,80}. In fact, some of these Asgard, particularly Lokiarchaeota, encode for histones with long, K-rich N-terminal tails but that bear no similarity with eukaryotic histones and are, therefore, most probably the result of convergent evolution. Interestingly, Lokiarchaeota genomes also frequently encode histone modifiers such as SET methyltransferases and MOZ_SAS acetylases. However, overall our results suggest that extensive usage of hPTMs is an eukaryotic innovation (Fig. 6a). Similarly, while we find the majority of catalytic domains of hPTM writers, hPTM erasers and chromatin remodellers in Archaea and even Bacteria, these appear only scattered in a small fraction of the examined taxa. In contrast, hPTM reader domains and histone chaperones are eukaryotic innovations, further supporting the idea that the functional readout of hPTMs and the role for histone variants in defining chromatin states are both exclusive to eukaryotes (Fig. 6a).

The origin of eukaryotes represents a major evolutionary transition in the history of life⁸¹. Thanks to sequencing and comparative analysis of archaeal and eukaryotic genomes, we also have a detailed reconstruction of the massive innovation in gene repertoires that occurred at the origin of eukaryotes. This gene innovation in the Last Eukaryotic Common Ancestor (LECA) includes cytoskeletal proteins and associated motors like myosins^{82,83} and kinesins⁸⁴, vesicle trafficking apparatus⁸⁵, splicing machinery⁸⁶, ubiquitin signalling systems⁸⁷ and a large repertoire of sequence-specific transcription factors³⁷. Combining parsimony analysis and knowledge on gene function in extant lineages (mostly vertebrates, yeast and plants), our results allow us to reconstruct a complex LECA repertoire of hPTMs and associated writing, eraser and reader gene families (Fig. 6b,c). We infer 23 to 29

highly-conserved lysine acetylations in canonical histones (e.g., H3K9ac and H3K27ac) and a repertoire of 65 and 20 histone acetylase and deacetylase families, respectively. With the exception of H4K16ac⁶⁴, most histone acetylations are thought to exert a generic, perhaps additive, effect on the opening of chromatin²². As such, acetylation marks like H3K27ac have been found to be enriched in promoters of active genes in diverse eukaryotes⁴². In contrast, histone methylations often have very specific readouts and they can be linked both to active and repressive chromatin states. We infer between 13 and 25 conserved methylated lysine residues in LECA histones, including marks typically associated to active promoters (H3K4me1/me2/me3), gene bodies (H3K36me3, H3K79me1/2, H4K20me1), and repressive chromatin states (H3K9me2/me3, H3K27me3, H4K20me3)^{88,89}. Finally, we also infer the existence of five histone variants in the LECA (cenH3, H3.3, H2A.Z, macroH2A and H2A.X), as well 33 chromatin remodellers (e.g., EP400/SWR1 and INO80, involved in loading and removal of H2A.Z, respectively) and 25 histone chaperones (e.g., ASF1A/B and NPM1/2/3). This indicates that, in addition to an extensive repertoire of hPTMs, the regulation of nucleosomal histone composition was also an important feature in the LECA.

Chromatin evolution after the origin of eukaryotes is characterized by an expansion of lineage-specific histone variants harboring unique hPTMs and a net expansion in the number of reader gene families, as opposed to the relatively static catalytic gene families (writers, erasers and remodellers). This is particularly relevant as it suggests extensive remodelling of chromatin networks during eukaryote evolution, that is, changes in the coupling of particular hPTMs to specific functional chromatin states. An example of such changing state-definitions comes from looking at the hPTMs associated to TEs in different organisms: H3K9me3+H4K20me3 in animals, H3K27me3 in some plants⁹⁰, H3K79me2+H4K20me3 in the brown multicellular algae *Ectocarpus siliculosus*⁴³, and H3K9me3+H3K27me3 in the ciliate *Paramecium tetraurelia*⁹¹. In the context of the histone code hypothesis^{3,20,92–94}, our findings indicate that, while there is an ancient core of conserved hPTMs across eukaryotes, evidence for a universal code/functional-readout is limited, with perhaps the exception of the highly conserved configuration of ancient hPTMs around active promoters across many eukaryotes⁴². Another interesting observation related to the evolution of chromatin networks is the capture of chromatin reader domains by TEs. We find evidence of this phenomenon in a number of species with a scattered phylogenetic distribution, suggesting that it is a recurrent process and that it often leads to the successful propagation of the TE in the host genome. We hypothesize that this process facilitates the targeting of TEs to specific chromatin states, as it has been described in the case of MBD DNA methylation readers captured by TEs^{95,96}.

In the future, a broader phylogenetic understanding of the genome-wide distribution of hPTMs, as well as the direct interrogation of hPTM binders in different species^{97–99}, will be crucial to further clarify questions such as the ancestral role of specific hPTM and the co-option of ancient hPTMs into novel functions.

Methods

Eukaryotic cell culture and tissue sources

Capsaspora owczarzaki strain ATCC30864 filopodial cells were grown axenically in 5 ml flasks with ATCC medium 1034 (modified PYNFH medium) in an incubator at 23°C (Sebé-Pedrés et al., 2013a).

Corallochytrium limacisporum strain India was axenically grown in Difco Marine Broth medium at 23°C, *Creolimax fragrantissima* strain CH2 was axenically grown in Difco Marine Broth medium at 12°C, *Spizellomyces punctatus* strain DAOM BR117 was axenically grown in (0,5% yeast extract, 3% glycerol, 1g/L K₂HPO₄, 0,5% EtOH) medium at 17°C, *Thecamonas trahens* strain ATCC50062 was grown in ATCC medium: 1525 Seawater 802 medium, *Chlamydomonas reinhardtii* strain CC-503 cw92 mt+ was axenically grown in Gibco TAP medium at 29°C, *Guillardia theta* strain CCMP2712 was axenically grown in L1+500uM NH₄Cl medium at 18°C, *Emiliania huxleyi* strain CCMP1516 was grown in L1-Si medium at 18°C, *Thalassiosira pseudonana* strain CCMP1335 was axenically grown in L1 medium at 18°C, *Bigelowiella natans* strain CCMP2755 was axenically grown in L1-Si medium at 23°C, *Naegleria gruberi* strain ATCC30224 was axenically grown in ATCC medium 1034 (modified PYNFH medium) at 29°C, *Gefionella okellyi* strain 249 was grown in 15% Water Complete Cereal Grass Media (WC-CGM3) at 18°C and *Fabomonas tropica* strain NYK3C was grown in L1 + YT medium at 18°C. All cells were grown in 250 ml culture flasks.

In addition, we used frozen tissues/cells from the following species: *Homo sapiens* (ES cells, courtesy of Cecilia Ballaré, CRG), *Physcomitrella patens* (strain Gransden 2004, vegetative stage, courtesy of Josep Casacuberta, CRAG-CSIC), *Sycon ciliatum* (adult sponges sampled from Bergen, Norway, courtesy of Maja Adamska, ANU) and *Phytophthora infestans* (strain T30-4, courtesy of Harold J.G.Meijer, Wageningen University).

Archaeal cell culture

Cultures of *Methanobrevibacter cuticularis* DSM 11139, *Methanospirillum stamsii* DSM 26304 and *Methanosarcina spelaei* DSM 26047 were purchased from the Deutsche Stammsammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Braunschweig, Germany. Cultures were grown in closed batch in 50mL of defined media in 120mL serum bottles (La-Pha-Pack, Langerwehe, Germany). Growth was monitored as OD (600 nm; Analytik Jena, Specord 200 plus). *Methanobrevibacter-cuticularis* was grown in modified *Methanobrevibacter cuticularis* medium DSMZ 734a (DSMZ 2014) omitting bovine rumen fluid, yeast extract and Na-resazurin at 1.5 bar overpressure H₂CO₂ (20 vol.-% CO₂ in H₂) at 37°C. As soon as a change in OD was observed, a constant agitation at 90rpm was applied. *Methanospirillum stamsii* was grown in modified *Methanobacterium* medium DSMZ 119 (DSMZ 2017) omitting sludge fluid, yeast extract and Na-resazurin at 1 bar overpressure H₂CO₂ (20 vol.-% CO₂ in H₂) at 29°C, under constant agitation at 90rpm. *Methanosarcina spelaei* was grown in modified *Methanosarcina barkeri* medium DSMZ 120a (DSMZ 2014) omitting yeast extract and Na-resazurin at 1.5 bar overpressure H₂CO₂ (20 vol.-% CO₂ in H₂) at 33°C, under constant agitation at 90rpm. All gases were obtained

from Air Liquide GmbH, Schwechat, Austria. *Nitrososphaera viennensis* EN76 was grown in continuous culture in a bioreactor as previously described¹⁰⁰.

Cells were harvested via centrifugation at 21,000xg 4°C 1h (Thermo scientific, Sorvall Lynx 4000 centrifuge), the supernatant discarded and the resulting pellet resuspended in 1ml of spent medium, followed by another round of centrifugation at 21,000xg 4°C for 1h (Eppendorf, Centrifuge 5424R). Pellets were stored at -70°C. All archaeal histones were extracted as described below.

Histone acid extraction

Starting material was a pellet of 50-100M cells (washed once with cold PBS) or a flash-frozen tissue homogenate in liquid nitrogen using a ceramic mortar grinder. Cells were washed first in 10ml of buffer I (10 mM TrisHCl pH 8, 10 mM MgCl₂, 0.4M Sucrose). After 5min incubation, samples were centrifuged at 8.000g for 20min at 4°C and supernatant was removed. The resulting pellet was resuspended in 1.5ml of Buffer II (10 mM TrisHCl pH 8, 10 mM MgCl₂, 0.25M Sucrose, 1% Triton X-100, 1% Igepal Ca-630) and incubated 15min on ice. In specific cases, cells at this stage were broken using a 2ml Dounce homogenizer (with Pestle B) or with a 20G syringe. Then samples were centrifuged at 15.000g for 10min at 4°C and supernatant was removed. The resulting pellet was then slowly resuspended in 300µL of Buffer III (10 mM TrisHCl pH 8, 2 mM MgCl₂, 1.7M Sucrose, 1% Triton X-100) and then resulting resuspended nuclei were layered on top of another 300µL of Buffer III. Sample was centrifuged at 20.000g for 1h at 4°C and supernatant was removed, resulting in a nuclear pellet ready for acid histone extraction. All buffers were supplemented with spermidine (1:1000), beta-mercaptoethanol (1:1000), protease inhibitors (1x cComplete cocktail Roche #11697498001, 1mM PMSF, 1:2000 Pepstatin), phosphatase inhibitors (1x phoSTOP cocktail Roche #4906845001) and deacetylase inhibitors (10mM Sodium butyrate).

For samples processed using a high-salt + HCl extraction protocol^{101,102}, the pellet was resuspended in 500µL of High Salt Extraction Buffer (20 mM TrisHCl pH 7.4, CaCl₂ 1M and protease, phosphatase and deacetylase inhibitors, same as above). Sample was incubated on ice for 30min and then pure HCl has added to a final 0.3N concentration (12.82µL to the initial 500µL). Samples were incubated for at least 2h on a rotor at 4°C and then centrifuged at 16.000g for 10min at 4°C to remove cellular/nuclear debris. The resulting supernatant containing solubilized histones was transferred to a clean 1.5ml tube and Trichloroacetic Acid (TCA) was added drop-wise to 25% final concentration (171µL TCA to an approximate initial 513µL sample) and left overnight at 4°C to precipitate histones. Samples were then centrifuged at 20.000g for 30min at 4°C and the supernatant removed. The pellet was then washed twice with 500µL of cold acetone and then dried for 20min at room temperature. Finally, clean histone pellets were resuspended in 30-50µL of ultrapure water. Protein concentration in the sample was measured using BCA and extraction was examined using an SDS-PAGE protein gel with Coomassie staining.

For samples processed using H₂SO₄¹⁰², the protocol was exactly the same except that 400µL 0.4N H₂SO₄ (freshly diluted) was used instead, with a similar incubation time of at least 2h at 4°C.

Histone chemical derivatization

Histones samples were quantified by the BCA method and 10 µg of each sample were derivatized with propionic anhydride, digested with trypsin and derivatized again with phenylisocyanate as previously described⁴⁹. Briefly, samples were dissolved in 9 µL of H₂O and 1 µL of triethyl ammonium bicarbonate was added to bring the pH to 8.5. The propionic anhydride was prepared by adding 1 µL of propionic anhydride to 99 µL of H₂O and 1 µL of propionic anhydride solution was added immediately to the samples with vortexing and incubation for 2 minutes. The reaction was quenched with 1 µL of 80mM hydroxylamine and samples were incubated at room temperature for 20 minutes. Tryptic digestion was performed for 3 h with 0.1 µg trypsin (Promega Sequencing Grade; Madison, WI) per sample. A 1% v/v solution of phenyl isocyanate (PIC) in acetonitrile was freshly prepared and 3 µl added to each sample (17 mM final concentration) and incubated for 60 min at 37 °C. Samples were acidified by adding 50 µL of 5% formic acid, vacuum dried and desalted with C18 ultramicrospin columns (The Nest Group, Inc, Southborough, MA).

Liquid Chromatography-Tandem Mass Spectrometry Sample Acquisition

A 2-µg aliquot of the peptide mixture was analyzed using a LTQ-Orbitrap Fusion Lumos mass spec-trometer (Thermo Fisher Scientific, San Jose, CA) coupled to an EASY-nLC 1000 (Thermo Fisher Scientific, San Jose, CA) with both collision induced dissociation (CID) and high energy collision dissociation (HCD) fragmentation.

Peptides were loaded directly onto the analytical column and were separated by reversed-phase chromatography using a 50-cm column with an inner diameter of 75 µm, packed with 2 µm C18 particles spectrometer (Thermo Scientific, San Jose, CA, USA) with a 90 min chromatographic gradient. The mass spectrometer was operated in positive ionization mode using a data dependent acquisition method. The “Top Speed” acquisition algorithm determined the number of selected precursor ions for fragmentation.

Mass-spectrometry Data Analysis

Acquired data were analyzed using the Proteome Discoverer software suite (v2.0, Thermo Fisher Scientific), and the Mascot search engine (v2.6, Matrix Science¹⁰³) was used for peptide identification using a double-search strategy. First, data were searched against each organism protein database plus the most common contaminants considering Propionylation on *N*-terminal, Propionylation on Lysines and Phenylisocyanate on *N*-terminal as variable modifications. Then a new database was generated with the proteins identified in the first search,, and a second search was done considering Propionylation on *N*-terminal, Propionylation on Lysines, Phenylisocyanate on *N*-terminal, Dimethyl lysine, trimethyl lysine, propionyl + methyl lysine, acetyl lysine, crotonyl lysine as variable modifications. Precursor ion mass tolerance of 7 ppm at the MS1 level was used, and up to 5 missed cleavages for trypsin were allowed. False discovery rate (FDR) in peptide identification was set to a maximum of 5%. The identified peptides were filtered by mascot ion score higher than 20 and only PTMs with a localization score ptmRS¹⁰⁴ higher than 45 were considered. The raw proteomics data have been deposited to the PRIDE¹⁰⁵ repository with the dataset identifier PXD031991.

Analysis of hPTM conservation

Identification of canonical and variant histones—We classified histone protein domains from a database of eukaryotic, prokaryotic and viral sequences (see details below) according to their similarity to known canonical (H2A, H2B, H3, H4) and variant histones (e.g., H2A.Z, macroH2A, cenH3 or H3.3), as well as other gene families with histone-like protein folds (e.g., the transcription factors DR1, DRAP1, NFYB/C, POLE3/4, SOS, TAF, or CHRAC). To that end, we used *diamond* to perform local alignments of each histone domain against (i) a set of curated histone variants obtained from HistoneDB 2.0⁴⁸, and (ii) annotated each domain according to the best hit in the reference database, which allowed us to classify histone fold-containing proteins as canonical histones (H2A, H2B, H3, H4) or their main variants (H2A.Z, macroH2A and cenH3). This best-hit strategy performs well in distinguishing canonical histones from each other, as well as each canonical histone from its main variants (H3 from cenH3, and H2A from H2A.Z and macroH2A; Extended Data Fig. 1a).

Then, we built a graph of pairwise similarity between histones, with edges weighted by the alignment bitscore (discarding edges with bitscore < 20). We created visualisations of each connected component in this graph using the spring layout algorithm implemented in the *networkx* 2.4 Python library (100 iterations, weighted by alignment bitscore)¹⁰⁶. We selected the four connected components in the graph that matched the four canonical eukaryotic histones (H2A, H2B, H3, H4; discarding edges with bitscore < 20), retrieved the protein sequences for each of them, aligned them using *mafft* (E-INS-i mode, 1,000 iterations)¹⁰⁷, and built phylogenetic trees with *IQ-TREE* 2.1.0 (-fast mode)¹⁰⁸.

Identification of hPTM homology—We retrieved the protein sequences of the canonical histones identified in each of the 26 species and we used them for the proteomic analysis of hPTMs, and aligned them using *mafft* (*G-INS-i* mode, up to 10,000 refinement iterations). For this subset of species, histone class identity was cross-referenced with the HistoneDB search tool. Then, we manually aligned the peptides mapping onto these proteins to identify the position of each hPTM along a consensus alignment. In the case of H3, H4, and macroH2A, the majority of alignment positions were conserved across most eukaryotes in our dataset, and we used a consensus numbering scheme. In the case of H2A, H2A.Z, and H2B, non-conserved insertions and deletions at the N-terminal tail precluded the use of a paneukaryotic numbering scheme. Instead, we reported hPTM positions based on the human homolog (if possible), or relative to taxonomically restricted conserved positions. In cases where position-wise homology could not be established, we grouped multiple amino-acids into stretches of unclear homology, which we report separately from conserved positions (question mark symbols in Fig. 1). The complete list of hPTMs and their position-wise coordinates relative to the consensus alignment is available in Supplementary Data 3.

Furthermore, we also reported the presence (in any position) of modifications in less-conserved histone variants, as well as the linker histone H1.

In addition to the 19 used in our proteomics survey, we also included previously published hPTM data from the following species (Supplementary Data 1c): the brown alga *Ectocarpus siliculosus*⁴³, the diatom *Phaeodactylum tricornutum*¹⁰⁹, the ciliate *Tetrahymena*

thermophila^{46,110–112}, the ascomycete *Neurospora crassa*¹¹³, *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*⁴⁶, and the plant *Arabidopsis thaliana*^{114–116}. When available in public repositories, we re-analysed these datasets using the strategy described above. Finally, we also complemented our own proteomics data using previously published hPTM data from *Homo sapiens*^{46,117–120} and *Capsaspora owczarzakii*⁴².

Comparative genomics analysis of chromatin-associated proteins

Data retrieval—We identified homologs of gene families associated with eukaryotic chromatin, using a database of predicted proteomes from a selection of eukaryotic species from all major supergroups ($n = 172$ species; see Supplementary Data 1 for their taxonomic classification and data sources), as well as archaeal and viral peptides available in the NCBI *nr* peptide collection (as of 25th of April, 2020) and bacterial peptides available in RefSeq (release 99, 11th May, 2020). The database of viral sequences was complemented with peptides from 501 genomes of nucleocytoplasmic large DNA viruses¹²¹.

Gene family searches—We defined 61 gene classes associated with eukaryotic chromatin, based on HMM models obtained from the Pfam database (release 33.0)¹²². This list included canonical and linker histones ($n = 2$ families), chromatin-specific lysine acetylases ($n = 5$), deacetylases ($n = 2$), methyltransferases ($n = 2$), demethylases ($n = 2$), chromatin readers ($n = 16$), remodellers ($n = 1$) and chaperones ($n = 13$), as well as multiple families associated with the Polycomb complexes ($n = 18$). The complete list of gene families, including the associated HMM models, is available in Supplementary Data 4.

For each gene family, we retrieved all homologs from the eukaryotic, archaeal, bacterial and viral databases using the *hmmsearch* tool from the *HMMER* 3.3 toolkit¹²³ and the gathering threshold defined in each Pfam HMM model. We recorded the taxonomic profile of each homolog.

Orthology identification—We aimed to identify groups of orthologs within each of the 61 chromatin-associated gene families using targeted phylogenetic analyses. We followed the following strategy for each of the 59 sets of eukaryotic genes. First, we partitioned each set into one or more homology groups based on pairwise local sequence alignments using *diamond* 0.9.36.137 (high sensitivity all-to-all search)¹²⁴, followed by clustering of the resulting pairwise alignments graph with *MCL* 14.137 (*--abc* mode)¹²⁵, using low inflation values (see Supplementary Data 4) to favour inclusive groupings. Second, we performed multiple sequence alignments of each homology group with *mafft* 7.471¹⁰⁷ under the E-INS-i mode (optimised for multiple conserved regions), running up to 10,000 refinement iterations. Third, we trimmed the resulting multiple sequence alignments using *clip-kit* 0.1 (*kpic-gappy* mode)¹²⁶. Fourth, we built phylogenetic trees for each trimmed alignment using *IQ-TREE* 2.1.0¹⁰⁸, selecting the best-fitting evolutionary model using its *ModelTest* module (according to the Bayesian Information Criterion) and using 1,000 UFBS bootstrap supports¹²⁷. Each tree was run for up to 10,000 iterations until convergence was attained (at the 0.999 correlation coefficient threshold, and for at least 200 iterations).

Then, we parsed the species composition of each gene tree in order to identify groups of orthologous proteins using the *POSSVM* pipeline¹²⁸. Specifically, we used the species

overlap algorithm¹²⁹ implemented in the *ETE* toolkit 3.1.1¹³⁰, which identifies pairs of orthologous genes in a phylogenetic tree by examining the species composition of each subtree, and classifying internal nodes as paralogy nodes (if there is overlap in the species composition between each of its two descendant subtrees) or orthology nodes (if there is no overlap). Pairs of genes linked by an orthology node are then recorded as orthology pairs. In our analysis, we used an overlap threshold=0 (i.e. any species composition overlap between the two descendant subtrees is classified as a paralogy event). The resulting list of pairwise orthology relationships between genes was clustered into groups of orthologs (orthogroups) using *MCL*. We further annotated each orthogroup with a string denoting the gene symbols of the human proteins therein (if any).

Overall, we classified 51,426 proteins from 61 gene classes (defined by protein structural domains), divided into 242 gene trees and 1,713 gene families (orthogroups). The source peptide sequences and gene trees used for these analyses are available in Supplementary Data 7 and 8.

Ancestral reconstruction of gene content—We inferred the presence, gain and loss of each orthogroup along the eukaryotic tree of life, using a phylogenetic birth-and-death model¹³¹ implemented in *Count*¹³². This tool takes a numeric profile of gene family presence/absence in extant species (172 in our dataset) and a phylogenetic tree defining their evolutionary relationships, and infers the probabilities of gain and loss of each family at each ancestral node along the tree.

First we trained the probabilistic model in *Count*. As a training set, we used a random sample of 1,000 PFAM domains annotated in the 172 species of interest (restricting the sampling to domains present in at least 5% of species). The final model consists of gain, loss and transfer rates with two Γ categories each, and a constant duplication rate (given that we only recorded gene presence/absence, duplication events are not included in our downstream analyses). This model was obtained in three sequential rounds of training, so as to sequentially add zero, one and two Γ categories to each evolutionary rate. Each round consisted of up to 100 iterations, and stopped when the relative change in the model log-likelihood fell by 0.1% in two consecutive rounds. The final evolutionary rates and the Newick-formatted species tree used in this step are available in the Supplementary Data 1 and Extended Data Fig. 3a.

Second, we calculated the posterior probability of gain, loss and presence of each orthogroup in our dataset with *Count*. The aggregated counts of gains and losses of the various classes of chromatin-associated proteins (acetylases, deacetylases, methyltransferases, demethylases, readers and remodellers) along the eukaryotic tree were obtained by summing the probabilities of gain, presence or loss of all orthogroups of a given class at each ancestral node. To investigate the evolutionary histories of specific orthogroups at a given node in the tree, we applied a probability threshold of 0.9 (for presence) or 0.5 (to identify the most probable gain and loss node). The *Count* model was not able to calculate ancestral probabilities for a few orthogroups with widespread phylogenetic distributions, due to violations of the birth-and-death model (25 out of 1,713 families). In order to be able to report presence probabilities in the LECA for these orthogroups, we inferred their

presence in this ancestor using the Wagner parsimony procedure implemented in *Count* with a gain-to-loss penalty $g = 5$, and recorded their presence as binary values (0/1) accordingly.

Protein domain architecture analyses—We annotated the Pfam domains present in each protein from the gene classes listed in Supplementary Data 4, using *Pfamscan* 1.6-3 and the Pfam 33.0 database¹²². We visualized the networks of protein domain co-occurrence from the point of view of the core domain(s) that define each gene class, using the *networkx* Python library (version 2.4)¹⁰⁶. Specifically, we built a graph where each node represented ‘accessory’ domains (i.e. domains that co-occur with the ‘core’ domain that defines given gene class), node size reflected number of co-occurrences with the ‘core’ domain, and edges reflected co-occurrences between accessory domains. We identified communities of frequently co-occurring accessory domains using the label propagation algorithm implemented in *networkx* (*communities* submodule), which we used as a basis to manually annotate groups of co-occurring domains of interest (Fig. 5c). Network visualizations were created using the *NEATO* spring layout algorithm from the *Graphviz* 2.40.1 Python library¹³³.

In parallel, we also recorded the presence of Pfam domains within individual orthogroups, and their taxonomic distribution.

Prokaryotic roots of the eukaryotic chromatin machinery—We retrieved all eukaryotic domains from gene class shared with prokaryotes (Histones, Acetyltransf_1, GNAT_acetyltr_2, MOZ_SAS, Hist_deacetyl, SIR2, DOT1, SET, CupinJmjC, ING, MBT, PWWP and SNF2_N), collapsing identical sequences at 100% similarity with *CD-HIT* 4.8.1¹³⁴, and identified their closest homologs amongst the corresponding archaea and bacteria protein domain sets, using *diamond* local alignments (high sensitivity search). The archaeal and bacterial protein sets were also reduced with *CD-HIT* (at 95% and 90% sequence similarity, respectively). Each set of sequences was then partitioned into low-granularity homology clusters using the *MCL*-based strategy described above (inflation $I = 1.2$), and a phylogenetic tree was then constructed from each homology cluster with *IQ-TREE* (as described above).

Then, we mapped each eukaryotic gene to its orthogroup (obtained from eukaryotic-only analyses, see above) and used the distribution of phylogenetic distances from the prokaryotic+eukaryotic gene trees to classify them according to their similarity to (*i*) eukaryotic genes in other orthogroups, (*ii*) archaeal homologs, or (*iii*) bacterial homologs. Specifically, we used a majority-voting procedure in which we recorded the number of sequences of eukaryotic, archaeal or bacterial origin amongst the ten nearest neighbors of each gene (measuring intergenic distances as substitutions per site), and assigned the most common taxonomic group as the ‘closest’ homolog of that gene (minimum 50% agreement). This fraction is termed ‘Phylogenetic affinity score’ and reported in Supplementary Data 5. The pairwise distances were obtained from each gene tree using the cophenetic distance method in the *cophenetic.phylo* utility of the *ape* 5.4 R library¹³⁵.

Characterisation of fusions with transposon-associated domains—We retrieved all classified genes from our eukaryotic dataset that contained transposon-associated Pfam

domains (version 33.0), using a list compiled from^{68,136} (complete list in Supplementary Data 4), totaling 823 candidate fusions from 91 species (listed in Supplementary Data 6). We annotated these genes to their most similar known TE element by aligning them against the Dfam 3.3 database¹³⁷ using the *tblastn* program in *BLAST* 2.2.31¹³⁸.

We validated each candidate fusion using the following criteria: (i) contiguity of the gene model on the genome assembly, i.e., recording which genes were interrupted by poly-*N* stretches (which might indicate an incorrect gene model); (ii) evidence of expression in at least one sample from a range of publicly available transcriptomic experiments (from the NCBI SRA repository); (iii) evidence of contiguous expression, i.e., whether an expressed transcript had mapped reads along the entire region located between the ‘core’ and ‘TE-associated’ domains; (iv) we also recorded the number of exons per gene; and (v) located near any other candidate fusion gene in the genome.

The list of SRA experiments used for these validation steps is available in Supplementary Data 1. This list includes 64 out of 91 species for which transcriptomics datasets are publicly available, and covers 768 out of the 822 TE fusion candidates (93%). RNA-seq read mapping was performed with *bwa mem* 0.7.17-r1188¹³⁹ using the complete set of spliced transcripts of each species as the reference database. We used *bedtools* 2.29.2¹⁴⁰ to identify poly-*N* stretches in the genome assembly (assembly contiguity criterion). We identified regions of low coverage along the transcript sequence (expression contiguity criterion) using the *bedtools genomecov* utility, requiring that the coverage along both domains involved in each fusion and their intermediate regions be higher or equal to two reads.

Analysis of viral homologs—We investigated the homology of the viral chromatin-associated genes (which included 19 out of 61 families present in our survey) using joint phylogenetic analyses of protein domains from virus, prokaryotic and eukaryotic genes. We used the same method described above to investigate the prokaryotic roots of eukaryotic gene classes: we aligned viral domains against a database of cellular homologs (high sensitivity *diamond* search), followed by low-granularity *MCL* clustering (inflation $I=1.2$) and phylogenetic tree building (*IQ-TREE*). Then, we used the same majority-voting procedure described above to classify viral homologs according to their similarity to eukaryotic, archaeal or bacterial gene families based on their distribution of phylogenetic distances. For viral genes that were most similar to eukaryotic genes, we used the same procedure to map them to their closest eukaryotic orthogroup.


The complete list of viral genes and their phylogenetic annotation is available in Supplementary Data 6. Out of 2,163 viral genes in our dataset, 2,144 could be annotated as similar to a particular cellular group using this procedure (99.1%), and the majority of these genes had a high agreement in the annotations of their nearest neighbors (2,096 with 50% agreement; 1,449 with 90% agreement).

In the case of viral histones, we built a separate phylogeny with a few modifications in our protocol: (i) we used additional viral genes obtained from⁷¹ as a reference; (ii) we omitted the *CD-HIT* reduction and *MCL* partitioning steps, and jointly analyzed the entire set of

homologs instead; and (iii) in the phylogenetic reconstruction step, we used the approximate Bayes posterior probabilities¹⁴¹ implemented in *IQ-TREE*.

Identification of archaeal N-terminal histone tails—We retrieved all archaeal histone domains classified belonging to the HMfB-like connected component in Fig. 1b, and retained those that fulfilled the following criteria: (i) contained a complete Cbfd_NFYB_HMF domain according to the *hmmscan* search (defined as an alignment starting at least at the 10th position of the HMM model, and up to the 55th position; the HMM model contains 65 positions); and (ii) the predicted tail (*N*-terminal to the core domain boundaries defined by *hmmscan*) was at least 10 residues long. 84 genes passed these filters, including three *N*-terminal containing histones previously identified by Henneman *et al.*⁵⁵. A complete list is available in Supplementary Data 2. We manually examined the sequences of archaeal tails and aligned four sets of similar histones with *mafft G-INS-i* (Extended Data Fig. 1d). Alignments were plotted using the *msa* 1.24.0 library in *R*¹⁴².


Extended Data



Extended Data Fig. 1. Histone classification and evolution.

a, Primary and secondary alignments of histone-fold containing proteins classified as canonical H2A, H2B, H3 and H4, based on identity to reference sequences in HistoneDB⁴⁸. Pie plots represent the number of alignments to HistoneDB-annotated sequences, for the entire dataset (prokaryotic, eukaryotic and viral sequences, large pie plots in the inset) and the eukaryotic subset (smaller plots in the inset). For those proteins that align to more


than one canonical histone or major variant (macroH2A, H2A.Z or cenH3), the scatter plots represent the relative identity between the primary (horizontal axis) and secondary alignment(s) (vertical axis). **b**, Aggregated counts of histone gene pairs, classified according to histone type and orientation. **c**, Presence of histone variants (left) and number of collinear pairs of histone-encoding genes (right) per species, classified according to their histone types and relative orientation (head-to-head, hh; head-to-tail, ht; and tail-to-tail, tt). Source data available in Supplementary Data 2. Histone variant classification is based on the highest-scoring HMM profile from HistoneDB. Asterisks colors in the macroH2A column indicate species where histone-less Macro domains orthologous to the macroH2A genes are found (see panel d). Lighter colors in the variant classification indicate ambiguously classified histones (i.e. cases in which the highest-scoring HMM profile exhibited a low bitscore, defined as a probability below 0.05 in the profile-wise distribution function of scaled bitscores; or cases in which the first-to-second ratio between high scoring profiles was below 1.01). **d**, Alignments of putatively conserved histone *N*-tails in archaea. Conserved amino-acids are color-coded according to chemical properties. Dots next to species names are color-coded according to taxonomy (same as Fig. 2c). **e**, Phylogenetic analysis of the Macro motif of macroH2A histones across eukaryotes, highlighting the macroH2A ortholog group (green), and, within this group, Macro-containing genes lacking histone domains (orange), and their protein domain architectures.



Extended Data Fig. 2. Histone post-translational modifications.

a, Proteomics detection coverage (% of amino acids), number of hPTMs and number of hPTMs per covered position, for the best-covered histone in each species in our proteomics survey. b, Number of samples in which each histone-matching peptide with post-translational modifications (peptide spectral matches defined by Proteome Discoverer) has been identified, per species. For each species, we report the percentage of modified peptides found in more than one replicate. c, Number of samples in which histone-matching modified peptide has been identified, across all the samples from this study. The tree pie charts


represent these distributions for all hPTMs, acetylations, and methylations. **d**, Evidence of hPTM conservation in the major histone variants H2A.Z and macroH2A (conserved positions only), as well as any position in the linker histones H1.



Extended Data Fig. 3. Gene family counts.

a-c, Number of taxa within each lineage that contain chromatin-associated genes, for archaeal, bacterial (per phyla) or viral (per family) genomes. Numbers indicate the exact number of taxa. **d**, Number of genes encoding core domains that define chromatin-


associated gene families per eukaryotic genome/transcriptome. Numbers indicate exact number of proteins.



Extended Data Fig. 4. Evolutionary reconstruction and domain architecture conservation.

a, Species tree of eukaryotes used in the ancestral reconstruction analysis, with branch lengths calibrated to the gain/loss rates of Pfam domains (see Methods). Available in Supplementary Data 1. **b**, Conservation of archetypical protein domain architectures across orthogroups, in acetylases, deacetylases, methyltransferases, demethylases, remodellers and


chaperones. In each heatmap, we indicate the fraction of genes within an orthogroup (rows) that contain a specific protein domain (columns). Domains in bold are catalytic (black) or reader (purple) functions. At the right of each heat-map, we summarize the presence/absence profile of each orthogroup across eukaryotic lineages (as listed in Fig. 1a).



Extended Data Fig. 5. Evolution of the hPTM reader toolkit.


a, Pie chart representing the number of genes classified as part of the catalytic (acetylases, deacetylases, methyltransferases, demethylases, remodellers or chaperones) or reader

families, or as both. The barplot at the right shows the most common reader domains in genes classified with both reader and catalytic functions. **b**, Pie plot representing the number of reader domain-encoding genes classified according to whether they contain one type of reader domain (e.g., PHD) or more than one (e.g., PHD + PWWP). The barplot at the right shows the most common combinations of reader domains among genes with multiple reader domains. **c**, Summary of gene family gains per reader family, with example cases highlighted in selected nodes. Node size is proportional to number of gains at 90% probability.



Extended Data Fig. 6. Transposon-chromatin gene fusions.
a, Number of candidate fusion genes classified by the level of gene model validation evidence, based on contiguity of the gene model over the genome assembly (i.e. lack of poly-N stretches in the genomic region between the TE- and chromatin-associated domains), evidence of expression, and evidence of contiguous expression (see inset at the right). **b**, Summary of candidate gene fusions within each chromatin-associated gene family, divided by gene family. For each gene, we indicate their similarity to known TE families, presence of TE-associated domains, the evidence of gene model validity, and information on their

gene structure (whether they are monoexonic or are located in clusters with other fusion genes). Source data available in Supplementary Data 6. **c**, Number of species with at least one valid fusion, divided by gene family. **d**, Mapping positions of RNA-seq reads supporting candidate gene-transposon fusions (selected examples from Fig. 5e). For each fusion, we show reads spanning the region along the spliced transcript that fully covers the transposon-associated domains (highlighted in green), the chromatin-associated domains, and the inter-domain region. Uninterrupted stretches of mapped positions between domains indicate the validity of a domain co-occurrence. For clarity purposes, reads mapping entirely within a single domain have been excluded from this visualization.



Extended Data Fig. 7. Chromatin proteins in viruses.

a-c, Selected gene trees highlighting examples of eukaryotic- and prokaryotic-like viral homologs. **d**, Number of viral genes of each chromatin-associated gene family, classified according to their closest neighbours from cellular clades in gene tree analyses based on phylogenetic affinity scores (see Methods). Within each gene family, viral sequences are classified according to their PFAM domain architecture – the most common architecture being single-domain in most gene families except for remodellers and BIR readers. **e**, *Id.*,

but classifying viral genes according to their phylogenetic affinity to eukaryotic orthology groups. Source data available in Supplementary Data 6.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We want to thank Alex de Mendoza for critical input on the analysis of transposable element fusions. We also want to thank Josep Casacuberta for *Physcomitrella patens* samples, Harold J. G. Meijer for *Phytophthora infestans* samples, Maja Adamska for *Sycon ciliatum* samples, and Alistar Simpson for access to the *Geffionella okellyi* culture (made possible by his funding from NSERC, Canada). Research in A.S-P. group was supported by the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Programme Grant Agreement (851647) and the Spanish Ministry of Science and Innovation (PGC2018-098210-A-I00). We also acknowledge support of the Spanish Ministry of Science and Innovation to the EMBL partnership, the Centro de Excelencia Severo Ochoa and the CERCA Programme (Generalitat de Catalunya). C.N. is supported by an FPI PhD fellowship from the Spanish Ministry of Economy, Industry and Competitiveness (MEIC). X.G-B. is supported by a Juan de la Cierva fellowship (FJC2018-036282-I) from the Spanish Ministry of Economy, Industry and Competitiveness (MEIC). I.R-T. was supported by a European Research Council Grant (616960). B.F.L. was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC; RGPIN-2017-05411) and by the 'Fonds de Recherche Nature et Technologie', Quebec. P.L-G. and D.M. were supported by a Moore and Simons foundations grant (GBMF9739) and by European Research Council Advanced Grants (322669, 787904). Research in C.S. group was supported by the European Research Council (ERC) through project TACKLE (AdvGrant No. 695192).

Data Availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD031991.

Code Availability

Code for reproducing the analysis is available in our lab Github repository (<https://github.com/sebepedroslab/chromatin-evolution-analysis>).

References

1. Struhl K. Fundamentally different logic of gene regulation in eukaryotes and prokaryotes. *Cell*. 1999; 98: 1–4. [PubMed: 10412974]
2. Kornberg RD, Lorch Y. Primary Role of the Nucleosome. *Mol Cell*. 2020; 79: 371–375. [PubMed: 32763226]
3. Jenuwein T, Allis CD. Translating the Histone Code. *Science*. 2001; 293: 1074–1080. [PubMed: 11498575]
4. Berger SL. The complex language of chromatin regulation during transcription. *Nature*. 2007; 447: 407–12. [PubMed: 17522673]
5. Banaszynski, La; Allis, CD; Lewis, PW. Histone variants in metazoan development. *Dev Cell*. 2010; 19: 662–74. [PubMed: 21074717]
6. Allis CD, Jenuwein T. The molecular hallmarks of epigenetic control. *Nat Rev Genet*. 2016; 1
7. Sultana T, et al. The Landscape of L1 Retrotransposons in the Human Genome Is Shaped by Pre-insertion Sequence Biases and Post-insertion Selection. *Mol Cell*. 2019; 74: 555–570. e7 [PubMed: 30956044]
8. Gangadharan S, Mularoni L, Fain-Thornton J, Wheelan SJ, Craig NL. DNA transposon Hermes inserts into DNA in nucleosome-free regions in vivo. *Proc Natl Acad Sci*. 2010; 107: 21966–21972. [PubMed: 21131571]

9. Shinn P, et al. HIV-1 Integration in the Human Genome Favors Active Genes and Local Hotspots. *Cell*. 2002; 110: 521–529. [PubMed: 12202041]
10. Goodier JL. Restricting retrotransposons: a review. *Mob DNA*. 2016; 7: 16. [PubMed: 27525044]
11. Molaro A, Malik HS. Hide and seek: how chromatin-based pathways silence retroelements in the mammalian germline. *Curr Opin Genet Dev*. 2016; 37: 51–58. [PubMed: 26821364]
12. Malik HS, Henikoff S. Phylogenomics of the nucleosome. *Nat Struct Biol*. 2003; 10: 882–91. [PubMed: 14583738]
13. Talbert PB, Henikoff S. Histone variants--ancient wrap artists of the epigenome. *Nat Rev Mol Cell Biol*. 2010; 11: 264–75. [PubMed: 20197778]
14. Soboleva, Ta; Nekrasov, M; Ryan, DP; Tremethick, DJ. Histone variants at the transcription start-site. *Trends Genet*. 2014; 30: 199–209. [PubMed: 24768041]
15. Zink L-M, Hake SB. Histone variants: nuclear function and disease. *Curr Opin Genet Dev*. 2016; 37: 82–89. [PubMed: 26826795]
16. Weber CM, Henikoff S. Histone variants: dynamic punctuation in transcription. *Genes Dev*. 2014; 28: 672–82. [PubMed: 24696452]
17. Borg M, Jiang D, Berger F. Histone variants take center stage in shaping the epigenome. *Curr Opin Plant Biol*. 2021; 61 101991 [PubMed: 33434757]
18. Zentner GE, Henikoff S. Regulation of nucleosome dynamics by histone modifications. *Nat Struct Mol Biol*. 2013; 20: 259–66. [PubMed: 23463310]
19. Campos EI, Reinberg D. Histones: annotating chromatin. *Annu Rev Genet*. 2009; 43: 559–99. [PubMed: 19886812]
20. Strahl BD, Allis CD. The language of covalent histone modifications. *Nature*. 2000; 403: 41–45. [PubMed: 10638745]
21. Bannister AJ, Kouzarides T. Regulation of chromatin by histone modifications. *Cell Res*. 2011; 21: 381–95. [PubMed: 21321607]
22. Talbert PB, Henikoff S. The Yin and Yang of Histone Marks in Transcription. *Annu Rev Genomics Hum Genet*. 2021; 22: 147–170. [PubMed: 33781079]
23. Taverna SD, Li H, Ruthenburg AJ, Allis CD, Patel DJ. How chromatin-binding modules interpret histone modifications: lessons from professional pocket pickers. *Nat Struct Mol Biol*. 2007; 14: 1025–40. [PubMed: 17984965]
24. Musselman, Ca; Lalonde, M-E; Côté, J; Kutateladze, TG. Perceiving the epigenetic landscape through histone readers. *Nat Struct Mol Biol*. 2012; 19: 1218–27. [PubMed: 23211769]
25. Gurard-Levin, Za; Quivy, J-P; Almouzni, G. Histone Chaperones: Assisting Histone Traffic and Nucleosome Dynamics. *Annu Rev Biochem*. 2014; 83: 487–517. [PubMed: 24905786]
26. Burgess RJ, Zhang Z. Histone chaperones in nucleosome assembly and human disease. *Nat Struct Mol Biol*. 2013; 20: 14–22. [PubMed: 23288364]
27. Koster MJE, Snel B, Timmers HTM. Genesis of Chromatin and Transcription Dynamics in the Origin of Species. *Cell*. 2015; 161: 724–736. [PubMed: 25957681]
28. Hargreaves DC, Crabtree GR. ATP-dependent chromatin remodeling: genetics, genomics and mechanisms. *Cell Res*. 2011; 21: 396–420. [PubMed: 21358755]
29. Gornik SG, et al. Loss of nucleosomal DNA condensation coincides with appearance of a novel nuclear protein in dinoflagellates. *Curr Biol*. 2012; 22: 2303–12. [PubMed: 23159597]
30. Mattioli F, et al. Structure of histone-based chromatin in Archaea. *Science*. 2017; 357: 609–612. [PubMed: 28798133]
31. Warnecke T, Becker Ea, Facciotti MT, Nislow C, Lehner B. Conserved substitution patterns around nucleosome footprints in eukaryotes and Archaea derive from frequent nucleosome repositioning through evolution. *PLoS Comput Biol*. 2013; 9 e1003373 [PubMed: 24278010]
32. Ammar R, et al. Chromatin is an ancient innovation conserved between Archaea and Eukarya. *Elife*. 2012; 1 e00078 [PubMed: 23240084]
33. Rojec M, Hocher A, Merckenschlager M, Warnecke T. Chromatinization of *Escherichia coli* with archaeal histones. *bioRxiv*. 2019; 660035 doi: 10.1101/660035

34. Forbes AJ, et al. Targeted analysis and discovery of posttranslational modifications in proteins from methanogenic archaea by top-down MS. *Proc Natl Acad Sci U S A*. 2004; 101: 2678–83. [PubMed: 14976258]
35. Weidenbach K, et al. Deletion of the archaeal histone in *Methanosarcina mazei* Gö1 results in reduced growth and genomic transcription. *Mol Microbiol*. 2008; 67: 662–671. [PubMed: 18086209]
36. Talbert PB, Meers MP, Henikoff S. Old cogs, new tricks: the evolution of gene expression in a chromatin context. *Nat Rev Genet*. 2019; doi: 10.1038/s41576-019-0105-7
37. de Mendoza A, Sebe-Pedros A. Origin and evolution of eukaryotic transcription factors. *Curr Opin Genet Dev*. 2019; 59: 25–32.
38. Schwaiger M, et al. Evolutionary conservation of the eumetazoan gene regulatory landscape. *Genome Res*. 2014; 24: 639–650. [PubMed: 24642862]
39. Sebé-Pedrós A, et al. Early metazoan cell type diversity and the evolution of multicellular gene regulation. *Nat Ecol Evol*. 2018; 2: 1176–1188. [PubMed: 29942020]
40. Connolly LR, Smith KM, Freitag M. The *Fusarium graminearum* Histone H3 K27 Methyltransferase KMT6 Regulates Development and Expression of Secondary Metabolite Gene Clusters. *PLoS Genet*. 2013; 9 e1003916 [PubMed: 24204317]
41. Jamieson K, Rountree MR, Lewis Za, Stajich JE, Selker EU. Regional control of histone H3 lysine 27 methylation in *Neurospora*. *Proc Natl Acad Sci*. 2013; 110: 6027–6032. [PubMed: 23530226]
42. Sebé-Pedrós A, et al. The Dynamic Regulatory Genome of *Capsaspora* and the Origin of Animal Multicellularity. *Cell*. 2016; 165: 1224–1237. [PubMed: 27114036]
43. Bourdareau S, et al. Histone modifications during the life cycle of the brown alga *Ectocarpus*. *Genome Biol*. 2021; 22: 12. [PubMed: 33397407]
44. Wang SY, et al. Role of epigenetics in unicellular to multicellular transition in *Dictyostelium*. *Genome Biol*. 2021; 22: 134. [PubMed: 33947439]
45. Taverna SD, Coyne RS, Allis CD. Methylation of Histone H3 at Lysine 9 Targets Programmed DNA Elimination in *Tetrahymena* University of Virginia Health System. *Cell*. 2002; 110: 701–711. [PubMed: 12297044]
46. Garcia, Ba; , et al. Organismal differences in post-translational modifications in histones H3 and H4. *J Biol Chem*. 2007; 282: 7641–55. [PubMed: 17194708]
47. Drinnenberg IA, et al. EvoChromo: towards a synthesis of chromatin biology and evolution. *Development*. 2019; 146 dev178962 [PubMed: 31558570]
48. Draizen EJ, et al. HistoneDB 2.0: a histone database with variants—an integrated resource to explore histones and their variants. *Database*. 2016; 2016 baw014 [PubMed: 26989147]
49. Maile TM, et al. Mass Spectrometric Quantification of Histone Post-translational Modifications by a Hybrid Chemical Labeling Method. *Mol Cell Proteomics*. 2015; 14: 1148–1158. [PubMed: 25680960]
50. Li B, Carey M, Workman JL. The Role of Chromatin during Transcription. *Cell*. 2007; 128: 707–719. [PubMed: 17320508]
51. Rajagopal N, et al. Distinct and Predictive Histone Lysine Acetylation Patterns at Promoters, Enhancers, and Gene Bodies. *G3 Genes|Genomes|Genetics*. 2014; 4: 2051–2063. [PubMed: 25122670]
52. Koonin EV, Yutin N. The Dispersed Archaeal Eukaryome and the Complex Archaeal Ancestor of Eukaryotes. *Cold Spring Harb Perspect Biol*. 2014; 6 a016188-a016188 [PubMed: 24691961]
53. Sandman K, Reeve JN. Archaeal histones and the origin of the histone fold. *Curr Opin Microbiol*. 2006; 9: 520–5. [PubMed: 16920388]
54. Pereira SL, Grayling Ra, Lurz R, Reeve JN. Archaeal nucleosomes. *Proc Natl Acad Sci U S A*. 1997; 94: 12633–7. [PubMed: 9356501]
55. Henneman B, van Emmerik C, van Ingen H, Dame RT. Structure and function of archaeal histones. *PLOS Genet*. 2018; 14 e1007582 [PubMed: 30212449]
56. Imachi H, et al. Isolation of an archaeon at the prokaryote-eukaryote interface. *bioRxiv*. 2019; 726976 doi: 10.1101/726976

57. Spang A, et al. Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature*. 2015; 521: 173–179. [PubMed: 25945739]
58. Zaremba-Niedzwiedzka K, et al. Asgard archaea illuminate the origin of eukaryotic cellular complexity. *Nature*. 2017; 541: 353–358. [PubMed: 28077874]
59. Da Cunha V, Gaia M, Nasir A, Forterre P. Asgard archaea do not close the debate about the universal tree of life topology. *PLOS Genet*. 2018; 14 e1007215 [PubMed: 29596428]
60. Alva V, Lupas AN. Histones predate the split between bacteria and archaea. *Bioinformatics*. 2019; 35: 2349–2353. [PubMed: 30520969]
61. Allis CD, et al. New Nomenclature for Chromatin-Modifying Enzymes. *Cell*. 2007; 131: 633–636. [PubMed: 18022353]
62. Wu F, et al. Unique mobile elements and scalable gene flow at the prokaryote-eukaryote boundary revealed by circularized Asgard archaea genomes. *Nat Microbiol*. 2022; 7: 200–212. [PubMed: 35027677]
63. Schuettengruber B, Bourbon H-M, Di Croce L, Cavalli G. Genome Regulation by Polycomb and Trithorax: 70 Years and Counting. *Cell*. 2017; 171: 34–57. [PubMed: 28938122]
64. Dion MF, Altschuler SJ, Wu LF, Rando OJ. Genomic characterization reveals a simple histone H4 acetylation code. *Proc Natl Acad Sci*. 2005; 102: 5501–5506. [PubMed: 15795371]
65. de Jong J, et al. Chromatin Landscapes of Retroviral and Transposon Integration Profiles. *PLoS Genet*. 2014; 10 e1004250 [PubMed: 24721906]
66. Sultana T, Zamborlini A, Cristofari G, Lesage P. Integration site selection by retroviruses and transposable elements in eukaryotes. *Nat Rev Genet*. 2017; 18: 292–308. [PubMed: 28286338]
67. Gao X, Hou Y, Ebina H, Levin HL, Voytas DF. Chromodomains direct integration of retrotransposons to heterochromatin. *Genome Res*. 2008; 18: 359–369. [PubMed: 18256242]
68. Cosby RL, et al. Recurrent evolution of vertebrate transcription factors by transposase capture. *Science*. 2021; 371 eabc6405 [PubMed: 33602827]
69. Cordaux R, Udit S, Batzer MA, Feschotte C. Birth of a chimeric primate gene by capture of the transposase gene from a mobile element. *Proc Natl Acad Sci*. 2006; 103: 8101–8106. [PubMed: 16672366]
70. Fiedler M, et al. Decoding of Methylated Histone H3 Tail by the Pygo-BCL9 Wnt Signaling Complex. *Mol Cell*. 2008; 30: 507–518. [PubMed: 18498752]
71. Erives AJ. Phylogenetic analysis of the core histone doublet and DNA topo II genes of *Marseilleviridae*: evidence of proto-eukaryotic provenance. *Epigenetics Chromatin*. 2017; 10: 55. [PubMed: 29179736]
72. Liu Y, et al. Virus-encoded histone doublets are essential and form nucleosome-like structures. *Cell*. 2021; 184: 4237–4250. e19 [PubMed: 34297924]
73. Valencia-Sánchez MI, et al. The structure of a virus-encoded nucleosome. *Nat Struct Mol Biol*. 2021; 28: 413–417. [PubMed: 33927388]
74. Iyer LM, Balaji S, Koonin EV, Aravind L. Evolutionary genomics of nucleo-cytoplasmic large DNA viruses. *Virus Res*. 2006; 117: 156–184. [PubMed: 16494962]
75. Nagamine T. Apoptotic arms races in insect-baculovirus coevolution. *Physiol Entomol*. 2021; phen.12371 doi: 10.1111/phen.12371
76. Starrett GJ, et al. Adintoviruses: An Animal-Tropic Family of Midsize Eukaryotic Linear dsDNA (MELD) Viruses. *bioRxiv*. 2020; 697771 doi: 10.1101/697771
77. Hocher A, et al. Growth temperature is the principal driver of chromatinization in archaea. *bioRxiv*. 2021; 2021.07.08.451601 doi: 10.1101/2021.07.08.451601
78. Alpha-Bazin B, et al. Lysine-specific acetylated proteome from the archaeon *Thermococcus gammatolerans* reveals the presence of acetylated histones. *J Proteomics*. 2021; 232 104044 [PubMed: 33161166]
79. Eme L, Spang A, Lombard J, Stairs CW, Ettema TJG. Archaea and the origin of eukaryotes. *Nat Rev Microbiol*. 2017; 15: 711–723. [PubMed: 29123225]
80. Akil C, Robinson RC. Genomes of Asgard archaea encode profilins that regulate actin. *Nature*. 2018; 562: 439–443. [PubMed: 30283132]

81. Koonin EV. The origin and early evolution of eukaryotes in the light of phylogenomics. *Genome Biol.* 2010; 11: 209. [PubMed: 20441612]
82. Sebé-Pedrós A, Grau-Bové X, Richards TA, Ruiz-Trillo I. Evolution and Classification of Myosins, a Paneukaryotic Whole-Genome Approach. *Genome Biol Evol.* 2014; 6: 290–305. [PubMed: 24443438]
83. Richards TA, Cavalier-Smith T. Myosin domain evolution and the primary divergence of eukaryotes. *Nature.* 2005; 436: 1113–8. [PubMed: 16121172]
84. Wickstead B, Gull K, Richards T. Patterns of kinesin evolution reveal a complex ancestral eukaryote with a multifunctional cytoskeleton. *BMC Evol Biol.* 2010; 10: 110. [PubMed: 20423470]
85. Dacks JB, Field MC. Evolution of the eukaryotic membrane-trafficking system: origin, tempo and mode. *J Cell Sci.* 2007; 120: 2977–85. [PubMed: 17715154]
86. Collins L, Penny D. Complex Spliceosomal Organization Ancestral to Extant Eukaryotes. *Mol Biol Evol.* 2005; 22: 1053–1066. [PubMed: 15659557]
87. Grau-Bové X, Sebé-Pedrós A, Ruiz-Trillo I. The Eukaryotic Ancestor Had a Complex Ubiquitin Signaling System of Archaeal Origin. *Mol Biol Evol.* 2015; 32: 726–739. [PubMed: 25525215]
88. Kundaje A, et al. Integrative analysis of 111 reference human epigenomes. *Nature.* 2015; 518: 317–330. [PubMed: 25693563]
89. Ho JW, et al. Comparative analysis of metazoan chromatin organization. *Nature.* 2014; 512: 449–452. [PubMed: 25164756]
90. Montgomery SA, et al. Chromatin Organization in Early Land Plants Reveals an Ancestral Association between H3K27me3, Transposons, and Constitutive Heterochromatin. *Curr Biol.* 2020; 30: 573–588. e7 [PubMed: 32004456]
91. Frapporti A, et al. The Polycomb protein Ezi1 mediates H3K9 and H3K27 methylation to repress transposable elements in *Paramecium*. *Nat Commun.* 2019; 10: 2710 [PubMed: 31221974]
92. Lennartsson A, Ekwall K. Histone modification patterns and epigenetic codes. *Biochim Biophys Acta.* 2009; 1790: 863–8. [PubMed: 19168116]
93. Peterson CL, Laniel M-A. Histones and histone modifications. *Curr Biol.* 2004; 14: R546–51. [PubMed: 15268870]
94. Rando OJ. Combinatorial complexity in chromatin structure and function: revisiting the histone code. *Curr Opin Genet Dev.* 2012; 22: 148–155. [PubMed: 22440480]
95. de Mendoza A, Pflueger J, Lister R. Capture of a functionally active methyl-CpG binding domain by an arthropod retrotransposon family. *Genome Res.* 2019; 29: 1277–1286. [PubMed: 31239280]
96. De Mendoza A, et al. Recurrent acquisition of cytosine methyltransferases into eukaryotic retrotransposons. *Nat Commun.* 2018; 9: 1–11. [PubMed: 29317637]
97. Ji X, et al. Chromatin proteomic profiling reveals novel proteins associated with histone-marked genomic regions. *Proc Natl Acad Sci U S A.* 2015; 112: 3841–3846. [PubMed: 25755260]
98. Wierer M, Mann M. Proteomics to study DNA-bound and chromatin-associated gene regulatory complexes. *Hum Mol Genet.* 2016; 25: R106–R114. [PubMed: 27402878]
99. Villaseñor R, et al. ChromID identifies the protein interactome at chromatin marks. *Nat Biotechnol.* 2020; 38: 728–736. [PubMed: 32123383]
100. Stieglmeier M, et al. *Nitrososphaera viennensis* gen. nov., sp. nov., an aerobic and mesophilic, ammonia-oxidizing archaeon from soil and a member of the archaeal phylum Thaumarchaeota. *Int J Syst Evol Microbiol.* 2014; 64: 2738–2752. [PubMed: 24907263]
101. Tirichine L, et al. Histone extraction protocol from the two model diatoms *Phaeodactylum tricorutum* and *Thalassiosira pseudonana*. *Mar Genomics.* 2014; 13: 21–25. [PubMed: 24315927]
102. Shechter D, Dormann HL, Allis CD, Hake SB. Extraction, purification and analysis of histones. *Nat Protoc.* 2007; 2: 1445–57. [PubMed: 17545981]
103. Perkins DN, Pappin DJC, Creasy DM, Cottrell JS. Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis.* 1999; 20: 3551–3567. [PubMed: 10612281]

104. Taus T, et al. Universal and confident phosphorylation site localization using phosphoRS. *J Proteome Res.* 2011; 10: 5354–5362. [PubMed: 22073976]
105. Vizcaíno JA, et al. 2016 update of the PRIDE database and its related tools. *Nucleic Acids Res.* 2016; 44: D447–56. [PubMed: 26527722]
106. Hagberg, AA; Schult, DA; Swart, PJ. In: Varoquaux, G; Vaught, T; Millman, J, editors. Exploring Network Structure, Dynamics, and Function using NetworkX; Proceedings of the 7th Python in Science Conference; 2008. 11–15.
107. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol Biol Evol.* 2013; 30: 772–780. [PubMed: 23329690]
108. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol.* 2015; 32: 268–274. [PubMed: 25371430]
109. Veluchamy A, et al. An integrative analysis of post-translational histone modifications in the marine diatom *Phaeodactylum tricornutum*. *Genome Biol.* 2015; 16: 102. [PubMed: 25990474]
110. Ren Q, Gorovsky Ma. Histone H2A.Z acetylation modulates an essential charge patch. *Mol Cell.* 2001; 7: 1329–35. [PubMed: 11430834]
111. Allis CD, et al. hv1 is an evolutionarily conserved H2A variant that is preferentially associated with active genes. *J Biol Chem.* 1986; 261: 1941–1948. [PubMed: 3944120]
112. Fusauchi Y, Iwai K. Tetrahymena histone H2A. Acetylation in the N-terminal sequence and phosphorylation in the C-terminal sequence. *J Biochem.* 1984; 95: 147–154. [PubMed: 6706903]
113. Xiong L, Adhvaryu KK, Selker EU, Wang Y. Mapping of Lysine Methylation and Acetylation in Core Histones of *Neurospora crassa*. *Biochemistry.* 2010; 49: 5236–5243. [PubMed: 20433192]
114. Zhang K, Sridhar VV, Zhu J, Kapoor A, Zhu JK. Distinctive core histone post-translational modification patterns in *Arabidopsis thaliana*. *PLoS One.* 2007; 2
115. Johnson L, et al. Mass spectrometry analysis of *Arabidopsis* histone H3 reveals distinct combinations of post-translational modifications. *Nucleic Acids Res.* 2004; 32: 6511–6518. [PubMed: 15598823]
116. Bergmüller E, Gehrig PM, Gruissem W. Characterization of post-translational modifications of histone H2B-variants isolated from *Arabidopsis thaliana*. *J Proteome Res.* 2007; 6: 3655–3668. [PubMed: 17691833]
117. Beck HC, et al. Quantitative Proteomic Analysis of Post-translational Modifications of Human Histones. *Mol Cell Proteomics.* 2006; 5: 1314–1325. [PubMed: 16627869]
118. Goudarzi A, et al. Dynamic Competing Histone H4 K5K8 Acetylation and Butyrylation Are Hallmarks of Highly Active Gene Promoters. *Mol Cell.* 2016; 62: 169–180. [PubMed: 27105113]
119. Hake SB, et al. Expression patterns and post-translational modifications associated with mammalian histone H3 variants. *J Biol Chem.* 2006; 281: 559–568. [PubMed: 16267050]
120. Tan M, et al. Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. *Cell.* 2011; 146: 1016–1028. [PubMed: 21925322]
121. Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO. Dynamic genome evolution and complex virocell metabolism of globally-distributed giant viruses. *Nat Commun.* 2020; 11: 1–11. [PubMed: 31911652]
122. Punta M, et al. The Pfam protein families database. *Nucleic Acids Res.* 2012; 40: D290–301. [PubMed: 22127870]
123. Eddy SR. Accelerated profile HMM searches. *PLoS Comput Biol.* 2011; 7 e1002195 [PubMed: 22039361]
124. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nat Methods.* 2015; 12: 59–60. [PubMed: 25402007]
125. Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale detection of protein families. *Nucleic Acids Res.* 2002; 30: 1575–1584. [PubMed: 11917018]
126. Steenwyk JL, Buida TJ, Li Y, Shen X-X, Rokas A. ClipKIT: A multiple sequence alignment trimming software for accurate phylogenomic inference. *PLOS Biol.* 2020; 18 e3001007 [PubMed: 33264284]

127. Minh BQ, Nguyen MAT, von Haeseler A. Ultrafast approximation for phylogenetic bootstrap. *Mol Biol Evol.* 2013; 30: 1188–95. [PubMed: 23418397]
128. Grau-Bové X, Sebé-Pedrós A. Orthology Clusters from Gene Trees with Possvm. *Mol Biol Evol.* 2021; 38: 5204–5208. [PubMed: 34352080]
129. Huerta-Cepas J, Dopazo H, Dopazo J, Gabaldón T. The human phylome. *Genome Biol.* 2007; 8: R109. [PubMed: 17567924]
130. Huerta-Cepas J, Serra F, Bork P. ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic Data. *Mol Biol Evol.* 2016; 33: 1635–1638. [PubMed: 26921390]
131. Csörös M, Miklós, I. A Probabilistic Model for Gene Content Evolution with Duplication, Loss, and Horizontal Transfer BT - *Research in Computational Molecular Biology*. Apostolico, A, Guerra, C, Istrail, S, Pevzner, PA, Waterman, M, editors. Springer; Berlin Heidelberg: 2006. 206–220.
132. Csörös M. Count: evolutionary analysis of phylogenetic profiles with parsimony and likelihood. *Bioinformatics.* 2010; 26: 1910–2. [PubMed: 20551134]
133. Gansner ER, North SC. An open graph visualization system and its applications to software engineering. *Softw Pract Exp.* 2000; 30: 1203–1233.
134. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics.* 2012; 28: 3150–3152. [PubMed: 23060610]
135. Jombart T, Balloux F, Dray S. adephylo: new tools for investigating the phylogenetic signal in biological traits. *Bioinformatics.* 2010; 26: 1907–1909. [PubMed: 20525823]
136. Wells JN, Feschotte C. A Field Guide to Eukaryotic Transposable Elements. *Annu Rev Genet.* 2020; 54: 539–561. [PubMed: 32955944]
137. Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. The Dfam community resource of transposable element families, sequence models, and genome annotations. *Mob DNA.* 2021; 12: 2. [PubMed: 33436076]
138. Camacho C, et al. BLAST+: architecture and applications. *BMC Bioinformatics.* 2009; 10: 421. [PubMed: 20003500]
139. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics.* 2009; 25: 1754–1760. [PubMed: 19451168]
140. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinforma.* 2010; 26: 841–842.
141. Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O. Survey of Branch Support Methods Demonstrates Accuracy, Power, and Robustness of Fast Likelihood-based Approximation Schemes. *Syst Biol.* 2011; 60: 685–699. [PubMed: 21540409]
142. Bodenhofer U, Bonatesta E, Horejš-Kainrath C, Hochreiter S. msa: an R package for multiple sequence alignment. *Bioinformatics.* 2015; 31: 3997–3999. [PubMed: 26315911]




Figure 1. Diversity of post-translational modifications in eukaryotic canonical and variant histones.

a. Eukaryotic taxon sampling used in this study. Colored dots indicate the number of species used in the comparative histone proteomics reconstruction, with solid dots indicating new species added in this analysis. Numbers in brackets indicate the number of genomes/transcriptomes used in the comparative genomics analyses. Dashed lines indicate uncertain phylogenetic relationships. Complete list of sampled species in Supplementary Data 1. Silhouettes adapted from <http://phylopic.org/>. **b.** Networks of pairwise protein similarity between histone protein domains in eukaryotes, archaea and viruses. Each node represents

one histone domain, colored according to their best alignment in the HistoneDB database (see Methods). Edges represent local alignments (bitscore ≥ 20). **c**, Schematic representation of the hPTM proteomics strategy employed in this study. **d**, Conservation of hPTMs in eukaryotic histones. hPTM coordinates are reported according to the amino-acid position in human orthologs (if conserved). In H2A and H2B, question marks indicate the presence of hPTMs in stretches of lysine residues of uncertain homology. In species with previously reported hPTMs, we further indicate which variants were also identified in our reanalysis. Only positions with hPTMs conserved in more than one species are reported (full table and consensus alignments available in Supplementary Data 3). **e**, Maximum likelihood phylogenetic trees of the connected components in panel b, corresponding to eukaryotic histones (H3, H4, H2A, H2B). Canonical histones included in panel d and variant histones detected are highlighted in red. hPTMs detected in non-canonical histones are indicated. Bottom, distributions of pairwise phylogenetic distances between all proteins in each gene tree. Violin plots above each distribution represent the distribution of distances between reference histones present in the HistoneDB database and histones with proteomic evidence included in our study, for each of the main canonical (H3, H4, H2A, and H2B) and variant histones (H2A.Z and macroH2A). Dots in the violin plot distributions represent the median.




Figure 2. Archaeal histone diversity and post-translational modifications.

a, Distribution of histones (fraction of taxa in each lineage) and histone tails (presence/absence) across Archaea phyla. **b**, Summary of proteomics evidence of archaeal histones, including the presence of modifications, tails, coverage, fraction of lysines identified, and isoelectric points. Human Histone H3 and H4 are included for reference. The alignments at the bottom depict the position of lysine modifications in the globular part of *Methanospirillum stamsii* and *Methanobrevibacter cuticularis* HMfB histones (modified residues in bold). **c**, Archaeal HMfB histones with N-terminal tails (at least 10 aa before a complete globular domain), sorted by frequency of lysine residues in the tail and color-coded according to taxonomy (same as panel A). Amino-acid sequences shown for selected examples. The dotted line indicates the median frequency of lysines in canonical eukaryotic H3 and H4 histone tails. Source data available in Supplementary Data 2. **d**, Mass spectra of three modified archaeal peptides, representing the relative abundance of fragments at various mass-to-charge ratios (m/z). Spectra were annotated using IPSA. b and y ions and their losses of H₂O are marked in green and purple, respectively; precursor ions are marked in dark grey. Unassigned peaks are marked in light grey. Some labels have been omitted to facilitate readability.




Figure 3. Taxonomic distribution of chromatin-associated gene classes.

a, Summary of the seven classes of genes with chromatin-related activity covered in our survey: histone-specific hPTM writers (acetylases and methyltransferases), erasers (deacetylases and demethylases), readers, remodellers, and chaperones. **b**, Percentage of surveyed taxa containing homologs from each chromatin-associated gene class, for eukaryotes (top), archaea, bacteria, and viruses (bottom). Species-level tables are available in Extended Data Fig. 3. **c**, Number of eukaryotic genes classified in each of the chromatin-associated modification enzymes, readers, remodellers, and chaperones. **d**, Overlap between the taxon-level phylogenetic distribution of histones and chromatin-associated domains in

archaea and four bacterial phyla, measured using the Jaccard index. **e**, Number of genes encoding writer, eraser, reader and remodeller domains, per species.




Figure 4. Origin and evolution of chromatin-associated gene families.

a, Summary of phylogenetic affinities of the eukaryotic homologs of gene classes that are also present in prokaryotes. For each gene family, we evaluate whether it is phylogenetically closer to a majority (> 50%) of eukaryotic sequences from a different orthogroup (indicating intra-eukaryotic diversification), or to sequences from Bacteria or Archaea. **b**, Left, gene tree of eukaryotic and prokaryotic Sirtuin deacetylases, showcasing an example of a eukaryotic family that diversified within eukaryotes (SIRT6) and another one with close relatives in Asgard archaea (SIRT7). Right, gene tree of KAT14 acetylase, a eukaryotic orthogroup with

bacterial origins. Statistical supports (UF bootstrap) are shown at selected internal nodes of the highlighted clades. **c**, Evolutionary reconstruction of hPTM writer and eraser gene families, remodellers, and histone chaperones along the eukaryotic phylogeny, including the number of genes present in the last eukaryotic common ancestor (LECA). Barplots indicate the number of orthologs of each gene family present at the LECA (at 90% posterior probability; see Methods) and whether the presence of a given orthogroup at LECA is supported by its conservation in various early-branching eukaryotic lineages (Amorphea, Discoba, Diaphoretickes and others). The list of ancestral gene families below each plot is non-exhaustive. Two ancestral gene counts are provided: all families at presence probability above 90%, and, in brackets, the subset of these that is present in at least two of the main eukaryotic early-branching lineages (Amorphea, Diaphoretickes, and Discoba). Source data in Supplementary Data 5. **d-e**, Reconstructed evolutionary origins of the different subunits of the Polycomb repressive complexes (PRC2 and PRC1) and Trithorax-group complexes (KMT1 to 5). **f-h**, Side-by-side comparison of the presence of individual hPTM marks and various subunits of the Polycomb and Trithorax complexes, as well as other hPTM writers, responsible for their deposition.




Figure 5. Evolution of chromatin readers and capture of chromatin proteins by transposable elements and viruses.

a, Evolutionary reconstruction of reader gene families along the eukaryotic phylogeny, highlighting the number of gains along the eukaryotic phylogeny (at 90% posterior probability). The Euler diagram at the top shows the overlap between presence of chromatin-associated catalytic domains and readers. The barplot at the left indicates the number of orthologs of each gene family present at the LECA and whether their presence is supported by its conservation in various early-branching eukaryotic lineages (Amorphea, Discoba,

Diaphoretickes, and others). Pie plots at the right summarize the number of orthogroups from each gene family gained within selected lineages: Metazoa, Holomycota, Viridiplantae and SAR+Haptophyta. **b**, Number of reader or catalytic orthogroups gained at each node in the species tree, for selected species. Source data in Supplementary Data 5. **c**, Networks of protein domain co-occurrence for Chromo and PHD readers. Each node represents a protein domain that co-occurs with Chromo or PHD domains, and node size denotes the number of co-occurrences with either Chromo or PHD. Edges represent co-occurrences between domains. Groups of frequently co-occurring protein domains have been manually annotated and color-coded, which has revealed sub-sets of retrotransposon and DNA transposon-associated domains. **d**, Number of chromatin-related eukaryotic genes fused with transposons grouped by gene family (left), including the fraction that are classified as valid gene models based on expression and assembly data (centre); and the number of species where each type of fusion is found (right). The number of fusion events are colored according to their similarity with known DNA transposons (red) or retrotransposons (orange) from the Dfam database (see Methods). (*) The 'Chromo' category excludes genes containing other chromatin-associated protein domains such as SNF2_N (listed separately as 'Chromo+SNF2_N', which includes remodellers with the domain of unknown function DUF1087, which is also common in DNA transposons). **e**, Selected examples of transposon fusion domains classified by orthogroup, including their archetypical protein domain architecture, homology to transposon class, their phylogenetic distribution, and number of fusion genes. Only orthogroups with at least one valid gene model are listed. Source data available in Supplementary Data 6. **f**, Example tree of Chromo readers, highlighting genes with fused TE-associated domains and their consensus domain architectures. **g**, Fraction of viral genomes containing homologs from each chromatin gene family, for nucleocytoplasmic giant DNA virus families (top) and other taxa containing histone domains (Nudiviridae, Polydnviridae; bottom). **h**, Phylogenetic analysis of histone domains, with a focus on viral homologs. Statistical supports (approximate Bayes posterior probabilities) are shown for the deepest node of each canonical eukaryotic or archaeal histone clade. The inset table summarizes the presence of doublet histone genes per lineage. **i**, Number of viral homologs in each chromatin-associated gene family, classified according to their closest cellular homologs (eukaryotes, bacteria or archaea) in phylogenetic analyses (see Methods). Source data available in Supplementary Data 6.




Figure 6. Chromatin evolution and eukaryogenesis.

a, Summary of events in chromatin evolution prior to, during and after the origin of eukaryotes. **b**, Number of chromatin-related gene families and hPTM marks inferred to have been present at the LECA. Ancestral gene counts are indicated at >90% probability. For gene counts, numbers within bars indicate the subset of families present in at least two of the most deeply-sampled early-branching eukaryotic lineages (Amorpha, Diaphoretickes, and Discoba). For hPTMs, the ancestral counts have been inferred using Dollo parsimony assuming a Diaphoretickes – Amorphea split at the root of eukaryotes, and numbers within bars indicate the number of hPTMs whose ancestral presence is supported by more than one species at both sides of the root. **c**, hPTMs inferred to be present in the last eukaryotic common ancestor (LECA) based on Dollo parsimony. Only amino-acid positions conserved in all eukaryotes in our dataset are shown. Asterisks indicate modifications whose presence at the LECA is supported by just one species at either side of the root. The inferred LECA presence of known writing/erasing enzymes associated to these hPTM is indicated.