

RETRACTION

Open Access

Retraction: Does changing from a first generation antipsychotic (perphenazin) to a second generation antipsychotic (risperidone) alter brain activation and motor activity? A case report

Jan Øystein Berle^{*}, Else-Marie Løberg and Ole Bernt Fasmer

Retraction

The authors have retracted this article [1] as the fMRI data presented in the case report are incorrect. The activation data reported for session 1 are the activation data for session 2 and vice versa. As a result the discussion and conclusions of the case report are based on the wrong set of data and are no longer valid. The authors apologise for the error.

Received: 20 August 2013 Accepted: 20 August 2013
Published: 20 August 2013

References

1. Berle JO, Else-Marie L, Ole Bernt F: Does changing from a first generation antipsychotic (perphenazin) to a second generation antipsychotic (risperidone) alter brain activation and motor activity? A case report. *BMC Research Notes* 2013, **6**:182.

doi:10.1186/1756-0500-6-332

Cite this article as: Berle et al.: Retraction: Does changing from a first generation antipsychotic (perphenazin) to a second generation antipsychotic (risperidone) alter brain activation and motor activity? A case report. *BMC Research Notes* 2013 **6**:332.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



^{*} Correspondence: jaob@helse-bergen.no
Division of Psychiatry, Haukeland University Hospital, Bergen, Norway