

Research article

Open Access

RpoE fine tunes expression of a subset of SsrB-regulated virulence factors in *Salmonella enterica* serovar Typhimurium

Suzanne E Osborne¹ and Brian K Coombes*^{1,2}

Address: ¹Michael G. DeGroot Institute for Infectious Disease Research, Department of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, ON, Canada and ²Laboratory for Foodborne Zoonoses, Public Health Agency of Canada, Guelph, ON, Canada

Email: Suzanne E Osborne - osborns@mcmaster.ca; Brian K Coombes* - coombes@mcmaster.ca

* Corresponding author

Published: 26 February 2009

Received: 15 October 2008

BMC Microbiology 2009, 9:45 doi:10.1186/1471-2180-9-45

Accepted: 26 February 2009

This article is available from: <http://www.biomedcentral.com/1471-2180/9/45>

© 2009 Osborne and Coombes; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The survival of *Salmonella enterica* within the intracellular host niche requires highly co-ordinated expression of virulence effectors predominantly regulated by the SsrAB two-component regulatory system. *S. enterica* serovar Typhimurium mutants lacking the *ssrAB* genes are avirulent in mice, highlighting the importance of this regulatory system *in vivo*. Mutants lacking the gene encoding the alternative sigma factor σ^E (*rpoE*) are also highly attenuated for intracellular survival, pointing to a potential connection with the SsrAB regulatory system.

Results: In this study we demonstrate that RpoE is involved in fine-tuning the expression of a subset of SsrB-regulated genes found in the *Salmonella* pathogenicity island-2 (SPI-2) genetic locus that encodes a horizontally acquired type III secretion system, and unlinked genes integrated into this regulon that are required for virulence in host animals.

Conclusion: These data point to a potential connection between the virulence phenotype of strains lacking *ssrB* and *rpoE*, and highlight new transcriptional regulation that might be essential for appropriate temporal and spatial control of the virulence-associated type III secretion system during host infection.

Background

Salmonella enterica are enteric pathogens that acquired a type III secretion system (T3SS) through horizontal gene transfer of a genomic island termed *Salmonella* Pathogenicity Island 2 (SPI-2) [1,2]. The SPI-2-encoded T3SS and its translocated effectors modify the intracellular host niche for *Salmonella* replication [3-5]. SPI-2 also has genes, *ssrA* and *ssrB*, which code for SsrAB, a two-component regulatory system needed for expression of the T3SS [6,7]. SsrB regulates the expression of SPI-2 encoded substrate effectors including *ssaB*, as well as several integrated virulence effectors such as *sseL* [8] and *srfN* [9] that are encoded elsewhere on the chromosome but that have

integrated into the SsrB regulon. Mutants lacking *ssrAB* are unable to survive within macrophages and are avirulent in mice [1].

Alternative sigma factors coordinate gene expression in response to environmental cues sensed by the bacterium. Sigma factors have a specific recognition motif at the -35 and -10 positions and function to concentrate RNA polymerase at a subset of promoters [10]. One alternative sigma factor, RpoE (σ^E) responds to envelope stress at the cell surface. Release of σ^E from its inner membrane anchored anti-sigma factor, RseA, leads to induction of genes required to maintain cell envelope integrity [11].

SsrB-regulated translocated effectors protect *S. Typhimurium* against host cell defences such as oxidative stress and antimicrobial peptides that perturb bacterial membrane integrity and provide a stimulus for σ^E release [4,12-15]. Although proficient at cellular invasion, *rpoE* or *ssrB* mutants are highly attenuated for intracellular survival in both cultured cells and animal hosts [16]. In addition, the expression of *rpoE* and *ssrB* is up-regulated within macrophages [17]. Links between RpoE and virulence gene expression is evident in other bacterial systems as well. Deletion of *rseA* in *Yersinia pseudotuberculosis* causes elevated virulence effector synthesis and secretion [18], establishing links between alternative sigma factors and virulence-specific regulators. Taken together, a connection between σ^E and SsrB is suggested from the available literature, however the role of σ^E in activating SsrB-regulated genes has not been studied.

We tested the hypothesis that RpoE is involved in expression of genes that use the SsrB response regulator for activation. By testing six promoters representing four classes of SsrB-regulated promoters ((i) two type III secretion structural operons in SPI-2, (ii) the effector operon in SPI-2, (iii) two effector genes unlinked with SPI-2, and (iv) an integrated virulence gene unlinked with SPI-2) we demonstrate that RpoE elicits an effect on a subset of SsrB-regulated genes. This effect was bidirectional depending on the promoter and was downstream of *ssrB* expression itself, since deletion of *rpoE* had no effect on SsrB levels in the mutant cells. These data help unite the virulence phenotypes of strains lacking SsrB and RpoE, and highlight new transcriptional regulation that might be essential for appropriate temporal and spatial control of the virulence-associated type III secretion system during host infection.

Results

Deletion of *rpoE* affects a subset of SsrB-regulated virulence genes

Salmonella virulence gene expression is coordinated *in vivo* and may be regulated, in part, by alternative sigma factor(s) in order to quickly respond to the host environment. To date, no sigma factor has been identified as regulating SsrB-dependent virulence genes. To start, we first screened four alternative sigma factor mutants of *S. Typhimurium* (*rpoS*, *rpoN*, *rpoE*, *rpoH*) for their ability to express a key virulence gene, *sseB*, that requires SsrB for expression and whose gene product is essential for intracellular pathogenesis. For an *rpoH* deletion, this strain was only viable at temperatures below 30°C. Since SPI-2 gene activation is integrated into a thermosensing circuit [19] we were unable to test the role of σ^H in this study (data not shown). In this screen, *rpoS* deletion resulted in a slight increase in SseB levels (Figure 1A) indicating a role for RpoS in the repression of SPI-2. Both *rpoE* and *rpoN* deletions resulted in decreased SseB levels with a more pro-

nounced effect in the *rpoE* deletion. Since we were predominantly interested in sigma factors that activate SPI-2 and which could be linked to the previous observation that *rpoE* mutants are highly attenuated *in vivo* we choose to focus on RpoE in the current study, which had the most influence on SseB levels in the cell.

An unmarked in-frame deletion of *rpoE* was then generated in *S. Typhimurium* strain SL1344 and we verified that this in-frame deletion had the same effect on SseB as




Figure 1

Figure 1

Loss of *rpoE* changes the abundance of virulence factors in *Salmonella*. (A) wild type (wt), $\Delta rpoE$, $\Delta rpoS$, and $\Delta rpoN$ *S. Typhimurium* 14028s were grown for 6 hours under SsrB-inducing conditions. Lysates were probed by western blot for SseB, a component of the T3SS needle complex. (B) Western blot was performed as above and lysates probed for SseB in wild type (wt) *S. Typhimurium* SL1344, $\Delta rpoE$ and in $\Delta rpoE$ complemented with pVWSK29 carrying full length *rpoE* with endogenous promoters. (C) Wild type *S. Typhimurium* SL1344 and $\Delta rpoE$ cells were immunoblotted as above and lysates probed for SseL-2HA, SrfN-2HA and SifA-2HA which were expressed from their endogenous promoters in pVWSK29. Blots were probed for DnaK as a control. The experiment was performed three times with similar results.

the *rpoE::cat* mutant used previously (Figure 1B). A low-copy plasmid containing full-length *rpoE* and the three endogenous promoters that can drive its expression [20] was able to restore wild type levels of SseB to $\Delta rpoE$ cells (Figure 1B) demonstrating that the results were specific to the *rpoE* deletion. In these complementation experiments, attempts were made to examine the levels of SseB secreted into the culture supernatant [21], however consistent with previous observations [22,23] perturbations to the *rpoE* pathway increased cell lysis resulting in contamination of secreted fractions with cytosolic proteins which precluded accurate interpretation (data not shown).

In order to examine the effect of σ^E (*rpoE*) on the expression of a broad range of SsrB-regulated virulence genes, we tested whether or not the effect of *rpoE* deletion was specific to *sseB* or if it extended to other SsrB-regulated genes. To do this we examined the levels of SseL-2HA, SifA-2HA and SrfN-2HA expressed from their endogenous promoters under SPI-2 inducing conditions (Figure 1C). Consistent with the results for SseB, there was a decrease in SifA-2HA levels in $\Delta rpoE$ compared to wild type, although deletion of *rpoE* did not have an effect on SseL-2HA. Relative to its expression in wild type cells, the level of SrfN-2HA was reproducibly increased in the $\Delta rpoE$ cells, suggesting a role for σ^E in the repression of SrfN, although it is unlikely that this is through a direct mechanism.

***RpoE* is involved in transcriptional activity of a subset of virulence genes**

In order to confirm the effect of σ^E on the expression of a broad range of SsrB-regulated virulence genes, we used wild type and $\Delta rpoE$ cells and integrated into the chromosome individually six single-copy transcriptional fusions representing promoters for four classes of SsrB-dependent genes or operons ((i) type III secretion effector operon

(*sseA*); (ii) structural operon I (*ssaB*); (iii) structural operon II (*ssaG*); (iv and v) effectors encoded outside of SPI-2 (*sseL* and *sifA*); and (vi) integrated virulence genes unlinked to SPI-2 (*srfN*) [9]. Transcriptional fusions to *lacZ* of the *sseA*, *ssaB*, *ssaG*, *sseL*, *sifA* and *srfN* promoters (mapped previously; [9,24] were integrated into the chromosome of wild type and $\Delta rpoE$ cells and then grown in the SsrB-activating medium LPM. The activity of each promoter was measured using a β -galactosidase assay during exponential growth. Although $\Delta rpoE$ was observed to have a slightly prolonged lag phase relative to wild type cells under these experimental conditions, at later time points the mutant grew similarly to wild type. To account for any differences in growth kinetics of the cultures, all data was normalized to the optical density at 600 nm of the culture, which permitted direct comparisons.

In wild type cells, promoter activity from all the transcriptional fusions was high, as expected, because LPM medium is highly inducing for SsrB activity [21]. In contrast, promoter activity for *sseA*, *ssaB*, and *sifA* decreased in the *rpoE* mutant compared to wild type cells (Figure 2A, B and 2D), whereas promoter activity from the *ssaG* and *srfN* reporters was upregulated in the *rpoE* mutant (Figure 2C and 2F). β -galactosidase activity observed from the *sseL* reporter was unaltered in the *rpoE* deletion compared to that in wild type cells (Figure 2E). These data are consistent with the protein levels detected for these gene products. Together, these data indicate that σ^E can have a variable and bidirectional effect on SsrB-regulated virulence genes.

The effect of *RpoE* on virulence genes is downstream of *ssrB* expression

The variable effects of *rpoE* deletion on SsrB-regulated effectors suggested that RpoE might direct transcription downstream of *ssrB* expression. To test this, we replaced the *ssrB* gene in $\Delta rpoE$ and wild type cells with an *ssrB::FLAG* allele [19] and examined the levels of SsrB protein in the strains by western blot. There was no change in the levels of SsrB-FLAG between wild type and $\Delta rpoE$ cells (Figure 3), indicating that the effects of RpoE on the four classes of virulence gene promoters examined here was not mediated through changes to SsrB protein levels. Together these data establish a role for RpoE in the fine-tuning of virulence gene expression in *S. Typhimurium*.

Discussion

In this work we found that the alternative sigma factor, σ^E , is involved in fine tuning the expressing of a subset of SsrB-regulated virulence genes required for *Salmonella* pathogenesis. Although the effect of *rpoE* deletion on promoter activity in some cases was mild, we have previously shown that gene regulators providing only modest transcriptional input have a profound influence on bacterial




Figure 2

The transcriptional activity of SsrB-regulated virulence genes is affected by an *rpoE* deletion. Wild type and $\Delta rpoE$ cells carrying single-copy chromosomal transcriptional reporters of (A) $P_{sseA}::pP_{sseA-lacZ}$, (B) $P_{ssaB}::pP_{ssaB-lacZ}$, (C) $P_{ssaG}::pP_{ssaG-lacZ}$, (D) $P_{sifA}::pP_{sifA-lacZ}$, (E) $P_{sseL}::pP_{sseL-lacZ}$ and (F) $P_{srfN}::pP_{srfN-lacZ}$ were grown in LPM (pH 5.8). At the indicated time β -galactosidase activity was measured and expressed as relative light units (RLU) normalized to optical density of the culture. Wild type and $\Delta rpoE$ cells lacking the transcriptional reporters were used as controls in each experiment. Data are the means with standard error from triplicate determinations from three independent experiments.




Figure 3
The effect of RpoE on SsrB-regulated genes is downstream of *ssrAB* expression. The *ssrB* gene in wild type and $\Delta rpoE$ cells was replaced with an *ssrB-FLAG* allele in its native location on the chromosome. Cells were grown under SsrB-activating conditions for six hours and lysates were probed by western blot to detect SsrB-FLAG and intracellular DnaK as a control. The data shown are representative of two experiments performed independently with identical results.

fitness in a host animal [25]. In cases where the regulator is deleted, the loss of genetic fine-tuning causes incongruous changes in the timing and magnitude of virulence gene expression, leading to fitness loss and strong attenuation. We predict that RpoE confers a similar fine-tuning effect on *Salmonella* virulence gene expression that is required for optimal within-host fitness during infection.

When we examined the -10 and -35 positions of the promoters studied here relative to the transcriptional start sites identified previously [24], these promoters did not appear to contain σ^E consensus sequences. Instead they appeared to have consensus sites for σ^{70} . Although a bioinformatics screen identified σ^E consensus sequences upstream of the SPI-2 genes *ssaU*, *ssaJ*, *sscA* and *ssaC* [26], these genes were not tested for σ^E -dependence in the present study because the identified consensus sites are in coding sequence within operons, and as a result may not be directly relevant. Due to the high degree of conservation in σ factor binding sequences, σ^E may not be directly regulating SsrB-dependent promoters. The lack of a canonical σ^E sequence at these promoters suggests that

another regulatory gene may be epistatic to σ^E or that these promoters encode functional, but non-canonical σ^E -binding sites due to their horizontal acquisition and gradual integration into the σ^E regulatory network. This integration may help *Salmonella* coordinate expression of the virulence-associated T3SS in response to host factors that compromise bacterial membrane integrity (Figure 4). This mechanism would activate a restorative σ^E pathway, which is consistent with the enhanced susceptibility of *rpoE* mutants to oxidative stress and antimicrobial peptides [13,15,16], both of which perturb membrane integrity *in vivo*. Although there is no evidence that σ^E can directly repress transcription, the negative effect on two promoters observed here might be due to an intermediate RpoE-regulated repressor or compensatory effect where loss of *rpoE* increases the relative abundance of another sigma factor that can directly activate the *ssaG* and *srfN* promoters. Future work will be required to resolve these possibilities.

Our results indicate that *rpoE* deletion has no effect on SsrB levels under SPI-2 inducing conditions suggesting that the σ^E pathway regulates effector expression downstream of *ssrAB* transcription. Unlinking *ssrAB* transcription from the σ^E regulon would be advantageous to the cell to prevent commitment to a virulence gene expression program in response to envelope stress not associated with infection. The results from this study demonstrate that σ^E has the ability to affect expression of SsrB-regulated virulence genes and offers potential insight into the virulence attenuation of *rpoE* mutants. Although when considered individually, each promoter was modestly affected by deletion of *rpoE*, the cumulative effects of mild rewired inputs on multiple virulence promoters has been shown to severely compromise in-host fitness and virulence ability [25].

Conclusion

Based on these and other data [4,12-15], the genetic interaction between σ^E and a subset of SsrB-regulated genes may serve to coordinate the spatial and temporal activation of virulence genes in a host setting, likely in response to membrane damage resulting from oxidative antimicrobial systems and membrane-targeted host defence peptides.

Methods

Strains and Growth Conditions

Bacteria were propagated at 37°C with aeration in Luria-Bertani (LB) broth. *S. enterica* serovar Typhimurium (*S. Typhimurium*) strain 14028s with inactivating mutations in *rpoE*, *rpoS*, *rpoN* and *rpoH* were provided by Ferric Fang (University of Washington, Seattle, WA) [27]. $\Delta rpoH$ was grown at 30°C and $\Delta rpoN$ was supplemented with 2 mM L-glutamine. An unmarked, in-frame deletion of *rpoE* was




Figure 4
Model for σ^E -dependent regulation of the SsrB regulon. Membrane-targeting host defences including reactive oxygen stress and antimicrobial peptides cause an accumulation of unfolded outer membrane proteins (OMPs) and stimulate the cleavage of the anti-sigma factor RseA consequently releasing σ^E into the cytoplasm where it directs RNA polymerase to a subset of SPI-2 promoters. RpoE can positively or negatively regulate SsrB-regulated genes including integrated virulence genes unlinked with SPI-2 but has no effect on some effector genes such as *sseL*. This regulatory pathway may have evolved to coordinate virulence gene expression with host infection by responding to host-specific defence pathways that perturb the bacterial outer membrane.

made in *S. Typhimurium* strain SL1344 by λ Red recombination [28] using primers BKC187 and BKC188. Mutants were screened for loss of *rpoE* using primers BKC193 and BKC194. To generate an *ssrB::FLAG* allele in $\Delta rpoE$, the *ssrB::FLAG* allele from wild type SL1344 [19] was transduced into $\Delta rpoE$ by P22-mediated transduction. All plasmids and strains used in this work are described in Table 1. Primer sequences for mutant and plasmid construction are listed in Table 2.

Plasmid Construction

A complementation construct for *rpoE* was generated that included the three endogenous promoters (*rpoEp1*, *rpoEp2*, *rpoEp3*) identified in *Salmonella* [20] plus the wild type *rpoE* gene. This construct was cloned into the low-copy plasmid pWSK29 using primers SEO095 and SEO096 as a *Sall* and *XbaI* fragment. Constructs were verified by sequencing and transformed into *S. Typhimurium* SL1344 $\Delta rpoE$ and selected on LB agar with appropriate antibiotics. The promoters for *ssaB* (SEO005

and SEO006), *ssaG* (SEO011 and SEO012), *sifA* (SEO205 and SEO206), *sseL* (BKC185 and BKC186) and *srfN* (BKC183 and BKC184) were cloned into pIVET5n [29] to generate single-copy transcriptional fusions to *lacZ*. Reporters were transformed into *E. coli* SM10 λpir , conjugated into SL1344 and merodiploid cells were selected on LB agar with appropriate antibiotics. Transcriptional fusions, including a previously constructed reporter for the *sseA* promoter [30], were integrated into the chromosome of wild type and $\Delta rpoE$ cells using homologous recombination. The promoters we chose use the SsrB response regulator for expression of the downstream gene or operon, and include both SPI-2-encoded and non-SPI-2-encoded virulence effectors representing structural apparatus genes and effector substrates of the type III secretion system [8,30-35]

Chemiluminescent β -galactosidase Assay

Reporter strains were inoculated from an overnight culture into culture medium (LPM pH 5.8) that induces SsrB-

Table 1: Strains and plasmids used in this study

Strain or plasmid	Genotype or description ^a	Reference
Plasmids		
pKD46	repA101ts, oriR101, araC-P _{araB} -gam-bet-exo, bla	[28]
pCP20	FLP ⁺ , λ cI857 ⁺ λ p _R Rep ^{ts} , Amp ^R , Cm ^R	[28]
pKD3	pANTSy derivative, FRT-flanked cat from pSCI40, Cm ^R	[28]
prpoE	rpoE with endogenous promoters in pWSK29, Amp ^R	This work
psrfN-2HA	srfN with endogenous promoter and C-term. tandem HA fusion in pWSK129, Kan ^R	Our collection
psseL-2HA	sseL with endogenous promoter and C-term. tandem HA fusion in pWSK129, Kan ^R	[8]
psseL-2HA	sifA with endogenous promoter and internal tandem HA fusion in pACYC184	[37]
pIVET5n	tnpR-lacZ, sacB, R6K ori, bla, Amp ^R	[29]
pP _{sseA} -lacZ	sseA promoter fused to tnpR-lacZ in pIVET5n, Amp ^R	[29]
pP _{sseL} -lacZ	sseA promoter fused to tnpR-lacZ in pIVET5n, Amp ^R	This work
pP _{srfN} -lacZ	srfN promoter fused to tnpR-lacZ in pIVET5n, Amp ^R	This work
pP _{sifA} -lacZ	sifA promoter fused to tnpR-lacZ in pIVET5n, Amp ^R	This work
pP _{ssaB} -lacZ	ssaB promoter fused to tnpR-lacZ in pIVET5n, Amp ^R	This work
pP _{ssaG} -lacZ	ssaG promoter fused to tnpR-lacZ in pIVET5n, Amp ^R	This work
Strains		
SL1344	wild type <i>S. enterica</i> sv. Typhimurium, Sm ^R	[38]
14028s	wild type <i>S. enterica</i> sv. Typhimurium	[27]
14028s rpoE::cat	rpoE deletion, Cm ^R	[27]
140282 rpoS::Tn10dCm	rpoS deletion, Cm ^R	[27]
14028s rpoN::Ap	rpoN deletion, Amp ^R	[27]
14028s rpoH::PCN	rpoH deletion, temperature sensitive, Amp ^R	[27]
SL1344 ΔrpoE	Unmarked, in-frame deletion of rpoE, Sm ^R	This work
SL1344 ssrB::3FLAG	ssrB-FLAG replacement allele on chromosome	This work
SL1344 P _{sseA} ::pP _{sseA} -lacZ	Merodiploid containing integrated P _{sseA} -lacZ reporter	[25]
SL1344 P _{sseL} ::pP _{sseL} -lacZ	Merodiploid containing integrated P _{sseL} -lacZ reporter	This work
SL1344 P _{srfN} ::pP _{srfN} -lacZ	Merodiploid containing integrated P _{srfN} -lacZ reporter	This work
SL1344 P _{sifA} ::pP _{sifA} -lacZ	Merodiploid containing integrated P _{sifA} -lacZ reporter	This work
SL1344 P _{ssaB} ::pP _{ssaB} -lacZ	Merodiploid containing integrated P _{ssaB} -lacZ reporter	This work
SL1344 P _{ssaG} ::pP _{ssaG} -lacZ	Merodiploid containing integrated P _{ssaG} -lacZ reporter	This work
ΔrpoE P _{sseA} ::pP _{sseA} -lacZ	Merodiploid containing integrated P _{sseA} -lacZ reporter	This work
ΔrpoE P _{sseL} ::pP _{sseL} -lacZ	Merodiploid containing integrated P _{sseL} -lacZ reporter	This work
ΔrpoE P _{srfN} ::pP _{srfN} -lacZ	Merodiploid containing integrated P _{srfN} -lacZ reporter	This work
ΔrpoE P _{sifA} ::pP _{sifA} -lacZ	Merodiploid containing integrated P _{sifA} -lacZ reporter	This work
ΔrpoE P _{ssaB} ::pP _{ssaB} -lacZ	Merodiploid containing integrated P _{ssaB} -lacZ reporter	This work
ΔrpoE P _{ssaG} ::pP _{ssaG} -lacZ	Merodiploid containing integrated P _{ssaG} -lacZ reporter	This work

^aSm, streptomycin; Amp, ampicillin; Cm, chloramphenicol; Kan, kanamycin

Table 2: Primers used in this study

Primer	Sequence ^{ab}
BKC183	<u>GCTCGAG</u> TTTACCGCGCCGGATAACATACCG
BKC184	CACCAATT <u>G</u> TTAGCTGAATGAAGCAACCGTTGCCAG
BKC185	<u>CCTCGAG</u> CGATTGCCGTCAAAGGTATTC
BKC186	CACCAATT <u>G</u> TTACTGATCACCGTTCTCTACGGCGCT
BKC187	GGTTTGGGGAGACATTACCTCGGATGAGCGAGCAGTTAGTGTAGGCTGGAGCTGCTTCG
BKC188	CCTAATACCTTTTCCAGTATCCCGCTATCGTCAACGC CATGAATATCCTCCTTA
BKC195	CAGCTCACTATCCAACGTTT
BKC196	TGCTTGCTCATAGTGCGGCTT
BKC205	CACCAATT <u>G</u> TTACTATAGTAATCGGCATATTA
BKC206	CCG <u>CTCGAG</u> TATGGCGCTGATCGCCAC
SEO005	CCG <u>CTCGAG</u> GTGCCATCCTTTGCCGTTT
SEO006	CACCAATT <u>G</u> TTACATGAATCCCTCCTCAGACAT
SEO011	CCG <u>CTCGAG</u> ATATGGAGAGTGGTAGAATAG
SEO012	CACCAATT <u>G</u> TTATAATTGTGCAATATCCATAA
SEO095	GCTCTAGATCAACGCCTGATAAGCGGTT
SEO096	ACGCGT <u>CGAC</u> CCCACTTTATTGCGATTCCA

^aunderlined nucleotides indicate restriction enzyme sites; XhoI (CTCGAG), MfeI (CAATTG)

^bbolded nucleotides indicate complementarity to *rpoE*

dependent gene expression [21,36]. Cultures were propagated at 37°C for 7 hours and samples were taken hourly to measure β-galactosidase activity using a chemiluminescence assay described previously [25]. Data was expressed as relative light units (RLU) and was normalized to the optical density (OD_{600 nm}) of the parent culture.

Immunoblotting

To examine the protein levels of SseB, SseL, SrfN and SifA under SPI-2 inducing conditions, we used plasmids *psifA-2HA*, *psseL-2HA* and *psrfN-2HA* that were published previously (Table 1) [8,37]. These constructs express the given gene under the control of the endogenous promoter. Wild type and Δ*rpoE* cells were transformed with these plasmids and grown in LPM pH 5.8 at 37°C for 6 hours. Whole cell lysates were collected and analyzed by immunoblotting using anti-SseB (1:1000) [21] and anti-HA (1:1000, Covance) antibodies. Blots were probed for DnaK (1:3500, Stressgen) as a control.

Authors' contributions

SEO designed and performed research, interpreted data and wrote the paper. BKC designed and interpreted research and wrote the paper. Both authors read and approved the final manuscript.

Acknowledgements

We would like to thank Jose Puente for providing λ Red recombination plasmids, Ferric Fang for providing sigma factor mutants in the 14028s strain background, and members of the Coombes laboratory for helpful comments on this work. This work was funded by a grant to BKC from the Canadian Institute for Health Research (MOP-82704). SEO is the recipient of an Ontario Graduate Scholarship. BKC is the recipient of a New Investigator Award from the CIHR, a Young Investigator Award from the American Society of Microbiology, and an Early Researcher Award from the Ontario Ministry of Research and Innovation.

References

1. Shea JE, Hensel M, Gleeson C, Holden DW: **Identification of a virulence locus encoding a second type III secretion system in *Salmonella typhimurium*.** *Proc Natl Acad Sci USA* 1996, **93**:2593-2597.
2. Ochman H, Soncini FC, Solomon F, Groisman EA: **Identification of a pathogenicity island required for *Salmonella* survival in host cells.** *Proc Natl Acad Sci USA* 1996, **93**:7800-7804.
3. Cirillo DM, Valdivia RH, Monack DM, Falkow S: **Macrophage-dependent induction of the *Salmonella* pathogenicity island 2 type III secretion system and its role in intracellular survival.** *Mol Microbiol* 1998, **30**:175-188.
4. Hensel M: ***Salmonella* pathogenicity island 2.** *Mol Microbiol* 2000, **36**:1015-1023.
5. Hensel M, Shea JE, Waterman SR, Mundy R, Nikolaus T, Banks G, Vazquez-Torres A, Gleeson C, Fang FC, Holden DW: **Genes encoding putative effector proteins of the type III secretion system of *Salmonella* pathogenicity island 2 are required for bacterial virulence and proliferation in macrophages.** *Mol Microbiol* 1998, **30**:163-174.
6. Garmendia J, Beuzon CR, Ruiz-Albert J, Holden DW: **The roles of SsrA-SsrB and OmpR-EnvZ in the regulation of genes encoding the *Salmonella typhimurium* SPI-2 type III secretion system.** *Microbiology* 2003, **149**:2385-2396.

7. Worley MJ, Ching KH, Heffron F: **Salmonella SsrB activates a global regulon of horizontally acquired genes.** *Mol Microbiol* 2000, **36**:749-761.
8. Coombes BK, Lowden MJ, Bishop JL, Wickham ME, Brown NF, Duong N, Osborne S, Gal-Mor O, Finlay BB: **SseL is a Salmonella-specific translocated effector integrated into the SsrB-controlled salmonella pathogenicity island 2 type III secretion system.** *Infect Immun* 2007, **75**:574-580.
9. Osborne S, Walthers D, Tomljenovic AM, Mulder D, Silphaduang U, Duong N, Lowden M, Wickham ME, Waller R, Kenney LJ, et al.: **Pathogenic adaptation of intracellular bacteria by rewiring a cis-regulatory input function.** *Proc Natl Acad Sci USA* 2009 in press.
10. Browning DF, Busby SJ: **The regulation of bacterial transcription initiation.** *Nat Rev Microbiol* 2004, **2**:57-65.
11. Alba BM, Gross CA: **Regulation of the Escherichia coli sigma-dependent envelope stress response.** *Mol Microbiol* 2004, **52**:613-619.
12. Vazquez-Torres A, Xu Y, Jones-Carson J, Holden DW, Lucia SM, Din-aucr MC, Mastroeni P, Fang FC: **Salmonella pathogenicity island 2-dependent evasion of the phagocyte NADPH oxidase.** *Science* 2000, **287**:1655-1658.
13. Testerman TL, Vazquez-Torres A, Xu Y, Jones-Carson J, Libby SJ, Fang FC: **The alternative sigma factor sigmaE controls antioxidant defences required for Salmonella virulence and stationary-phase survival.** *Mol Microbiol* 2002, **43**:771-782.
14. Bader MW, Sanowar S, Daley ME, Schneider AR, Cho U, Xu W, Klevit RE, Le Moual H, Miller SI: **Recognition of antimicrobial peptides by a bacterial sensor kinase.** *Cell* 2005, **122**:461-472.
15. Crouch ML, Becker LA, Bang IS, Tanabe H, Ouellette AJ, Fang FC: **The alternative sigma factor sigma is required for resistance of Salmonella enterica serovar Typhimurium to anti-microbial peptides.** *Mol Microbiol* 2005, **56**:789-799.
16. Humphreys S, Stevenson A, Bacon A, Weinhardt AB, Roberts M: **The alternative sigma factor, sigmaE, is critically important for the virulence of Salmonella typhimurium.** *Infect Immun* 1999, **67**:1560-1568.
17. Eriksson S, Lucchini S, Thompson A, Rhen M, Hinton JC: **Unravelling the biology of macrophage infection by gene expression profiling of intracellular Salmonella enterica.** *Mol Microbiol* 2003, **47**:103-118.
18. Carlsson KE, Liu J, Edqvist PJ, Francis MS: **Extracytoplasmic-stress-responsive pathways modulate type III secretion in Yersinia pseudotuberculosis.** *Infect Immun* 2007, **75**:3913-3924.
19. Duong N, Osborne S, Bustamante VH, Tomljenovic AM, Puente JL, Coombes BK: **Thermosensing coordinates a cis-regulatory module for transcriptional activation of the intracellular virulence system in Salmonella enterica serovar Typhimurium.** *J Biol Chem* 2007, **282**:34077-34084.
20. Miticka H, Rowley G, Rezuchova B, Homerova D, Humphreys S, Farn J, Roberts M, Kormanec J: **Transcriptional analysis of the rpoE gene encoding extracytoplasmic stress response sigma factor sigmaE in Salmonella enterica serovar Typhimurium.** *FEMS Microbiol Lett* 2003, **226**:307-314.
21. Coombes BK, Brown NF, Valdez Y, Brumell JH, Finlay BB: **Expression and secretion of Salmonella pathogenicity island-2 virulence genes in response to acidification exhibit differential requirements of a functional type III secretion apparatus and SsaL.** *J Biol Chem* 2004, **279**:49804-49815.
22. Nitta T, Nagamitsu H, Murata M, Izu H, Yamada M: **Function of the sigma(E) regulon in dead-cell lysis in stationary-phase Escherichia coli.** *J Bacteriol* 2000, **182**:5231-5237.
23. Kabir MS, Yamashita D, Koyama S, Oshima T, Kurokawa K, Maeda M, Tsunedomi R, Murata M, Wada C, Mori H, et al.: **Cell lysis directed by sigmaE in early stationary phase and effect of induction of the rpoE gene on global gene expression in Escherichia coli.** *Microbiology* 2005, **151**:2721-2735.
24. Walthers D, Carroll RK, Navarre WW, Libby SJ, Fang FC, Kenney LJ: **The response regulator SsrB activates expression of diverse Salmonella pathogenicity island 2 promoters and counters silencing by the nucleoid-associated protein H-NS.** *Mol Microbiol* 2007, **65**:477-493.
25. Coombes BK, Wickham ME, Lowden MJ, Brown NF, Finlay BB: **Negative regulation of Salmonella pathogenicity island 2 is required for contextual control of virulence during typhoid.** *Proc Natl Acad Sci USA* 2005, **102**:17460-17465.
26. Rhodius VA, Suh WC, Nonaka G, West J, Gross CA: **Conserved and variable functions of the sigmaE stress response in related genomes.** *PLoS Biol* 2006, **4**:e2.
27. Bang IS, Frye JG, McClelland M, Velayudhan J, Fang FC: **Alternative sigma factor interactions in Salmonella: sigma and sigma promote antioxidant defences by enhancing sigma levels.** *Mol Microbiol* 2005, **56**:811-823.
28. Datsenko KA, Wanner BL: **One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products.** *Proc Natl Acad Sci USA* 2000, **97**:6640-6645.
29. Brown NF, Vallance BA, Coombes BK, Valdez Y, Coburn BA, Finlay BB: **Salmonella pathogenicity island 2 is expressed prior to penetrating the intestine.** *PLoS Pathog* 2005, **1**:e32.
30. Coombes BK, Brown NF, Kujat-Choy S, Vallance BA, Finlay BB: **SseA is required for translocation of Salmonella pathogenicity island-2 effectors into host cells.** *Microbes Infect* 2003, **5**:561-570.
31. Beuzon CR, Meresse S, Unsworth KE, Ruiz-Albert J, Garvis S, Waterman SR, Ryder TA, Boucrot E, Holden DW: **Salmonella maintains the integrity of its intracellular vacuole through the action of SifA.** *EMBO J* 2000, **19**:3235-3249.
32. Brumell JH, Tang P, Zaharik ML, Finlay BB: **Disruption of the Salmonella-containing vacuole leads to increased replication of Salmonella enterica serovar typhimurium in the cytosol of epithelial cells.** *Infect Immun* 2002, **70**:3264-3270.
33. Ruiz-Albert J, Mundy R, Yu XJ, Beuzon CR, Holden DW: **SseA is a chaperone for the SseB and SseD translocon components of the Salmonella pathogenicity-island-2-encoded type III secretion system.** *Microbiology* 2003, **149**:1103-1111.
34. Zurawski DV, Stein MA: **SseA acts as the chaperone for the SseB component of the Salmonella Pathogenicity Island 2 translocon.** *Mol Microbiol* 2003, **47**:1341-1351.
35. Rytkonen A, Poh J, Garmendia J, Boyle C, Thompson A, Liu M, Freemont P, Hinton JC, Holden DW: **SseL, a Salmonella deubiquitinase required for macrophage killing and virulence.** *Proc Natl Acad Sci USA* 2007, **104**:3502-3507.
36. Deiwick J, Nikolaus T, Erdogan S, Hensel M: **Environmental regulation of Salmonella pathogenicity island 2 gene expression.** *Mol Microbiol* 1999, **31**:1759-1773.
37. Brumell JH, Goosney DL, Finlay BB: **SifA, a type III secreted effector of Salmonella typhimurium, directs Salmonella-induced filament (Sif) formation along microtubules.** *Traffic* 2002, **3**:407-415.
38. Wray C, Sojka WJ: **Experimental Salmonella typhimurium infection in calves.** *Res Vet Sci* 1978, **25**:139-143.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

