

SCIENTIFIC REPORTS



OPEN

Cyanamide is biosynthesized from L-canavanine in plants

Tsunashi Kamo¹, Sakae Sakurai², Tatsuya Yamanashi³ & Yasushi Todoroki^{2,4}

Received: 09 January 2015

Accepted: 17 April 2015

Published: 27 May 2015

Cyanamide had long been recognized as a synthetic compound but more recently has been found as a natural product from several leguminous plants. This compound's biosynthetic pathway, as yet unelaborated, has attracted attention because of its utility in many domains, such as agriculture, chemistry, and medicine. We noticed that the distribution of L-canavanine in the plant kingdom appeared to include that of cyanamide and that the guanidino group structure in L-canavanine contained the cyanamide skeleton. Here, quantification of these compounds in *Vicia* species suggested that cyanamide was biosynthesized from L-canavanine. Subsequent experiments involving L-[guanidineimino-¹⁵N₂]canavanine addition to young *Vicia villosa* seedlings resulted in significant incorporation of ¹⁵N-label into cyanamide, verifying its presumed biosynthetic pathway.

Cyanamide (NH₂CN) was first synthesized from ammonia and cyanogen chloride by Cannizzaro and Cloëz in 1851¹. The scheme for its mass production as a calcium salt was established in Germany in the late 19th century. Cyanamide is multifunctional for agricultural purposes because it serves in the soil as an insecticide, fungicide, and herbicide for a time after application, then decomposes to urea in the soil, and finally is absorbed by crops as fertilizer. Maier-Greiner *et al.* isolated and characterized cyanamide hydratase from a soil fungus *Myrothecium verrucaria*^{2,3}. The high specificity of this enzyme was surprising because the substrate had not been previously found in nature³. In 2003, natural cyanamide was first isolated from hairy vetch, *Vicia villosa*⁴. The distribution of this compound in the plant kingdom seems highly limited, having only been detected in four leguminous species among more than 550 species tested^{5,6}. Several studies have suggested that cyanamide in plants might function in the context of chemical ecology; for example, livestock feeding on vetch-dominant meadows often die of vetch-disease^{7–9}, probably due to cyanamide toxicity^{6,10}.

L-Canavanine is the guanidinoxy structural analogue of L-arginine. When assimilated, it is used for peptide extension in place of L-arginine, resulting in protein malfunction^{11,12}, which explains its toxicity to a wide range of organisms¹³. A variety of leguminous species accumulate this toxic compound in their seeds for protection from herbivores as well as nitrogen storage^{14,15}. L-Canavanine distribution in the plant kingdom is broader than cyanamide; it has been found in more than 1500 members of the Faboideae, a subfamily of the Fabaceae^{16,17}.

L-Canavanine's biosynthetic pathway has already been established and described in jack bean to involve the conversion of L-canavaninosuccinic acid to L-canavanine and fumaric acid¹⁸. In contrast, cyanamide biosynthesis has remained totally undescribed. Although there have been no reports that indicate a biosynthetic relationship between L-canavanine and cyanamide, we focused here on the following information: First, canavanine has been detected in many *Vicia* species¹⁶, some of which also contain cyanamide^{5,6}. Second, L-canavanine's chemical structure consists of cyanamide and another amino acid, L-canaline, and it is possible to actually synthesize L-canavanine from them in the presence of zinc cations¹⁹. And third, L-canavanine is stored in seeds, whereas cyanamide starts to accumulate after germination⁶. Thus, it is conceivable that in plants L-canavanine is the biosynthetic precursor of cyanamide. In the present study, we conducted several experiments to examine the validity of this hypothesis.

¹Biodiversity Division, National Institute for Agro-Environmental Sciences, 3-1-3 Kan-nondai, Tsukuba, 305-8604 Ibaraki, Japan. ²Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, 422-8529 Shizuoka, Japan. ³Faculty of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, 422-8529 Shizuoka, Japan. ⁴Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, 422-8529 Shizuoka, Japan. Correspondence and requests for materials should be addressed to T.K. (email: tkamo@affrc.go.jp)




Figure 1. Concentration of cyanamide and L-canavanine in the leaves of leguminous plants. Tested species with sampling day after sowing in parentheses; 1, *Lens culinaris* (11 d); 2, *Robinia pseudoacacia* (22 d); 3, *Vicia angustifolia* (11 d); 4, *V. amoena* (18 d); 5, *V. benghalensis* (10 d); 6, *V. cracca* (14 d); 7, *V. hirsuta* (11 d); 8, *V. pseudo-orobus* (22 d); 9, *V. tetrasperma* (13 d); 10, *V. unijuga* (18 d); 11, *V. villosa* subsp. *villosa* (10 d); and 12, *V. villosa* subsp. *varia* (8 d). First, second and third leaves sampled when third leaf developed; black and grey bars, cyanamide and L-canavanine, respectively; and error bars, standard deviation ($n = 4$).




Figure 2. Changes in cyanamide and L-canavanine content in *V. villosa* subsp. *varia* seedlings. Black and grey circles, cyanamide and L-canavanine, respectively; arrow, germination; and error bars, standard error ($n = 6$).

Results and Discussion

We quantified cyanamide and L-canavanine in the leaves of leguminous plants (Fig. 1). In addition to *Vicia* species, two other species were also analyzed; *Robinia pseudoacacia* as a lone species so far tested that contains cyanamide other than *Vicia* species⁵, and *Lens culinaris* for its placement, by recent molecular phylogeny studies, within a monophyletic group of the genus *Vicia*^{20,21}. Notably, L-canavanine has been detected in all species that contain cyanamide, supporting the idea that there is a link between cyanamide biosynthesis and L-canavanine.

Figure 2 presents the changes in cyanamide and L-canavanine content in young seedlings of *V. villosa* subsp. *varia*. Cyanamide was absent in the seed and began to accumulate when the first leaf appeared at 5–6 day after sowing. In contrast, L-canavanine was abundant in the seed and started to decrease after leaves developed. This synchrony was thought-provoking, particularly with the intriguing observation that the total molar content of the two compounds per individual plant remained nearly constant throughout the monitored period. This suggested that L-canavanine was a cyanamide biosynthetic precursor. It is unclear whether L-canavanine is biosynthesized at the early stage of development.

The plant organ in which cyanamide was biosynthesized was investigated by quantifying the two compounds in each part of the seedling: cotyledon, root, epicotyl, and leaf (Table S1). Since *V. villosa* is a plant with hypogeal germination, the cotyledons stay below the ground. More than half of the total L-canavanine was present in the cotyledons, which was a reasonable localization because it is contained in the seeds (Fig. 2). Nearly 40% of the total L-canavanine, probably transferred from the cotyledons, was present in the epicotyl and leaves. In contrast, almost all the cyanamide was detected in the epicotyl and leaves. Considering that this compound started to accumulate when leaves emerged, cyanamide appeared to be biosynthesized in leaves. Thus, it was highly likely that L-canavanine stored in the seed was transported to the seedling shoot after germination and then converted to cyanamide in leaves.

We confirmed the conversion of L-canavanine to cyanamide by administration of ¹⁵N-labelled compounds to 4-day-old shoots of *V. villosa* subsp. *varia*. After 48 h of incubation, cyanamide was isolated from the leaves and analyzed by gas chromatography-mass spectrometer (GC-MS). Compared with

Treatment	Relative area of the isotopic [M] ⁺ ions at <i>m/z</i> 42 to 45 (%) [*]			
	42	43	44	45
(Theoretical value)	100.0	1.9	0.0	0.0
Control	100.0	1.9	0.1	0.0
[¹⁵ N ₂]Ammonium nitrate	100.0	1.7	0.1	0.0
[¹⁵ N ₂]Urea	100.0	1.8	0.1	0.0
L-[Guanidineimino- ¹⁵ N ₂]arginine	100.0	1.9	0.1	0.0
L-[Guanidineimino- ¹⁵ N ₂]canavanine	100.0	15.1	98.4	0.8

Table 1. Incorporation of ¹⁵N-labels into cyanamide in shoots of 4-day-old seedlings of *V. villosa* subsp. *varia*. Labelled compounds administered at 2.0 mM for 48 h to *V. villosa* shoot (leaves plus epicotyl) excised from roots and cotyledons. ^{*}Cyanamide isolated from leaves analyzed by GC-MS after preparative HPLC.




Figure 3. Cyanamide biosynthesis from L-canavanine.

the control, the isotopic molecular ion peak at *m/z* 44 (the [M + 2]⁺ ion) was clearly enhanced when L-[guanidineimino-¹⁵N₂]canavanine was administered (Table 1; Figures S1 and S2). This indicated that this substrate was converted to [¹⁵N₂]cyanamide in shoots (Fig. 3). Administration of [¹⁵N₂]ammonium nitrate and [¹⁵N₂]urea showed no effect on the abundance of key isotopic molecular ion peaks, negating the possibility that L-canavanine degradative products, such as urea or ammonia, were used for cyanamide biosynthesis. ¹⁵N-Label from L-[guanidineimino-¹⁵N₂]arginine was also not incorporated into cyanamide, which illustrated a clear difference in incorporation between L-canavanine and L-arginine and implied high substrate specificity by the enzyme responsible for cyanamide production.

Plant arginase can convert L-canavanine into L-canaline and urea, but L-arginine is a better substrate for this enzyme (Figure S3)^{22,23}. In addition to this conversion, L-canavanine is also decomposed to smaller molecules or coupled with other compounds by enzymatic reactions in which L-arginine is the favored primary substrate²⁴. In contrast, some canavanine-resistant microorganisms benefit from mechanisms that specifically function on this toxic amino acid; it is cleaved to L-homoserine and guanidine in *Streptococcus faecalis* and *S. equinus* and hydrolyzed to L-homoserine and hydroxyguanidine in *Pseudomonas* sp. (Figure S4)^{24,25}. The former route reportedly also functions in canavanine-resistant insects, such as the tobacco budworm *Heliothis virescens*²⁶. Our results here indicated that a novel enzymatic activity specific for L-canavanine was present in some *Vicia* species. Although the amino acid formed in this cleavage is yet to be clarified, the conversion from L-canavanine to cyanamide is reasonably explainable by a general acid-base catalysis mechanism (Figure S5).

In conclusion, we demonstrated that cyanamide was biosynthesized from L-canavanine in plants. As cyanamide has a short history since it was first isolated from a natural source⁴, the biosynthesis and metabolism of natural cyanamide has not been fully explained. This finding will serve, in future studies, as a crucial step for the isolation and characterization of the responsible enzyme.

Methods

General. GC-MS was performed on a QP5050A system (Shimadzu Corp., Kyoto, Japan) using an Equity-5 column (0.25 mm i.d. × 30 m, 0.25 μm film thickness; Supelco, Inc., Bellefonte, PA, USA). The HPLC consisted of a 626 pump with 996 photodiode array detector (Waters Corp., Milford, MA) equipped with a C-18 column (Inertsil ODS-3, 5 μm, 250 × 4.6 mm ID; GL Sciences, Inc., Tokyo, Japan). ¹H and ¹³C NMR spectra were recorded with tetramethylsilane as the internal standard using JNM-EX270 (270 MHz) and JNM-LA500 (500 MHz) NMR spectrometers (JEOL Ltd., Tokyo, Japan). High resolution mass spectra were obtained with a JMS-T100LC AccuTOF mass spectrometer (JEOL). The seeds used in the present study have been previously described⁶. They were planted at a depth of 5 mm in pots (8 × 8 × 6 cm deep) containing sand and then incubated in an illuminated growth chamber (FLI-301N; Tokyo, Rikakikai, Co., Ltd., Tokyo, Japan) under a 16-h light/8-h dark cycle at 22 °C.

¹⁵N-Labelled chemicals. [¹⁵N₂]Urea (98+ atom% ¹⁵N) was purchased from Sigma-Aldrich Corp. (Milwaukee, WI, USA). L-[Guanidineimino-¹⁵N₂]arginine hydrochloride (min 98 atom% ¹⁵N), [¹⁵N₂]

ammonium nitrate (min 98+ atom% $^{15}\text{NH}_4$, min 98+ atom% $^{15}\text{NO}_3$) and [^{13}C , $^{15}\text{N}_2$]cyanamide (99 atom% ^{13}C , 98 atom% ^{15}N , 50wt% aqueous solution) were purchased from Isotec (Miamisburg, OH, USA). L-[Guanidineimino- $^{15}\text{N}_2$]canavanine was synthesized from [$^{15}\text{N}_2$]urea (Figure S6).

Quantification of cyanamide and L-canavanine in leguminous species. Fresh plant materials (20–50 mg fresh weight) were extracted with 1 mL ethanol, according to a published procedure⁶, and the extracts used for quantification of cyanamide and L-canavanine. The leaves obtained from a single seedling was used for one lot of the extraction. For cyanamide quantification by GC-MS, 50 μL of [^{13}C , $^{15}\text{N}_2$] cyanamide solution (10 $\mu\text{g}/\text{mL}$ in acetonitrile) was added as internal standard to the ethanol extract (50 μL ; GC-MS conditions⁶). For L-canavanine quantification by HPLC, the ethanol extract (160 μL) was mixed with 20 μL triethylamine and 20 μL phenyl isothiocyanate and then placed at room temperature for 20 min before concentrating the solution to dryness in vacuo^{27,28}. The residue was dissolved in 100 μL ethanol, and a 10- μL aliquot injected into the HPLC (HPLC conditions²⁹). For cyanamide and L-canavanine quantification in 12 leguminous species (Fig. 1), leaves were sampled when the third leaf developed. The species used are listed here with the sampling day after sowing in parentheses: *Lens culinaris* (11 d), *Robinia pseudoacacia* (22 d), *Vicia angustifolia* (11 d), *V. amoena* (18 d), *V. benghalensis* (10 d), *V. cracca* (14 d), *V. hirsuta* (11 d), *V. pseudo-orobus* (22 d), *V. tetrasperma* (13 d), *V. unijuga* (18 d), *V. villosa* subsp. *villosa* (10 d), and *V. villosa* subsp. *varia* (8 d).

Synthesis of L-[guanidineimino- $^{15}\text{N}_2$]canavanine. L-Canaline was prepared using a modified previously-described method³⁰. A solution of benzyl L-2-[(carbobenzyloxy)amino]-4-(benzamidoxy) butanoate (1.26 g, 2.72 mmol) in 4 M HCl (30 mL) was refluxed for 5 h and the solvent removed in vacuo. The residue was dissolved in ethanol (1.5 mL), precipitated with diethyl ether (6 mL) and filtered. The resultant white cake was washed with diethyl ether (2 \times 6 mL) and dissolved in ethanol (1.5 mL). Triethylamine (10 mL) was added to produce a white precipitate, which was concentrated by solvent removal in vacuo. The residue was dissolved in water (15 mL), charcoal added, and the mixture stirred for 10 min. Next, it was filtered and concentrated in vacuo to yield a white solid (0.99 g), which was then washed with ethanol (2 \times 10 mL) to remove the triethylamine hydrochloride. The residual oil containing a white solid was purified by silica gel column chromatography with ethanol/water stepwise to obtain L-canaline (177 mg, 1.32 mmol, 49% yield) as a colorless oil. ^1H NMR (270 MHz, D_2O): δ 3.69 (m, 3H), 2.00 (m, 2H); HRMS (m/z): [M-H]⁻ calcd. for $\text{C}_4\text{H}_9\text{N}_2\text{O}_3$, 133.0613; found, 133.0609 (ESI-TOF, negative mode).

L-[Guanidineimino- $^{15}\text{N}_2$]canavanine was prepared from L-canaline and [$^{15}\text{N}_2$]-O-methylisourea methylsulfate by a modified previously-described method^{31,32}. [$^{15}\text{N}_2$]-O-Methylisourea methylsulfate was obtained from [$^{15}\text{N}_2$]urea using the same method as for preparation of the non-labelled compound³³. Cupric oxide (239 mg) was added to a stirred solution of L-canaline (177 mg, 1.32 mmol) in water (2 mL) and the solution refluxed for 10 min to obtain a copper/L-canaline complex. The resultant deep blue solution was next stirred at 60 °C for 2 h and at room temperature for 4 d. After filtration, the filtrate was concentrated to a small volume (~2 mL) in vacuo and then [$^{15}\text{N}_2$]-O-methylisourea methylsulfate (440 mg, 2.30 mmol) dissolved in water (1.2 mL) added at 0 °C. After addition of 2 M NaOH aq. (1.2 mL), the solution was stirred for 16 d at room temperature, acidified (pH ~1) with 1 M HCl (3.6 mL) to pale blue, and treated with hydrogen sulfide. After filtration to remove the resulting black precipitate (CuS), the filtrate was heated at 90 °C for 5 h to remove hydrogen sulfide and concentrated in vacuo to produce a crude oil containing a pale yellow solid (603 mg). Flavianic acid dehydrate (460 mg, 1.31 mmol) in water (0.6 mL) was added to the crude product in water (3 mL) and held at 5 °C for 17 h. After suction filtration, the resulting yellow crystals were collected and dissolved in water (10 mL) at ~90 °C. Barium hydroxide octahydrate (310 mg) dissolved in hot water (10 mL) was added to the solution, which was held at 5 °C for 16 h and then filtered. The filtrate was next treated with 5% sulfuric acid (0.6 mL) and filtered through filter paper to remove the white precipitate. The yellow filtrate was then treated with charcoal for 30 min, filtered, and concentrated in vacuo. The residual colorless oil (48 mg) was purified by strong cation exchange resin column chromatography (PoraPak Rxn CX, Waters Corp., Milford, MA, USA) with 5% 3 M ammonia solution in methanol. The eluate was concentrated in vacuo and recrystallized in ethanol/water (25/1, v/v) to obtain L-[guanidineimino- $^{15}\text{N}_2$]canavanine (2.0 mg, 11 μmol , 0.9% yield). ^1H NMR (500 MHz, D_2O): δ 3.83 (m, 2H), 3.73 (t, J = 5.8 Hz, 1H), 2.13 (m, 1H), 2.06 (m, 1H); ^{13}C NMR (125 MHz, D_2O): δ 176.2, 159.8, 70.9, 54.4, 31.2; HRMS (m/z): [M + H]⁺ calcd. for $\text{C}_5\text{H}_{13}\text{N}_2^{15}\text{N}_2\text{O}_3$, 179.0928; found, 179.0923 (ESI-TOF, positive mode).

Administration of ^{15}N -labelled compounds to *Vicia villosa* subsp. *varia* seedlings. A 4-day-old *V. villosa* subsp. *varia* shoot (leaves plus epicotyl) excised from the roots and cotyledons was inserted into each tube containing 0.8 mL of a solution of ^{15}N -labelled compound (2.0 mM). The shoots were next incubated for 48 h in the illuminated growth chamber under a 16-h light/8-h dark cycle at 22 °C, with the solution replaced at 24 h. The leaves (~30 mg fresh weight) were used for cyanamide analysis (for the extraction procedure, see the literature⁶). Out of 900 μL of ethanol extract, a 300 μL aliquot was used for purification. After the addition of 60 μL of water, the ethanol extract was concentrated in vacuo and 20 μL of the resulting water solution (ca. 30 μL) then purified by HPLC to yield a cyanamide-containing

fraction (t_R 3.3–3.9 min), which was then concentrated to dryness and dissolved in 50 μ L ethyl acetate. A portion of the sample solution was finally subjected to GC-MS, using previously described conditions⁶, except the splitless injection mode was used with a 2 min sampling time at 200 kPa, and the sample injection volume was at 2.5 μ L.

References

- Cannizzaro, S. & Cloëz, S. Recherche sur les amides cyaniques. *Compt. Rendus* **32**, 62–64 (1851).
- Maier-Greiner, U. H., Klaus, C. B. A., Estermaier, L. M. & Hartmann, G. R. Herbicide resistance in transgenic plants through degradation of the phytotoxin to urea. *Angew. Chem. Int. Ed.* **30**, 1314–1315, 10.1002/anie.199113141 (1991).
- Maier-Greiner, U. H. *et al.* Isolation and properties of a nitrile hydratase from the soil fungus *Myrothecium verrucaria* that is highly specific for the fertilizer cyanamide and cloning of its gene. *Proc. Natl. Acad. Sci. USA* **88**, 4260–4264, 10.1073/pnas.88.10.4260 (1991).
- Kamo, T., Hiradate, S. & Fujii, Y. First isolation of natural cyanamide as a possible allelochemical from hairy vetch *Vicia villosa*. *J. Chem. Ecol.* **29**, 275–283, 10.1023/A:1022621709486 (2003).
- Kamo, T. *et al.* Limited distribution of natural cyanamide in higher plants: occurrence in *Vicia villosa* subsp. *varia*, *V. cracca*, and *Robinia pseudo-acacia*. *Phytochemistry* **69**, 1166–1172, 10.1016/j.phytochem.2007.11.004 (2008).
- Kamo, T., Takemura, T., Wasano, N., Fujii, Y. & Hiradate, S. Quantification of cyanamide in young seedlings of *Vicia* species, *Lens culinaris*, and *Robinia pseudo-acacia* by gas chromatography-mass spectrometry. *Biosci. Biotechnol. Biochem.* **76**, 1416–1418, 10.1271/bbb.120262 (2012).
- Odriozola, E., Paloma, E., Lopez, T. & Campero, C. An outbreak of *Vicia villosa* (hairy vetch) poisoning in grazing Aberdeen Angus bulls in Argentina. *Vet. Hum. Toxicol.* **33**, 278–280 (1991).
- Pancieria, R. J., Johnson, L. & Osburn, B. I. A disease of cattle grazing hairy vetch pasture. *J. Am. Vet. Med. Assoc.* **148**, 804–808 (1966).
- Peet, R. L. & Cardner, J. J. Poisoning of cattle by hairy or woolly-pod vetch, *Vicia villosa* subspecies *dasycarpa*. *Aust. Vet. J.* **63**, 381–382, 10.1111/j.1751-0813.1986.tb02910.x (1986).
- Onda, K. *et al.* Contact dermatitis in dairy cattle caused by calcium cyanamide. *Vet. Rec.* **163**, 418–422, 10.1136/vr.163.14.418 (2008).
- Rosenthal, G. A. L-Canavanine: a higher plant insecticidal allelochemical. *Amino Acids* **21**, 319–330, 10.1007/s007260170017 (2001).
- Igloi, G. L. & Schiefermayr, E. Amino acid discrimination by arginyl-tRNA synthetases as revealed by an examination of natural specificity variants. *FEBS J.* **276**, 1307–1318, 10.1111/j.1742-4658.2009.06866.x (2009).
- Rosenthal, G. A. The biological effects and mode of action of L-canavanine, a structural analogue of L-arginine. *Q. Rev. Biol.* **52**, 155–178 (1977).
- Rosenthal, G. A. & Dahlman, D. L. L-Canavanine and protein synthesis in the tobacco hornworm *Manduca sexta*. *Proc. Natl. Acad. Sci. USA* **83**, 14–18 (1986).
- Rosenthal, G. A. Nitrogen allocation for L-canavanine synthesis and its relationship to chemical defense of the seed. *Biochem. Sys. Ecol.* **5**, 219–220 (1977).
- Bell, E. A., Lackey, J. A. & Polhill, R. M. Systematic significance of canavanine in the Papilionoideae (Faboideae). *Biochem. Sys. Ecol.* **6**, 201–212, 10.1016/0305-1978(78)90008-X (1978).
- Rosenthal, G. A. Metabolism of L-canavanine and L-canaline in leguminous plants. *Plant Physiol.* **94**, 1–3 (1990).
- Rosenthal, G. A. L-Canavanine metabolism in jack bean, *Canavalia ensiformis* (L.) DC. (Leguminosae). *Plant Physiol.* **69**, 1066–1069 (1982).
- Ozinskas, A. J. & Rosenthal, G. A. L-Canavanine synthesis by zinc-mediated guanidination of L-canaline with cyanamide. *Bioorg. Chem.* **14**, 157–162, 10.1016/0045-2068(86)90025-8 (1986).
- Endo, Y. *et al.* Molecular phylogeny of *Vicia* sect. *Amurensis* (Leguminosae). *J. Jpn. Bot.* **85**, 337–349 (2010).
- Schaefer, H. *et al.* Systematics, biogeography, and character evolution of the legume tribe Fabaeae with special focus on the middle-Atlantic island lineages. *BMC Evol. Biol.* **12**, 250–268, 10.1186/1471-2148-12-250 (2012).
- Kitagawa, M. & Tomiyama, T. A new amino-compound in the jack bean and a corresponding new ferment. (I). *J. Biochem.* **11**, 265–271 (1929).
- Damodaran, M. & Karayanan, K. G. A. A comparative study of arginase and canavanase. *Biochem. J.* **34**, 1449–1459 (1940).
- Kalyankar, G. D., Ikawa, M. & Snell, E. E. The enzymatic cleavage of canavanine to homoserine and hydroxyguanidine. *J. Biol. Chem.* **233**, 1175–1178 (1958).
- Kihara, H., Prescott, J. M. & Snell, E. E. The bacterial cleavage of canavanine to homoserine and guanidine. *J. Biol. Chem.* **217**, 497–504 (1955).
- Berge, M. A., Rosenthal, G. A. & Dahlman, D. L. Tobacco budworm, *Heliothis virescens* [Noctuidae] resistance to L-canavanine, a protective allelochemical. *Pestic. Biochem. Physiol.* **25**, 319–326 (1986).
- Bidlingmeyer, B. A., Cohen, S. A. & Tarvin, T. L. Rapid analysis of amino acids using pre-column derivatization. *J. Chromatogr. B* **336**, 93–104, 10.1016/S0378-4347(00)85133-6 (1984).
- Frye, E. B., Degenhardt, T. P., Thorpe, S. R. & Baynes, J. W. Role of the Maillard reaction in aging of tissue proteins: Advanced glycation end product-dependent increase in imidazolium cross-links in human lens proteins. *J. Biol. Chem.* **273**, 18714–18719, 10.1074/jbc.273.30.18714 (1998).
- Kamo, T., Tokuoaka, Y. & Miyazaki, M. Quantification of canavanine, 2-aminoethanol, and cyanamide in *Aphis craccivora* and its host plants, *Robinia pseudoacacia* and *Vicia angustifolia*: effects of these compounds on larval survivorship of *Harmonia axyridis*. *J. Chem. Ecol.* **38**, 1552–1560, 10.1007/s10886-012-0220-9 (2012).
- Ozinskas, A. J. & Rosenthal, G. A. Synthesis of L-canaline and γ -functional 2-aminobutyric acid derivatives. *J. Org. Chem.* **51**, 5047–5050, 10.1021/jo00376a001 (1986).
- Crooks, P. A., Rosenthal, G. A., inventors; University of Kentucky Research Foundation, assignee. Canavanine analogs and their use as chemotherapeutic agents. United States patent US 5, 859,295. 1999 Jan 12.
- Nyberg, D. D. & Christensen, B. E. The synthesis of DL-canaline, DL-canavanine and related compounds. *J. Am. Chem. Soc.* **79**, 1222–1226, 10.1021/ja01562a050 (1957).
- Greindl, T., Scherr, G., Schneider, R., Mundinger, K., inventors; Basf Ag, assignee. Verfahren zur Herstellung von substituierten Guanidinderivaten. German patent DE 19,748,695. 1999 May 6.

Author Contributions

T.K. conceived, designed and performed the research, analyzed the data, and wrote the manuscript. S.S., T.Y. and Y.T. synthesized the ^{15}N -labelled canavanine. All authors have approved the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Kamo, T. *et al.* Cyanamide is biosynthesized from L-canavanine in plants. *Sci. Rep.* 5, 10527; doi: 10.1038/srep10527 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>