

Review Article

Halomonas as a chassis

Jian-Wen Ye¹ and  Guo-Qiang Chen^{2,3,4,5}

¹School of Biology and Biological Engineering, South China University of Technology, Guangzhou 510006, China; ²School of Life Sciences, Tsinghua University, Beijing 100084, China; ³Center for Synthetic and Systems Biology, Tsinghua University, Beijing 100084, China; ⁴MOE Key Laboratory for Industrial Biocatalysts, Dept Chemical Engineering, Tsinghua University, Beijing 100084, China; ⁵Tsinghua-Peking Center for Life Sciences, Tsinghua University, Beijing 100084, China

Correspondence: Guo-Qiang Chen (chengq@mail.tsinghua.edu.cn)



With the rapid development of systems and synthetic biology, the non-model bacteria, *Halomonas* spp., have been developed recently to become a cost-competitive platform for producing a variety of products including polyesters, chemicals and proteins owing to their contamination resistance and ability of high cell density growth at alkaline pH and high salt concentration. These salt-loving microbes can partially solve the challenges of current industrial biotechnology (CIB) which requires high energy-consuming sterilization to prevent contamination as CIB is based on traditional chassis, typically, *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas putida* and *Corynebacterium glutamicum*. The advantages and current status of *Halomonas* spp. including their molecular biology and metabolic engineering approaches as well as their applications are reviewed here. Moreover, a systematic strain engineering streamline, including product-based host development, genetic parts mining, static and dynamic optimization of modularized pathways and bioprocess-inspired cell engineering are summarized. All of these developments result in the term called next-generation industrial biotechnology (NGIB). Increasing efforts are made to develop their versatile cell factories powered by synthetic biology to demonstrate a new biomanufacturing strategy under open and continuous processes with significant cost-reduction on process complexity, energy, substrates and fresh water consumption.

Introduction

Due to the sustainable and environment-friendly advantages of industrial biotechnology [1–4], it has been developed for decades to produce numerous bio-based products such as bioplastics [5–7], medicines [8], chemicals, food additives [9,10] and biofuels [11–13], with an aim to gradually replace the traditional petrol-based industry. However, the long-standing challenges of contamination risk, high sterile energy input and high fresh water consumption involved in bioprocesses of current industrial biotechnology (CIB) still reduce its competitiveness in spite of the rapid development of synthetic biology [14,15].

To make industrial biotechnology competitive to chemical industry, suitable microorganisms should be constructed to solve the existing difficulties of CIB [16]. Recently, the salt-tolerating bacteria, *Halomonas* spp., are becoming the attractive candidate hosts for microbial cell factory engineering due to their strong metabolism of diverse substrates and fast growth under high salt and high pH conditions, making possible the contamination free, non-food raw materials- and seawater-consuming fermentation processes [17,18]. In contrast with the traditional model microorganisms, such as *Escherichia coli* [14,19], *Bacillus subtilis* [20], *Pseudomonas putida* [21] and others [22], the development of molecular manipulation tools and methods for *Halomonas* spp. is difficult and less effective [23]. Recent attempts have been successful to establish genetic engineering approaches [24,25] including genome editing and gene expression control [26] to construct *Halomonas* spp. into a diverse bioproduct producers.

It is important to note that, *Halomonas bluephagenesis*, one of the well-studied salt-loving wildtype microbe able to accumulate poly-3-hydroxybutyrate (PHB), has been reprogrammed to be a low-cost chassis for various biosyntheses conducted under seawater-based unsterile open fermentation [27]. This *Halomonas*-based biotechnology, namely, the ‘next-generation industrial biotechnology’ (NGIB), has

Received: 10 February 2021
Revised: 24 March 2021
Accepted: 26 March 2021

Version of Record published:
26 July 2021

exemplified successful cases in pilot-scale polyhydroxyalkanoate (PHA) productions with significant cost reduction [28], possessing promising advantages that CIB cannot have. Besides, several predominant *Halomonas* strains of industrial potential, including *Halomonas smyrnensis* AAD6 [29–31], *Halomonas* sp. KM-1 [32–34] and *Halomonas* sp. HAL1 [35], are recently developed for producing different metabolic targets, demonstrating *Halomonas* as powerful chassis in biomanufacturing based on NGIB.

The state-of-the-art developments of *Halomonas* spp. used as microbial production chassis are reviewed. Meanwhile, future perspectives and critical comments involving upstream strain engineering and downstream process optimization have been discussed in details aiming to provide constructive thoughts and possible solutions for further improvements of NGIB based on the current achievements.

Properties of *Halomonas* spp.

Halomonas spp. are Gram-negative bacteria belonging to the family of Halophiles that prefer to grow in saline environments (commonly referred to NaCl or KCl), such as salt lakes and marshes, oceans or other saline areas on earth [36]. According to their preference of salt concentration under optimal growth, *Halomonas* spp. are commonly divided into two subtypes, moderate (3–15% NaCl w/v) and extreme (>20% NaCl w/v) *Halomonas* [18,36]. Most *Halomonas* spp. can survive in a wide range of temperature, reaching up to 50°C, and at alkaline conditions with a pH value over 10 [37]. Due to the fast-growth at such extreme environments, *Halomonas* spp. are grown with less contamination risks compared with non-halophilic microorganisms. *Halomonas* spp. are becoming favored hosts for developing contamination-free fermentation processes without strict sterilization [16,27].

Halomonas are reported to possess two typical osmotic (salt-resistant) regulatory mechanisms enabling flourished growth under saline conditions: (I) accumulation of inorganic ions, such as K⁺, so as to balance the extracellular osmotic pressure of NaCl [38]; (II) production of water soluble and compatible solutes, also termed osmolytes, including ectoine, hydroxyectoine, betaine and several amino acids such as glycine, valine and proline to form an intracellular barrier resisting the influx of NaCl from saline environments [39]. For most *Halomonas* spp., mechanism-II is the preferred strategy to maintain the intra- and extra-cellular osmotic balance [40]. Interestingly, the compatible solutes ectoine and hydroxyectoine, are highly value-added compounds acting as protective agent for proteins or cells [41]; they have been commercially used in cosmetics, organ transplantation and medicinal areas. Thus, most *Halomonas* spp. are high-performing ectoine producers due to their possession of natural ectoine synthesis pathways [42–46], that can also be cloned into other hosts for enhanced ectoine production [47–49]. In addition, many *Halomonas* spp. have been reported to accumulate intracellular inclusion body, typically, PHB, one of the members of biodegradable polyesters, PHAs [350–5250–52]. Moreover, increasing interests on biosurfactants, bioemulsifiers, some proteins from *Halomonas* [53–58], have turned *Halomonas* spp. into platforms for diverse bioproductions (Figure 1).

Genetic manipulations in *Halomonas* spp.

Due to the recent growing interest in *Halomonas* spp. used as microbial chassis, many efforts have been made to develop genetic tools, such as expression vectors [59,60], promoters [24,61], ribosome-binding sites (RBSs), inducible systems [25], genome editing tools [26] and pathway tuning approaches [42,62,63], allowing genetic manipulations of *Halomonas* spp. possible for strain improvements [64]. For a given module of gene over-expression in model microbes such as *E. coli* and *P. putida*, several parts are necessary including vectors containing relative antibiotic-resistant gene(s) and replicon, promoters, RBSs, coding sequences and terminators [65]. Insulators are sometimes needed to minimize the unwanted sequence context between two different gene parts [66]. Generally, transformation of broad host range genetic parts into *Halomonas* spp. is an efficient strategy to screen suitable gene expression elements [23]. However, native plasmids isolated from Gram-negative halophiles are also good choice for cloning uses [67]. Till now, many expression vectors including pSEVA plasmids [68], shuttle vectors pWL102 and pUBP2 [69] and other broad host range or native plasmids are found to be usable in *Halomonas* spp. Accordingly, antibiotics such as chloramycetin (Cm) and spectinomycin (Spe) have been proven effective for *Halomonas* spp [70]. More importantly, a *porin* constitutive promoter library consisting of a wide range of transcriptional strengths, a novel type of T7-like inducible system and expression variances cross different expression systems were developed and characterized in *Halomonas* spp. for gene expression tuning [25,61,63] (Figure 3A). Moreover, recent efforts on whole-genome sequencing (WGS) of over 450 *Halomonas* spp. strains (Supplementary Table S1), including *H. bluephagenesis* (also termed *Halomonas* TD01), *Halomonas smyrnensis* AAD6 [71,72], *Halomonas* sp. KM-1 [73], *Halomonas* sp. HAL1 [74] and so on, provides plentiful genomic information for boosting the developments of endogenous genetic parts mining, valuable pathways identification, metabolic networks modeling [72,75] etc., which offers fundamental insights into rational microbial cell factory engineering based on *Halomonas* sp. [35,75–77].




Figure 1. Overview of *Halomonas* engineering for biotechnological industry

Many systems and synthetic biology tools and approaches, for example CRISPR/Cas9-based gene editing, omics profiling, parts mining, static and dynamic optimization methods, have been developed for *Halomonas* spp. It is thus advanced, the genetic reprogramming of *Halomonas* spp. allowing construction of high-performance *Halomonas* cell factories for production of a variety of chemicals, polyesters and proteins. A cost-effective NGIB has been developed based on extremophilic bacteria especially *Halomonas* spp. for bioproduction in various scales. Abbreviation: CRISPR, clustered regularly interspaced short palindromic repeats.

Genome editing tools including homologous recombination and clustered regularly interspaced short palindromic repeats (CRISPR)/Cas9 have been commonly used for site-specific mutagenesis in many microorganisms [78]. However, there were less studies reported in *Halomonas* spp. [26,79] except some successful cases based on double cross-overs homologous recombination using lethal genes or helper plasmids as selection pressures. An essential gene-deficient mutant can be used as the host (in the presence of the gene encoded in a plasmid), for example, the deletion of gene *pryF* encoding orotidine-5'-phosphate decarboxylase, can significantly improve the selection pressure during mutagenesis, especially for essential gene deletion and large deoxyribonucleic acid (DNA) fragment integration [80,81]. Recently, a CRISPR/Cas9-based gene editing tool was established in *H. bluephagenesis* in the authors' lab for engineering chromosomes, such as gene knock-down for morphology control [80], bypass deletion for product flux enhancement [82], target module integration on to the chromosome [63]. All of these efforts help turn *Halomonas* spp. into a reprogrammable chassis comparable with other model microbes (Figure 1).




Figure 2. Metabolic pathways for diverse bioproductions by engineered *Halomonas*

Halomonas spp. have been engineered to produce different chemicals and polyesters from glucose, fatty acids, gluconate or other structure-related carbon sources (light gray). Three major metabolic modules involved in glucose metabolism, PHA biosynthesis (light blue), and productions of amino acids and their derivatives (light orange) are summarized according to the published studies. Crosses in red represent bypasses (letters in light gray) required to be knockout.

Table 1 Key achievements of bioproductions by *Halomonas*

Strains	Productions	Titer	References
<i>H. bluephagenesis</i> (<i>Halomonas</i> TD01)	Ectoine	28 g/l	[42]
	Threonine	33 g/l	[54]
	3-Hydroxypropionate	154 g/l	[87]
	5-Minolevulinic acid	0.7 g/l	[53]
<i>H. smymensis</i> AAD6	Levan	18.06 g/l	[31]
<i>Halomonas</i> sp. KM-1	Pyruvate	63.3 g/l	[34]

Currently, conjugation is still the most commonly used genetic transformation method for *Halomonas* spp. due to the unsolved difficulty of electroporation or chemical transformation of expression vectors [79]. Therefore, many high throughput-dependent methods are not useful for *Halomonas* spp. engineering, resulting in challenges of attempts in large dataset mining and analysis when leveraging ‘Design-Build-Test-Learn’ cycle [83].

Bioproductions by recombinant *Halomonas* spp.

Due to the natural accumulation capability of PHAs and osmolytes by many isolated *Halomonas* spp., metabolic engineering on *Halomonas* spp. has attracted growing attention for enhanced production of PHA, ectoine and their derived products, levan, exopolymers and so on (Figure 1 and Table 1) [29,84]. A lot of studies have been devoted to produce diverse PHA, including PHB [27,50,85], copolyester of 3-hydroxybutyrate and 3-hydroxyvalerate (PHBV) [82], copolyester of 3-hydroxybutyrate and 4-hydroxybutyrate (P34HB) [70] from glucose, sucrose [56]




Figure 3. Streamlined engineering of *Halomonas* as a chassis

(A) Genetic tools development includes promoter engineering, novel type T7-like inducible system, expression correlation cross vectors and/or different genomic locus. (B) Stimulus response-based pathway tuning approach enables fast and exquisite gene expression optimization on chromosome-based system in corporation of GFP-mediated transcription mapping approach. (C) Innovative strain engineering strategies boosts the industrial performance of engineered cells by enhancing oxygen availability, increasing the cell volume, achieving self-flocculation at the end of the growth and production processes. (D) Strain engineering pipeline of *Halomonas* for industrial purposes. Abbreviation: GFP, green fluorescent protein.

and waste gluconate [28]. *Halomonas boliviensis* [46,50], *Halomonas campaniensis* [52] and *H. bluephagenesis* [28] performed well in PHA accumulation. *H. boliviensis* can utilize diverse substrates to produce high molecular weight PHB, reaching up to 1100 kilo-Daltons (kDa) [50]. *H. campaniensis*, a moderate halophile, was engineered to produce over 70 wt% PHB using kitchen waste-like mixed substrates conducted under continuous and open fermentation over a period of 65 days [52]. Notably, *H. bluephagenesis*, a predominant PHA producer isolated from Idyngo Lake, Xinjiang/China, is able to accumulate more than 90 g/l biomass containing over 80 wt% of PHB [27]. Further engineering effects have made it a diverse PHA copolymer producer, such as PHBV [82], P34HB [70] and copolyester of 3-hydroxybutyrate and 3-hydroxyhexanoate (PHBHHx) [86], when grown on glucose and/or structure-related carbon sources serving as precursors of the non-3HB monomer (Figure 2). It is important to note

that the pilot-scale P34HB production by engineered *H. bluephagenesis* has demonstrated its success as an industrial chassis for NGIB [6]. In addition, recombinant *H. bluephagenesis* has displayed proven ability for productions of bio-surfactant and bio-emulsifier (PHA surface binding proteins PhaR and PhaP) [23,57], ectoine [42], L-threonine [54], 5-minolevulinic acid (ALA) [53], 3-hydroxypropionate (3HP) [87] and many more to come, all under open unsterile conditions (Figure 2).

Metabolic engineering in *Halomonas* spp.

Metabolic engineering generally requires exquisite expression tuning of target pathways to channel the metabolic flux towards metabolic targets leveraging different static optimization approaches in addition to the commonly used methods, including enzyme engineering, promoter engineering, RBS optimization, gene over-expression (on/off-control only) and bypass knockout (Figure 2). Recently, an approach termed high-resolution gene expression control was developed for *H. bluephagenesis* based on the combination of two isopropyl- β -D-thiogalactopyranoside (IPTG)-induced systems with different dynamic ranges [63]. Similar to the length measuring tools with different scales of measuring ranges, this approach allows precise transcription tuning of target genes on chromosome directly in cross-magnitude scopes in corporation of green fluorescent protein (GFP)-mediated transcriptional mapping strategy [63]. On the basis of these successes, two orthogonal inducible systems induced by acyl homoserine lactone (AHL) and IPTG, respectively, were used to fine-tune two individual expression modules simultaneously, resulting in over 12-folds improvement of ectoine titer by engineered *H. bluephagenesis* [42]. This attempt gives a successful example for high throughput-independent strain engineering paradigm without large size library construction and labor-intensive screening process (Figure 3B), which are helpful for non-model bacterial engineering with low efficient transformation methods, such as conjugation.

In addition to the metabolic engineering approaches, several strategies have been employed to achieve strain improvements for *Halomonas* spp.: (I) morphology engineering to enlarge cell shapes for enhanced intracellular substances (such as PHA or proteins) accumulation and for benefiting sedimentation during the separation process [80]; (II) introduction of bacterial hemoglobin, *Vitreoscilla* hemoglobin (VHb) for enhanced oxygen uptake leading to higher cell mass and more product formation as well as aeration energy saving [88]; (III) self-flocculation for convenient separation and wastewater-less bioprocessing by recycling the supernatant of fermented broth after heat treatment and membrane processing with a recycling rate of 70–85% [89]; (IV) control of redox potential nicotinamide adenine dinucleotide plus hydrogen (NADH)/nicotinamide adenine dinucleotide plus (NAD⁺) by supplementing acetate to improve PHA and biomass accumulation (Figure 3C) [90]. All of these strategies extend ability of *Halomonas* spp.

Halomonas spp. have been successfully engineered in previous studies including cell engineering, pathway optimization and process scale-up (Figure 3D). Firstly, DNA sequence-based fundamental understanding of the selected *Halomonas* sp. and the genetic tools mining thereof is the cornerstone for strain engineering. Generally, the genome-wide sequencing and annotation can provide us a predictable overview of functional gene sets and metabolic networks [91]. Growth and characterization are important to obtain cultural protocols for various engineered purposes in different scales, including 96 deep well-plates, shake flasks and bench-top bioreactors, their results serve as reference-standards for growth profiling and pre-knowledge of recombinant cells without performing the scale-up studies [63], these are especially important as *Halomonas* spp. have NaCl concentrations and pH preferences.

Most importantly, optimization of genetic parts, promoter, RBS [49] and tools, promoter library construction [92], inducible systems [93], effective CRISPR-based gene editing approach [94] and alternative expression vectors, have been demonstrated to strongly improve the programmability of *Halomonas*. Screening on chromosomal locus with low expression variance can provide applicable genomic sites for efficient integrations of multiple expression modules [95]. Secondly, assembling catalytic enzymes to rewire the endogenous flux towards metabolic targets is commonly employed to achieve prototype success of target products from 0 to 1, namely pathway construction. Further optimization leveraging static regulation, mainly refers to bypass deletion and flux tuning, and dynamic regulation [96] allowing gene expression control over time and levels, is a proven strategy to generate significant breakthrough from 1 to 100. Thirdly, a rigorous scale-up test including medium and feeding strategy optimization can obtain iterative bioprocess of refinement. Accordingly, many fundamental insights can also be uncovered during the growth assays in corporation of omics profiling [97] and adaptive evolution [98], directing the redesign of cell factory by implementing ‘Design-Build-Test’ cycle, this is also true for *Halomonas* spp.

Many bioprocess-inspired strategies are useful in real cases, such as morphology control and self-flocculating cells aiming to make continuous fermentation and easier downstream separation possible. For compiling these three parts into a closed loop streamline, the interactive relationship among ‘host engineering’, ‘downstream engineering’ and

'scale-up' is usually missed (Figure 3D), however, they should be combined into strain engineering so that the engineering concept is used throughout the production. The NGIB concept based on extremophilic bacteria proposed by the authors' lab has demonstrated the engineering concept up- and down-stream of the bioproduction [64]. Recent efforts have also exemplified a successful case in rebuilding the yeast system for obtaining a metabolic target-dependent (or -preferred) host to realize enhanced production of fatty acids before the execution of pathway optimization [98].

Opportunities and challenges

Compared with the well-studied chassis especially *E. coli* and *C. glutamicum*, *Halomonas* spp. displaying several advantages including contamination-resistance, fast growth, seawater-based media and wide range substrates utilization, added with rich engineering tools and approaches developed recently, have become a promising platform for NGIB. A variety of products have been successfully produced with some of them scaled up to at least 5000 liters bioreactors under open unsterile conditions. Since *Halomonas* spp. can be engineered as convenient as *E. coli*, their potentials will be at least as promising as *E. coli* which has been employed to produce many bio-based products in various scales. *Halomonas* spp.-based NGIB has provided a versatile low-cost platform of biomanufacturing to meet the increasing demand of sustainable development that chemical industry and CIB cannot realize.

More efforts and attempts should be made to strengthen the sciences and technological sites of NGIB for overcoming the accompanying challenges. For example, the development of high throughput genetic transformation methods to enable generation of large size datasets from clones [49], establishments of multigene pathway tuning strategy [93] and dynamic control systems [99], exploration of high cell density cultivation and induction technology [100], enhancement of substrates to product conversion efficiency, controllable cell morphology changes for better growth, production and downstream processing. The coming joint efforts offer more possibilities for *Halomonas* as a chassis comparable with *E. coli* yet with more advantages.

Summary

- *Halomonas* spp. as chassis are able to grow rapidly in saline and alkaline environments allowing contamination-free cultivation under open conditions.
- *Halomonas* spp. display advantages including fast growth, seawater-based medium and a wide substrate range as well as simple cultivation processes.
- Similar to *E. coli*, *Halomonas* spp. can be engineered to produce diverse products including several PHAs, chemicals, food additives and proteins.
- *H. bluephagenesis* is one of the most promising hosts as it allows convenient genetic manipulations and process scale-up under open unsterile conditions.
- NGIB based on *Halomonas* spp. as chassis provides a competitive solution for overcoming the challenges of high energy and fresh water consumption based on CIB.

Competing Interests

The authors declare that there are no competing interests associated with the manuscript.

Funding

This work was supported by the Ministry of Science and Technology [grant number 2018YFA0900200 (to G.-Q.C.); the National Natural Science Foundation of China [grant numbers 31430003, 21761132013, 31870859 (to G.-Q.C.), 32001029 (to J.-W.Y.)]; the Tsinghua University-INDITEX Sustainable Development Fund [grant number TISD201907 (to J.-W.Y.)]; and the National Natural Science Foundation of China [grant numbers 31961133017, 31961133018, 31961133019], which are part of MIX-UP, a joint NSFC and EU H2020 collaboration. In Europe, MIX-UP has received funding from the European Union's Horizon 2020 Research and Innovation Program [grant number 870294].

Author Contribution

J.-W.Y. and G.-Q.C. wrote the paper and drafted the figures.

Abbreviations

AHL, acyl homoserine lactone; CIB, current industrial biotechnology; CRISPR, clustered regularly interspaced short palindromic repeats; DNA, deoxyribonucleic acid; IPTG, isopropyl- β -D-thiogalactopyranoside; NGIB, next-generation industrial biotechnology; P34HB, poly(3-hydroxybutyrate-co-4-hydroxybutyrate); PHA, polyhydroxyalkanoate; PHB, poly-3-hydroxybutyrate; PHBV, poly(3-hydroxybutyrate-co-3-hydroxyvalerate); RBS, ribosome-binding site.

References

- Murphy, A.C. (2011) Metabolic engineering is key to a sustainable chemical industry. *Nat. Prod. Rep.* **8**, 1406–1425, <https://doi.org/10.1039/c1np00029b>
- Olguin, E.J., Giuliano, G., Porro, D., Tuberosa, R. and Salamini, F. (2012) Biotechnology for a more sustainable world. *Biotechnol. Adv.* **30**, 931–932, <https://doi.org/10.1016/j.biotechadv.2012.06.001>
- Philp, J.C., Ritchie, R.J. and Allan, J.E.M. (2013) Biobased chemicals: the convergence of green chemistry with industrial biotechnology. *Trends Biotechnol.* **31**, 219–222, <https://doi.org/10.1016/j.tibtech.2012.12.007>
- Becker, J., Rohles, C.M. and Wittmann, C. (2018) Metabolically engineered *Corynebacterium glutamicum* for bio-based production of chemicals, fuels, materials, and healthcare products. *Metab. Eng.* **50**, 122–141, <https://doi.org/10.1016/j.ymben.2018.07.008>
- Yu, L.P., Wu, F.Q. and Chen, G.-Q. (2019) Next-generation industrial biotechnology-transforming the current industrial biotechnology into competitive processes. *Biotechnol. J.* **14**, 1800437, <https://doi.org/10.1002/biot.201800437>
- Chen, G.-Q. and Jiang, X. (2018) Next generation industrial biotechnology based on extremophilic bacteria. *Curr. Opin. Biotechnol.* **50**, 94–100, <https://doi.org/10.1016/j.copbio.2017.11.016>
- Park, S.J., Kim, T.W., Kim, M.K., Lee, S.Y. and Lim, S.C. (2011) Advanced bacterial polyhydroxyalkanoates: towards a versatile and sustainable platform for unnatural tailor-made polyesters. *Biotechnol. Adv.* **30**, 1196–1206, <https://doi.org/10.1016/j.biotechadv.2011.11.007>
- Pollard, D.J. and Woodley, J.M. (2007) Biocatalysis for pharmaceutical intermediates: the future is now. *Trends Biotechnol.* **25**, 66–73, <https://doi.org/10.1016/j.tibtech.2006.12.005>
- Fang, H., Li, D., Kang, J., Jiang, P., Sun, J. and Zhang, D. (2018) Metabolic engineering of *Escherichia coli* for de novo biosynthesis of vitamin B12. *Nat. Commun.* **9**, 4917, <https://doi.org/10.1038/s41467-018-07412-6>
- Fernandesab, P., Cruza, A., Angelovaac, B., Pinheiroa, H. and Cabral, J.M. (2003) Microbial conversion of steroid compounds: recent developments. *Enzyme Microb. Technol.* **32**, 688–705, [https://doi.org/10.1016/S0141-0229\(03\)00029-2](https://doi.org/10.1016/S0141-0229(03)00029-2)
- Ingram, L.O., Gomez, P.F., Lai, X., Moniruzzaman, M., Wood, B.E., Yomano, L.P. et al. (2000) Metabolic engineering of bacteria for ethanol production. *Biotechnol. Bioeng.* **58**, 204–214, [https://doi.org/10.1002/\(SICI\)1097-0290\(19980420\)58:2<3%3c204::AID-BIT13%3e3.0.CO;2-C](https://doi.org/10.1002/(SICI)1097-0290(19980420)58:2<3%3c204::AID-BIT13%3e3.0.CO;2-C)
- Steen, E.J., Chan, R., Prasad, N., Myers, S., Petzold, C.J., Redding, A. et al. (2008) Metabolic engineering of *Saccharomyces cerevisiae* for the production of n-butanol. *Microb. Cell Fact.* **7**, 36, <https://doi.org/10.1186/1475-2859-7-36>
- Dutta, D., De, D., Chaudhuri, S. and Bhattacharya, S.K. (2005) Hydrogen production by *Cyanobacteria*. *Microb. Cell Fact.* **4**, 36, <https://doi.org/10.1186/1475-2859-4-36>
- Dueber, J.E., Wu, G.C., Malmirchegini, G.R., Moon, T.S., Petzold, C.J., Ullal, A.V. et al. (2009) Synthetic protein scaffolds provide modular control over metabolic flux. *Nat. Biotechnol.* **27**, 753–759, <https://doi.org/10.1038/nbt.1557>
- Park, J.H., Lee, S.Y., Kim, T.Y. and Kin, H.U. (2008) Application of systems biology for bioprocess development. *Trends Biotechnol.* **26**, 404–412, <https://doi.org/10.1016/j.tibtech.2008.05.001>
- Chen, Z. and Wan, C. (2017) Non-sterile fermentations for the economical biochemical conversion of renewable feedstocks. *Biotechnol. Lett.* **39**, 1765–1777, <https://doi.org/10.1007/s10529-017-2429-8>
- Kivisto, A., Santala, V. and Karp, M. (2013) Non-sterile process for biohydrogen and 1,3-propanediol production from raw glycerol. *Int. J. Hydrog. Energy* **38**, 11749–11755, <https://doi.org/10.1016/j.ijhydene.2013.06.119>
- Oren, A. (2010) Industrial and environmental applications of halophilic microorganisms. *Environ. Technol.* **31**, 825–834, <https://doi.org/10.1080/09593330903370026>
- Tyo, K.E.J., Ajikumar, P.K. and Stephanopoulos, G. (2009) Stabilized gene duplication enables long-term selection-free heterologous pathway expression. *Nat. Biotechnol.* **27**, 760–765, <https://doi.org/10.1038/nbt.1555>
- Yztürk, S., Jälek, P. and Jzdamar, T.H. (2016) Fed-batch biomolecule production by *Bacillus subtilis*: A state of the art review. *Trends Biotechnol.* **34**, 329–345, <https://doi.org/10.1016/j.tibtech.2015.12.008>
- Loeschcke, A. and Thies, S. (2015) *Pseudomonas putida*-a versatile host for the production of natural products. *Appl. Microb. Biotechnol.* **99**, 6197–6214, <https://doi.org/10.1007/s00253-015-6745-4>
- Beckera, J., Zelder, O., Häfner, S., Schröder, H. and Wittmann, C. (2011) From zero to hero-Design-based systems metabolic engineering of *Corynebacterium glutamicum* for L-lysine production. *Metab. Eng.* **13**, 159–168, <https://doi.org/10.1016/j.ymben.2011.01.003>
- Fu, X.Z., Tan, D., Aibaidula, G., Wu, Q., Chen, J.C. and Chen, G.-Q. (2014) Development of *Halomonas* TD01 as a host for open production of chemicals. *Metab. Eng.* **23**, 78–91, <https://doi.org/10.1016/j.ymben.2014.02.006>
- Li, T., Li, T., Ji, W., Wang, Q., Zhang, H., Chen, G.-Q. et al. (2016) Engineering of core promoter regions enables the construction of constitutive and inducible promoters in *Halomonas* sp. *Biotechnol. J.* **11**, 219–227, <https://doi.org/10.1002/biot.201400828>

- 25 Zhao, H., Zhang, H.M., Chen, X.B., Li, T., Wu, Q., Ouyang, Q. et al. (2017) Novel T7-like expression systems used for *Halomonas*. *Metab. Eng.* **39**, 128–140, <https://doi.org/10.1016/j.ymben.2016.11.007>
- 26 Qin, Q., Ling, C., Zhao, Y.Q., Yang, T., Yin, J., Guo, Y.Y. et al. (2018) CRISPR/Cas9 editing genome of extremophile *Halomonas* spp. *Metab. Eng.* **47**, 219–229, <https://doi.org/10.1016/j.ymben.2018.03.018>
- 27 Tan, D., Xue, Y.S., Aibaidula, G. and Chen, G.Q. (2011) Unsterile and continuous production of polyhydroxybutyrate by *Halomonas* TD01. *Bioresour. Technol.* **102**, 8130–8136, <https://doi.org/10.1016/j.biortech.2011.05.068>
- 28 Ye, J.W., Huang, W.Z., Wang, D.S., Chen, F.Y., Yin, J., Li, T. et al. (2018) Pilot scale-up of poly(3-hydroxybutyrate-co-4-hydroxybutyrate) production by *Halomonas bluephagenesis* via cell growth adapted optimization process. *Biotechnol. J.* **13**, e1800074, <https://doi.org/10.1002/biot.201800074>
- 29 Hande, K.S., Ozlem, A., Gonca, O., Kazim, Y.A. and Ebru, T.O. (2015) Effective stimulating factors for microbial levan production by *Halomonas smyrnensis* AAD6. *J. Biosci. Bioeng.* **119**, 455–463, <https://doi.org/10.1016/j.jbiosc.2014.09.019>
- 30 Souha, T., Gül, G.H., Mehmet, S.E., Ceyda, K., Seval, G., Zehra, S.C. et al. (2018) *Halomonas smyrnensis* as a cell factory for co-production of PHB and levan. *Int. J. Biol. Macromol.* **118**, 1238–1246, <https://doi.org/10.1016/j.ijbiomac.2018.06.197>
- 31 Erkorkmaz, B.A., Krtel, O., Duru, Ö.A. and Öner, E.T. (2018) Development of a cost-effective production process for *Halomonas* levan. *Biopro. Biosyst.* **41**, 1247–1259, <https://doi.org/10.1007/s00449-018-1952-x>
- 32 Jin, Y.X., Shi, L.H. and Yoshikazu, K. (2013) Metabolomics-based component profiling of *Halomonas* sp. KM-1 during different growth phases in poly(3-hydroxybutyrate) production. *Bioresour. Technol.* **140**, 73–79, <https://doi.org/10.1016/j.biortech.2013.04.059>
- 33 Yoshikazu, K. and Aiba, S. (2010) Poly(3-hydroxybutyrate) production by isolated *Halomonas* sp. KM-1 using waste glycerol. *Biosci. Biotechnol. Biochem.* **74**, 175–177, <https://doi.org/10.1271/bbb.90459>
- 34 Yoshikazu, K., Taku, N., Isao, M. and Jun, T. (2016) Efficient production and secretion of pyruvate from *Halomonas* sp. KM-1 under aerobic conditions. *AMB Express* **6**, 22, <https://doi.org/10.1186/s13568-016-0195-y>
- 35 Chen, F., Cao, Y., Wei, S., Li, Y., Li, X., Wang, Q. et al. (2015) Regulation of arsenite oxidation by the phosphate two-component system PhoBR in *Halomonas* sp. HAL1. *Front. Microbiol.* **6**, 923, <https://doi.org/10.3389/fmicb.2015.00923>
- 36 Setati, M.E. (2010) Diversity and industrial potential of hydrolase-producing halophilic/halotolerant eubacteria. *Afr. J. Biotechnol.* **9**, 1555–1560, <https://doi.org/10.5897/AJB10.051>
- 37 Yin, J., Chen, J.C., Wu, Q. and Chen, G.-Q. (2015) Halophiles, coming stars for industrial biotechnology. *Biotechnol. Adv.* **33**, 1433–1442, <https://doi.org/10.1016/j.biotechadv.2014.10.008>
- 38 Oren, A. (2002) Diversity of halophilic microorganisms: environments, phylogeny, physiology, and applications. *Ind. Microbiol. Biotechnol.* **28**, 56–63, <https://doi.org/10.1038/sj/jim/7000176>
- 39 Oren, A. (2008) Microbial life at high salt concentrations: phylogenetic and metabolic diversity. *Saline Syst.* **4**, 2, <https://doi.org/10.1186/1746-1448-4-2>
- 40 Quillaguamán, J., Guzmán, H., Van-Thuoc, D. and Hatti-Kaul, R. (2010) Synthesis and production of polyhydroxyalkanoates by halophiles: current potential and future prospects. *Appl. Microbiol. Biotechnol.* **85**, 1687–1696, <https://doi.org/10.1007/s00253-009-2397-6>
- 41 Pastor, J., Salvador, M., Argandoña, M., Bernal, V., Reina-Bueno, M., Csonka, L. et al. (2010) Ectoines in cell stress protection: uses and biotechnological production. *Biotechnol. Adv.* **28**, 782–801, <https://doi.org/10.1016/j.biotechadv.2010.06.005>
- 42 Ma, H., Zhao, Y.Q., Huang, W.Z., Zhang, L.Z., Wu, F.Q., Ye, J.W. et al. (2020) Rational flux-tuning of *Halomonas bluephagenesis* for co-production of bioplastic PHB and ectoine. *Nat. Commun.* **11**, 3311, <https://doi.org/10.1038/s41467-020-17223-3>
- 43 Tanimura, K., Nakayama, H., Tanaka, T. and Kondo, A. (2013) Ectoine production from lignocellulosic biomass-derived sugars by engineered *Halomonas elongate*. *Bioresour. Technol.* **142**, 523–529, <https://doi.org/10.1016/j.biortech.2013.05.004>
- 44 Schwibbert, K., Marin-Sanguino, A., Bagyan, I., Heidrich, G., Lentzen, G., Seitz, H. et al. (2011) A blueprint of ectoine metabolism from the genome of the industrial producer *Halomonas elongata* DSM 2581T. *Environ. Microbiol.* **13**, 1973–1994, <https://doi.org/10.1111/j.1462-2920.2010.02336.x>
- 45 Chen, P.W., Cui, Z.Y., Ng, H.S. and Lan, J.C. (2020) Exploring the additive bio-agent impacts upon ectoine production by *Halomonas salina* DSM5928 T using corn steep liquor and soybean hydrolysate as nutrient supplement. *J. Biosci. Bioeng.* **130**, 195–199, <https://doi.org/10.1016/j.jbiosc.2020.03.011>
- 46 Guzmán, H., Van-Thuoc, D., Martín, J., Hatti-Kaul, R. and Quillaguamán, J. (2009) A process for the production of ectoine and poly(3-hydroxybutyrate) by *Halomonas boliviensis*. *Appl. Microbiol. Biotechnol.* **84**, 1069–1077, <https://doi.org/10.1007/s00253-009-2036-2>
- 47 Chen, J., Liu, P., Chu, X., Chen, J., Zhang, H., David, C.R. et al. (2020) Metabolic pathway construction and optimization of *Escherichia coli* for high-level ectoine production. *Curr. Microbiol.* **77**, 1412–1418, <https://doi.org/10.1007/s00284-020-01888-6>
- 48 Ning, Y.K., Wu, X.J., Zhang, C., Xu, Q., Chen, N. and Xie, X. (2016) Pathway construction and metabolic engineering for fermentative production of ectoine in *Escherichia coli*. *Metab. Eng.* **36**, 10–18, <https://doi.org/10.1016/j.ymben.2016.02.013>
- 49 Gideon, G., Demian, D., Lukas, J., Michael, K., Eun, J.J., Sung, S.Y. et al. (2019) Metabolic engineering of *Corynebacterium glutamicum* for high-level ectoine production: design, combinatorial assembly, and implementation of a transcriptionally balanced heterologous ectoine pathway. *Biotechnol. J.* **14**, e1800417, <https://doi.org/10.1002/biot.201800417>
- 50 Quillaguamán, J., Hashim, S., Bento, F., Mattiasson, B. and Hatti-Kaul, R. (2005) Poly(β -hydroxybutyrate) production by a moderate halophile, *Halomonas boliviensis* LC1 using starch hydrolysate as substrate. *J. Appl. Microbiol.* **99**, 151–157, <https://doi.org/10.1111/j.1365-2672.2005.02589.x>
- 51 Quillaguamán, J., Guzmán, H., Van-Thuoc, D. and Hatti-Kaul, R. (2010) Synthesis and production of polyhydroxyalkanoates by halophiles: current potential and future prospects. *Appl. Microbiol. Biotechnol.* **85**, 1687–1696, <https://doi.org/10.1007/s00253-009-2397-6>
- 52 Yue, H., Ling, C., Yang, T., Chen, X., Chen, Y., Deng, H. et al. (2014) A seawater based open and continuous process for polyhydroxyalkanoates production by recombinant *Halomonas campaniensis* LS21 grown in mixed substrates. *Biotechnol. Biofuels* **7**, 108, <https://doi.org/10.1186/1754-6834-7-108>

- 53 Li, T., Guo, Y., Qiao, G.Q. and Chen, G.-Q. (2016) Microbial synthesis of 5-aminolevulinic acid and its coproduction with polyhydroxybutyrate. *ACS Synth. Biol.* **5**, 1264–1274, <https://doi.org/10.1021/acssynbio.6b00105>
- 54 Du, H.T., Zhao, Y.Q., Wu, F.Q., Ouyang, P.F., Chen, J.C., Jiang, X.R. et al. (2020) Engineering *Halomonas bluephagenesis* for L-Threonine production. *Metab. Eng.* **60**, 119–127, <https://doi.org/10.1016/j.ymben.2020.04.004>
- 55 Coronado, M.J., Vargas, C., Hofemeister, J., Ventosa, A. and Nieto, J.J. (2006) Production and biochemical characterization of an α -amylase from the moderate halophile *Halomonas meridiana*. *FEMS Microb. Lett.* **183**, 67–71
- 56 Shivanand, P., Mugeraya, G. and Kumar, A. (2013) Utilization of renewable agricultural residues for the production of extracellular halostable cellulase from newly isolated *Halomonas* sp. strain PS47. *Ann. Microbiol.* **63**, 1257–1263, <https://doi.org/10.1007/s13213-012-0583-8>
- 57 Lan, L.H., Zhao, H., Chen, J.C. and Chen, G.-Q. (2016) Engineering *Halomonas* spp. as a low-cost production host for production of bio-surfactant protein PhaP. *Biotechnol. J.* **11**, 1595–1604, <https://doi.org/10.1002/biot.201600459>
- 58 Ma, H.K., Liu, M.M., Li, S.Y., Wu, Q., Chen, J.C. and Chen, G.-Q. (2013) Application of polyhydroxyalkanoate (PHA) synthesis regulatory protein PhaR as a bio-surfactant and bactericidal agent. *J. Biotechnol.* **166**, 34–41, <https://doi.org/10.1016/j.jbiotec.2013.04.017>
- 59 Vargas, C., Fernández-Castillo, R., Cánovas, D., Ventosa, A. and Nieto, J.J. (1995) Isolation of cryptic plasmids from moderately halophilic eubacteria of the genus *Halomonas*: characterization of a small plasmid from *H. elongata* and its use for shuttle vector construction. *Mol. Gen. Genet.* **246**, 411–418, <https://doi.org/10.1007/BF00290444>
- 60 Vargas, C., Tegos, G., Vartholomatos, G., Drainas, C., Ventosa, A. and Nieto, J.J. (1999) Genetic organization of the mobilization region of the plasmid pHE1 from *Halomonas elongate*. *Syst. Appl. Microbiol.* **22**, 520–529
- 61 Shen, R., Yin, J., Ye, J.W., Xiang, R.J., Ning, Z.Y., Huang, W.Z. et al. (2018) Promoter engineering for enhanced P(3HB-co-4HB) production by *Halomonas bluephagenesis*. *ACS Synth. Biol.* **7**, 1897–1906, <https://doi.org/10.1021/acssynbio.8b00102>
- 62 Zhang, X., Lin, Y.N., Wu, Q., Wang, Y. and Chen, G.-Q. (2019) Synthetic biology and genome-editing tools for improving PHA metabolic engineering. *Trends Biotechnol.* **38**, 689–700, <https://doi.org/10.1016/j.tibtech.2019.10.006>
- 63 Ye, J.W., Hu, D.K., Yin, J., Huang, W.Z., Xiang, R.J., Zhang, L.Z. et al. (2020) Stimulus response-based fine-tuning of polyhydroxyalkanoate pathway in *Halomonas*. *Metab. Eng.* **57**, 85–95, <https://doi.org/10.1016/j.ymben.2019.10.007>
- 64 Chen, G.-Q. and Jiang, X. (2018) Next generation industrial biotechnology based on extremophilic bacteria. *Curr. Opin. Biotechnol.* **50**, 94–100, <https://doi.org/10.1016/j.copbio.2017.11.016>
- 65 Nielsen, A.A.K., Der, B.S., Shin, J., Vaidyanathan, P., Paralanov, V., Strychalski, E.A. et al. (2016) Genetic circuit design automation. *Science* **352**, 7341, <https://doi.org/10.1126/science.aac7341>
- 66 Lou, C.B., Stanton, B., Chen, Y.J., Munsky, B. and Voigt, C.A. (2012) Ribozyme-based insulator parts buffer synthetic circuits from genetic context. *Nat. Biotechnol.* **30**, 1137–1142, <https://doi.org/10.1038/nbt.2401>
- 67 Argandoña, M., Vargas, C., Reina-Bueno, M., Rodríguez-Moya, J., Salvador, M. and Nieto, J. (2012) An extended suite of genetic tools for use in bacteria of the *Halomonadaceae*: an overview. *Methods Mol. Biol.* **824**, 167–201, https://doi.org/10.1007/978-1-61779-433-9_9
- 68 Martínez-García, E., Goñi-Moreno, A., Bartley, B., McLaughlin, J., Sánchez-Sampedro, L., Pozo, H.P.D. et al. (2020) SEVA 3.0: an update of the Standard European Vector Architecture for enabling portability of genetic constructs among diverse bacterial hosts. *Nucleic Acids Res.* **48**, 3395, <https://doi.org/10.1093/nar/gkaa114>
- 69 Han, J., Lu, Q., Zhou, L., Zhou, J. and Xiang, H. (2007) Molecular characterization of the *phaE_{CHm}* genes, required for biosynthesis of poly(3-hydroxybutyrate) in the extremely halophilic archaeon *Haloarcula marismortui*. *Appl. Environ. Microbiol.* **73**, 6058–6065, <https://doi.org/10.1128/AEM.00953-07>
- 70 Ye, J.W., Hu, D.K., Che, X.M., Jiang, X.R., Li, T., Chen, J.C. et al. (2018) Engineering of *Halomonas bluephagenesis* for low cost production of poly(3-hydroxybutyrate-co-4-hydroxybutyrate) from glucose. *Metab. Eng.* **47**, 143–152, <https://doi.org/10.1016/j.ymben.2018.03.013>
- 71 Elif, S., Zeliha, E., Muzzaffer, A., Aris, C., Neslihan, A., Ebru, T.O. et al. (2012) Draft genome sequence of *Halomonas smyrnensis* AAD6. *J. Bacteriol.* **194**, 5690–5691, <https://doi.org/10.1128/JB.00559-12>
- 72 Elif, D., Tugba, O., Muzaffer, A., Zeliha, E., Ebru, T.O., Duran, U. et al. (2015) Genomic analysis reveals the biotechnological and industrial potential of levan producing halophilic extremophile, *Halomonas smyrnensis* AAD6T. *SpringerPlus* **4**, 393, <https://doi.org/10.1186/s40064-015-1184-3>
- 73 Yoshikazu, K., Kazunori, K. and Yasushi, S. (2012) Draft genome sequence of *Halomonas* sp. strain KM-1, a moderately halophilic bacterium that produces the bioplastic poly(3-Hydroxybutyrate). *J. Bacteriol.* **194**, 2738–2739, <https://doi.org/10.1128/JB.00237-12>
- 74 Lin, Y., Fan, H., Hao, X., Johnstone, L., Hu, Y., Wei, G. et al. (2012) Draft genome sequence of *Halomonas* sp. strain HAL1, a moderately halophilic arsenite-oxidizing bacterium isolated from gold-mine soil. *J. Bacteriol.* **194**, 199–200, <https://doi.org/10.1128/JB.06359-11>
- 75 Ates, O., Arga, K.Y. and Oner, E.T. (2013) The stimulatory effect of mannitol on levan biosynthesis: Lessons from metabolic systems analysis of *Halomonas smyrnensis* AAD6(T). *Biotechnol. Prog.* **29**, 1386–1397, <https://doi.org/10.1002/btpr.1823>
- 76 Aydin, B., Ozer, T., Oner, E.T. and Arga, K.Y. (2018) The genome-based metabolic systems engineering to boost levan production in a halophilic bacterial model. *OMICS* **22**, 198–209, <https://doi.org/10.1089/omi.2017.0216>
- 77 Kawata, Y., Shi, L.H., Kawasaki, K. and Shigeri, Y. (2012) Taxonomic characterization and metabolic analysis of the *Halomonas* sp. KM-1, a highly bioplastic poly(3-hydroxybutyrate)-producing bacterium. *J. Biosci. Bioeng.* **113**, 456–460, <https://doi.org/10.1016/j.jbiosc.2011.11.018>
- 78 Gao, J., Gao, N., Zhai, X. and Zhou, Y.J. (2021) Recombination machinery engineering for precise genome editing in methylotrophic yeast *Ogataea polymorpha*. *iScience* **24**, 102168, <https://doi.org/10.1016/j.isci.2021.102168>
- 79 Tan, D., Wu, Q., Chen, J.C. and Chen, G.-Q. (2014) Engineering *Halomonas* TD01 for low cost production of polyhydroxyalkanoates. *Metab. Eng.* **26**, 34–47, <https://doi.org/10.1016/j.ymben.2014.09.001>
- 80 Jiang, X.R., Yao, Z.H., Chen, G.-Q. et al. (2017) Controlling cell volume for efficient PHB production by *Halomonas*. *Metab. Eng.* **44**, 30–37, <https://doi.org/10.1016/j.ymben.2017.09.004>

- 81 Liu, H., Han, J., Liu, X., Zhou, J. and Xiang, H. (2011) Development of *pyrF*-based gene knockout systems for genome-wide manipulation of the archaea *Haloflex mediterranei* and *Haloarcula hispaica*. *J. Genet. Genomics* **38**, 261–269, <https://doi.org/10.1016/j.jgg.2011.05.003>
- 82 Chen, Y., Chen, X.Y., Du, H.T., Zhang, X., Ma, Y.M. and Chen, J.C. (2019) Chromosome engineering of the TCA cycle in *Halomonas bluephagenesis* for production of copolymers of 3-hydroxybutyrate and 3-hydroxyvalerate (PHBV). *Metab. Eng.* **54**, 69–82, <https://doi.org/10.1016/j.ymben.2019.03.006>
- 83 Hamedirad, M., Chao, R., Weisberg, S., Lian, J., Sinha, S. and Zhao, H. (2019) Towards a fully automated algorithm driven platform for biosystems design. *Nat. Commun.* **10**, 5150, <https://doi.org/10.1038/s41467-019-13189-z>
- 84 Gutierrez, T., Morris, G., Ellis, D., Mulloy, B. and Aitken, M.D. (2020) Production and characterisation of a marine *Halomonas* surface-active exopolymer. *Appl. Microbiol. Biotechnol.* **104**, 1063–1076, <https://doi.org/10.1007/s00253-019-10270-x>
- 85 Fernandez-Castillo, R., Rodriguez-Valera, F., Gonzalez-Ramos, J. and Ruiz-Berraquero, F. (1986) Accumulation of poly(beta-hydroxybutyrate) by halobacteria. *Appl. Environ. Microbiol.* **51**, 214–216, <https://doi.org/10.1128/AEM.51.1.214-216.1986>
- 86 Yu, L.P., Yan, X., Zhang, X., Chen, X.B., Wu, Q., Jiang, X.R. et al. (2020) Biosynthesis of functional polyhydroxyalkanoates by engineered *Halomonas bluephagenesis*. *Metab. Eng.* **59**, 119–130, <https://doi.org/10.1016/j.ymben.2020.02.005>
- 87 Jiang, X.R., Yan, X., Yu, L.P., Liu, X.Y. and Chen, G.-Q. (2021) Hyperproduction of 3-hydroxypropionate by *Halomonas bluephagenesis*. *Nat. Commun.* **12**, 1513, <https://doi.org/10.1038/s41467-021-21632-3>
- 88 Ouyang, P.F., Wang, H., Ivan, H. and Wu, Q. (2017) Increasing oxygen availability for improving poly(3-hydroxybutyrate) production by *Halomonas*. *Metab. Eng.* **45**, 20–31, <https://doi.org/10.1016/j.ymben.2017.11.006>
- 89 Ling, C., Qiao, G.Q., Shuai, B.W., Song, K.N., Yao, W.X., Jiang, X.R. et al. (2019) Engineering self-flocculating *Halomonas campaniensis* for wastewaterless open and continuous fermentation. *Biotechnol. Bioeng.* **116**, 805–815
- 90 Ling, C., Qiao, G.Q., Shuai, B.W., Karel, O., Yin, J., Xiang, R.J. et al. (2018) Engineering NADH/NAD⁺ ratio in *Halomonas bluephagenesis* for enhanced production of polyhydroxyalkanoates (PHA). *Metab. Eng.* **49**, 275–286, <https://doi.org/10.1016/j.ymben.2018.09.007>
- 91 Yim, H., Haselbeck, R., Niu, W., Pujol-Baxley, C., Burgard, A., Boldt, J. et al. (2011) Metabolic engineering of *Escherichia coli* for direct production of 1,4-butanediol. *Nat. Chem. Biol.* **7**, 445–452, <https://doi.org/10.1038/nchembio.580>
- 92 Zhou, H., Vonk, B., Roubos, J.A., Bovenberg, R.A.L. and Voigt, C.A. (2015) Algorithmic co-optimization of genetic constructs and growth conditions: application to 6-ACA, a potential nylon-6 precursor. *Nucleic Acids Res.* **43**, 10560–10570, <https://doi.org/10.1093/nar/gkv1071>
- 93 Meyer, A.J., Segall-Shapiro, T.H., Glassey, E., Zhang, J. and Voigt, C.A. (2018) *Escherichia coli* “Marionette” strains with 12 highly optimized small-molecule sensors. *Nat. Chem. Biol.* **15**, 196–205, <https://doi.org/10.1038/s41589-018-0168-3>
- 94 Wang, Y., Cheng, H., Liu, Y., Liu, Y., Wen, X., Zhang, K. et al. (2021) *In-situ* generation of large numbers of genetic combinations for metabolic reprogramming via CRISPR-guided base editing. *Nat. Commun.* **12**, 678, <https://doi.org/10.1038/s41467-021-21003-y>
- 95 Goormans, A.R., Snoeck, N., Decadt, T., Vermeulen, K., Peters, C., Coussement, P. et al. (2020) Comprehensive study on *Escherichia coli* genomic expression: does position really matter? *Metab. Eng.* **62**, 10–19, <https://doi.org/10.1016/j.ymben.2020.07.007>
- 96 Xu, X., Li, X., Liu, Y., Zhu, Y., Li, J., Du, G. et al. (2020) Pyruvate-responsive genetic circuits for dynamic control of central metabolism. *Nat. Chem. Biol.* **16**, 1261–1268, <https://doi.org/10.1038/s41589-020-0637-3>
- 97 Björkerötha, J., Campbell, K., Malinaa, C., Yu, R., Bartolomeo, F.D. and Nielsen, J. (2020) Proteome reallocation from amino acid biosynthesis to ribosomes enables yeast to grow faster in rich media. *Proc. Natl. Acad. Sci. U.S.A.* **17**, 21804–21812, <https://doi.org/10.1073/pnas.1921890117>
- 98 Yu, T., Zhou, Y.J.J., Huang, M., Liu, Q., Pereira, R., David, F. et al. (2018) Reprogramming yeast metabolism from alcoholic fermentation to lipogenesis. *Cell* **174**, 1549–1558, <https://doi.org/10.1016/j.cell.2018.07.013>
- 99 Gupta, A., Reizman, I.M.B., Reisch, C.R. and Prather, K.L.J. (2017) Dynamic regulation of metabolic flux in engineered bacteria using a pathway-independent quorum-sensing circuit. *Nat. Biotechnol.* **35**, 273–279, <https://doi.org/10.1038/nbt.3796>
- 100 Ren, Y.L., Ling, C., Ivan, H., Wu, Q. and Chen, G.-Q. (2018) Construction of *Halomonas bluephagenesis* capable of high cell density growth for efficient PHA production. *Appl. Microbiol. Biotechnol.* **102**, 4499–4510, <https://doi.org/10.1007/s00253-018-8931-7>