

Research article

A ~35 kDa polypeptide from insect cells binds to yeast ACS like elements in the presence of ATP

Suman K Dhar*¹, Neelima Mondal², Rajesh K Soni¹ and Gauranga Mukhopadhyay¹

Address: ¹Special Centre for Molecular Medicine, Jawaharlal Nehru University, New Delhi-110067, India and ²Brigham and Women's Hospital, Harvard Medical School, Boston, USA

E-mail: Suman K Dhar* - skdhar2002@yahoo.co.in; Neelima Mondal - nmondal@rics.bwh.harvard.edu; Rajesh K Soni - rajeshsoni@lycos.com; Gauranga Mukhopadhyay - garunga@hotmail.com

*Corresponding author

Published: 19 August 2002

Received: 28 March 2002

BMC Biochemistry 2002, 3:23

Accepted: 19 August 2002

This article is available from: <http://www.biomedcentral.com/1471-2091/3/23>

© 2002 Dhar et al; licensee BioMed Central Ltd. This article is published in Open Access: verbatim copying and redistribution of this article are permitted in all media for any non-commercial purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: The *S. cerevisiae* origin recognition complex binds to the ARS consensus sequence in an ATP dependent fashion. Recently, the yeast Cdc6 has been reported to have DNA binding activity. Conservation of replication proteins among different species strongly supports their functional similarity. Here we report the results of an investigation into the DNA binding activity of human Cdc6 protein. Cdc6 was expressed and purified from baculovirus infected Sf9 (*Spodoptera frugiperda*) insect cells as GST fusion protein (GST-Cdc6) and its DNA binding activity was tested.

Results: Partially purified fractions containing GSTCdc6 or GST showed an ACS binding activity in an ATP dependent manner. However, further purification revealed the presence of a putative 35 kDa insect cell protein (p35) which was found responsible for the DNA binding activity. A close match to the 9/11 bases of the ARS consensus sequence was sufficient for p35 binding activity. A DNA fragment from the human c-myc origin region containing yeast ACS like elements also showed p35 binding activity.

Conclusions: We have identified a *Spodoptera frugiperda* protein with ATP dependent DNA binding activity to ACS like elements. ACS like elements have been reported to be essential for ORC binding and replication initiation in yeast but their role in higher eukaryotes still remains elusive. Like the ARS consensus sequence elements of yeast, ACS like elements found in c-myc and lamin beta 2 origin regions may play similar roles in replication and indicate a conserved role for this DNA motif among eukaryotes.

Background

Proteins recognizing specific DNA sequences play an important role in the regulation of gene expression and in DNA replication. Nearly all eukaryotic genes transcribed by RNA Polymerase II for instance, contain the conserved

TATA box which is present upstream of the transcription start site. The TATA-box binding protein, a ~30 kDa component of the TFIID complex binds specifically to the heptanucleotide A and T residues [1] and forms the core of the transcription initiation complex. Additionally, many specific transcription factors bind to the upstream promoter




Figure 1
Purification of GST-Cdc6 and its DNA binding activity. (A) SDS-PAGE of GST-Cdc6 purified by glutathione agarose beads purification technique from Sf9 insect cells infected with baculovirus expressing GST-Cdc6. The polypeptide was visualised by comassie stain. Molecular mass markers are indicated on the right. (B) Gel retardation assays were performed in the presence of reduced glutathione eluted GST or GST-Cdc6 fusion protein. *S. cerevisiae* ARS1 fragment (240 bp) was end labelled using γ - ^{32}P ATP and incubated with GST or GST-Cdc6 as described in the materials and methods. Both GST and GST-Cdc6 showed strong band shifts.

in a sequence specific manner and regulate gene expression. For example in *Drosophila*, a heat-shock transcription factor (HSTF) can bind to consensus heat-shock response elements [2] and regulates expression of heat/stress inducible genes. Promoter activity in later branching eukaryotes is greatly modulated by enhancer and repressor sequences which have no activity of their own but are the targets of DNA binding proteins or protein complexes which can remodel the promoter chromatin to make it more or less accessible to RNA polymerase. Biochemical assays have shown that the action of ATP-dependent chromatin remodelling activities increase the accessibility of DNA within chromatin templates. *S. cerevisiae* SWI/SNF [3], Ino80 complex [4], *Drosophila* NURF [5] are examples of some high molecular weight chromatin remodelling factors which can facilitate transcription by binding to chromatinized DNA templates. However, none of the above chromatin remodelling factors binds to specific DNA sequences.

Unlike transcription, the role of sequence specific DNA binding proteins in eukaryotic DNA replication is not well characterized. In higher eukaryotes finding of specific DNA sequences essential for DNA replication has been elusive so far. In yeast *Saccharomyces cerevisiae*, a six-protein origin recognition complex binds to ARS consensus sequence (ACS) in a sequence specific manner [6]. Individual ORC subunits have not been demonstrated to show DNA binding activity *in vitro*. Recently, in an *in vitro* study ScCdc6 has been shown to bind double stranded DNA [7]. The minimal requirement for the binding of Cdc6 to DNA has been mapped within its N-terminal 47-amino acid sequence. *Saccharomyces pombe* ORC4 subunit has been reported to contain DNA binding activity by using its N-terminal AT hook region [8]. Neither ScCdc6 nor SpORC4 showed any sequence specific DNA binding activity. Recombinant six protein *Drosophila* ORC (DmORC) binds to ACE region of the *Drosophila* chorion gene [9]. *In vivo*, DmORC co-localized with the amplified chorion gene locus. In *Xenopus*, biochemical analysis of replication and cell cycle events using egg extracts has helped to understand the mechanism of eukaryotic DNA replication [10]. However, two dimensional gel electrophoresis analysis of the rDNA locus showed that replication initiated at all sites tested [11]. All six human homologs of yeast and *Drosophila* ORC subunits have been cloned and characterized [12,13]. Other replication proteins like Cdc6, Cdt1, MCMs, Cdc45 that are essential for initiation of DNA replication have also been reported [12]. Conservation of replication factors among higher eukaryotes suggests that functionally they may play similar roles.

In an attempt to identify DNA binding activity of human Cdc6, it was expressed and purified as a GST-Cdc6 fusion

protein from baculovirus infected Sf9 insect cells. Partially purified fractions (reduced glutathione eluate) containing GSTCdc6 or GST showed an ACS binding activity in an ATP dependent manner. The GSTCdc6 protein fraction contained both the GSTCdc6 and a 35 kDa *S. frugiperda* protein. The DNA binding activity was confined to a 35 kDa polypeptide. It was latter found that the p35 has an intrinsic affinity to GST. This polypeptide bound to yeast ACS like elements in the presence of ATP. 9/11 matches to ARS consensus sequence were found to be essential for this DNA binding activity both by gel shift assay as well as by in vitro foot printing assay. A DNA fragment containing 9/11 matches from human c-myc replication origin region also showed p35 binding activity suggesting that this polypeptide has intrinsic DNA binding activity. The implications of this DNA binding activity are discussed here.

Results

Partially purified protein fractions containing GSTCdc6 or GST contain an ACS binding activity

We infected Sf9 insect cells with the baculovirus expressing GSTCdc6. Cells were harvested 48 hours post infection and the proteins were extracted according to the procedures described in materials and methods. The GSTCdc6 protein was partially purified by pull down on glutathione beads (Fig. 1A). The partially purified protein was used in DNA binding assays with a 240 bp DNA fragment containing all three conserved boxes (A, B1 and B2) of the ARS consensus sequences (Fig. 7A & 7B). As a control, we used GST alone, which was purified using the same strategy used for GSTCdc6 purification. A DNA protein complex was formed in both the cases as evidenced by the retarded mobility of the free 32 P phosphate labelled probe (Fig 1B). The specificity of the DNA binding was examined in a competition reaction by increasing the amount of unlabeled DNA fragment containing ACS like elements. It was determined that the DNA-Protein complex could be competed efficiently by increasing amount of unlabelled ACS like DNA (20 \times , 50 \times , 100 \times and 200 \times respectively) (Fig 2). A \sim 350 bp DNA fragment from pBlue-Script KS⁺ (*Hinf*I digested and subsequently gel purified) of similar base composition did not compete with the complex formation in the Electrophoretic Mobility Shift Assay when added at a similar concentration indicating a degree of specificity in the DNA-Protein complex formation (Fig. 2).

An unidentified \sim 35 kDa protein from baculovirus infected insect cells is responsible for DNA binding activity

Partially purified fractions containing GSTCdc6 or GST showed DNA binding activity. To further fractionate the proteins present in the partially purified GSTCdc6 fraction the glutathione column was washed with 200 mM and 300 mM sodium chloride prior to GSTCdc6 elution with reduced glutathione. Western blot analysis using anti GST




Figure 2
Specificity of the ARS1 binding. Gel retardation assays were performed in the absence (-) or presence (+) of GST-Cdc6 using radiolabeled ARS1 probe as described in Fig. 1B. The binding reaction was competed either with cold ARS1 fragment (lanes 3–6) or a 350 bp fragment from the plasmid BlueScript KS(+) digested with *Hinf*I followed by gel purification (lanes 7–10). Free probe and the band shifts are indicated by arrow.

antibodies (Fig. 3A) revealed that neither 200 mM nor 300 mM fractions contained GSTCdc6 (Fig. 3A). GSTCdc6 protein was present only in the proteins eluted by reduced glutathione. Proteins released by different salt fractionation were separated on an SDS-PAGE and visualized by silver stain (Figure 3B). A prominent band of molecular mass \sim 35 kDa was visualised both in the 200 mM and 300 mM salt eluate.

To check whether the DNA binding activity was due to the presence of \sim 35 kDa band or GSTCdc6 itself, gel shift assays were performed either using 50 ng \sim 35 kDa protein obtained from the salt wash or GSTCdc6 by itself (Fig. 3C). Salt eluate gave a strong band shift which was identical with the band shift found with GSTCdc6 found in Fig. 2. GSTCdc6 eluted from the beads following high salt wash failed to give any band shift suggesting that the \sim 35 kDa polypeptide was responsible for the DNA binding activity.

To further test whether the presence of p35 is absolutely required for DNA binding activity, the 300 mM salt eluate was dialysed against low salt buffer H/0.15 and then passed through Superose 12 gel filtration column. Each fraction was checked for DNA binding activity by gel retardation assay using a 32 P labelled DNA fragment containing ACS elements. DNA binding activity was found only




Figure 3
Identification of p35. (A) Western blot analysis of salt eluate and reduced GSH eluate using anti GST antibody. GST-Cdc6 beads were washed using wash buffer containing NaCl (200 mM and 300 mM respectively). Beads following salt wash were treated with reduced glutathione and released proteins were run in SDS-PAGE along with salt eluate followed by western blot analysis. GST-Cdc6 was found only in GSH eluate. (B) Silver staining of salt eluate: Salt eluates were run in SDS-PAGE and proteins were visualised by silver stain. A distinct band of p35 was found both in 200 mM and 300 mM salt eluate. Molecular mass markers are shown on the right. (C) DNA binding activity of p35. Gel shift assays were performed using salt wash (300 mM NaCl eluate) and GSH eluate following 300 mM salt wash of GST-Cdc6 beads. ARS 240 bp DNA fragment was used as probe. 300 mM salt eluate gave a strong band shift whereas GSH eluate failed to do so though it contained GST-Cdc6 as shown in figure 3A.

in high molecular weight fraction (~670 kDa) (Fig. 4A). Proteins present in the gel filtration fractions in the high molecular weight range (fractions 13–17) were separated by SDS-PAGE followed by silver stain. Surprisingly, p35 was found to be present only in the fraction 15 which contains the DNA binding activity (Fig. 4B). The presence of p35 in the same fraction containing the DNA binding ac-




Figure 4
Purification of p35 by gel filtration chromatography. (A) Gel retardation assay. 300 mM salt eluate (as shown in fig. 3B) was dialysed against Buffer H/0.15 and passed through Superose-12 Gel filtration column. Fractions 13–25 were tested for ARS 240bp DNA binding activity. Only fraction 15 showed a band shift compared to the free probe. Molecular mass markers are shown on the top. (B) Silver stain of gel filtration fractions. 0.25 ml of each gel filtration fractions (fraction 13–17) were precipitated using trichloro acetic acid (TCA). The samples were run in SDS-PAGE and proteins were visualised by silver stain. p35 was found only in the fraction 15.

tivity strongly suggests that p35 is responsible for the binding activity.

DNA binding activity is ATP dependent

One of the hallmarks of yeast ORC binding to yeast ARS consensus sequences is its ATP dependence [6]. We were interested to see whether the DNA binding activity of the ~35 kDa protein is ATP dependent or not. Gel shift assays were performed either in the absence or in the presence of increasing amount of ATP in the reaction mixture (Fig. 5). In the absence of ATP, a very weak binding was observed whereas with increasing amount of ATP strong binding was detected. There is a threshold of ATP concentration (6 mM and onwards) which stimulated the binding remarkably. A nonhydrolysable ATP analog, ATPγS was used in the binding reaction to see whether ATP hydrolysis is required for this binding. With increasing amount of ATPγS, the band shift was completely inhibited suggesting that ATP hydrolysis is required for this DNA binding activity.




Figure 5
ATP hydrolysis is required for the DNA binding activity of p35. Gel shift assays were performed using ^{32}P labelled ARS fragment either in the presence of ATP (lanes 3–7) or ATP γ S (lanes 8–12) or both (lanes 13–14). A strong band shift was observed in the presence of ATP whereas ATP γ S completely inhibited the binding. The inhibition can be reversed by adding back ATP in the reaction mixture.

This was confirmed by adding ATP back in the reaction mixture when ATP γ S was already present in the reaction. Under these reaction conditions, increasing amount of ATP again stimulated the DNA binding activity even in the presence of ATP γ S suggesting that ATP γ S can be competed with ATP and it is the ATP hydrolysis which is essential for this DNA binding activity (Fig. 5, lanes 13 and 14).

p35 binds to A, B1 and B2 boxes of ARS consensus sequences as revealed by footprinting assay

After establishing the fact that a protein of approximately 35 kDa binds to DNA fragment containing ARS consensus sequences, the exact site of binding of the protein on the DNA was mapped by copper-phenanthroline footprinting assay. Unlike DNaseI, 1,10-phenanthroline-cuprous complex is a small chemical probe which can demark the boundaries of the protected region clearly. The protein was bound to a 5' ^{32}P labelled 240 bp DNA fragment containing yeast ARS consensus sequence and separated from the free DNA by gel shift assay. The gel was then treated with copper-phenanthroline reagent as described in materials and methods. The bound and unbound DNA was purified and allowed to run in a sequencing gel (Fig. 6). On the T-rich strand (bottom strand) two protected regions were observed. The first region entirely covered the A box of the ARS consensus sequence and the second region covered the overlapping regions of B1 and B2. Therefore the 35 kDa protein has a strong affinity to bind to A, B1 and B2 boxes of the ARS consensus sequences. The long stretch of protection could be due to multimeric form of p35 or

could be due to the formation of a higher order nucleoprotein complex.

p35 binds to A, B1 and B2 boxes and mutation in these boxes abolish DNA binding activity

Previous studies of the structure of ARS1 in both plasmid and chromosome contexts have shown that it contains one essential DNA element, A, that includes a perfect (11/11) match to the ARS consensus sequence (ACS, Fig. 7A), and three additional elements, B1, B2, and B3, with 9 out of 11 bases match to the ACS that are also important for ARS function [14]. We were interested to see whether A, B1 and B2 boxes were sufficient to allow p35 binding activity. We used the p21N protein (N terminal 150 bases of coding region of p21) [15] as a control. This DNA when incubated with \sim 35 kDa protein does not form a DNA protein complex (Fig 7D). The DNA fragment was divided into four subfragments (a, b, c and d; Fig. 7B) and subcloned in the middle of p21N fragment. Fragment 'a' does not contain any ACS sequence whereas fragments b, c and d contain at least one ACS like element. All the ACS containing sub fragments (b, c, d) showed a mobility shift (Fig. 7D), which suggests that at least one ACS like element (either 11/11 match or 9/11 match to ARS consensus sequence) is essential and sufficient for p35 binding activity. This was further confirmed by using a subfragment d mutated at the ACS motif (Fig. 7C). Subfragment d as shown in Fig 7D lane 10 can bind strongly to the 35 kDa protein, however mutation altering the A and Ts of the core ACS sequence to G and Cs (Fig 7C) to generate the dmut oligonucleotide results in the abolition of its ability to bind p35. Therefore the ACS sequence is essential for the p35-DNA protein complex formation.

35 kDa polypeptide binds to ARS consensus sequence found in c-myc origin of replication

In *S. cerevisiae*, ARS elements have been implicated to be important both for ORC binding and origin function [6]. We looked for the availability of such sequences in known human origins of replication like c-myc, lamin beta 2 and Dnmt1. An origin of replication was mapped previously by nascent strand abundance analysis within 2.0 kb zone immediately upstream of c-myc gene [16]. Detailed analysis of the 2.0 kb upstream sequences revealed the presence of two ACS like elements separated by 148 nucleotides (AAAAGATAAAG and AAAAGAAAAA). A 300 bp DNA fragment containing both the ACS elements was amplified by polymerase chain reaction and the product was used subsequently for p35 binding studies.

A strong band shift was observed (Fig. 8) which could be competed out using a 50 bp long double stranded oligo containing two ACS like elements (oligo 'b', Fig. 7B). Increasing amount of \sim 65 bp long unrelated (non specific) double stranded oligo did not have any effect on this




Figure 6
Copper-Phenanthroline footprint analysis of ARS DNA fragment bound to p35. ARS DNA fragment was 5' end labelled at the T rich strand and incubated with p35 (300 mM salt eluate) and finally subjected to gel retardation assay. The gel was then treated with copper-phenanthroline mixture and DNA from bound (B) and unbound (U) fractions were extracted from gel, denatured and loaded in a sequencing gel. A+G ladder was also loaded to figure out the position of footprint. The relative positions of A, B1 and B2 boxes were indicated. The protected regions are shown by solid lines and asteric (*). The hypersensitive site is denoted by arrowhead.

binding activity suggesting that binding of p35 to c-myc origin region is specific. Interestingly, analysis of Lamin beta 2 origin of replication region also revealed the presence of two ACS (9/11 match) like elements [17]. Further studies are required to find out whether p35 also binds to lamin beta 2 origin region.

p35-ACS interaction is sensitive to high salt, temperature and EDTA

The stability of p35-ACS DNA complex was further tested either by changing NaCl concentration in the reaction mixture, or by shifting reaction temperature or by adding EDTA. The DNA binding activity was found to be sensitive to NaCl concentration (Fig. 9). Strong band shift was obtained up to 0.2 M NaCl. NaCl concentration at 0.5 M and above completely inhibited the binding activity. Higher temperature also showed a drastic effect on the binding activity. Normal DNA binding activity was observed up to 42°C. Temperature higher than 42°C completely abolished the binding activity suggesting that the off rate of p35 from DNA is much faster at higher temperature. Finally inclusion of EDTA in the reaction mixture inhibited the binding reaction suggesting that the divalent cations are essential for this binding activity.

Discussion

Few proteins have been reported in the literature, which are capable of binding to DNA in a sequence specific ATP dependent manner. Although transcription factors bind to specific DNA sequences, the binding activity is not dependent on ATP hydrolysis. In contrast, chromatin remodelling factors like SWI/SNF, ISW1, BRG1 facilitate transcription from chromatinised templates in the presence of ATP [18]. However, these factors do not bind to specific DNA sequences. In eukaryotic DNA replication, sequence specific ATP dependent DNA binding activity has been demonstrated in yeast *S. cerevisiae* where ORC, a six polypeptide complex binds to yeast ARS consensus sequence in an ATP dependent manner [6]. The binding sites for other ORCs are not very clear at present. DmORC binds the critical elements of well-characterized, chromosome III amplification domain (ACE3 and ori- β , though the precise sequence recognized by DmORC within ACE3 and ori- β have not been identified [9]. Studies of both ScORC [19] and DmORC [20] indicate the ATP binding by Orc1p is required for DNA binding. However ATP hydrolysis is not required for DNA binding for both the cases suggesting that ATP hydrolysis may be required for further downstream processes. Chromatin immunoprecipitation (ChIP) studies have demonstrated the association of SpORC with *S. pombe* origins [21] and human ORC with the EBV *OriP* [22-24]. Whether ORC binds to these sequences directly or indirectly with the help of other proteins are subject to in vitro DNA binding assays using purified ORC proteins.




Figure 7

At least one ACS like element is required for p35 binding. (A) ARS consensus sequence. (B) 240 bp yeast *S. cerevisiae* ARS1 fragment containing several functional elements required for ARS activity (Boxes A, B1, B2 and B3 respectively). Several oligos spanning the entire ARS1 as indicated in the figure were designed and subsequently cloned into p21N fragment. Double stranded oligo 'a' (ARSa) does not contain any ARS like element. Oligo 'b' (ARSb) contains one 11/11 match and one 9/11 match to ACS element. Oligo 'c' (ARSc) and oligo 'd' (ARSd) contain one 9/11 match to ACS element each. In the fragment dmut, 'A's and 'T's within the core ARS were mutated to 'G's and 'C's. (C) Nucleotide sequences of ARSd and ARSdmut. (D) Gel shift assay was performed either using ARSwT (lanes 1,8), or p21N (lanes 2,3 and 9) or the constructs derived from p21N by cloning fragments a-d and dmut (lanes 4-7, 10,11) as described in (B). p35 does not bind to p21N or p21N + ARSa or p21N + ARSdmut. All the other DNA fragments showed DNA binding activity.




Figure 8
p35 binds to DNA fragment containing ACS like elements from c-myc origin region. A 300 bp DNA fragment containing two ACS like elements (10/11 matches) was PCR amplified using primers specific for the zone of replication from c-myc locus. PCR product was end labelled using γ -³²P ATP and used for gel shift assays. Competition reactions were performed by using different quantities (50×, 100× and 200×) of 50 bp long double stranded oligos (oligo 'b', Fig. 7B) in the reaction mixtures corresponding to lanes 3–5. Lanes 6–8 contained 65 bp double stranded DNA fragment (plasmid BlueScript KS(+)) digested with *Hin*I followed by gel purification).

In this study, we report a ~35 kDa protein from the baculovirus infected Sf9 insect cells that binds to yeast ACS sequences in an ATP dependent fashion. p35 was purified as high salt (300 mM NaCl) eluate from the GST-Cdc6 beads. GST-Cdc6 eluted from the beads following high salt wash failed to show any DNA binding activity (Fig. 3C) whereas high salt eluate containing only p35 showed strong DNA binding activity suggesting that p35 not Cdc6 is responsible for the binding activity. This experiment was repeated several times and always the protein preparations containing p35 showed DNA binding activity.

p35 has an intrinsic affinity to GST moieties. Sf9 insect cells were infected with baculovirus expressing GST alone. The cell lysate was allowed to bind to GST beads. High salt eluate (300 mM NaCl) from GST beads was tested for

DNA binding activity. Surprisingly, we observed a very similar band shift as obtained previously using high salt eluate from GSTCdc6 (data not shown). Further, high salt eluate from GSTORC2 and GSTORC4 (GST fusion protein containing human origin recognition complex subunit 1 and 2 respectively) also showed DNA binding activity (data not shown). Therefore, it can be concluded that p35 interacts with GST and high salt concentration is required to disrupt this interaction. The nature and the specificity of the interaction between GST and p35 are not clear at this moment. It is important to note that majority of the p35 bound to GST or GST fusion proteins are released mostly at high salt concentration (300 mM) allowing us to get rid of most of the impurities by stringent washing of the GST beads with buffer containing 250 mM NaCl.




Figure 9
Effect of salt concentration, temperature and EDTA on p35 gel shift assay. Gel retardation assay was performed either using no extra salt or different concentration (lanes 1–5) of NaCl. Band shift was completely inhibited at 0.5 M NaCl concentration and above. Effect of temperature on binding reaction was tested by incubating the reaction mixture (lanes 6–9) at different temperature. Binding reaction was completely inhibited at 55°C and above. All the other reactions were carried out at room temperature. Effect of bivalent cations were tested by incubating the reaction mixture in the presence of 50 mM and 100 mM EDTA (lanes 10–11). Presence of EDTA completely inhibited the binding reaction.

A weak DNA binding activity was found till 4 mM ATP concentration (Fig. 5). A strong stimulation was obtained at 6 mM ATP concentration. It is possible that p35 is purified as ATP bound form but an associated weak ATPase activity does not allow it to give a strong binding activity. It is suggested that 6 mM ATP concentration may be suffi-

cient to overcome this inhibitory effect. ATP hydrolysis is essential for DNA binding activity of p35 since ATPγ S, a nonhydrolysable analog of ATP completely inhibited the binding activity (Fig. 5). It is possible that ATP is required for strand opening which needs to be further explored. An insect cell factor, polyhedrin promoter binding protein

has been reported previously, capable of binding to AT rich DNA sequence [25]. However, the reported DNA binding activity was unusual in a sense that the activity was heat and salt concentration resistant. 100 mM EDTA concentration did not affect the DNA binding activity. The DNA binding activity reported in this study was found to be temperature, EDTA and salt concentration sensitive (Fig. 9) suggesting that this polypeptide is completely different from the polyhedrin promoter binding protein.

p35 showed a strong DNA binding affinity towards ACS like elements. This was confirmed first by using unlabeled specific competitor DNA which completely abolished the binding of ^{32}P labelled probe. Competition using unrelated DNA did not affect the DNA binding activity. We took p21N, which normally does not bind to p35, to further test the binding specificity. Introduction of a single ACS like element (9/11 match) in p21N (ARSc and ARSd) showed a strong band shift (Fig. 7D) suggesting that only one ACS like element is sufficient for p35 binding. This was further confirmed by making mutations in the ARSd fragment. The resulting ARSdmut did not show any p35 binding activity suggesting that the ACS like element itself but not the adjacent sequences are essential for the p35 binding activity. However, a systematic mutational analysis of ACS like elements will be required to explore the exact binding specificity of p35 towards ACS like elements. ACS elements are normally AT rich. However, p35 did not bind to random AT rich sequences. p35 did not bind to p21N + ARSa which contains ARSa oligo (78% AT rich) with no ACS like element. Surprisingly, p35 showed strong binding activity in the presence of the oligo ARSb (64% AT rich), ARSc (66% AT rich) and ARSd (74% AT rich) respectively (Figure 7D). ARSb, c and d contain at least one ACS like element (Fig. 7B). Finally the copper phenanthroline footprint analysis confirmed that p35 binds to A and B1-B2 boxes of the ARS1 DNA fragment. At this moment, the function of p35 is not very clear. It may play major role(s) in the transcription of certain insect cell genes. It may as well be responsible for DNA replication. The fact that it binds to yeast ACS and to a DNA fragment from c-myc origin of replication region containing yeast ACS like elements in an ATP dependent manner further strengthen the hypothesis. *S. cerevisiae* origin recognition complex (ORC) binds to ARS consensus sequences in an ATP dependent fashion and this binding is essential for both origin function and activity. It is interesting to note that a huge six protein origin recognition complex binds to yeast ARS1 whereas p35, a small protein is showing same kind of protection as evidenced by footprint analysis (Fig. 6). We believe that p35 forms an oligomeric structure or it maintains a multimeric form which may explain the wide footprint over the ARS1 fragment. The presence of p35 in high molecular weight fraction (~670 kDa) following superose 12 gel filtration chroma-

tography strengthen this hypothesis and clearly suggests that p35 forms an oligomeric structure.

ARS consensus sequence has been found near the vicinity of c-myc, lamin beta 2 and Dnmt1 replication origin [26]. Therefore, Identification and characterization of this protein from insect cells and finding its human counterpart will greatly help in elucidating its possible function in DNA replication.

Conclusions

The data presented here leads to the identification and characterization of a polypeptide from insect cells with ATP dependent DNA binding activity. This is an important and unique observation. In *S. cerevisiae*, ACS elements have been reported to be essential for ORC binding and replication initiation. Yeast ACS like elements found in c-myc and lamin beta 2 origin region may play similar roles in replication initiation. However, it is also possible that p35 is a transcription factor which may facilitate transcription of some insect cell genes. Further characterization of p35 from insect cells and finding its human homolog will be very helpful to dissect its functional role in replication and/or transcription.

Materials and Methods

Plasmid construction

Cloning of human Cdc6 cDNA is described elsewhere. Coding sequence of human Cdc6 was cloned in pFastBac-GST vector (Life Technologies, Inc.) to express GST fusion protein. A 240 bp DNA fragment from *S. cerevisiae* ARS1 chromosomal DNA replication origin containing all the key elements including boxes A, B1, B2 and B3 was subcloned in pBlueScript KS(+) between *EcoRI* and *HindIII*. Subsequently *EcoRI-HindIII* fragment was end labelled using $\gamma^{32}\text{P}$ ATP and used either for gel shift assay or copper phenanthroline footprint assay. p21N, (N terminal ~150 bp of p21) was previously cloned in pBlueScript KS(+) between *EcoRI* and *HindIII* sites.

Complementary oligos corresponding to ARSa, b, c, d and dmut (Fig. 7B) were synthesized (~50 bases in length) and subsequently annealed to get double stranded oligos. p21N/KS(+) construct contains only one *StuI* site which is present within the p21N insert. All the annealed double stranded oligos were cloned into the *StuI* site using blunt end ligation. The sequences of ARSa, b, c and d are followed:

ARSa: ttagtttttcggtttactaatcgtaatagaaatgtagaacaataaaatgt

ARSb: tctaaaatacaaatctagaaaatacgaacgaaaagtttccggacgtc-cgt

ARSd: cggacgtcggttcagtgtttggatgaattttatgatgagtcattat

ARSd: tgagtcattattggataaagaatcgtaaaaactgctttaacgataaaa

Plasmid containing 2.5 kb DNA fragment from c-myc origin region was a kind gift from Michael Leffak, Wright State University Ohio. Forward and reverse PCR primers (5'-gaagaaaaactctcttttc-3' and 5'-atttgctgggtgaaaaatg-3' respectively) were used to amplify 300 bp region containing two ACS like elements.

Expression of GSTCdc6 and GST in insect cells and purification

Baculoviruses were produced from the recombinant pFB-GST plasmid using Bac-to-Bac expression system (Life Technologies Inc.). Sf9 cells (Invitrogen) were infected with the pFB-GSTCdc6 or pFBGSTbaculovirus according to the manufacturers' recommendations. Cells were harvested 48 hours post-infection. The cell pellet was washed once in cold phosphate-buffered saline and subsequently resuspended in hypotonic lysis buffer (10 mM Tris.Cl, pH 7.9, 10 mM KCl, 1.5 mM MgCl₂, 1 mM phenylmethylsulfonyl fluoride, 2 µg/ml pepstatin, 2 µg/ml leupeptin, 5 µg/ml aprotinin, 1 mM dithiothreitol). The cell suspension was homogenized in a Dounce homogenizer using a B-type pestle followed by centrifugation at 3000 rpm for 7 min. The pellet containing the nuclei was lysed in buffer H/0.15 (50 mM HEPES/KOH, pH 7.5, 150 mM KCl, 0.02% Nonidet P-40, 5 mM magnesium acetate, 1 mM EDTA, 1 mM EGTA, 10% glycerol, 1 mM phenylmethylsulfonyl fluoride, 2 µg/ml pepstatin, 2 µg/ml leupeptin, 5 µg/ml aprotinin, 1 mM dithiothreitol). The resulting suspension was subjected to ammonium sulphate precipitation (starting with 10% followed by 30% and finally 50%). The pellet after the 50% ammonium sulphate cut was resuspended in buffer H/0.0 (no salt) and then dialyzed overnight against buffer H/0.15. The dialyzed sample was then bound to GST beads (Sigma) and washed three times with buffer H*/0.15 (containing 150 mM NaCl instead of 150 mM KCl). Proteins were eluted using reduced glutathione elution buffer (50 mM Tris.Cl, pH 8.0, 20 mM reduced glutathione, 0.01% Nonidet P-40, 100 mM NaCl).

Immunoblotting and Silver stain

Anti-GST polyclonal antibodies were purchased from Santa Cruz Biotechnologies. Western blotting technique was carried out using standard protocol. The silver stain protocol is described elsewhere [27].

Gel retardation assay

Gel retardation assay was performed with slight modification of the protocol used by Mukhopadhyay et al [28]. The DNA fragments to be used for gel retardation assay were endlabeled with $\gamma^{32}\text{P}$ ATP. The binding reactions were performed in 20 µl of T buffer (50 mM Tris-HCl, pH 7.4, 50 mM KCl, 50 mM NaCl, 10 mM MgCl₂, 0.1 mM

EDTA, 0.5 mM DTT, 30 µg/ml BSA) supplemented with 5 mM ATP and 6% (v/v) glycerol for most of the reactions. The mixture was incubated at 37°C for 10 min and loaded directly on a 5% polyacrylamide gel in TBE buffer (89 mM Tris base, 89 mM boric acid, 2.5 mM EDTA, pH 8.3). The gel was run at 150 V for 2 hours, dried and autoradiographed.

Copper-phenanthroline footprint assay

The Copper-phenanthroline footprint assay was performed essentially by using the protocol described by Kuwabara et al [29]. The DNA-protein gel is run in the absence of free radical scavengers as described in gel retardation assay. The gel is placed in 200 ml of 50 mM Tris-HCl, pH 8.0. The gel is further incubated for 10 minutes in a solution containing equal volume of solution A (40 mM 1,10 Phenanthroline monohydrate in 100% EtOH and 9 mM Cupric sulphate mixed with equal volume followed by 1:10 dilution with water) and solution B (0.5% 3-Merceptopropionic acid in water). Finally the gel is soaked in solution C (28 mM 2,9 Dimethyl-1,10 Phenanthroline in 100% EtOH) for 2 min. The gel is washed twice in deionised water. After the pre-treatment of the gel, it is autoradiographed and the retarded band is cut from the gel and placed in an eppendorf tube. The DNA is eluted from the gel slice, denatured and loaded in a sequencing gel. The sequencing gel is fixed, dried and the bands were visualised by autoradiography.

Authors' contributions

SKD designed, performed and co-ordinated the whole study. NM participated in insect cell culture and helped in making and amplification of baculovirus expressing GST-Cdc6 and GST. RKS helped in making the figures and drafting the manuscript. GM helped in analysing the data and critically reviewed the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors acknowledge Dr Anindya Dutta, Harvard Medical School for conceiving the idea and providing full support to carry out this work. Authors greatly acknowledge Carol Gilchrist, University of Virginia for thorough editing of the manuscript. David Garcia Quintana helped to amplify the c-myc origin of replication by polymerase chain reaction. Jeffrey D Parvin helped to perform the Copper-phenanthroline footprint assay. RKS is thankful to Council of Scientific and Industrial Research, India for providing a fellowship. SKD acknowledges University Grant Commission, India for providing funds (University Excellence programme).

References

1. Guarente L, Bermingham-McDonogh O: **Conservation and evolution of transcriptional mechanisms in eukaryotes.** *Trends Genet* 1992, **8**:27-32
2. Dynan WS, Tjian R: **Control of eukaryotic messenger RNA synthesis by sequence-specific DNA-binding proteins.** *Nature* 1985, **316**:774-8
3. Peterson CL, Tamkun JW: **The SWI-SNF complex: a chromatin remodeling machine?** *Trends Biochem Sci* 1995, **20**:143-6
4. Shen X, Mizuguchi G, Hamiche A, Wu C: **A chromatin remodeling complex involved in transcription and DNA processing.** *Nature* 2000, **406**:541-4

5. Mizuguchi G, Wu C: **Nucleosome remodeling factor NURF and in vitro transcription of chromatin.** *Methods Mol Biol* 1999, **119**:333-42
6. Bell SP, Stillman B: **ATP-dependent recognition of eukaryotic origins of DNA replication by a multiprotein complex.** *Nature* 1992, **357**:128-34
7. Feng L, Wang B, Driscoll B, Jong A: **Identification and characterization of *Saccharomyces cerevisiae* Cdc6 DNA-binding properties.** *Mol Biol Cell* 2000, **11**:1673-85
8. Lee JK, Moon KY, Jiang Y, Hurwitz J: **The *Schizosaccharomyces pombe* origin recognition complex interacts with multiple AT-rich regions of the replication origin DNA by means of the AT-hook domains of the spOrc4 protein.** *Proc Natl Acad Sci U S A* 2001, **98**:13589-94
9. Austin RJ, Orr-Weaver TL, Bell SP: **Drosophila ORC specifically binds to ACE3, an origin of DNA replication control element.** *Genes Dev* 1999, **13**:2639-49
10. Blow JJ: **Control of chromosomal DNA replication in the early *Xenopus* embryo.** *EMBO J* 2000, **20**:3293-7
11. Hyrien O, Mechali M: **Chromosomal replication initiates and terminates at random sequences but at regular intervals in the ribosomal DNA of *Xenopus* early embryos.** *EMBO J* 1993, **12**:4511-20
12. Kelly TJ, Brown GW: **Regulation of chromosome replication.** *Annu Rev Biochem* 2000, **69**:829-80
13. Dhar SK, Dutta A: **Identification and characterization of the human ORC6 homolog.** *J Biol Chem* 2000, **275**:34983-8
14. Marahrens Y, Stillman B: **A yeast chromosomal origin of DNA replication defined by multiple functional elements.** *Science* 1992, **255**:817-23
15. Chen J, Jackson PK, Kirschner MW, Dutta A: **Separate domains of p21 involved in the inhibition of Cdk kinase and PCNA.** *Nature* 1995, **374**:386-8
16. Vassilev L, Johnson EM: **An initiation zone of chromosomal DNA replication located upstream of the c-myc gene in proliferating HeLa cells.** *Mol Cell Biol* 1990, **10**:4899-904
17. Dimitrova DS, GM, Demarchi F, Biamonti G, Riva S, Falaschi A: **In vivo protein-DNA interactions at human DNA replication origin.** *Proc Natl Acad Sci U S A* 1996, **93**:1498-503
18. Havas K, Whitehouse I: **ATP-dependent chromatin remodeling activities.** *Cell Mol Life Sci* 2001, **58**:673-82
19. Klemm RD, Austin RJ, Bell SP: **Coordinate binding of ATP and origin DNA regulates the ATPase activity of the origin recognition complex.** *Cell* 1997, **88**:493-502
20. Chesnokov I, Remus D, Botchan M: **Functional analysis of mutant and wild-type *Drosophila* origin recognition complex.** *Proc Natl Acad Sci U S A* 2001, **8**:11997-2002
21. Ogawa Y, Takahashi T, Masukata H: **Association of fission yeast Orp1 and Mcm6 proteins with chromosomal replication origins.** *Mol Cell Biol* 1999, **19**:7228-36
22. Dhar SK, Yoshida K, Machida Y, Khaira P, Chaudhuri B, Wohlschlegel JA, Leffak M, Yates J, Dutta A: **Replication from oriP of Epstein-Barr virus requires human ORC and is inhibited by geminin.** *Cell* 2001, **106**:287-96
23. Chaudhuri B, Xu H, Todorov I, Dutta A, Yates JL: **Human DNA replication initiation factors, ORC and MCM, associate with oriP of Epstein-Barr virus.** *Proc Natl Acad Sci U S A* 2001, **98**:10085-9
24. Schepers A, Ritzi M, Bousset K, Kremmer E, Yates JL, Harwood J, Difley JF, Hammerschmidt W: **Human origin recognition complex binds to the region of the latent origin of DNA replication of Epstein-Barr virus.** *EMBO J* 2001, **20**:4588-602
25. Burma S, Mukherjee B, Jain A, Habib S, Hasnain SE: **An unusual 30-kDa protein binding to the polyhedrin gene promoter of *Autographa californica* nuclear polyhedrosis virus.** *J Biol Chem* 1994, **269**:2750-7
26. Araujo FD, Knox JD, Ramchandani S, Pelletier R, Bigey P, Price G, Zyzf M, Zannis-Hadjopoulos M: **Identification of initiation sites for DNA replication in the human dnmt1 (DNA-methyltransferase) locus.** *J Biol Chem* 1999, **274**:9335-41
27. Dunn MJ, Giacca M: **Detection of proteins in polyacrylamide gels using an ultrasensitive silver staining technique.** *Methods Mol Biol* 1994, **32**:113-8
28. Mukhopadhyay G, Carr KM, Kaguni JM, Chatteraj DK: **JM, DK. Open-complex formation by the host initiator, DnaA, at the origin of P1 plasmid replication.** *EMBO J* 1993, **12**:4547-54
29. Kuwabara MD, Sigman DS: **Footprinting DNA-protein complexes in situ following gel retardation assays using 1,10-phenanthroline-copper ion: *Escherichia coli* RNA polymerase-lac promoter complexes.** *Biochemistry* 1987, **26**:7234-8

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com