

POSTER PRESENTATIONS

Open Access

Plasmodium falciparum cytoadherence to ICAM-1 is associated with cerebral malaria

Lucy B Ochola¹, Bethsheba R Siddondo¹, Harold Ocholla¹, Siana Nyka³, Eva N Kimani¹, Thomas N Williams¹, Johnstone O Makale¹, Anne Liljander⁴, Britta C Urban^{1,2}, Pete Bull¹, Tadge Szeszak², Kevin Marsh¹, Alister G Craig^{2*}

From Parasite to Prevention: Advances in the understanding of malaria
Edinburgh, UK. 20-22 October 2010

The pathology of severe malaria is in part related to the pro-inflammatory nature of the host response but a number of other factors are also thought to be involved, including the interaction between infected erythrocytes and endothelium. This phenotype involves a range of host receptors and the parasite-derived variant antigen PfEMP1, which is expressed on the surface of the infected erythrocyte membrane. Previous studies have suggested a role for ICAM-1 in the pathology of cerebral malaria, although these were inconclusive. In this study we measured the binding to CD36 and ICAM-1 of patient isolates from varying clinical syndromes under static and flow conditions. We also used mutant ICAM-1 proteins to characterise the key contact residues on ICAM-1 and produce a detailed binding phenotype. Our results show that increased binding to CD36 is associated with uncomplicated malaria while ICAM-1 adhesion under flow conditions is raised in parasites from cerebral malaria cases. The pattern of ICAM-1 binding has also been investigated using mutant ICAM-1 proteins and indicates that isolates from severe malaria are biased towards a binding signature also seen with ITO4, a laboratory isolate selected for binding on human endothelium with similar receptor expression to that seen in the brain.

Author details

¹KEMRI/Wellcome Trust Research Programme, P. O. Box 230, 80108 Kilifi, Kenya. ²Liverpool School of Tropical Medicine, Pembroke Place, Liverpool L3 5QA, UK. ³Dares Salaam University College of Education, Faculty of Science, P.O. Box 2329, Dares Salaam, Tanzania. ⁴Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden.

Published: 20 October 2010

²Liverpool School of Tropical Medicine, Pembroke Place, Liverpool L3 5QA, UK

Full list of author information is available at the end of the article

doi:10.1186/1475-2875-9-S2-P27

Cite this article as: Ochola et al.: *Plasmodium falciparum* cytoadherence to ICAM-1 is associated with cerebral malaria. *Malaria Journal* 2010, 9 (Suppl 2):P27.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

