

Redefining Liver Fibrosis Risk Assessment in Indians with Type 2 Diabetes: New FIB-4 Score Cutoff for Optimizing Sequential Assessment with Transient Elastography

Rajat Deb, Soumik Goswami, Nilanjan Sengupta, Arjun Baidya, Vibhu R. Khare, Joydip Datta, Mousumi Das¹, Debes Ray¹

Departments of Endocrinology and ¹Biochemistry, Nil Ratan Sircar Medical College and Hospital, Kolkata, West Bengal, India

Abstract

Introduction: Liver fibrosis in metabolic dysfunction-associated steatotic liver disease (MASLD) is strongly related to hepatic and extrahepatic outcomes. Clinically Significant Liver Fibrosis (CSLF) screening using FIB-4 score is mandated in all T2D patients. The existing FIB-4 cutoff of 1.3 is derived from Western studies and could differ for Indians. Hence, we aimed to determine the FIB-4 cutoff to rule out Transient Elastography (TE) proven CSLF among Indians with T2D. **Methods:** 551 individuals with T2D underwent laboratory tests for FIB-4 calculation and transient elastography (TE) to detect CSLF defined as Liver Stiffness Measurement (LSM) $\geq$ 8kPa. The Receiver Operative Characteristic (ROC) curve was used to determine the optimum cutoff value of FIB4 to rule out CSLF. **Results:** 129 (23.4%) of 551 T2D patients in our cohort had CSLF. We found that a FIB-4 of 1.5 rules out CSLF with a sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and Diagnostic Accuracy (DA) of 82.9%, 79.9%, 55.7%, 93.8%, and 80.6%, respectively, compared with a FIB4 of 1.3 which has values of 91.5%, 67.3%, 46.1%, 96.2%, and 72.9%, respectively. **Conclusion:** A FIB-4 cutoff of 1.5 rather than 1.3 is suggested for Indian subjects with T2DM and needs to be validated in further large multicenter prospective studies, preferably with histopathology as the gold standard.

Keywords: Clinically significant liver fibrosis, FIB4 score, MASLD, transient elastography, type 2 diabetes mellitus

INTRODUCTION

Metabolic dysfunction-associated steatotic liver disease (MASLD) encompasses a spectrum from simple hepatic steatosis to nonalcoholic steatohepatitis (NASH), liver fibrosis, cirrhosis, and hepatocellular carcinoma in individuals with Type 2 diabetes mellitus (T2D).^[1] Fibrosis is a critical prognostic factor in MASLD, linked to adverse liver-related outcomes, cardiovascular complications, and mortality.^[2] Current guidelines recommend screening for Clinically Significant Liver Fibrosis (CSLF) using the Fibrosis Index Based on Four Factors (FIB-4) score as an accessible and cost-effective initial assessment.^[1,3,4] The Fibrosis-4 (FIB-4) score is calculated based on age, aspartate aminotransferase (AST), alanine aminotransferase (ALT), and platelet count—variables that are readily available and commonly measured.^[5] Transient elastography (TE) serves as the standard of care for confirming hepatic fibrosis severity when FIB-4 scores suggest the need for further evaluation.^[6,7] Currently, a FIB-4 score threshold of 1.3, derived from Western population studies, is used to

exclude CSLF in individuals with T2D.^[1,3,4] This study aims to determine the optimal FIB-4 cutoff value for excluding TE-confirmed CSLF in Indian patients with T2D.

MATERIALS AND METHODS

Study population

This cross-sectional descriptive study evaluated 551 consecutive patients with type 2 diabetes mellitus (T2DM) attending the Department of Endocrinology from March to December 2023.

Address for correspondence: Dr. Rajat Deb,
Department of Endocrinology, Nil Ratan Sircar Medical College and Hospital,
Kolkata - 700 014, West Bengal, India.
E-mail: rajatdeb2411.rd@gmail.com

Submitted: 05-Nov-2024

Revised: 12-Jan-2025

Accepted: 27-Feb-2025

Published: 29-Apr-2025

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Deb R, Goswami S, Sengupta N, Baidya A, Khare VR, Datta J, *et al.* Redefining liver fibrosis risk assessment in Indians with type 2 diabetes: New FIB-4 score cutoff for optimizing sequential assessment with transient elastography. *Indian J Endocr Metab* 2025;29:237-41.

Access this article online

Quick Response Code:



Website:
<https://journals.lww.com/indjem/>

DOI:
10.4103/ijem.ijem_457_24

Participants were aged between 35 and 65 years to align with FIB-4 score efficacy, as it performs less accurately in individuals outside this age range.^[8] Exclusion criteria included male patients consuming over 30 g of alcohol per day and female patients consuming more than 20 g per day, as well as patients with secondary causes of hepatic steatosis (e.g., chronic corticosteroid use), positive hepatitis B surface antigen or antihepatitis C virus antibody, and histories of other chronic liver diseases or thyroid dysfunction. Patients on pharmacotherapy for MASLD, including Pioglitazone or Saroglitazar, were also excluded. The diagnosis of T2DM among participants was based on the American Diabetes Association (ADA) criteria.^[9] These included fasting plasma glucose levels ≥ 126 mg/dL, 2-hour plasma glucose ≥ 200 mg/dL during an oral glucose tolerance test, HbA1c $\geq 6.5\%$, or random plasma glucose ≥ 200 mg/dL in individuals with classic symptoms of hyperglycemia. Participants with both newly diagnosed and long-standing T2DM were included, as no minimum duration of diabetes was required for inclusion in the study. In addition, participants managing their diabetes through lifestyle interventions alone, without pharmacotherapy, were eligible for inclusion. While detailed pharmacotherapy data were not collected, it was confirmed that none of the participants were on GLP-1 receptor agonists (GLP-1RA), Pioglitazone, or Saroglitazar, although some were on sodium-glucose cotransporter-2 inhibitors (SGLT-2i) [Figure 1].

The study protocol received approval from the Institutional Ethics Committee. Written informed consent was obtained from all participants before enrolment. The study was conducted in accordance with the principles of the Declaration of Helsinki.

Sample size calculation

The calculation of sample size was based on prevalence and sensitivity; the estimated sample size required for this study was around 101 patients with liver fibrosis, assuming an unlimited sample size and 5% precision.

The formula used to estimate the sample size was –

$$N = \frac{Z^2 \times \text{Sensitivity} \times (1 - \text{sensitivity})}{L^2 \times (1 - \text{prevalence})}$$

$$N = \frac{(1.96)^2 \times 0.95 \times (1 - 0.95)}{(0.05)^2 \times (1 - 0.28)} = 101$$

Z = Standard normal variant (1.96)

L = Precision 5% at 95% CI

Sensitivity of FIB4,^[10] = 95%

Prevalence of CSLF in T2DM,^[11] = 28%

Based on prevalence from previous studies, we needed to screen only 375 T2D patients to achieve our sample size target but we succeeded in screening as many as 600 T2D patients for our study.

Clinical assessment

Comprehensive clinical assessments included anthropometric measurements of body weight and height. Body mass index (BMI) was calculated as weight (kg) divided by height




Figure 1: Derivation of the study cohort. LSM, liver stiffness measurement; T2D, type 2 diabetes; CSLF, clinically significant liver fibrosis

squared (m^2). Blood for complete blood count (CBC) and liver function test (LFT) were collected after TE and analyzed on the same day in the institute's central laboratory.

Liver stiffness measurements

Liver stiffness was assessed using Transient Elastography (TE), which served as the gold standard in this study for evaluating the diagnostic accuracy of the FIB-4 score. A single operator performed the TE procedure on the right liver lobe, utilizing the FibroScan® Mini+ 430 model by Echosens through the intercostal spaces, with participants positioned flat on their backs and their right arm maximally abducted. Liver stiffness results were reported in kiloPascals (kPa) and liver stiffness measurement (LSM) of ≥ 8.0 kPa was used as the threshold for Clinically Significant Liver Fibrosis (CSLF), while $\text{LSM} \geq 12.0$ kPa indicated advanced fibrosis. These thresholds were chosen based on previous studies, which have demonstrated high positive predictive values (PPVs) for the presence of clinically significant fibrosis.^[12]

Calculation of FIB4 score

The FIB4 score was calculated by using the formula:^[5]

$$\text{FIB-4} = \frac{\text{Age (years)} \times \text{AST (U/L)}}{\text{Platelets (} 10^9 / \text{L)} \times \text{ALT}^{1/2} (\text{U/L})}$$

Statistical analysis

Data were analyzed using the SPSS 20.0 software (IBM, NY, USA). Categorical variables were expressed as frequencies and percentages, while quantitative variables were reported as

means with standard deviations or as medians with interquartile ranges for non-normally distributed data. Continuous data were presented as mean $\pm$ SD and results of categorical measurements were expressed in terms of frequency and percentage. The Receiver Operative Characteristic (ROC) curve was used to determine the modified cutoff points of the FIB4 index that could best discriminate between the absence and presence of significant fibrosis.

Ethical aspect

The study protocol received approval from the Institutional Ethics Committee (reg no. ECR/609/Instt/WB/2014/RR20) of Nil Ratan Sircar Medical College and Hospital, Kolkata Memo No. NRSMEC/IEC/06/2023, approved on 11.01.2023. Written informed consent was obtained from participants for their involvement in the study and the use of data for research and educational purposes. The study adhered to the current version of the World Medical Association Declaration of Helsinki (2013).

RESULTS

A total of 551 T2D patients were included in the analysis after applying the inclusion and exclusion criteria, while 49 participants were excluded for various reasons [detailed in Figure 1]. The cohort's mean age was 49.9 ± 7.7 years, with a predominance of female participants (58.6%). The average BMI was 24.9 ± 3.9 kg/m², with 21.2% of participants categorized as overweight and 47.1% as obese. The average liver stiffness measurement (LSM) was 7.3 ± 6.1 kPa, indicative of varying degrees of hepatic involvement.

Baseline characteristics of participants

129 (23.4%) of 551 T2D patients in our cohort had CSLF. The detailed baseline characteristics of all participants, including demographics, biochemical parameters, and comorbidities, are summarized in Table 1.

ROC curve analysis and FIB-4 cutoff determination

ROC analysis demonstrated an area under the curve (AUROC) of 90.10% [Figure 2] for the FIB-4 score in discriminating CSLF. The proposed FIB-4 cutoff value of 1.5 offered a balance between sensitivity (82.9%) and specificity (79.9%), resulting in a positive predictive value (PPV) of 55.7% and a negative predictive value (NPV) of 93.8%, with a diagnostic accuracy of

80.6%. For reference, the performance of conventional cutoff values and their comparative sensitivity, specificity, PPV, and NPV values are presented in Table 2.

DISCUSSION

The global rise in MASLD prevalence has increased the need for effective, noninvasive screening tools for assessing liver fibrosis, particularly in patients with T2D, a group with elevated liver disease risk. This study aimed to validate and potentially adjust the FIB-4 score cutoff to better suit the Indian population with T2D, where Western-derived thresholds may not be fully applicable.

The new FIB-4 cutoff of 1.5 proposed in this study aligns closely with findings from *Torres et al.*,^[10] who reported a 1.49 cutoff for MRE-based fibrosis screening in a Brazilian cohort with MASLD and T2D. This higher cutoff improved specificity in both studies, reducing false positives and limiting unnecessary testing. Notably, studies from Western populations often recommend a lower threshold, around 1.3, for similar screening purposes.^[13] This discrepancy may stem from population differences in metabolic profiles, as Asian populations, including the Indian demographic, are known to develop liver-related complications at lower BMI thresholds and with distinct metabolic risk factors compared with Western counterparts.^[14,15] These findings support the need for population-specific adjustments in diagnostic criteria.

The proposed FIB-4 cutoff of 1.5 offers a more targeted approach, particularly valuable in settings with limited resources, where access to advanced tests like TE and MRE may be restricted. By enhancing specificity, this cutoff helps reserve such resources for patients at higher risk of CSLF, reducing the clinical and financial burden associated with unnecessary testing. This aligns with existing guidelines suggesting that only T2D patients exceeding the FIB-4 threshold should undergo further testing.^[1,3,16]

This study's strength lies in its large, Indian T2D cohort, making it one of the first to propose an adjusted FIB-4 threshold

Table 1: Baseline characteristics of all participants, with CSLF and without CSLF

Characteristics	All participants	LSM <8	LSM $\geq$ 8
No. of participants	551	422	129
Age (years)	49.9 $\pm$ 7.7	49.8 $\pm$ 7.7	50.2 $\pm$ 7.8
BMI (kg/m ²)	24.9 $\pm$ 3.9	24.5 $\pm$ 3.7	26.6 $\pm$ 4.3
Diabetes Duration (years)	7.4 $\pm$ 5.3	7.6 $\pm$ 5.1	6.9 $\pm$ 5.9
Platelet count ($\times 10^9$ /L)	215.8 $\pm$ 77.0	233.2 $\pm$ 74.3	158.9 $\pm$ 55.6
AST (U/L)	28.2 $\pm$ 16.9	24.2 $\pm$ 10.3	41.2 $\pm$ 25.8
ALT (U/L)	27.9 $\pm$ 19.1	25.6 $\pm$ 16.3	35.9 $\pm$ 24.6

Data are presented as mean ($\pm$ SD). BMI, body mass index; AST, Aspartate aminotransferase; ALT, Alanine aminotransferase;




Figure 2: ROC curve of FIB4 score for patients with T2D

Table 2: Analysis and comparison between the sensitivity, specificity, PPV, and NPV of proposed cutoff points and those already used. AUROC, area under receiver operating characteristic curve; PPV, positive predictive value; NPV, negative predicted value

Cutoff	AUROC	Sensitivity	Specificity	PPV	NPV
Conventional					
<1.3	90.10%	91.5%	67.3%	46.1%	96.2%
Modified					
<1.2	90.10%	96.1%	59.0%	41.8%	98.0%
<1.4	90.10%	86.8%	73.9%	50.5%	94.8%
<1.5 [#]	90.10%	82.9%	79.9%	55.7%	93.8%
<1.6	90.10%	76.7%	85.1%	61.1%	92.3%
<1.7	90.10%	71.3%	88.6%	65.7%	90.9%

[#]proposed cutoff value with optimum sensitivity, specificity, NPV, and PPV

specific to this population. Our study has several limitations that warrant consideration. Being cross-sectional, the study was designed to evaluate associations rather than establish causal relationships between FIB-4 cutoff values and the risk of clinically significant liver fibrosis (CSLF). In addition, as the data were collected from a single tertiary care center, the generalizability of our findings to the broader Indian population may be limited, especially given regional variations in metabolic profiles and healthcare access. Although transient elastography (TE) is a validated non-invasive method for assessing liver fibrosis, it is not equivalent to liver biopsy (the gold standard for diagnosing CSLF) and the absence of histopathological confirmation limits the definitive validation of our findings.^[17,18] While our sample size was sufficient for addressing the primary objective, the study was not powered for subgroup analyses, such as stratification by BMI categories, limiting insights into the role of obesity in CSLF risk. Information regarding the use of anti-hyperglycemic drugs among study participants was limited. While none of the participants were on GLP-1 receptor agonists (GLP-1RA), Pioglitazone, or Saroglitazar; some participants were on sodium-glucose cotransporter-2 inhibitors (SGLT-2i), the exact number was not recorded. The potential effects of SGLT-2i on transient elastography findings could not be fully accounted for. Despite these limitations, our study offers valuable preliminary insights into a population-specific FIB-4 cutoff for assessing CSLF in Indian patients with type 2 diabetes, and future multicenter studies involving diverse populations and incorporating histopathology as the gold standard addressing these limitations could further validate and refine our findings.

CONCLUSION

A FIB-4 cutoff of 1.5 rather than 1.3 is suggested for Indian subjects with T2DM and needs to be validated in further large multicenter prospective studies, preferably with histopathology as the gold standard.

Acknowledgement

We want to thank all study participants for agreeing to take part in the study. We would also like to thank the scientific committee and institutional ethical committee for approval

of the study. No funding or financial sponsorship was taken from external sources. Resources and facilities available in our institute were used for the current study.

Authors' contribution

Rajat Deb and Soumik Goswami were involved in the conception and design of the study, along with the drafting of initial manuscript. Rajat Deb, Vibhu Ranjan Khare, Joydip Datta, and Mousumi Das acquisition of data and its analysis. Nilanjan Sengupta, Arjun Baidya, and Debes Ray contributed to the revision of article and final approval of the version to be published.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Data availability statement

The data supporting this study is not publicly available due to ethical considerations and ongoing analysis. However, data can be accessed upon request by contacting the corresponding author via email.

REFERENCES

1. Rinella ME, Neuschwander-Tetri BA, Siddiqui MS, Abdelmalek MF, Caldwell S, Barb D, *et al.* AASLD practice guidance on the clinical assessment and management of nonalcoholic fatty liver disease. *Hepatology* 2023;77:1797-835.
2. Targher G, Byrne CD, Tilg H. NAFLD and increased risk of cardiovascular disease: Clinical associations, pathophysiological mechanisms, and pharmacological implications. *Gut* 2020;69:1691-705.
3. Wattacheril JJ, Abdelmalek MF, Lim JK, Sanyal AJ. AGA clinical practice update on the role of noninvasive biomarkers in the evaluation and management of nonalcoholic fatty liver disease: Expert review. *Gastroenterology* 2023;165:1080-8.
4. Cusi K, Isaacs S, Barb D, Basu R, Caprio S, Garvey WT, *et al.* American Association of Clinical Endocrinology clinical practice guideline for the diagnosis and management of nonalcoholic fatty liver disease in primary care and endocrinology clinical settings: Co-sponsored by the American Association for the Study of Liver Diseases (AASLD). *Endocr Pract* 2022;28:528-62.
5. Sterling RK, Lissen E, Clumeck N, Sola R, Correa MC, Montaner J, *et al.* Development of a simple noninvasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology* 2006;43:1317-25.

6. Talwalkar JA, Kurtz DM, Schoenleber SJ, West CP, Montori VM. Ultrasound- based transient elastography for the detection of hepatic fibrosis: Systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2007;5:1214-20.
7. Friedrich-Rust M, Poynard T, Castera L. Critical comparison of elastography methods to assess chronic liver disease. *Nat Rev Gastroenterol Hepatol* 2016;13:402-11.
8. McPherson S, Hardy T, Dufour JF, Petta S, Romero-Gomez M, Allison M, *et al.* Age as a confounding factor for the accurate non-invasive diagnosis of advanced NAFLD fibrosis. *Am J Gastroenterol* 2017;112:740-51.
9. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33(Suppl 1):S62-9.
10. Torres L, Schuch A, Longo L, Valentini BB, Galvão GS, Luchese E, *et al.* New FIB-4 and NFS cutoffs to guide sequential non-invasive assessment of liver fibrosis by magnetic resonance elastography in NAFLD. *Ann Hepatol* 2023;28:100774.
11. Kuchay MS, Choudhary NS, Mishra SK, Bano T, Gagneja S, Mathew A. Prevalence of clinically relevant liver fibrosis due to non-alcoholic fatty liver disease in Indian individuals with type 2 diabetes. *JGH Open* 2021;5:915-22.
12. Koehler EM, Plompen EP, Schouten JN, Hansen BE, Darwish Murad S, Taimr P, *et al.* Presence of diabetes mellitus and steatosis is associated with liver stiffness in a general population: The Rotterdam study. *Hepatology* 2016;63:138-47.
13. Xiao G, Zhu S, Xiao X, Yan L, Yang J, Wu G. Comparison of laboratory tests, ultrasound, or magnetic resonance elastography to detect fibrosis in patients with nonalcoholic fatty liver disease: A meta-analysis. *Hepatology* 2017;66:1486-501.
14. Misra A, Soares MJ, Mohan V, Anoop S, Abhishek V, Vaidya R, *et al.* Body fat, metabolic syndrome and hyperglycemia in South Asians. *J Diabetes Complications* 2018;32:1068-75.
15. Shah AG, Lydecker A, Murray K, Tetri BN, Contos MJ, Sanyal AJ, NASH Clinical Research Network. Comparison of noninvasive markers of fibrosis in patients with nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol* 2009;7:1104-12.
16. European Association for the Study of the Liver (EASL), European Association for the Study of Diabetes (EASD), European Association for the Study of Obesity (EASO). EASL–EASD–EASO clinical practice guidelines on the management of metabolic dysfunction-associated steatotic liver disease (MASLD). *J Hepatol* 2024;81:492-542.
17. Wong VW, Vergniol J, Wong GL, Foucher J, Chan HL, Le Bail B, *et al.* Diagnosis of fibrosis and cirrhosis using liver stiffness measurement in nonalcoholic fatty liver disease. *Hepatology* 2010;51:454-62.
18. Siddiqui MS, Vuppalanchi R, Van Natta ML, Hallinan E, Kowdley KV, Abdelmalek M, *et al.* Vibration-controlled transient elastography to assess fibrosis and steatosis in patients with nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol* 2019;17:156-63.e2.