

## Technical Note

# Limited vertical dural opening for lesions of the vermis, 4<sup>th</sup> ventricle, and distal PICA segments

Patrick Graupman<sup>1,2</sup>, Archie Defillo<sup>1</sup>, Leslie Nussbaum<sup>1</sup>, Eric S. Nussbaum<sup>1</sup>

<sup>1</sup>National Brain Aneurysm Center, Saint Joseph's Hospital, West 10<sup>th</sup> Street, <sup>2</sup>Gillette Children's Specialty Healthcare, East University Ave, St. Paul, Minnesota, USA

E-mail: \*Patrick Graupman - [pgraupman@gillettechildrens.com](mailto:pgraupman@gillettechildrens.com); Archie Defillo - [adefillo@healtheast.org](mailto:adefillo@healtheast.org); Leslie Nussbaum - [lnussbaum@comcast.net](mailto:lnussbaum@comcast.net); Eric S. Nussbaum - [lnussbaum@comcast.net](mailto:lnussbaum@comcast.net)

\*Corresponding author

Received: 24 September 12

Accepted: 28 September 12

Published: 27 November 12

### This article may be cited as:

Graupman P, Defillo A, Nussbaum L, Nussbaum ES. Limited vertical dural opening for lesions of the vermis, 4<sup>th</sup> ventricle, and distal PICA segments. *Surg Neurol Int* 2012;3:141. Available FREE in open access from: <http://www.surgicalneurologyint.com/text.asp?2012/3/1/141/103881>

Copyright: © 2012 Graupman P. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Background:** Lesions of the vermis and 4<sup>th</sup> ventricle are commonly addressed through a midline suboccipital approach. Most neurosurgeons use either a Y-shaped or a curvilinear dural opening in this setting. Although these approaches offer a wide intraoperative surgical exposure, in occasion, the dural opening is difficult to repair primarily, often necessitating the use of a patch, which may increase the risk for development of CSF fistula. We are describing our experience with a limited, vertical, midline, dural opening for approaches to the vermis, tentorium, 4<sup>th</sup> ventricle, and distal posterior-inferior cerebellar artery (PICA) segments as an alternative to the classic Y-shaped or curvilinear incision.

**Methods:** We report our experience with a limited vertical midline durotomy in five patients with posterior fossa lesions. The lesions treated included a PICA dissecting aneurysm, three metastatic lesions (located in the vermian, floor of the 4<sup>th</sup> ventricle, and undersurface of the tentorium cerebelli), and one intra-axial tumor (ependymoma). All patients were positioned prone, and the lesions were accessed without difficulty through a limited, vertical, midline durotomy.

**Results:** Mass lesions and vascular abnormalities located from the midline as far lateral as the outlet foramina of the 4<sup>th</sup> ventricle can be accessed comfortably via a limited midline dural opening when combined with microsurgical techniques, and the use of a frameless Stealth Station Neuronavigation System (SSNS) [Medtronic Sofamor Danek, Inc., Memphis, TN]. By doing this, simple primary dural closure was achieved with a single running absorbable suture without tension in each case.

**Conclusion:** In our experience, a suboccipital linear dural opening appears to be as effective as the more traditional Y-shaped incision, yet allows for quicker and easier primary dural repair.

**Key Words:** Dural opening, median suboccipital approach, posterior fossa

### Access this article online

#### Website:

[www.surgicalneurologyint.com](http://www.surgicalneurologyint.com)

#### DOI:

10.4103/2152-7806.103881

#### Quick Response Code:



## INTRODUCTION

Surgical approaches to the posterior fossa are designed to achieve adequate intraoperative exposure with minimal

brain retraction and manipulation. The combination of intraoperative microsurgical techniques with the SSNS allows surgeons to precisely navigate and dissect within the posterior fossa corridors and subarachnoid

spaces. Traditionally, neurosurgeons have used either a “Y”-shaped or a curvilinear dural opening to access the posterior fossa.<sup>[1-5]</sup> In this report, we describe the usefulness of a simple vertical, linear durotomy to access the midline cerebellar regions, facilitating dural repair. With this technique, exposure of the vermian, tentorial, 4<sup>th</sup> ventricle structures and distal PICA segments was achieved without undue brain retraction or manipulation. Dural repair was simpler and quicker than with the more traditional Y-shaped dural opening.

## MATERIALS AND METHODS

We report our experience with a limited vertical midline durotomy in five patients with post-fossa lesions. The lesions treated included a PICA dissecting aneurysm, three metastatic lesions (located in the vermian, floor of the 4<sup>th</sup> ventricle, and undersurface of the tentorium cerebelli), and one intra-axial tumor (ependymoma). All patients were positioned prone, and the lesions were accessed without difficulty through a limited, vertical, midline durotomy.

## RESULTS

### Illustrative cases

#### Case 1

A 72-year-old female with previously diagnosed metastatic lung cancer presented with a newly found, enlarging, vermian lesion that failed stereotactic radiosurgical treatment. A decision was made to proceed with surgical resection, and a midline suboccipital approach was planned. A limited linear durotomy was performed and the SSNS was utilized to localize the tumor. The vermis was entered via a paramedian approach, sparing the vertically oriented vermian veins. Following gross total resection of the lesion, the dura was closed without difficulty. The postoperative course was uneventful, and there was no evidence of cerebrospinal fluid (CSF) leak or accumulation.

#### Case 2

A 24-year-old male was found to have a large 4<sup>th</sup> ventricular tumor, requiring surgical resection. Following a midline suboccipital exposure, the dura was opened using a limited, midline incision. The cerebellar tonsils were separated from the tumor region inferiorly, and PICA segments were mobilized and dissected away laterally. The floor of the 4<sup>th</sup> ventricle was entered. The tumor, which was fairly vascular in nature, was dissected using its natural cleavage plane and removed as a single specimen. Primary dural repair was performed, and the postoperative course was without incident [Figure 1a-d].


#### Case 3

A 66-year-old male presented with a coma producing subarachnoid hemorrhage due to a ruptured dissecting


aneurysm of the right PICA. He underwent microsurgical wrapping and remote distal outflow occlusion of the aneurysm. Surgically, the aneurysm was accessed via a midline suboccipital craniotomy, with bone work involving exposure and removal of the C1 arch. Dural opening was performed in the midline, using a linear incision, followed by the lifting or “tenting” of the dural edges superiorly. The right PICA in the 4<sup>th</sup> ventricular region was identified and followed proximally. The aneurysm was extremely friable in nature, which prompted the decision to wrap it. The distal outflow occlusion was carried out beyond the PICA tonsillar point, just after the last evident brainstem perforators. There were no postoperative complications.

### Technique description

The patient positioning on the operation table is crucial. We prefer the prone position for exposing the posterior fossa via a median suboccipital approach, because the cerebellum falls away from the brainstem surface, opening the subvermian surgical corridor. A midline skin lineal incision is made using the external occipital protuberance and the spinous process of C2 as a landmark [Figure 2]. The paraspinal muscles and the avascular plane of the ligamentum nuchae are divided in the midline with diathermy coagulation. Separating the fascia in this manner is of paramount importance, since this creates a fascial plane, which is easily approachable to closure, preventing future CSF leakage. The occipital bone is exposed from the superior nuchal line to the region of the foramen magnum. Partial removal of the arch of C1 is not often needed, but may allow additional exposure to the cervicomedullary junction after dural opening. During the bone work, it is crucial to avoid injuring the vertebral arteries and dura mater. After performing the craniotomy, a midline, vertical durotomy is made extending from the



**Figure 1: (a and b) Pre-operative sagittal and coronal MRI views demonstrating a heterogeneous, nodular mass that fills the 4<sup>th</sup> ventricle and extends through the foramen magnum into the upper cervical canal. (c) Postoperative sagittal MRI showing gross total resection of the lesion. (d) Intraoperative photomicrograph shows the minimally invasive durotomy, with sutures “tenting” the dural edges**



**Figure 2: Illustration demonstrating the traditionally “Y”-shaped versus the straight incision. A midline, vertical, dural opening is made extending from the region of the foramen magnum to below the bony margin superiorly**

region of the foramen magnum to below the bony margin superiorly. The free dural edges are “tenting” with sutures [Figure 3]. At this point, the microscope is brought into the field, allowing us to open the cervicomedullary cistern.


## DISCUSSION

There are several benefits of Y-shaped dural opening when performing a medial suboccipital approach, which consist of wide surgical field exposure, avoidance of injury to the occipital sinus, good control of the sinusoid vessels around the foramen magnum and easy access to the cerebellomedullary cistern for CSF drainage.<sup>[1,3,4]</sup>

Our description demonstrates that a lineal dural opening may provide the same intraoperative benefits as the Y-shaped incision. Surgical dissection of the posterior fossa corridors and CSF space can be achieved without significant brain manipulation and retraction.

Our indications for this kind of practice include elective lesions located in the vicinity of the cerebellar vermis, floor of the 4<sup>th</sup> ventricle, cortical, telovelar, tonsilomedullary and lateral medullary segments of the PICA.<sup>[1-5]</sup>

Although the amount of exposure may be slightly diminished by a midline vertical dural opening, in our experience, this exposure has been more than sufficient for lesions as far lateral as the foramina of Luschka.



**Figure 3: Graphic picture showing the lifting or “tenting” of the free dural edges with sutures, and exposure of the vermician, tentorial, 4<sup>th</sup> ventricle structures, and distal PICA**

When choosing a linear durotomy, we have used intraoperative navigation techniques to carefully guide our opening as well as the surgical trajectory. By working through the natural arachnoid planes and releasing CSF as needed, adequate exposure can be achieved without significant brain manipulation and retraction. The vermician, tentorial, 4<sup>th</sup> ventricle structures, and distal PICA segments, can be accessed without difficulty [Figure 3].

## CONCLUSION

In our experience, a suboccipital linear dural opening appears to be as effective as the more traditional Y-shaped incision, yet allows for quicker and easier primary dural repair.

## REFERENCES

1. Algird A, Morcos JJ, Macdonald RL. Vertebral and Posterior inferior Cerebellar Artery Aneurysms. In: Macdonald RL, editor: *Neurosurgical Operative Atlas*. 2<sup>nd</sup> ed. Vascular Neurosurgery. New York: Thieme; 2008. p. 81-5.
2. Koos WT, Spetzler RF, Lang J, editors. *Color Atlas of Microneurosurgery*. Vol 2. Cerebrovascular Lesions. Stuttgart, Germany: Georg Thieme-Verlag; 1997.
3. Lawton MT, Vates EG, Spetzler RF. Surgical Approaches for Posterior Circulation Aneurysms. In: Winn RH, editor: *Youmans Neurological Surgery*. 5<sup>th</sup> ed. Philadelphia: Elsevier Saunders; 2004. p. 1971-2005.
4. Perneczky A, Reisch, R. *Keyhole Approaches in Neurosurgery*. Vol 1. Concept and Surgical Technique. New York: Springer Wien; 2008.
5. Wright DC, Wilson CB. Surgical treatment of basilar aneurysms. *Neurosurgery* 1979;5:325-33.