

2-(1*H*-Benzotriazol-1-yl)-3-(2,6-dichlorophenyl)-1-phenylpropan-1-ol

Özden Özel Güven,^a Seval Çapanlar,^a Simon J. Coles^b and Tuncer Hökelek^{c*}

^aDepartment of Chemistry, Zonguldak Karaelmas University, 67100 Zonguldak, Turkey, ^bDepartment of Chemistry, Southampton University, SO17 1BJ Southampton, England, and ^cDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 12 August 2011; accepted 24 August 2011

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.042; wR factor = 0.108; data-to-parameter ratio = 17.9.

The asymmetric unit of the title compound, $\text{C}_{21}\text{H}_{17}\text{Cl}_2\text{N}_3\text{O}$, contains two crystallographically independent molecules with similar conformations. The benzotriazole ring is oriented at dihedral angles of 30.61 (5) and 43.36 (5)°, respectively, to the phenyl and dichlorophenyl rings in one molecule, and 32.25 (5) and 41.04 (5)° in the other. The dihedral angles between the phenyl and dichlorophenyl rings are 66.38 (7) and 66.14 (6)° in the two molecules. An intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond links the benzotriazole ring and phenylpropanol unit in each molecule. In the crystal, weak intermolecular $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds link the molecules into chains along the a axis. $\pi-\pi$ stacking between the dichlorophenyl rings [centroid-centroid distances = 3.809 (1) and 3.735 (1) Å] may further stabilize the crystal structure.

Related literature

For the biological activity of azole compounds, see: Cozzi *et al.* (1994) and of triazole derivatives, see: Jin *et al.* (2006). For related structures, see: Özel Güven *et al.* (2007, 2008, 2010).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{17}\text{Cl}_2\text{N}_3\text{O}$
 $M_r = 398.28$
 Triclinic, $P\bar{1}$

$a = 9.3894$ (2) Å
 $b = 9.4947$ (2) Å
 $c = 21.2687$ (3) Å

$\alpha = 91.415$ (2)°
 $\beta = 92.324$ (2)°
 $\gamma = 90.406$ (1)°
 $V = 1893.90$ (6) Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.36$ mm⁻¹
 $T = 120$ K
 $0.4 \times 0.4 \times 0.3$ mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 2007)
 $T_{\min} = 0.866$, $T_{\max} = 0.897$

40396 measured reflections
 8697 independent reflections
 7206 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.108$
 $S = 1.07$
 8697 reflections

487 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.43$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.73$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1A}\cdots\text{N2}$	0.82	2.56	3.1254 (19)	127
$\text{O1}'-\text{H1B}\cdots\text{N2}'$	0.82	2.52	3.0881 (19)	128
$\text{C19}-\text{H19}\cdots\text{N3}^{\text{i}}$	0.93	2.54	3.420 (2)	158
$\text{C19}'-\text{H19}'\cdots\text{N3}^{\text{ii}}$	0.93	2.53	3.410 (2)	159

Symmetry codes: (i) $x + 1, y, z$; (ii) $x, y + 1, z$.

Data collection: COLLECT (Nonius, 1998); cell refinement: DENZO (Otwinowski & Minor, 1997) and COLLECT; data reduction: DENZO and COLLECT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON (Spek, 2009).

The authors acknowledge Zonguldak Karaelmas University Research Fund (project No: 2010-13-02-06).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5297).

References

- Cozzi, P., Giordani, A., Menichincheri, M., Pillan, A., Pinciroli, V., Rossi, A., Tonani, R., Volpi, D., Tamburin, M., Ferrario, R., Fusar, D. & Salvati, P. (1994). *J. Med. Chem.* **37**, 3588–3604.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Jin, Z., Hu, Y., Huo, A., Tao, W., Shao, L., Liu, J. & Fang, J. (2006). *J. Organomet. Chem.* **691**, 2340–2345.
- Nonius (1998). COLLECT. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Özel Güven, Ö., Bayraktar, M., Coles, S. J. & Hökelek, T. (2010). *Acta Cryst.* **E66**, o959.
- Özel Güven, Ö., Erdoğan, T., Çaylak, N. & Hökelek, T. (2007). *Acta Cryst.* **E63**, o3638.
- Özel Güven, Ö., Tahtacı, H., Coles, S. J. & Hökelek, T. (2008). *Acta Cryst.* **E64**, o1254.
- Sheldrick, G. M. (2007). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2011). E67, o2510 [doi:10.1107/S1600536811034738]

2-(1*H*-Benzotriazol-1-yl)-3-(2,6-dichlorophenyl)-1-phenylpropan-1-ol

Ö. Özel Güven, S. Çapanlar, S. J. Coles and T. Hökelek

Comment

Azole compounds have important biological activities. In literature 2-(1*H*-imidazol-1-yl)-1,3-diphenylpropan-1-one and its derivatives have been reported that they show both high and selective thromboxane A₂ receptor antagonist and thromboxane A₂ synthase inhibitory activity (Cozzi *et al.*, 1994). Some 1*H*-1,2,4-triazole derivatives have been known as antifungal and plant growth regulatory agents (Jin *et al.*, 2006). Crystal structures of similar compounds like ketone having benzimidazole and furan rings (Özel Güven *et al.*, 2007) and alcohols having 1,2,4-triazole and benzotriazole rings have been reported (Özel Güven *et al.*, 2008; Özel Güven *et al.*, 2010). Now, we report herein the crystal structure of the title benzotriazole derivative, (I).

The asymmetric unit of the title compound (Fig. 1) contains two crystallographically independent molecules, in which they differ slightly in the orientations of the 2-6-dichlorophenyl units and the bond lengths and angles are generally within normal ranges. The intramolecular O—H···N hydrogen bonds (Table 1) link the benzotriazole rings and phenylpropan units. In each molecule, the planar benzotriazole rings [A (N1-N3/C9-C14) and A' (N1'-N3'/C9'-C14')] are oriented with respect to the phenyl [B (C3-C8) and B' (C3'-C8')] and 2-6-dichlorophenyl [C (C16-C21)] and C' (C16'-C21')] rings at dihedral angles of A/B = 30.61 (5), A/C = 43.36 (5) ° and A'/B' = 32.25 (5), A'/C' = 41.04 (5) °. The dihedral angles between phenyl rings are B/C = 66.38 (7) and B'/C' = 66.14 (6) °. Atoms C1 and C1' are -0.044 (2) and 0.086 (2) Å away from the planes of the benzotriazole rings, respectively.

In the crystal structure, intermolecular C—H···N hydrogen bonds (Table 1) link the molecules into chains along the *a*-axis (Fig. 2). The π - π contacts between the 2-6-dichlorophenyl rings, Cg3—Cg3ⁱ and Cg3ⁱⁱ—Cg3ⁱⁱⁱ, [symmetry codes: (i) 2 - *x*, 2 - *y*, - *z*, (ii) 1 - *x*, 1 - *y*, 1 - *z*, where Cg3 and Cg3' are the centroids of the rings C (C16-C21)] and C' (C16'-C21'), respectively], may further stabilize the structure, with centroid-centroid distances of 3.809 (1) and 3.735 (1) Å, respectively.

Experimental

The title compound, (I), was synthesized by the reduction of 2-(1*H*-benzotriazol-1-yl)-3-(2,6-dichlorophenyl)-1-phenylpropane-1-one. Sodiumborohydride (38.21 mg, 1.01 mmol) was added to a solution of 2-(1*H*-benzotriazol-1-yl)-3-(2,6-dichlorophenyl)-1-phenylpropane-1-one (200 mg, 0.505 mmol) in ethanol (15 ml). The mixture was refluxed for 5 h. The solvent was removed and the mixture was neutralized with dilute HCl, and then refluxed for 30 min. After cooling the mixture, it was alkalized with dilute NaOH. The precipitate was filtered and washed with ethanol. The filtrate was extracted with chloroform, then the organic phase was dried and solvents were removed by rotary evaporator. The residue was purified with column chromatography using hexane: ethyl acetate (7:3) mixture as solvent. The product was crystallized from benzene to obtain colorless single crystals suitable for X-ray analysis (yield; 82 mg, 41%).

Refinement

H atoms were positioned geometrically with O—H = 0.82 for hydroxy H, and C—H = 0.93, 0.97 and 0.98 Å for aromatic, methylene and methine H-atoms, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C}, \text{O})$, where $k = 1.5$ for hydroxy H-atoms and $k = 1.2$ for all other H-atoms.

Figures




Fig. 1. The molecular structure of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H-atoms have been omitted for clarity.




Fig. 2. A view of the crystal packing of the title compound. The intermolecular C—H...N hydrogen bonds are shown as dashed lines [H-atoms not involved in hydrogen bonding have been omitted for clarity].

2-(1H-Benzotriazol-1-yl)-3-(2,6-dichlorophenyl)-1-phenylpropan-1-ol

Crystal data

$\text{C}_{21}\text{H}_{17}\text{Cl}_2\text{N}_3\text{O}$

$M_r = 398.28$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.3894$ (2) Å

$b = 9.4947$ (2) Å

$c = 21.2687$ (3) Å

$\alpha = 91.415$ (2)°

$\beta = 92.324$ (2)°

$\gamma = 90.406$ (1)°

$V = 1893.90$ (6) Å³

$Z = 4$

$F(000) = 824$

$D_x = 1.397$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 14651 reflections

$\theta = 2.9$ – 27.5 °

$\mu = 0.36$ mm⁻¹

$T = 120$ K

Block, colorless

$0.4 \times 0.4 \times 0.3$ mm

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

φ and ω scans

Absorption correction: multi-scan

8697 independent reflections

7206 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

$\theta_{\text{max}} = 27.6$ °, $\theta_{\text{min}} = 3.0$ °

$h = -12 \rightarrow 12$

(SADABS; Sheldrick, 2007)

$T_{\min} = 0.866$, $T_{\max} = 0.897$

40396 measured reflections

$k = -12 \rightarrow 11$

$l = -27 \rightarrow 27$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.108$

$S = 1.07$

8697 reflections

487 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0504P)^2 + 0.9534P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.43 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.73 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.77473 (4)	1.06219 (5)	0.104106 (19)	0.02207 (10)
C12	0.97499 (5)	0.66618 (5)	-0.05798 (2)	0.03245 (12)
O1	0.42613 (13)	0.61760 (13)	-0.02537 (6)	0.0256 (3)
H1A	0.4009	0.6380	0.0101	0.038*
N1	0.60715 (15)	0.74669 (14)	0.06575 (6)	0.0162 (3)
N2	0.50172 (15)	0.83379 (15)	0.08432 (7)	0.0200 (3)
N3	0.49164 (16)	0.82583 (16)	0.14548 (7)	0.0206 (3)
C1	0.64956 (17)	0.73778 (17)	0.00021 (7)	0.0149 (3)
H1	0.7061	0.6522	-0.0048	0.018*
C2	0.52030 (18)	0.72524 (18)	-0.04605 (8)	0.0193 (3)
H2	0.4698	0.8152	-0.0462	0.023*
C3	0.57040 (18)	0.69279 (18)	-0.11180 (8)	0.0197 (4)
C4	0.5683 (2)	0.7957 (2)	-0.15719 (9)	0.0284 (4)
H4	0.5318	0.8842	-0.1479	0.034*
C5	0.6205 (3)	0.7673 (2)	-0.21634 (10)	0.0366 (5)

supplementary materials

H5	0.6183	0.8366	-0.2465	0.044*
C6	0.6758 (2)	0.6356 (2)	-0.23049 (10)	0.0338 (5)
H6	0.7117	0.6171	-0.2699	0.041*
C7	0.6774 (2)	0.5318 (2)	-0.18587 (9)	0.0295 (4)
H7	0.7136	0.4433	-0.1955	0.035*
C8	0.6248 (2)	0.55996 (19)	-0.12650 (9)	0.0237 (4)
H8	0.6260	0.4901	-0.0966	0.028*
C9	0.66758 (18)	0.68088 (17)	0.11643 (8)	0.0160 (3)
C10	0.7773 (2)	0.58173 (18)	0.12292 (8)	0.0215 (4)
H10	0.8286	0.5502	0.0889	0.026*
C11	0.8040 (2)	0.5342 (2)	0.18290 (9)	0.0258 (4)
H11	0.8748	0.4676	0.1894	0.031*
C12	0.7273 (2)	0.5834 (2)	0.23490 (9)	0.0262 (4)
H12	0.7481	0.5472	0.2744	0.031*
C13	0.6226 (2)	0.6834 (2)	0.22842 (8)	0.0237 (4)
H13	0.5740	0.7172	0.2628	0.028*
C14	0.59246 (18)	0.73200 (17)	0.16758 (8)	0.0173 (3)
C15	0.74540 (17)	0.86394 (17)	-0.01401 (8)	0.0157 (3)
H15A	0.7664	0.8609	-0.0583	0.019*
H15B	0.6948	0.9506	-0.0055	0.019*
C16	0.88295 (17)	0.86400 (17)	0.02488 (8)	0.0166 (3)
C17	0.90693 (18)	0.94709 (18)	0.07957 (8)	0.0183 (3)
C18	1.03235 (19)	0.9434 (2)	0.11631 (9)	0.0238 (4)
H18	1.0437	0.9998	0.1526	0.029*
C19	1.1407 (2)	0.8542 (2)	0.09808 (9)	0.0289 (4)
H19	1.2255	0.8510	0.1222	0.035*
C20	1.12324 (19)	0.7704 (2)	0.04434 (10)	0.0276 (4)
H20	1.1960	0.7113	0.0319	0.033*
C21	0.99607 (19)	0.77537 (19)	0.00931 (9)	0.0225 (4)
Cl1'	0.43627 (5)	1.25631 (4)	0.394633 (19)	0.02133 (10)
Cl2'	0.83607 (5)	1.47909 (5)	0.56172 (2)	0.02874 (12)
O1'	0.88141 (13)	0.93042 (12)	0.52629 (6)	0.0217 (3)
H1B	0.8569	0.8991	0.4912	0.032*
N1'	0.74653 (15)	1.09289 (14)	0.43580 (6)	0.0143 (3)
N2'	0.65658 (15)	0.98414 (14)	0.41974 (7)	0.0184 (3)
N3'	0.65629 (16)	0.96282 (15)	0.35860 (7)	0.0201 (3)
C1'	0.76151 (17)	1.14764 (16)	0.50055 (7)	0.0134 (3)
H1'	0.8488	1.2049	0.5043	0.016*
C2'	0.77665 (18)	1.02758 (17)	0.54764 (8)	0.0163 (3)
H2'	0.6849	0.9782	0.5494	0.020*
C3'	0.81778 (18)	1.08827 (17)	0.61265 (8)	0.0167 (3)
C4'	0.95523 (19)	1.14282 (18)	0.62497 (8)	0.0207 (4)
H4'	1.0218	1.1385	0.5938	0.025*
C5'	0.9927 (2)	1.2033 (2)	0.68340 (9)	0.0251 (4)
H5'	1.0840	1.2402	0.6911	0.030*
C6'	0.8942 (2)	1.2090 (2)	0.73039 (9)	0.0283 (4)
H6'	0.9194	1.2499	0.7695	0.034*
C7'	0.7588 (2)	1.1537 (2)	0.71895 (9)	0.0278 (4)
H7'	0.6932	1.1569	0.7505	0.033*

C8'	0.7202 (2)	1.09329 (19)	0.66023 (8)	0.0224 (4)
H8'	0.6289	1.0562	0.6528	0.027*
C9'	0.80557 (17)	1.14485 (17)	0.38334 (7)	0.0150 (3)
C10'	0.90362 (19)	1.25377 (18)	0.37405 (8)	0.0200 (4)
H10'	0.9412	1.3100	0.4072	0.024*
C11'	0.9407 (2)	1.2722 (2)	0.31273 (9)	0.0272 (4)
H11'	1.0059	1.3427	0.3042	0.033*
C12'	0.8824 (2)	1.1869 (2)	0.26232 (9)	0.0309 (4)
H12'	0.9107	1.2029	0.2217	0.037*
C13'	0.7850 (2)	1.08096 (19)	0.27169 (8)	0.0262 (4)
H13'	0.7465	1.0257	0.2384	0.031*
C14'	0.74682 (19)	1.06059 (17)	0.33400 (8)	0.0180 (3)
C15'	0.63528 (17)	1.24411 (16)	0.51530 (7)	0.0146 (3)
H15C	0.5466	1.1929	0.5066	0.017*
H15D	0.6403	1.2706	0.5597	0.017*
C16'	0.63594 (18)	1.37529 (16)	0.47679 (8)	0.0160 (3)
C17'	0.55197 (18)	1.39105 (17)	0.42163 (8)	0.0167 (3)
C18'	0.5560 (2)	1.51185 (18)	0.38579 (8)	0.0213 (4)
H18'	0.4990	1.5184	0.3492	0.026*
C19'	0.6460 (2)	1.62216 (18)	0.40542 (9)	0.0237 (4)
H19'	0.6494	1.7032	0.3818	0.028*
C20'	0.7307 (2)	1.61274 (18)	0.45971 (9)	0.0238 (4)
H20'	0.7903	1.6871	0.4731	0.029*
C21'	0.72545 (19)	1.49006 (18)	0.49406 (8)	0.0193 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0183 (2)	0.0301 (2)	0.0176 (2)	0.00322 (16)	0.00066 (16)	-0.00500 (16)
Cl2	0.0273 (2)	0.0317 (3)	0.0388 (3)	0.00458 (19)	0.0124 (2)	-0.0105 (2)
O1	0.0194 (6)	0.0270 (7)	0.0307 (7)	-0.0052 (5)	0.0077 (5)	-0.0054 (5)
N1	0.0161 (7)	0.0160 (7)	0.0170 (7)	0.0044 (5)	0.0052 (5)	0.0029 (5)
N2	0.0157 (7)	0.0215 (7)	0.0233 (8)	0.0046 (6)	0.0058 (6)	-0.0001 (6)
N3	0.0177 (7)	0.0252 (8)	0.0190 (7)	0.0010 (6)	0.0038 (6)	-0.0031 (6)
C1	0.0155 (8)	0.0155 (8)	0.0139 (8)	0.0033 (6)	0.0034 (6)	0.0018 (6)
C2	0.0154 (8)	0.0177 (8)	0.0244 (9)	-0.0001 (6)	0.0000 (7)	-0.0025 (7)
C3	0.0174 (8)	0.0211 (8)	0.0202 (9)	-0.0008 (7)	-0.0030 (7)	-0.0031 (7)
C4	0.0334 (11)	0.0256 (10)	0.0256 (10)	0.0024 (8)	-0.0051 (8)	-0.0002 (8)
C5	0.0459 (13)	0.0398 (12)	0.0240 (10)	-0.0039 (10)	-0.0013 (9)	0.0067 (9)
C6	0.0345 (11)	0.0463 (12)	0.0205 (10)	-0.0060 (9)	0.0039 (8)	-0.0059 (8)
C7	0.0299 (11)	0.0288 (10)	0.0295 (10)	-0.0005 (8)	0.0044 (8)	-0.0115 (8)
C8	0.0251 (9)	0.0219 (9)	0.0239 (9)	-0.0022 (7)	0.0029 (7)	-0.0040 (7)
C9	0.0177 (8)	0.0145 (8)	0.0158 (8)	-0.0008 (6)	0.0018 (6)	0.0006 (6)
C10	0.0256 (9)	0.0210 (9)	0.0183 (8)	0.0065 (7)	0.0037 (7)	0.0012 (7)
C11	0.0294 (10)	0.0244 (9)	0.0238 (9)	0.0067 (8)	-0.0019 (8)	0.0052 (7)
C12	0.0298 (10)	0.0321 (10)	0.0167 (9)	-0.0015 (8)	-0.0021 (7)	0.0052 (7)
C13	0.0231 (9)	0.0308 (10)	0.0170 (8)	-0.0040 (7)	0.0038 (7)	-0.0035 (7)
C14	0.0150 (8)	0.0186 (8)	0.0184 (8)	-0.0027 (6)	0.0029 (6)	-0.0023 (6)

supplementary materials

C15	0.0155 (8)	0.0176 (8)	0.0140 (8)	0.0000 (6)	0.0010 (6)	0.0005 (6)
C16	0.0139 (8)	0.0198 (8)	0.0165 (8)	-0.0003 (6)	0.0028 (6)	0.0038 (6)
C17	0.0134 (8)	0.0232 (9)	0.0185 (8)	-0.0002 (6)	0.0032 (6)	0.0040 (7)
C18	0.0175 (9)	0.0347 (10)	0.0191 (9)	-0.0029 (7)	-0.0022 (7)	0.0059 (7)
C19	0.0139 (9)	0.0434 (12)	0.0298 (10)	0.0001 (8)	-0.0012 (7)	0.0144 (9)
C20	0.0148 (9)	0.0320 (10)	0.0373 (11)	0.0061 (7)	0.0081 (8)	0.0102 (8)
C21	0.0174 (9)	0.0256 (9)	0.0250 (9)	0.0005 (7)	0.0072 (7)	0.0028 (7)
C11'	0.0268 (2)	0.0159 (2)	0.0207 (2)	-0.00349 (16)	-0.00538 (17)	0.00179 (15)
C12'	0.0305 (3)	0.0212 (2)	0.0332 (3)	-0.00356 (17)	-0.01093 (19)	-0.00507 (18)
O1'	0.0232 (6)	0.0154 (6)	0.0258 (7)	0.0046 (5)	-0.0037 (5)	-0.0032 (5)
N1'	0.0161 (7)	0.0123 (6)	0.0146 (7)	-0.0029 (5)	0.0016 (5)	-0.0021 (5)
N2'	0.0200 (7)	0.0137 (7)	0.0211 (7)	-0.0041 (5)	-0.0005 (6)	-0.0042 (5)
N3'	0.0253 (8)	0.0154 (7)	0.0189 (7)	-0.0010 (6)	-0.0030 (6)	-0.0025 (5)
C1'	0.0147 (8)	0.0121 (7)	0.0133 (7)	-0.0023 (6)	0.0008 (6)	-0.0014 (6)
C2'	0.0158 (8)	0.0136 (8)	0.0193 (8)	-0.0012 (6)	-0.0009 (6)	0.0027 (6)
C3'	0.0186 (8)	0.0140 (8)	0.0173 (8)	0.0019 (6)	-0.0024 (6)	0.0040 (6)
C4'	0.0184 (9)	0.0237 (9)	0.0200 (9)	0.0011 (7)	-0.0011 (7)	0.0009 (7)
C5'	0.0218 (9)	0.0266 (9)	0.0262 (10)	-0.0010 (7)	-0.0061 (7)	-0.0016 (7)
C6'	0.0340 (11)	0.0314 (10)	0.0189 (9)	0.0071 (8)	-0.0046 (8)	-0.0028 (7)
C7'	0.0307 (10)	0.0322 (10)	0.0210 (9)	0.0047 (8)	0.0052 (8)	0.0021 (8)
C8'	0.0194 (9)	0.0243 (9)	0.0236 (9)	0.0008 (7)	0.0014 (7)	0.0046 (7)
C9'	0.0152 (8)	0.0149 (8)	0.0148 (8)	0.0025 (6)	0.0014 (6)	0.0005 (6)
C10'	0.0203 (9)	0.0204 (8)	0.0192 (8)	-0.0046 (7)	0.0017 (7)	-0.0008 (6)
C11'	0.0325 (11)	0.0261 (10)	0.0235 (9)	-0.0057 (8)	0.0074 (8)	0.0045 (7)
C12'	0.0500 (13)	0.0278 (10)	0.0157 (9)	0.0001 (9)	0.0088 (8)	0.0034 (7)
C13'	0.0429 (12)	0.0204 (9)	0.0148 (8)	0.0000 (8)	-0.0025 (8)	-0.0017 (7)
C14'	0.0217 (9)	0.0145 (8)	0.0177 (8)	0.0025 (6)	-0.0018 (7)	-0.0005 (6)
C15'	0.0170 (8)	0.0129 (7)	0.0140 (8)	0.0005 (6)	0.0022 (6)	0.0007 (6)
C16'	0.0199 (8)	0.0114 (7)	0.0170 (8)	0.0022 (6)	0.0045 (6)	-0.0008 (6)
C17'	0.0207 (8)	0.0125 (7)	0.0170 (8)	0.0000 (6)	0.0041 (6)	-0.0009 (6)
C18'	0.0289 (10)	0.0178 (8)	0.0176 (8)	0.0037 (7)	0.0051 (7)	0.0033 (6)
C19'	0.0334 (10)	0.0126 (8)	0.0259 (9)	0.0000 (7)	0.0105 (8)	0.0040 (7)
C20'	0.0272 (10)	0.0125 (8)	0.0321 (10)	-0.0037 (7)	0.0082 (8)	-0.0026 (7)
C21'	0.0201 (9)	0.0165 (8)	0.0212 (9)	0.0003 (6)	0.0012 (7)	-0.0036 (6)

Geometric parameters (Å, °)

C11—C17	1.7467 (17)	C11'—C17'	1.7435 (17)
C12—C21	1.7500 (19)	C12'—C21'	1.7460 (18)
O1—C2	1.435 (2)	O1'—C2'	1.433 (2)
O1—H1A	0.8200	O1'—H1B	0.8200
N1—N2	1.3591 (19)	N1'—C1'	1.461 (2)
N1—C1	1.466 (2)	N1'—C9'	1.367 (2)
N1—C9	1.365 (2)	N2'—N1'	1.3576 (19)
N2—N3	1.312 (2)	N2'—N3'	1.311 (2)
N3—C14	1.381 (2)	N3'—C14'	1.381 (2)
C1—C2	1.533 (2)	C1'—C2'	1.540 (2)
C1—C15	1.536 (2)	C1'—C15'	1.541 (2)
C1—H1	0.9800	C1'—H1'	0.9800

C2—H2	0.9800	C2'—C3'	1.519 (2)
C3—C2	1.519 (2)	C2'—H2'	0.9800
C3—C4	1.390 (3)	C3'—C4'	1.400 (2)
C3—C8	1.397 (2)	C3'—C8'	1.393 (2)
C4—C5	1.389 (3)	C4'—C5'	1.387 (3)
C4—H4	0.9300	C4'—H4'	0.9300
C5—H5	0.9300	C5'—H5'	0.9300
C6—C5	1.388 (3)	C6'—C7'	1.382 (3)
C6—H6	0.9300	C6'—C5'	1.389 (3)
C7—C6	1.385 (3)	C6'—H6'	0.9300
C7—H7	0.9300	C7'—H7'	0.9300
C8—C7	1.395 (3)	C8'—C7'	1.395 (3)
C8—H8	0.9300	C8'—H8'	0.9300
C9—C10	1.405 (2)	C9'—C10'	1.403 (2)
C9—C14	1.399 (2)	C9'—C14'	1.397 (2)
C10—C11	1.378 (3)	C10'—C11'	1.379 (2)
C10—H10	0.9300	C10'—H10'	0.9300
C11—H11	0.9300	C11'—C12'	1.418 (3)
C12—C11	1.415 (3)	C11'—H11'	0.9300
C12—H12	0.9300	C12'—H12'	0.9300
C13—C12	1.377 (3)	C13'—C12'	1.378 (3)
C13—C14	1.403 (2)	C13'—H13'	0.9300
C13—H13	0.9300	C14'—C13'	1.404 (2)
C15—H15A	0.9700	C15'—H15C	0.9700
C15—H15B	0.9700	C15'—H15D	0.9700
C16—C15	1.505 (2)	C16'—C15'	1.508 (2)
C16—C17	1.399 (2)	C16'—C17'	1.398 (2)
C16—C21	1.405 (2)	C16'—C21'	1.404 (2)
C17—C18	1.388 (2)	C17'—C18'	1.394 (2)
C18—C19	1.390 (3)	C18'—C19'	1.387 (3)
C18—H18	0.9300	C18'—H18'	0.9300
C19—H19	0.9300	C19'—H19'	0.9300
C20—C19	1.380 (3)	C20'—C19'	1.380 (3)
C20—H20	0.9300	C20'—C21'	1.392 (2)
C21—C20	1.383 (3)	C20'—H20'	0.9300
C2—O1—H1A	109.5	C2'—O1'—H1B	109.5
N2—N1—C1	121.83 (13)	N2'—N1'—C1'	121.39 (13)
N2—N1—C9	110.25 (13)	N2'—N1'—C9'	110.31 (13)
C9—N1—C1	127.81 (13)	C9'—N1'—C1'	128.08 (13)
N3—N2—N1	108.75 (13)	N3'—N2'—N1'	108.85 (13)
N2—N3—C14	108.35 (13)	N2'—N3'—C14'	108.13 (13)
N1—C1—C2	111.92 (13)	N1'—C1'—C2'	111.37 (12)
N1—C1—C15	110.06 (13)	N1'—C1'—C15'	110.12 (13)
N1—C1—H1	107.6	N1'—C1'—H1'	107.8
C2—C1—C15	111.85 (13)	C2'—C1'—C15'	111.68 (13)
C2—C1—H1	107.6	C2'—C1'—H1'	107.8
C15—C1—H1	107.6	C15'—C1'—H1'	107.8
O1—C2—C3	111.67 (14)	O1'—C2'—C3'	111.31 (13)
O1—C2—C1	108.97 (14)	O1'—C2'—C1'	109.00 (13)

supplementary materials

O1—C2—H2	108.9	O1'—C2'—H2'	108.9
C1—C2—H2	108.9	C1'—C2'—H2'	108.9
C3—C2—C1	109.52 (13)	C3'—C2'—C1'	109.66 (13)
C3—C2—H2	108.9	C3'—C2'—H2'	108.9
C4—C3—C2	120.78 (16)	C4'—C3'—C2'	119.74 (15)
C4—C3—C8	119.18 (17)	C8'—C3'—C2'	121.20 (15)
C8—C3—C2	119.99 (16)	C8'—C3'—C4'	119.05 (16)
C3—C4—H4	119.8	C3'—C4'—H4'	119.8
C5—C4—C3	120.48 (18)	C5'—C4'—C3'	120.41 (17)
C5—C4—H4	119.8	C5'—C4'—H4'	119.8
C4—C5—H5	119.9	C4'—C5'—C6'	120.11 (18)
C6—C5—C4	120.11 (19)	C4'—C5'—H5'	119.9
C6—C5—H5	119.9	C6'—C5'—H5'	119.9
C5—C6—H6	120.0	C5'—C6'—H6'	120.0
C7—C6—C5	119.98 (19)	C7'—C6'—C5'	119.91 (18)
C7—C6—H6	120.0	C7'—C6'—H6'	120.0
C6—C7—C8	120.03 (18)	C6'—C7'—C8'	120.28 (18)
C6—C7—H7	120.0	C6'—C7'—H7'	119.9
C8—C7—H7	120.0	C8'—C7'—H7'	119.9
C3—C8—H8	119.9	C3'—C8'—C7'	120.22 (17)
C7—C8—C3	120.21 (18)	C3'—C8'—H8'	119.9
C7—C8—H8	119.9	C7'—C8'—H8'	119.9
N1—C9—C10	133.02 (15)	N1'—C9'—C10'	133.11 (15)
N1—C9—C14	104.37 (14)	N1'—C9'—C14'	104.09 (14)
C14—C9—C10	122.61 (15)	C14'—C9'—C10'	122.80 (15)
C9—C10—H10	122.1	C9'—C10'—H10'	122.1
C11—C10—C9	115.72 (16)	C11'—C10'—C9'	115.79 (16)
C11—C10—H10	122.1	C11'—C10'—H10'	122.1
C10—C11—C12	122.25 (17)	C10'—C11'—C12'	121.91 (18)
C10—C11—H11	118.9	C10'—C11'—H11'	119.0
C12—C11—H11	118.9	C12'—C11'—H11'	119.0
C11—C12—H12	119.2	C11'—C12'—H12'	119.0
C13—C12—C11	121.64 (17)	C13'—C12'—C11'	122.00 (17)
C13—C12—H12	119.2	C13'—C12'—H12'	119.0
C12—C13—C14	117.06 (16)	C12'—C13'—C14'	116.67 (17)
C12—C13—H13	121.5	C12'—C13'—H13'	121.7
C14—C13—H13	121.5	C14'—C13'—H13'	121.7
N3—C14—C9	108.29 (14)	N3'—C14'—C9'	108.62 (15)
N3—C14—C13	130.99 (16)	N3'—C14'—C13'	130.53 (16)
C9—C14—C13	120.68 (16)	C9'—C14'—C13'	120.83 (16)
C1—C15—H15A	109.2	C1'—C15'—H15C	109.3
C1—C15—H15B	109.2	C1'—C15'—H15D	109.3
C16—C15—C1	111.91 (13)	C16'—C15'—C1'	111.63 (13)
C16—C15—H15A	109.2	C16'—C15'—H15C	109.3
C16—C15—H15B	109.2	C16'—C15'—H15D	109.3
H15A—C15—H15B	107.9	H15C—C15'—H15D	108.0
C17—C16—C15	123.48 (15)	C17'—C16'—C15'	123.46 (15)
C17—C16—C21	115.01 (16)	C17'—C16'—C21'	115.44 (15)
C21—C16—C15	121.47 (15)	C21'—C16'—C15'	121.08 (15)

C16—C17—C11	119.61 (13)	C16'—C17'—C11'	120.24 (12)
C18—C17—C11	117.19 (14)	C18'—C17'—C11'	116.96 (14)
C18—C17—C16	123.21 (16)	C18'—C17'—C16'	122.80 (16)
C17—C18—C19	118.98 (18)	C17'—C18'—H18'	120.4
C17—C18—H18	120.5	C19'—C18'—C17'	119.14 (17)
C19—C18—H18	120.5	C19'—C18'—H18'	120.4
C18—C19—H19	119.8	C18'—C19'—H19'	119.7
C20—C19—C18	120.31 (17)	C20'—C19'—C18'	120.56 (16)
C20—C19—H19	119.8	C20'—C19'—H19'	119.7
C19—C20—C21	119.15 (17)	C19'—C20'—C21'	118.90 (16)
C19—C20—H20	120.4	C19'—C20'—H20'	120.6
C21—C20—H20	120.4	C21'—C20'—H20'	120.6
C16—C21—C12	118.25 (14)	C20'—C21'—C16'	123.15 (16)
C20—C21—C12	118.42 (14)	C20'—C21'—C12'	118.11 (14)
C20—C21—C16	123.33 (17)	C16'—C21'—C12'	118.74 (13)
C1—N1—N2—N3	-176.70 (14)	N2'—N1'—C1'—C2'	45.82 (19)
C9—N1—N2—N3	-0.29 (19)	N2'—N1'—C1'—C15'	-78.65 (17)
N2—N1—C1—C2	-46.4 (2)	C9'—N1'—C1'—C2'	-140.11 (16)
N2—N1—C1—C15	78.61 (18)	C9'—N1'—C1'—C15'	95.42 (18)
C9—N1—C1—C2	137.85 (17)	N2'—N1'—C9'—C10'	-179.53 (17)
C9—N1—C1—C15	-97.12 (18)	N2'—N1'—C9'—C14'	-0.33 (17)
N2—N1—C9—C10	179.28 (18)	C1'—N1'—C9'—C10'	5.9 (3)
N2—N1—C9—C14	0.33 (18)	C1'—N1'—C9'—C14'	-174.94 (15)
C1—N1—C9—C10	-4.6 (3)	N3'—N2'—N1'—C9'	0.62 (18)
C1—N1—C9—C14	176.47 (15)	N3'—N2'—N1'—C1'	175.64 (13)
N1—N2—N3—C14	0.12 (18)	N1'—N2'—N3'—C14'	-0.63 (17)
N2—N3—C14—C9	0.09 (19)	N2'—N3'—C14'—C9'	0.42 (18)
N2—N3—C14—C13	-177.81 (18)	N2'—N3'—C14'—C13'	179.01 (18)
N1—C1—C2—O1	-48.50 (17)	N1'—C1'—C2'—O1'	48.02 (17)
N1—C1—C2—C3	-170.92 (13)	N1'—C1'—C2'—C3'	170.11 (13)
C15—C1—C2—O1	-172.54 (13)	C15'—C1'—C2'—O1'	171.61 (13)
C15—C1—C2—C3	65.04 (17)	C15'—C1'—C2'—C3'	-66.30 (17)
N1—C1—C15—C16	63.47 (17)	N1'—C1'—C15'—C16'	-65.81 (17)
C2—C1—C15—C16	-171.46 (13)	C2'—C1'—C15'—C16'	169.90 (13)
C4—C3—C2—O1	133.27 (17)	O1'—C2'—C3'—C4'	49.0 (2)
C4—C3—C2—C1	-105.93 (19)	O1'—C2'—C3'—C8'	-132.25 (16)
C8—C3—C2—O1	-49.4 (2)	C1'—C2'—C3'—C4'	-71.69 (19)
C8—C3—C2—C1	71.4 (2)	C1'—C2'—C3'—C8'	107.05 (17)
C8—C3—C4—C5	-0.4 (3)	C2'—C3'—C4'—C5'	177.60 (15)
C2—C3—C4—C5	176.99 (18)	C8'—C3'—C4'—C5'	-1.2 (3)
C2—C3—C8—C7	-176.87 (17)	C2'—C3'—C8'—C7'	-177.86 (16)
C4—C3—C8—C7	0.5 (3)	C4'—C3'—C8'—C7'	0.9 (3)
C3—C4—C5—C6	-0.3 (3)	C3'—C4'—C5'—C6'	0.6 (3)
C7—C6—C5—C4	0.8 (3)	C7'—C6'—C5'—C4'	0.3 (3)
C8—C7—C6—C5	-0.6 (3)	C5'—C6'—C7'—C8'	-0.5 (3)
C3—C8—C7—C6	0.0 (3)	C3'—C8'—C7'—C6'	0.0 (3)
N1—C9—C10—C11	-176.93 (18)	N1'—C9'—C10'—C11'	178.29 (18)
C14—C9—C10—C11	1.9 (3)	C14'—C9'—C10'—C11'	-0.8 (3)
N1—C9—C14—N3	-0.25 (18)	N1'—C9'—C14'—N3'	-0.05 (18)

supplementary materials

N1—C9—C14—C13	177.90 (16)	N1'—C9'—C14'—C13'	-178.80 (16)
C10—C9—C14—N3	-179.34 (16)	C10'—C9'—C14'—N3'	179.25 (15)
C10—C9—C14—C13	-1.2 (3)	C10'—C9'—C14'—C13'	0.5 (3)
C9—C10—C11—C12	-0.8 (3)	C9'—C10'—C11'—C12'	0.4 (3)
C13—C12—C11—C10	-0.9 (3)	C10'—C11'—C12'—C13'	0.2 (3)
C14—C13—C12—C11	1.6 (3)	C14'—C13'—C12'—C11'	-0.5 (3)
C12—C13—C14—N3	177.08 (18)	N3'—C14'—C13'—C12'	-178.28 (18)
C12—C13—C14—C9	-0.6 (3)	C9'—C14'—C13'—C12'	0.2 (3)
C17—C16—C15—C1	-99.63 (18)	C17'—C16'—C15'—C1'	98.63 (18)
C21—C16—C15—C1	78.26 (19)	C21'—C16'—C15'—C1'	-79.95 (19)
C15—C16—C17—C11	-2.1 (2)	C15'—C16'—C17'—C11'	1.2 (2)
C15—C16—C17—C18	177.91 (16)	C15'—C16'—C17'—C18'	-178.60 (15)
C21—C16—C17—C11	179.90 (12)	C21'—C16'—C17'—C11'	179.83 (12)
C21—C16—C17—C18	-0.1 (2)	C21'—C16'—C17'—C18'	0.1 (2)
C15—C16—C21—C12	1.3 (2)	C15'—C16'—C21'—C20'	179.40 (16)
C15—C16—C21—C20	-178.87 (16)	C17'—C16'—C21'—C20'	0.7 (2)
C17—C16—C21—C12	179.38 (12)	C15'—C16'—C21'—C12'	-0.5 (2)
C17—C16—C21—C20	-0.8 (3)	C17'—C16'—C21'—C12'	-179.16 (12)
C11—C17—C18—C19	-179.39 (14)	C11'—C17'—C18'—C19'	179.82 (13)
C16—C17—C18—C19	0.6 (3)	C16'—C17'—C18'—C19'	-0.4 (3)
C17—C18—C19—C20	-0.2 (3)	C17'—C18'—C19'—C20'	0.0 (3)
C21—C20—C19—C18	-0.6 (3)	C21'—C20'—C19'—C18'	0.7 (3)
C12—C21—C20—C19	-179.01 (14)	C19'—C20'—C21'—C12'	178.75 (14)
C16—C21—C20—C19	1.2 (3)	C19'—C20'—C21'—C16'	-1.1 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1A \cdots N2	0.82	2.56	3.1254 (19)	127
O1'—H1B \cdots N2'	0.82	2.52	3.0881 (19)	128
C19—H19 \cdots N3 ⁱ	0.93	2.54	3.420 (2)	158
C19'—H19' \cdots N3 ⁱⁱⁱ	0.93	2.53	3.410 (2)	159

Symmetry codes: (i) $x+1, y, z$; (ii) $x, y+1, z$.

Fig. 1




Fig. 2

