





Complete Genome Sequence of Human Coronavirus Strain 229E Isolated from Plasma Collected from a Haitian Child in 2016

Tania S. Bonny,^{a,b*}  Kuttichantran Subramaniam,^c Thomas B. Waltzek,^c Maha A. Elbadry,^{a,b} Valery Madsen Beau De Rochars,^{b,d} Taina Telisma,^e Mohammed Rashid,^b J. Glenn Morris, Jr.,^{b,f}  John A. Lednicky^{a,b}

Department of Environmental and Global Health, College of Public Health and Health Professions, University of Florida, Gainesville, Florida, USA^a; Emerging Pathogens Institute, University of Florida, Gainesville, Florida, USA^b; Department of Infectious Diseases and Immunology, College of Veterinary Medicine, University of Florida, Gainesville, Florida, USA^c; Department of Health Services Research, College of Public Health and Health Professions, University of Florida, Gainesville, Florida, USA^d; Christianville Foundation School Clinic, Gressier, Haiti^e; Department of Medicine, College of Medicine, University of Florida, Gainesville, Florida, USA^f

ABSTRACT Human coronavirus strain 229E (HCoV-229E) and human alphaherpesvirus 1 were isolated from the plasma of a Haitian child in 2016 with suspected arbovirus diseases. To our knowledge, this is the first description of HCoV-229E in human plasma, which is the focus of this article.

Human coronavirus strain 229E (HCoV-229E), one of the causative agents of the common cold, is increasingly associated with more severe respiratory infections in children, elders, and individuals with underlying medical conditions (1–4). The virus also has neuroinvasive and neurotropic properties (5). The presence of HCoV-229E within clinical samples (e.g., nasal or throat swabs, nasopharyngeal aspirate, bronchoalveolar lavage, or saliva) is typically confirmed by molecular (e.g., reverse transcription-PCR [RT-PCR] targeting of virus-specific genes) or serological (immunofluorescence assay targeting viral antigen) methods (3, 6–8). Primary isolation of HCoV-229E in cell culture is technically challenging, and few have succeeded at isolating the virus from respiratory specimens (3, 6, 9, 10). The isolation of HCoV-229E from human plasma, to our knowledge, has never been reported.

During a suspected arbovirus outbreak in March 2016, HCoV-229E and human alphaherpesvirus 1 (information to be presented elsewhere) were isolated from the plasma of a child presenting with acute undifferentiated febrile illness at a school clinic in rural Haiti. The study protocol was approved by the University of Florida Institutional Review Board (IRB) and the Haitian National IRB, with written consent obtained from parents/guardians of all study participants. The patient's plasma tested negative for chikungunya, dengue, and zika genomic RNAs by RT-PCR (11). As the potential causative agent(s) was unknown, virus isolation was attempted in several human and animal cell lines. Virus-induced cytopathic effects (CPE) were observed in VERO E6, LLC-MK2, and MRC-5 by 14 days postinoculation of the cells. The CPE first consisted of either cell vacuolation or the formation of syncytia, followed by cell lysis or detachment of clumped cells from the cell monolayers, suggesting mixed infections. Infected cells and spent cell growth media were tested for the presence of viruses at various time points postinoculation using molecular assays. HCoV-229E viral genomic RNA (vRNA) was detected using a GenMark multiplex PCR eSensor XT-8 respiratory viral panel, and its identity was supported by the generation of correct-sized PCR amplicons using RT-PCR assays targeting HCoV-229E-specific genes (12–14). Whole-genome sequencing

Received 23 October 2017 **Accepted** 24 October 2017 **Published** 22 November 2017

Citation Bonny TS, Subramaniam K, Waltzek TB, Elbadry MA, Beau De Rochars VM, Telisma T, Rashid M, Morris JG, Jr, Lednicky JA. 2017. Complete genome sequence of human coronavirus strain 229E isolated from plasma collected from a Haitian child in 2016. *Genome Announc* 5:e01313-17. <https://doi.org/10.1128/genomeA.01313-17>.

Copyright © 2017 Bonny et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to John A. Lednicky, jlednicky@php.ufl.edu.

* Present address: Tania S. Bonny, Department of Microbiology, University of Dhaka, Dhaka, Bangladesh.

of HCoV-229E vRNA from Vero E6 cells was accomplished by Sanger sequencing following previously described methods (15) to attain the consensus sequence. In parallel, next-generation sequencing (NGS) with a cDNA library prepared using the NEBNext Ultra RNA library prep kit, and sequencing performed using a version 3 chemistry 600-cycle kit on a MiSeq platform (Illumina), produced the same consensus sequence.

Whole-genome sequence analyses of the Haitian HCoV-229E isolate, designated strain 229E/Haiti-1/2016, revealed close genetic relatedness (>99% nucleotide identity) to several American HCoV-229E strains reported in 2015 (the GenBank accession numbers of the three HCoV-229E sequences with the highest nucleotide similarities as of 22 July 2017 are KY983587, KY684760, and KY967357). Compared to the American strains, the Haitian HCoV-229E genome has unique nucleotide polymorphisms that result in changes within the deduced amino acid sequences of the replicase polyprotein 1ab, spike, accessory, membrane, envelope, and nucleocapsid proteins.

Accession number(s). The complete genome sequence of HCoV-229E strain 229E/Haiti-1/2016 has been deposited in the GenBank database under accession number [MF542265](https://www.ncbi.nlm.nih.gov/nuccore/MF542265).

ACKNOWLEDGMENTS

These studies were funded in part by internal funds from the University of Florida Emerging Pathogens Institute and by a grant to J.G.M. from DHHS (NIH R01 AI126357-01S1).

REFERENCES

- van der Hoek L. 2007. Human coronaviruses: what do they cause? *Antivir Ther* 12:651–658.
- Walsh EE, Shin JH, Falsey AR. 2013. Clinical impact of human coronaviruses 229E and OC43 infection in diverse adult populations. *J Infect Dis* 208:1634–1642. <https://doi.org/10.1093/infdis/jit393>.
- Pene F, Merlat A, Vabret A, Rozenberg F, Buzyn A, Dreyfus F, Cariou A, Freymuth F, Lebon P. 2003. Coronavirus 229E-related pneumonia in immunocompromised patients. *Clin Infect Dis* 37:929–932. <https://doi.org/10.1086/377612>.
- Gorse GJ, O'Connor TZ, Hall SL, Vitale JN, Nichol KL. 2009. Human coronavirus and acute respiratory illness in older adults with chronic obstructive pulmonary disease. *J Infect Dis* 199:847–857. <https://doi.org/10.1086/597122>.
- Desforges M, Le Coupanec A, Stodola JK, Meessen-Pinard M, Talbot PJ. 2014. Human coronaviruses: viral and cellular factors involved in neuroinvasiveness and neuropathogenesis. *Virus Res* 194:145–158. <https://doi.org/10.1016/j.virusres.2014.09.011>.
- Farsani SM, Dijkman R, Jebbink MF, Goossens H, Ieven M, Deijs M, Molenkamp R, van der Hoek L. 2012. The first complete genome sequences of clinical isolates of human coronavirus 229E. *Virus Genes* 45:433–439. <https://doi.org/10.1007/s11262-012-0807-9>.
- Chibo D, Birch C. 2006. Analysis of human coronavirus 229E spike and nucleoprotein genes demonstrates genetic drift between chronologically distinct strains. *J Gen Virol* 87:1203–1208. <https://doi.org/10.1099/vir.0.81662-0>.
- Kim YG, Yun SG, Kim MY, Park K, Cho CH, Yoon SY, Nam MH, Lee CK, Cho YJ, Lim CS. 2017. Comparison between saliva and nasopharyngeal swab specimens for detection of respiratory viruses by multiplex reverse transcription-PCR. *J Clin Microbiol* 55:226–233. <https://doi.org/10.1128/JCM.01704-16>.
- Dijkman R, Jebbink MF, Koekkoek SM, Deijs M, Jónsdóttir HR, Molenkamp R, Ieven M, Goossens H, Thiel V, van der Hoek L. 2013. Isolation and characterization of current human coronavirus strains in primary human epithelial cell cultures reveal differences in target cell tropism. *J Virol* 87:6081–6090. <https://doi.org/10.1128/JVI.03368-12>.
- Matoba Y, Aoki Y, Tanaka S, Yahagi K, Itagaki T, Matsuzaki Y, Mizuta K. 2015. Picornavirus-like cytopathic effects on RD-18S cell lines were induced by human coronavirus 229E not picornaviruses. *Jpn J Infect Dis* 68:78–79. <https://doi.org/10.7883/yoken.JJID.2014.487>.
- Waggoner JJ, Gresh L, Mohamed-Hadley A, Ballesteros G, Davila MJ, Tellez Y, Sahoo MK, Balmaseda A, Harris E, Pinsky BA. 2016. Single-reaction multiplex reverse transcription PCR for detection of Zika, chikungunya, and dengue viruses. *Emerg Infect Dis* 22:1295–1297. <https://doi.org/10.3201/eid2207.160326>.
- Vabret A, Mouthon F, Mourez T, Gouarin S, Petitjean J, Freymuth F. 2001. Direct diagnosis of human respiratory coronaviruses 229E and OC43 by the polymerase chain reaction. *J Virol Methods* 97:59–66. [https://doi.org/10.1016/S0166-0934\(01\)00343-3](https://doi.org/10.1016/S0166-0934(01)00343-3).
- Myint S, Johnston S, Sanderson G, Simpson H. 1994. Evaluation of nested polymerase chain methods for the detection of human coronaviruses 229E and OC43. *Mol Cell Probes* 8:357–364. <https://doi.org/10.1006/mcpr.1994.1052>.
- Stewart JN, Mounir S, Talbot PJ. 1992. Detection of coronaviruses by the polymerase chain reaction, p 316–327. *In* Becker Y, Darai G (ed). *Diagnosis of human viruses by polymerase chain reaction technology, frontiers of virology*, vol 1. Springer, Berlin, Germany.
- Bonny TS, Yezli S, Lednický JA. 8 September 2017. Isolation and identification of *human coronavirus 229E* from frequently touched environmental surfaces of a university classroom that is cleaned daily. *Am J Infect Control*. <https://doi.org/10.1016/j.ajic.2017.07.014>.