




Complete Genome Sequence of *Staphylococcus pseudintermedius* Type Strain LMG 22219

Mohamed A. Abouelkhair,^{a,b} Matthew C. Riley,^a David A. Bemis,^a

 Stephen A. Kania^a

Department of Biomedical and Diagnostic Sciences, Knoxville, Tennessee, USA^a; Faculty of Veterinary Medicine, University of Sadat City, Menoufia, Egypt^b

ABSTRACT We report the first complete genome sequence of LMG 22219 (=ON 86^T = CCUG 49543^T), the *Staphylococcus pseudintermedius* type strain isolated from feline lung tissue. This sequence information will facilitate phylogenetic comparisons of staphylococcal species and other bacteria at the genome level.

Here, we present the first complete genome sequence of LMG 22219 (=ON 86^T = CCUG 49543^T), the *Staphylococcus pseudintermedius* type strain isolated from lung tissue of a cat (1). *S. pseudintermedius* is a Gram-positive opportunistic bacterial pathogen (2) commonly associated with canine pyoderma and occasionally isolated from human infections (3–6). It was first defined as a unique species by Devriese et al. (1). This species is classified as a member of the *Staphylococcus intermedius* group, which also includes *S. intermedius* and *S. delphini*, as defined by Takahashi et al. (7). *S. pseudintermedius* is characterized by nonpigmented colonies surrounded by double-zone hemolysis on Columbia sheep blood agar. It is catalase-positive and coagulates rabbit plasma but is clumping-factor-negative in slide coagulase testing. *S. pseudintermedius* is positive for DNase, β -glucosidase, arginine dihydrolase, urease, nitrate reduction, pyrrolidonyl arylamidase, and ONPG (β -galactosidase). It does not produce β -glucuronidase and is susceptible to 8 $\mu\text{g ml}^{-1}$ acriflavine and to novobiocin. It is also resistant to deferroxamine (1).

Whereas assays exist to differentiate *S. pseudintermedius* from other *Staphylococcus intermedius* group species, it may be misidentified in veterinary and human medicine (6, 8). Clinical treatment of infections relies upon accurate pathogen identification (9) and the availability of the complete genome for this organism may allow for genetic validation of phenotypic testing, as well as identification of new genetic targets for interspecies comparisons. Only genome sequences of *S. intermedius* and *S. delphini* (10) type strains are currently available, and consequently the elucidation of the genome sequence of the *S. pseudintermedius* type strain LMG 22219 will facilitate comparisons of different species and strains based on genetic analysis.

DNA was extracted and a library prepared using the Nextera XT library preparation kit in accordance with the manufacturer's protocol. Sequencing was performed using Illumina MiSeq version 2 (Illumina Inc., USA), and *de novo* assembly was performed using Geneious version 9.1.6 (11). Automated annotation of the assembled contigs was performed using the NCBI Prokaryote Genome Annotation Pipeline (http://www.ncbi.nlm.nih.gov/genome/annotation_prok).

A total of 1,432,416 MiSeq paired-end reads were used to generate 50 contigs with >100 \times average depth of coverage. The LMG 22219 genome comprises 2,523,112 bp with 37.6% GC content, 2,242 predicted coding sequences, and 81 RNAs.

Received 7 December 2016 Accepted 11 December 2016 Published 16 February 2017

Citation Abouelkhair MA, Riley MC, Bemis DA, Kania SA. 2017. Complete genome sequence of *Staphylococcus pseudintermedius* type strain LMG 22219. *Genome Announc* 5:e01651-16. [10.1128/genomeA.01651-16](https://doi.org/10.1128/genomeA.01651-16).

Copyright © 2017 Abouelkhair et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stephen A. Kania, skania@utk.edu.

Accession number(s). This whole-genome sequence has been deposited at DDBJ/ENA/GenBank under the accession number [MLGE00000000](#). The version described in this paper is the first version, MLGE00000000.1.

ACKNOWLEDGMENTS

We thank the University of Tennessee Institute of Agriculture Center of Excellence in Livestock Diseases and Human Health for funding and support.

REFERENCES

- Devriese LA, Vancanneyt M, Baele M, Vaneechoutte M, De Graef E, Snauwaert C, Cleenwerck I, Dawyndt P, Swings J, Decostere A, Haesebrouck F. 2005. *Staphylococcus pseudintermedius* sp. nov., a coagulase positive species from animals. *Int J Syst Evol Microbiol* 55:1569–1573. <https://doi.org/10.1099/ijss.0.63413-0>.
- Van Hoovels L, Vankeerberghen A, Boel A, Van Vaerenbergh K, De Beenhouwer H. 2006. First case of *Staphylococcus pseudintermedius* infection in a human. *J Clin Microbiol* 44:4609–4612.
- Börjesson S, Gómez-Sanz E, Ekström K, Torres C, Grönlund U. 2015. *Staphylococcus pseudintermedius* can be misdiagnosed as *Staphylococcus aureus* in humans with dog bite wounds. *Eur J Clin Microbiol Infect Dis* 34:839–844. <https://doi.org/10.1007/s10096-014-2300-y>.
- Frank LA, Kania SA, Hnilica KA, Wilkes RP, Bemis DA. 2003. Isolation of *Staphylococcus schleiferi* from dogs with pyoderma. *J Am Vet Med Assoc* 222:451–454. <https://doi.org/10.2460/javma.2003.222.451>.
- Lee J, Murray A, Bendall R, Gaze W, Zhang L, Vos M. 2015. Improved detection of *Staphylococcus intermedius* group in a routine diagnostic laboratory. *J Clin Microbiol* 53:961–963. <https://doi.org/10.1128/JCM.02474-14>.
- Sasaki T, Kikuchi K, Tanaka Y, Takahashi N, Kamata S, Hiramatsu K. 2007. Reclassification of phenotypically identified *Staphylococcus intermedius* strains. *J Clin Microbiol* 45:2770–2778. <https://doi.org/10.1128/JCM.00360-07>.
- Takahashi T, Satoh I, Kikuchi N. 1999. Phylogenetic relationships of 38 taxa of the genus *Staphylococcus* based on 16S rRNA gene sequence analysis. *Int J Syst Bacteriol* 49:725–728. <https://doi.org/10.1099/00207713-49-2-725>.
- Weese JS. 2013. Phenotypic identification of *Staphylococcus intermedius* may be inaccurate. *J Clin Microbiol* 51:734. <https://doi.org/10.1128/JCM.02736-12>.
- Starlander G, Börjesson S, Grönlund-Andersson U, Tellgren-Roth C, Melhus A. 2014. Cluster of infections caused by methicillin-resistant *Staphylococcus pseudintermedius* in humans in a tertiary hospital. *J Clin Microbiol* 52:3118–3120. <https://doi.org/10.1128/JCM.00703-14>.
- Ben Zakour NL, Beatson SA, van den Broek AHM, Thoday KL, Fitzgerald JR. 2012. Comparative genomics of the *Staphylococcus intermedius* group of animal pathogens. *Front Cell Infect Microbiol* 2:44. <https://doi.org/10.3389/fcimb.2012.00044>.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.