

RESEARCH

Open Access

# The absence of myelin basic protein promotes neuroinflammation and reduces amyloid $\beta$ -protein accumulation in Tg-5xFAD mice

Ming-Hsuan Ou-Yang and William E Van Nostrand\*

## Abstract

**Background:** Abnormal accumulation of amyloid  $\beta$ -protein ( $A\beta$ ) in the brain plays an important role in the pathogenesis of Alzheimer's disease (AD).  $A\beta$  monomers assemble into oligomers and fibrils that promote neuronal dysfunction. This assembly pathway is influenced by naturally occurring brain molecules, the  $A\beta$  chaperone proteins, which bind to  $A\beta$  and modulate its aggregation. Myelin basic protein (MBP) was previously identified as a novel  $A\beta$  chaperone protein and a potent inhibitor for  $A\beta$  fibril assembly *in vitro*.

**Methods:** In this study, we determined whether the absence of MBP would influence  $A\beta$  pathology *in vivo* by breeding MBP knockout mice ( $MBP^{-/-}$ ) with Tg-5xFAD mice, a model of AD-like parenchymal  $A\beta$  pathology.

**Results:** Through biochemical and immunohistochemical experiments, we found that bigenic Tg-5xFAD/ $MBP^{-/-}$  mice had a significant decrease of insoluble  $A\beta$  and parenchymal plaque deposition at an early age. The expression of transgene encoded human  $A\beta$ PP, the levels of C-terminal fragments generated during  $A\beta$  production and the intracellular  $A\beta$  were unaffected in the absence of MBP. Likewise, we did not find a significant difference in plasma  $A\beta$  or cerebrospinal fluid  $A\beta$ , suggesting these clearance routes were unaltered in bigenic Tg-5xFAD/ $MBP^{-/-}$  mice. However,  $MBP^{-/-}$  mice and bigenic Tg-5xFAD/ $MBP^{-/-}$  mice exhibited elevated reactive astrocytes and activated microglia compared with Tg-5xFAD mice. The  $A\beta$  degrading enzyme matrix metalloproteinase 9 (MMP-9), which is expressed by activated glial cells, was significantly increased in the Tg-5xFAD/ $MBP^{-/-}$  mice.

**Conclusions:** These findings indicate that the absence of MBP decreases  $A\beta$  deposition in transgenic mice and that this consequence may result from increased glial activation and expression of MMP-9, an  $A\beta$  degrading enzyme.

**Keywords:** Alzheimer's disease, Amyloid  $\beta$ -protein, Astrocyte, Chaperone molecules, Matrix metalloproteinases, Microglia, Myelin basic protein, Transgenic mice

## Background

One of the pathological hallmarks of Alzheimer's disease (AD) is the abnormal accumulation amyloid  $\beta$  ( $A\beta$ ) aggregates in brain.  $A\beta$  is a 38–43 peptide produced from the sequential proteolysis of the amyloid precursor protein ( $A\beta$ PP), a ubiquitously expressed type I membrane protein, by  $\beta$  secretase [1] and  $\gamma$  secretase [2,3]. The assembly of  $A\beta$  into soluble oligomeric forms and fibrils is proposed to have a causative role in AD through various mechanisms [4]. Soluble oligomers have been shown to correlate with synaptic plasticity and memory deficit

[5,6]. Fibrillar  $A\beta$  can promote oxidative stress and neuroinflammation, and is toxic to neuronal and vascular cells [7,8].

The assembly of  $A\beta$  is influenced by a number of naturally occurring brain factors, the ' $A\beta$  chaperone molecules' that bind and modulate the aggregation process of the peptide. One of the better known  $A\beta$  chaperones is the apolipoprotein E (apoE) family. The apoE2 and apoE3 isoforms can suppress fibrillar  $A\beta$  deposition, while apoE4 can promote fibril formation [9,10]. Studies using transgenic mice have demonstrated that by modulating the levels of these  $A\beta$  binding partners,  $A\beta$  deposition is delayed or enhanced [11]. Other examples of  $A\beta$  chaperones include apolipoprotein J [12,13], members of heat shock

\* Correspondence: William.VanNostrand@stonybrook.edu  
Departments of Neurosurgery & Medicine, Stony Brook University, Stony Brook, NY 11794-8122, USA

proteins [14,15],  $\alpha_1$ -anti-chymotrypsin [16], transthyretin [17,18], proteoglycans [19], and gangliosides [20,21].

Previously, we identified myelin basic protein (MBP) as a novel A $\beta$  chaperone that can potentially inhibit its fibrillar assembly [22]. MBP is best known as a major structural protein in the central nervous system (CNS) myelin sheath. It is also suggested to have a role in intracellular signaling through interactions with membrane actin and tubulin [23]. MBPs are products of the Golli (genes of the oligodendrocyte lineage)-MBP gene complex [24]. Four major MBP isoforms are products of alternative splicing of the Golli-MBP gene complex [25,26]. The expression of the different MBP isoforms by oligodendrocytes is developmentally regulated. The predominant MBP isoform in mature human beings is 18.5 kDa [27,28]. Although the 18.5 kDa MBP undergoes post-translational modifications to give rise to eight charge isomers, its ability to bind A $\beta$  appears to be solely sequence-dependent [22]. The strong binding of MBP to A $\beta$ 42 was demonstrated to inhibit A $\beta$  fibril assembly in a substoichiometric molar ratio *in vitro* [29].

Interestingly, a number of studies have reported a loss of myelin and breakdown of MBP in AD patients and mouse models of AD pathology. This loss of myelin is associated with AD risk factors (for example, aging, apoE4, traumatic brain injury) [30-34] and an increase of A $\beta$  peptides [35]. Immunolabeling of brains for A $\beta$  showed that the most susceptible areas for its deposition are in gray matter, where little MBP is present. Conversely, areas of white matter that are abundantly supplied with MBP (for example, corpus callosum, striatum) exhibit very little A $\beta$  deposition. Furthermore, other studies showed there was no myelin staining inside amyloid plaques [36]. Taken together, these findings suggest an inverse correlation between the levels of MBP and A $\beta$ . However, whether MBP can actually influence A $\beta$  accumulation *in vivo* remains unknown.

Here, we directly tested whether MBP could modulate A $\beta$  *in vivo* by removing endogenous MBP from a mouse model of AD-like A $\beta$  pathology. We took advantage of MBP<sup>-/-</sup> mice, known as *shiverer* mice, in which no functional MBP is produced due to a gene breakage from the middle of MBP exon II [37]. MBP<sup>-/-</sup> mice were crossed with human A $\beta$ PP transgenic mice Tg-5xFAD, a model of parenchymal plaque amyloid pathology [38]. We show that in the absence of endogenous mouse MBP there was a significant reduction in cerebral A $\beta$  levels and the amount of deposited fibrillar amyloid. The reduction in A $\beta$  was not due to changes in expression or processing of human A $\beta$ PP or in clearance through cerebrospinal fluid (CSF) or plasma pathways. However, in bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice there was a significant elevation in activated astrocytes and microglia as well as in the levels of the A $\beta$ -degrading enzyme MMP-9. Together, these

findings indicate that in the absence of MBP there is a marked reduction in A $\beta$  pathology in Tg-5xFAD mice but that this decrease is likely to result from increased degradation via elevated neuroinflammatory glial cells and associated MMP-9.

## Methods

### Animals

All work with mice followed National Institutes of Health guidelines and was approved by the Stony Brook University Institutional Animal Care and Use Committee. Tg-5xFAD mice were obtained from Jackson Laboratories. Tg-5xFAD mice coexpress human APP and human presenilin 1 with five familial AD mutations (APP K670N/M671L + I716V + V717I and PS1 M146L + L286V) and develop early-onset A $\beta$  accumulation and fibrillar A $\beta$  plaques in the brain, starting at about two months of age [38]. *Shiverer* MBP<sup>-/-</sup> mice were also obtained from Jackson Laboratories. *Shiverer* MBP<sup>-/-</sup> mice produce no functional MBP, owing to a gene breakage from the middle of MBP exon II [37]. Hemizygous Tg-5xFAD mice were successively bred with MBP<sup>+/-</sup> mice to obtain cohorts of wild-type mice, Tg-5xFAD mice, MBP<sup>-/-</sup> mice, and bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice. 10 to 12 mice of each genotype were collected at two months of age.

### Tissue preparation

Mice were overdosed with 2.5% Avertin followed by the collection of CSF, plasma and brain. CSF was obtained following a protocol adapted from [39]. Blood was collected through heart puncture with a 27½G needle in one-tenth volume of 3.8% sodium citrate to prevent coagulation. Blood was centrifuged at 8,000g for 5 min at room temperature to remove platelets and cellular components. Plasma samples were stored at -80°C until analysis. Brains were perfused with PBS and bisected along the midsagittal plane. One hemisphere was snap frozen and stored at -80°C. The other hemisphere was placed in 70% ethanol, followed by xylene treatment and embedding in paraffin for immunohistochemical and histological analyses.

### ELISA analysis of cerebral A $\beta$ peptides

The pools of A $\beta$ <sub>40</sub> and A $\beta$ <sub>42</sub> were determined by using a specific ELISA as previously described [40]. Sequential extraction of pulverized mouse forebrain tissues was as follows. To obtain a soluble fraction, tissue aliquots were homogenized with tris-buffered saline (TBS) (10  $\mu$ l/mg tissue) using a bullet blender and 0.5 mm glass beads (Next Advance, Inc.) followed by 20 min centrifugation at 8,000 g at 4°C. The supernatant was removed as the soluble fraction and the pellet was next extracted with TBS/1% Triton X-100 following the same procedure to obtain a membrane-associated fraction. Finally, the

resulting pellet was resuspended in 5M guanidine-HCl (pH 8.0), rotating at room temperature for 3 hours. After centrifugation, the supernatant was removed and kept as the insoluble fraction. Plasma was treated with 5M guanidine-HCl (pH 8.0) at room temperature for 30 min. For each fraction, a sandwich ELISA was performed, where A $\beta$ <sub>40</sub> and A $\beta$ <sub>42</sub> were captured using their respective carboxyl terminus-specific antibodies, m2G3 and m21F12, and biotinylated antibody m3D6, specific for human A $\beta$ , was used for detection [41].

#### Immunoblot analysis

The TBS/1% Triton X-100 extraction (membrane-associated fraction) was used to detect A $\beta$ PP and A $\beta$ PP C-terminal fragments (CTFs). Direct TBS/1% Triton X-100 extraction (total extraction) was used to detect GFAP. Protein concentration was determined using a BCA kit (Pierce). Equal amounts of total protein were separated on 4 to 12% Tris-Glycine (Invitrogen) or 16% Tricine (Invitrogen) for APP CTFs. Gels were transferred onto nitrocellulose membranes (Amersham Hybond-ECL). Membranes were blocked with 5% nonfat milk and incubated overnight at 4°C with anti-human A $\beta$ PP (mouse mAb P2-1, 1:1,000), anti-A $\beta$ PP-CTF (rabbit pAb, 1:1,000), anti-MMP-9 (Abcam ab38898 1:1,000), anti-neurospecific  $\beta$ -tubulin (Abcam ab18207 1:2,000), anti-GFAP (Chemicon MAB360 1:1,000). Secondary HRP conjugated anti-mouse or anti-rabbit was used at 1:5,000 dilution. Membranes were developed using ECL (Pierce) and signals were quantified with VersaDoc (BioRad Model 3000).

#### Immunohistochemical analysis

10  $\mu$ m paraffin sections were deparaffinated in xylene and rehydrated with ethanol. Sections were blocked in SuperBlock blocking buffer (Thermo #37515) with 0.3% triton X-100 and incubated overnight with diluted primary antibody in 1:10 SuperBlock/PBS containing 0.1% triton X-100 at 4°C. The following antibodies were used: A $\beta$  rabbit pAb anti-A $\beta$ 1-28 1:500), GFAP antibody (Chemicon MAB360 1:1,000), Keratan sulfate antibody (5D4) (Seikagaku Corp. 1:1,000), OC antibody (a gift from Dr. Charles Glabe, UC Irvine, 1:1,000). Antigen retrieval was done in antigen unmasking solution (Vector labs H-3301) 30 minutes at 90°C for 5D4 staining and 15 min in 88% formic acid for intraneuronal A $\beta$  (OC) staining before blocking. Sections were treated with Alexa Fluor 488 (Invitrogen) for fluorescence staining or biotinylated secondary antibodies followed by vectastain ABC kit (Vector Labs) for DAB staining.

#### Gelatin zymography

Pulverized brain aliquots were homogenized in TBS containing 1% Triton X-100 as described. After centrifugation, 400  $\mu$ l of supernatant was incubated with 50  $\mu$ l

50% pre-washed gelatin agarose beads and allowed to rotate overnight at 4°C. The beads were then pelleted by centrifugation. After removing the supernatant, the beads were washed in PBS and eluted with 50  $\mu$ l 1X gel loading dye. Half of the elution was separated on 7.5% SDS-PAGE containing 0.1% gelatin. Following electrophoresis, the gels were gently agitated in 2.5% Triton X-100 at room temperature. The buffer was changed every 40 min for three times. After briefly rinsed in the assay buffer, gels were incubated with shaking in assay buffer (50 mM Tris, 0.2M NaCl, 6.7mM CaCl<sub>2</sub>) at 37°C for 20 hours. Gels were stained with Coomassie blue and destained until clearing by gelatinases was visible.

#### Statistical analysis

Data were analyzed using the unpaired two-tailed Student's *t* test. Error bars represent standard error of the mean (SEM). Significance was taken when *P* value was less than 0.05.


## Results

### Significant reduction of insoluble cerebral A $\beta$ in Tg-5xFAD/MBP<sup>-/-</sup> mice

Previously, we identified MBP as a potent A $\beta$  fibrillogenesis inhibitor *in vitro* via a sequence-dependent interaction [22,29,42]. To investigate whether the absence of MBP could influence A $\beta$  pathology *in vivo*, we bred MBP<sup>-/-</sup> mice to Tg-5xFAD mice, a model of parenchymal AD-like A $\beta$  pathology. Age-matched Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice were collected at 2 months of age when thioflavin S-positive fibrillar plaques begin to appear in Tg-5xFAD mice [43]. Pulverized brain aliquots were sequentially extracted into soluble (s), membrane-associated (m) and insoluble (i) fractions for A $\beta$  ELISA analysis. Bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice had a significant reduction in the amount of insoluble A $\beta$ , with an eight-fold reduction in A $\beta$ <sub>40</sub> and a 30-fold reduction in A $\beta$ <sub>42</sub>. No significant differences were found in the levels of soluble A $\beta$  and membrane-associated A $\beta$  between Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice (Figure 1).

### parenchymal A $\beta$ deposition in Tg-5xFAD/MBP<sup>-/-</sup> mice

We next performed immunofluorescent labeling for A $\beta$  on brain sections from Tg-5xFAD and bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice using an anti-A $\beta$  N-terminal antibody. Even though A $\beta$  deposition just begins at this early age in Tg-5xFAD mice, there was a remarkable decrease in both number and the size of A $\beta$  plaques in bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice observed in the cortex, subiculum, and thalamus (Figure 2), which was consistent with the reduction of insoluble A $\beta$  from the ELISA analysis.





**The absence of MBP does not alter human AβPP protein levels or processing by α and β secretases in Tg-5xFAD mice**  
 The observed reduction in Aβ levels and deposition could be a consequence of decreased Aβ production or increased Aβ catabolism. To determine whether the

absence of MBP led to a decrease in human AβPP expression or processing in Tg-5xFAD mice, we performed quantitative immunoblotting on membrane-associated fractions using antibodies against human AβPP and the AβPP CTFs generated from α secretase (C83) and β secretase cleavages (C99) (Figure 3A). There were no significant differences in the levels of human AβPP, or in the levels of AβPP CTF cleavage products, between Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice (Figure 3B). This finding suggests that the reduction of Aβ in bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice was unlikely to have resulted from decreased Aβ production.

**The levels of intracellular Aβ are unaltered between Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice**

We next evaluated the level of intracellular Aβ (iAβ), since its accumulation is proposed to precede extracellular Aβ deposition and it is suggested as one of the first events in the progression of Aβ pathology [44,45]. The detection of iAβ has been controversial, owing to the cross-reaction of some Aβ antibodies with AβPP. To avoid this potential confound, we used a conformational antibody (OC), which is specific to a fibrillar epitope present in Aβ oligomers and fibrils [46]. We saw prominent iAβ-containing neurons in the cortical layer V that appeared comparable between Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice (Figures 4A,B). The numbers of cortical neurons that were positive with iAβ were counted (Figure 4C). At the age of two months, male mice had three-fold less iAβ positive neurons than female






**Figure 3** Absence of MBP does not alter AβPP expression or processing in Tg-5xFAD/MBP<sup>-/-</sup> mice. **(A)** Equal amount of total protein was separated on 4 to 12% or Tris-Glycine gel for AβPP or 16% Tricine for AβPP CTFs. **(B)** The chemiluminescence signals were quantified and presented as percentage of Tg-5xFAD. Data presented are the mean ± SEM of 11 or 12 mice per group.

mice, but there was no significant difference between Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice of the same sex. This result and the quantitative data from Figure 3 together indicate that the absence of MBP does not alter Aβ production and suggest that the events causing Aβ reduction in Tg-5xFAD/MBP<sup>-/-</sup> mice probably occur extracellularly, after Aβ is released.

#### Efflux of Aβ into plasma or CSF is unaltered in Tg-5xFAD/MBP<sup>-/-</sup> mice

Efflux into plasma or CSF represents major clearance pathways for Aβ in brain [47-49]. To determine whether the efflux of Aβ was affected by the absence of MBP, we performed ELISA analyses for Aβ on guanidine-extracted plasma samples and CSF samples obtained from the two



**Figure 4** No significant difference in the level of intraneuronal Aβ between Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice. To assess the level of intraneuronal Aβ, sections were pretreated with formic acid and incubated with an oligomer specific antibody, OC. Prominent cell body Aβ staining was observed in layer V of cortex in both **(A)** Tg-5xFAD and **(B)** Tg-5xFAD/MBP<sup>-/-</sup> mice. **(C)** Cortical neurons with positive iAβ were counted, no difference was observed between different genotypes. Scale bar, 100 μm. Data presented are the mean ± SEM of 4 or 5 mice per group.

groups of mice. There were similar levels of A $\beta$ 40 and A $\beta$ 42 in the plasma of Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice (Figure 5A). Likewise, in CSF there was no significant difference in the levels of A $\beta$ 40 and A $\beta$ 42 (Figure 5B). These findings suggest that there is no enhancement of plasma or CSF clearance of A $\beta$  in the absence of MBP.

#### Elevated neuroinflammatory cells in Tg-5xFAD/MBP<sup>-/-</sup> mice

Activated glial cells are known to participate in A $\beta$  clearance by producing a number of A $\beta$  degrading enzymes. Immunostaining for astrocytes using an antibody to GFAP, we found that bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice (Figure 6C) had extensive astrocyte staining compared with wild-type mice (Figure 6A) and Tg-5xFAD mice (Figure 6B). In addition, we observed a similar elevated astrocyte immunostaining pattern in the MBP<sup>-/-</sup> mice (Figure 6D) compared with wild-type mice, as reported previously [50,51]. The elevated GFAP was also confirmed by quantitative immunoblotting on brain homogenates from the different mice (Figure 6E). GFAP levels were increased three- to four-fold in Tg-5xFAD/MBP<sup>-/-</sup> mice and MBP<sup>-/-</sup> mice compared with Tg-5xFAD mice (Figure 6F).


Similarly, using a marker for activated microglia showed increased immunostaining in bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice (Figure 7C) compared with wild-type mice (Figure 7A) and Tg-5xFAD mice (Figure 7B). Again, we observed a similar elevated activated microglial immunostaining pattern in MBP<sup>-/-</sup> mice (Figure 7D). Together, these findings indicate that bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice have elevated levels of neuroinflammatory cells compared with Tg-5xFAD mice and that this effect appears to be associated with the absence of MBP, as MBP<sup>-/-</sup> mice exhibit comparable increases.

#### Increased expression of the A $\beta$ degrading enzyme MMP-9 in Tg-5xFAD/MBP<sup>-/-</sup> mice


Reactive astrocytes and activated microglia are known to express the A $\beta$ -degrading enzymes matrix metalloproteinase 2 and 9 (MMP-2 and MMP-9) [52-54]. To measure the activity of MMP-2 and MMP-9, gelatin zymography was performed using brain lysates prepared from Tg-5xFAD mice and bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice that were concentrated by filtration over gelatin agarose. Gelatin zymography showed a two-fold increase in MMP-9, but not MMP-2, in Tg-5xFAD/MBP<sup>-/-</sup> mice compared with Tg-5xFAD mice (Figure 8A,B). Immunoblotting showed a similar increase in MMP-9 protein levels in Tg-5xFAD/MBP<sup>-/-</sup> mice (Figure 8C).

#### Discussion

The assembly pathway of endogenous A $\beta$  is influenced by various A $\beta$  chaperone proteins, which can either promote or inhibit the aggregation of A $\beta$ . Previously, our *in vitro* work showed that MBP could strongly bind fibrillogenic forms of A $\beta$  and potently inhibit their assembly into fibrils [22,29]. Although MBP is largely embedded in myelin sheaths, it can be readily detected in the CSF of healthy individuals within the range of  $\mu$ g/l [55,56]. Furthermore, after brain injuries and myelin breakdown, the MBP is increased in the CSF [55,57,58]. Golli-MBP proteins are expressed in several cell types in the CNS, including oligodendrocytes, neurons, and microglia [59-61]. Recent studies have implicated Golli-MBP proteins as multifunctional intracellular scaffolds that can bind a number of intracellular proteins and small molecule ligands affecting diverse cellular processes [62]. These findings suggest that cells that express Golli-MBP proteins could affect intracellular, and possibly extracellular, A $\beta$  assembly, molecular interactions, and associated pathogenic effects. Thus with its abundance in the brain and its close proximity to A $\beta$  it is



**Figure 5** No significant difference in plasma and CSF A $\beta$  levels of Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice. ELISA measurements were performed for A $\beta$ 40 and A $\beta$ 42 in plasma samples (A) and CSF samples (B) collected from Tg-5xFAD and Tg-5xFAD/MBP<sup>-/-</sup> mice. The data presented are the mean  $\pm$  SEM of ten mice per group.



possible that MBP could influence A $\beta$  levels, especially at early stages of myelin breakdown.


In this present study, we sought to explore the consequences of removing endogenous MBP on A $\beta$  accumulation in transgenic mice. To do this, we generated Tg-5xFAD/MBP<sup>-/-</sup> mice by breeding MBP<sup>-/-</sup> mice to Tg-5xFAD mice, a model of early-onset parenchymal A $\beta$  pathology. Owing to the short life span of MBP<sup>-/-</sup> mice [63], the choice of an aggressive A $\beta$  depositing mouse model such as Tg-5xFAD was necessary to allow us to investigate A $\beta$  pathology at a young age. At weaning age, the bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice exhibited the same severe shivering phenotype that is characteristic of the MBP<sup>-/-</sup> mice, but died at a younger age before reaching three months, which we suspected to be a result of the rapid A $\beta$  accumulation from the FAD mutations.

Based on our earlier *in vitro* data, showing a potent inhibitory effect of MBP on A $\beta$  fibril assembly, we might have expected to see an increase in A $\beta$  accumulation and deposition in the absence of MBP. Conversely, in the absence of MBP the Tg-5xFAD mice exhibited significantly decreased A $\beta$  levels and A $\beta$  deposition in the brain at two months of age (Figure 1 and Figure 2, respectively). However, this finding was not unique to Tg-5xFAD/MBP<sup>-/-</sup> mice. We bred Tg-SwDI mice, another model of early-onset A $\beta$

accumulation and deposition, with MBP<sup>-/-</sup> mice. Like bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice, bigenic Tg-SwDI/MBP<sup>-/-</sup> mice also exhibited decreased A $\beta$  levels and A $\beta$  deposition at 2 to 3 months of age (data not shown).

A lowering of cerebral A $\beta$  levels can result from reduced expression of A $\beta$ PP and production of the peptide. However, the levels of A $\beta$ PP protein, A $\beta$ PP CTFs and the presence of intraneuronal A $\beta$  were similar in the brains of Tg-5xFAD mice and bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice, suggesting that the reductions in A $\beta$  were not the consequence of decreased production in the absence of MBP. Alternatively, A $\beta$  reductions in the brain can arise, owing to increased clearance through established efflux pathways. For example, through one route A $\beta$  initially released by neurons enters the interstitial fluid, which drains to the CSF [48]. Yet another mechanism involves active transport of A $\beta$  across the blood-brain barrier into the circulation that is mediated by known A $\beta$  receptors, including low-density lipoprotein receptor-related protein 1 (LRP1) and P-glycoprotein [64,65]. However, we found no increase in the levels of A $\beta$  in the CSF or plasma of bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice, arguing against increased A $\beta$  efflux in the absence of MBP.

Another recognized clearance mechanism of A $\beta$  in brain involves a broad class of A $\beta$ -degrading enzymes




that are largely released by activated neuroinflammatory cells [66]. In the bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice, we found markedly elevated staining for reactive astrocytes and activated microglia compared with Tg-5xFAD mice (Panels C of Figures 6 and 7). Once Tg-5xFAD mice age and develop numerous fibrillar Aβ plaques, they exhibit a robust neuroinflammatory response to these plaques that is characterized by reactive astrocytes and activated microglia [38]. Yet, in our study the young animals at two months of age are just beginning to develop amyloid plaques and neuroinflammatory cells are scarce or absent (Panels B of Figures 6 and 7). Furthermore, the increased immunostaining of activated glial markers in bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice was not observed around the few plaques that had developed. However, the increase in neuroinflammatory cells was similarly observed in the MBP<sup>-/-</sup> mice alone (Panels D of Figures 6 and 7); this is consistent with previous reports regarding these mice [51]. Activation of glial cells is a common

phenomenon in neurodegenerative diseases, including AD, multiple sclerosis, and traumatic brain injury. A number of other mouse models with deficiencies in myelination from different causes including *jimpy*, *MBP<sup>mla</sup>*, and *quaking* were all found to exhibit glial activation [67,68]. Although it is not well understood, the increase in neuroinflammatory cells observed in the MBP<sup>-/-</sup> mice appears to be a pleiotropic effect due to the loss of MBP and abnormal myelination, thereby disrupting normal interaction between glial cells. Indeed, it has been suggested that a proliferation of mixed-phenotype glial cells, which were found to be increased in the pathogenic white matter, contribute to this gliosis in MBP<sup>-/-</sup> mice [69].

In any case, reactive astrocytes and activated microglia both produce MMP-2 and MMP-9 in a number of CNS disorders, including AD and multiple sclerosis [70-73]. In regards to Aβ-degrading enzymes, MMP-2 and MMP-9 are distinctive in that both can degrade soluble Aβ peptides and fibrillar plaque Aβ [74-78]. While we did not





find elevated MMP-2 levels in the brains of bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice, we did observe a significant increase in MMP-9 levels, as assessed by zymography and immunoblotting (Figure 8). Although it is possible that other A $\beta$ -degrading enzymes could be elevated in Tg-5xFAD/MBP<sup>-/-</sup> mice, quantitative PCR analysis did not reveal increased expression of some of the more common enzymes including insulin-degrading enzyme, neprilysin, or angiotensin-converting enzyme (data not shown). This suggests that elevated MMP-9, produced by reactive astrocytes and activated microglia as a consequence of the absence of MBP, could contribute to the decreased A $\beta$  levels observed in Tg-5xFAD/MBP<sup>-/-</sup> mice.

A goal of this study was to investigate the potential consequences of MBP-A $\beta$  interactions in the brain. Based on our previous *in vitro* work showing that MBP strongly binds A $\beta$  and inhibits fibrillar assembly [22,29,42], one prediction is that in the absence of MBP there could be greater accumulation of fibrillar A $\beta$ . On the contrary, as shown here, in the complete absence of MBP there was a significant reduction in the accumulation of A $\beta$ . However, the A $\beta$  reduction observed in the bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice is more likely to be an indirect pleiotropic effect of the absence of MBP and proper myelination, leading to glial activation and increased A $\beta$  degrading enzymes, rather than a consequence of the direct loss of interaction between MBP and A $\beta$ . Recently, we identified specific residues in MBP that are essential for A $\beta$  binding and fibril assembly inhibition [79]. To overcome the significant

limitations of MBP<sup>-/-</sup> mice, future efforts are focused on utilizing novel mice that harbor specific mutations of the residues in MBP involved in A $\beta$  binding and fibril inhibition. In contrast with the MBP<sup>-/-</sup> mice, these new mutant MBP mice are largely expected to retain normal physiological functions of MBP but will be devoid of the ability to interact with A $\beta$ . Alternatively, efforts are also focused on characterizing novel transgenic mice that over-express biologically active fragment of MBP in brain. Together, the novel MBP knock-in mutant mice and MBP-expressing transgenic mice, crossed with A $\beta$ PP transgenic mice, will provide more suitable models for studying *in vivo* MBP-A $\beta$  relationships, thereby enabling us to gain insight into A $\beta$  assembly in brain and a potential therapeutic role of MBP as an A $\beta$  fibril assembly inhibitor.

### Conclusions

The primary findings of this study show that in the absence of MBP there is decreased accumulation and deposition of A $\beta$  in Tg-5xFAD mice. The decrease in A $\beta$  was not a consequence of reduced A $\beta$ PP expression or processing or of increased peptide clearance through plasma and CSF efflux pathways. However, there were elevated reactive astrocytes and microglia accompanied by increased expression of A $\beta$ -degrading enzyme MMP-9 in bigenic Tg-5xFAD/MBP<sup>-/-</sup> mice. The absence of MBP promotes a neuroinflammatory environment that can reduce A $\beta$  accumulation in the brains of transgenic mice.

## Abbreviations

A $\beta$ PP: amyloid precursor protein; A $\beta$ : amyloid beta protein; apoE: apolipoprotein E; MBP: myelin basic protein; C83: C-terminal fragment of A $\beta$ PP generated by  $\alpha$  secretase cleavage; C99: C-terminal fragment of A $\beta$ PP generated by  $\beta$  secretase cleavage; CNS: central nervous system; CSF: cerebrospinal fluid; CTF: C-terminal fragment; ELISA: enzyme-linked immunosorbent assay; GFAP: glial fibrillary acidic protein; iA $\beta$ : intracellular amyloid beta protein; MBP: myelin basic protein; MMP: matrix metalloproteinase; OC: a fibril specific, conformation dependent antibody used in intracellular A $\beta$  detection; PBS: phosphate-buffered saline; SEM: standard error of the mean; TBS: tris-buffered saline.

## Competing interests

Both authors declare that they have no competing interests.

## Authors' contributions

MO designed, performed experiments and analyzed the data. WEVN obtained funding, helped conceive the project and interpreted data. Both authors read and approved the final manuscript.

## Acknowledgements

This work was supported in part by National Institutes of Health grants R21-NS079951 and R21-AG039215. Antibody reagents for the A $\beta$  ELISAs were generously provided by Lilly Research Laboratories.

Received: 2 August 2013 Accepted: 27 September 2013

Published: 5 November 2013

## References

- Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiaz EA, Denis P, Teplow DB, Ross S, Amarante P, Loeloff R, Luo Y, Fisher S, Fuller J, Edenson S, Lile J, Jarosinski MA, Biere AL, Curran E, Burgess T, Louis JC, Collins F, Treanor J, Rogers G, Citron M:  $\beta$ -secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* 1999, **286**:735–741.
- De Strooper B, Saffig P, Craessaerts K, Vanderstichele H, Guhde G, Annaert W, Von Figura K, Van Leuven F: Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* 1998, **391**:387–390.
- Edbauer D, Winkler E, Regula JT, Pesold B, Steiner H, Haass C: Reconstitution of  $\gamma$ -secretase activity. *Nat Cell Biol* 2003, **5**:486–488.
- Benilova I, Karran E, De Strooper B: The toxic A $\beta$  oligomer and Alzheimer's disease: an emperor in need of clothes. *Nat Neurosci* 2012, **15**:349–357.
- Haass C, Selkoe DJ: Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid  $\beta$ -peptide. *Nat Rev Mol Cell Biol* 2007, **8**:101–112.
- Selkoe DJ: Soluble oligomers of the amyloid  $\beta$ -protein impair synaptic plasticity and behavior. *Behav Brain Res* 2008, **192**:106–113.
- Urbanc B, Cruz L, Le R, Sanders J, Ashe KH, Duff K, Stanley HE, Irizarry MC, Hyman BT: Neurotoxic effects of thioflavin S-positive amyloid deposits in transgenic mice and Alzheimer's disease. *Proc Natl Acad Sci USA* 2002, **99**:13990–13995.
- McLellan ME, Kajdasz ST, Hyman BT, Bacskai BJ: *In vivo* imaging of reactive oxygen species specifically associated with thioflavin S-positive amyloid plaques by multiphoton microscopy. *J Neurosci* 2003, **23**:2212–2217.
- Strittmatter WJ, Saunders AM, Schmechel D, Pericak-Vance M, Enghild J, Salvesen GS, Roses AD: Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc Natl Acad Sci USA* 1993, **90**:1977–1981.
- LaDu MJ, Falduto MT, Manelli AM, Reardon CA, Getz GS, Frail DE: Isoform-specific binding of apolipoprotein E to beta-amyloid. *J Biol Chem* 1994, **269**:23403–23406.
- Holtzman DM: Role of apoE/A $\beta$  interactions in the pathogenesis of Alzheimer's disease and cerebral amyloid angiopathy. *J Mol Neurosci* 2001, **17**:147–155.
- Matsubara E, Frangione B, Ghiso J: Characterization of apolipoprotein J-Alzheimer's A $\beta$  interaction. *J Biol Chem* 1995, **270**:7563–7567.
- DeMattos RB, O'Dell MA, Parsadanian M, Taylor JW, Harmony JA, Bales KR, Paul SM, Aronow BJ, Holtzman DM: Clusterin promotes amyloid plaque formation and is critical for neuritic toxicity in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci USA* 2002, **99**:10843–10848.
- Magrane J, Smith RC, Walsh K, Querfurth HW: Heat shock protein 70 participates in the neuroprotective response to intracellularly expressed  $\beta$ -amyloid in neurons. *J Neurosci* 2004, **24**:1700–1706.
- Evans CG, Wisen S, Gestwicki JE: Heat shock proteins 70 and 90 inhibit early stages of amyloid  $\beta$ -(1–42) aggregation *in vitro*. *J Biol Chem* 2006, **281**:33182–33191.
- Yamamoto N, Hirabayashi Y, Amari M, Yamaguchi H, Romanov G, Van Nostrand WE, Yanagisawa K: Assembly of hereditary amyloid  $\beta$ -protein variants in the presence of favorable gangliosides. *FEBS Lett* 2005, **579**:2185–2190.
- Schwarzman AL, Goldgaber D: Interaction of transthyretin with amyloid  $\beta$ -protein: binding and inhibition of amyloid formation. *Ciba Found Symp* 1996, **199**:146–160. discussion 160–144.
- Buxbaum JN, Ye Z, Reixach N, Friske L, Levy C, Das P, Golde T, Masliah E, Roberts AR, Bartfai T: Transthyretin protects Alzheimer's mice from the behavioral and biochemical effects of A $\beta$  toxicity. *Proc Natl Acad Sci USA* 2008, **105**:2681–2686.
- van Horsen J, Wesseling P, van den Heuvel LP, de Waal RM, Verbeek MM: Heparan sulphate proteoglycans in Alzheimer's disease and amyloid-related disorders. *Lancet Neurol* 2003, **2**:482–492.
- Yanagisawa K, Odaka A, Suzuki N, Ihara Y: GM1 ganglioside-bound amyloid  $\beta$ -protein (A $\beta$ ): a possible form of preamyloid in Alzheimer's disease. *Nat Med* 1995, **1**:1062–1066.
- Chi EY, Frey SL, Lee KY: Ganglioside G(M1)-mediated amyloid-beta fibrillogenesis and membrane disruption. *Biochemistry* 2007, **46**:1913–1924.
- Hoos MD, Ahmed M, Smith SO, Van Nostrand WE: Inhibition of familial cerebral amyloid angiopathy mutant amyloid  $\beta$ -protein fibril assembly by myelin basic protein. *J Biol Chem* 2007, **282**:9952–9961.
- Boggs JM: Myelin basic protein: a multifunctional protein. *Cell Mol Life Sci* 2006, **63**:1945–1961.
- Campagnoni AT, Pribyl TM, Campagnoni CW, Kampf K, Amur-Umarjee S, Landry CF, Handley VW, Newman SL, Garbay B, Kitamura K: Structure and developmental regulation of Golli-mbp, a 105-kilobase gene that encompasses the myelin basic protein gene and is expressed in cells in the oligodendrocyte lineage in the brain. *J Biol Chem* 1993, **268**:4930–4938.
- Kamholz J, de Ferra F, Puckett C, Lazzarini R: Identification of three forms of human myelin basic protein by cDNA cloning. *Proc Natl Acad Sci USA* 1986, **63**:4962–4966.
- Roth HJ, Kronquist KE, Kerlero De Rosbo N, Crandall BF, Campagnoni AT: Evidence for the expression of four myelin basic protein variants in the developing human spinal cord through cDNA cloning. *J Neurosci Res* 1987, **17**:321–328.
- Harauz G, Ishiyama N, Hill CM, Bates IR, Libich DS, Fares C: Myelin basic protein-diverse conformational states of an intrinsically unstructured protein and its roles in myelin assembly and multiple sclerosis. *Micron* 2004, **35**:503–542.
- de Ferra F, Engh H, Hudson L, Kamholz J, Puckett C, Molineaux S, Lazzarini RA: Alternative splicing accounts for the four forms of myelin basic protein. *Cell* 1985, **43**:721–727.
- Hoos MD, Ahmed M, Smith SO, Van Nostrand WE: Myelin basic protein binds to and inhibits the fibrillar assembly of A $\beta$ 42 *in vitro*. *Biochemistry* 2009, **48**:4720–4727.
- Bartzokis G, Lu PH, Geschwind DH, Edwards N, Mintz J, Cummings JL: Apolipoprotein E genotype and age-related myelin breakdown in healthy individuals: implications for cognitive decline and dementia. *Arch Gen Psychiatry* 2006, **63**:63–72.
- Liu MC, Akle V, Zheng W, Kitlen J, O'Steen B, Larner SF, Dave JR, Tortella FC, Hayes RL, Wang KK: Extensive degradation of myelin basic protein isoforms by calpain following traumatic brain injury. *J Neurochem* 2006, **98**:700–712.
- Bartzokis G, Lu PH, Geschwind DH, Tingus K, Huang D, Mendez MF, Edwards N, Mintz J: Apolipoprotein E affects both myelin breakdown and cognition: implications for age-related trajectories of decline into dementia. *Biol Psychiatry* 2007, **62**:1380–1387.
- Bartzokis G, Sultzer D, Lu PH, Nuechterlein KH, Mintz J, Cummings JL: Heterogeneous age-related breakdown of white matter structural integrity: implications for cortical 'disconnection' in aging and Alzheimer's disease. *Neurobiol Aging* 2004, **25**:843–851.
- Marner L, Nyengaard JR, Tang Y, Pakkenberg B: Marked loss of myelinated nerve fibers in the human brain with age. *J Comp Neurol* 2003, **462**:144–152.
- Roher AE, Weiss N, Kokjohn TA, Kuo YM, Kalback W, Anthony J, Watson D, Lehrs DC, Sue L, Walker D, Emmerling M, Goux W, Beach T: Increased A $\beta$

- peptides and reduced cholesterol and myelin proteins characterize white matter degeneration in Alzheimer's disease. *Biochemistry* 2002, **41**:11080–11090.
36. Mitew S, Kirkcaldie MT, Halliday GM, Shepherd CE, Vickers JC, Dickson TC: Focal demyelination in Alzheimer's disease and transgenic mouse models. *Acta Neuropathol* 2010, **119**:567–577.
  37. Molineaux SM, Engh H, de Ferra F, Hudson L, Lazzarini RA: Recombination within the myelin basic protein gene created the dysmyelinating shiverer mouse mutation. *Proc Natl Acad Sci USA* 1986, **83**:7542–7546.
  38. Oakley H, Cole SL, Logan S, Maus E, Shao P, Craft J, Guillozet-Bongaarts A, Ohno M, Disterhoft J, Van Eldik L, Berry R, Vassar R: Intraneuronal  $\beta$ -amyloid aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial Alzheimer's disease mutations: potential factors in amyloid plaque formation. *J Neurosci* 2006, **26**:10129–10140.
  39. Liu L, Duff K: A technique for serial collection of cerebrospinal fluid from the cisterna magna in mouse. *J Vis Exp* 2008, **21**:960.
  40. Davis J, Xu F, Deane R, Romanov G, Previti ML, Zeigler K, Zlokovic BV, Van Nostrand WE: Early-onset and robust cerebral microvascular accumulation of amyloid  $\beta$ -protein in transgenic mice expressing low levels of a vasculotropic Dutch/lowa mutant form of amyloid  $\beta$ -protein precursor. *J Biol Chem* 2004, **279**:20296–20306.
  41. DeMattos RB, Bales KR, Parsadanian M, O'Dell MA, Foss EM, Paul SM, Holtzman DM: Plaque-associated disruption of CSF and plasma amyloid- $\beta$  ( $\text{A}\beta$ ) equilibrium in a mouse model of Alzheimer's disease. *J Neurochem* 2002, **81**:229–236.
  42. Liao MC, Hoos MD, Aucoin D, Ahmed M, Davis J, Smith SO, Van Nostrand WE: N-terminal domain of myelin basic protein inhibits amyloid  $\beta$ -protein fibril assembly. *J Biol Chem* 2010, **285**:35590–35598.
  43. Oddo S, Billings L, Kesslak JP, Cribbs DH, LaFerla FM:  $\text{A}\beta$  immunotherapy leads to clearance of early, but not late, hyperphosphorylated tau aggregates via the proteasome. *Neuron* 2004, **43**:321–332.
  44. Gyure KA, Durham R, Stewart WF, Smialek JE, Troncoso JC: Intraneuronal  $\text{A}\beta$ -amyloid precedes development of amyloid plaques in Down syndrome. *Arch Pathol Lab Med* 2001, **125**:489–492.
  45. Wirths O, Multhaup G, Czech C, Blanchard V, Moussaoui S, Tremp G, Pradier L, Beyreuther K, Bayer TA: Intraneuronal  $\text{A}\beta$  accumulation precedes plaque formation in  $\beta$ -amyloid precursor protein and presenilin-1 double-transgenic mice. *Neurosci Lett* 2001, **306**:116–120.
  46. Kaye R, Head E, Sarsoza F, Saing T, Cotman CW, Necula M, Margol L, Wu J, Breydo L, Thompson JL, Rasool S, Gurlo T, Butler P, Glabe CG: Fibril specific, conformation dependent antibodies recognize a generic epitope common to amyloid fibrils and fibrillar oligomers that is absent in prefibrillar oligomers. *Mol Neurodegener* 2007, **2**:18.
  47. Shibata M, Yamada S, Kumar SR, Calero M, Bading J, Frangione B, Holtzman DM, Miller CA, Strickland DK, Ghiso J, Zlokovic BV: Clearance of Alzheimer's amyloid-ss(1–40) peptide from brain by LDL receptor-related protein-1 at the blood–brain barrier. *J Clin Invest* 2000, **106**:1489–1499.
  48. Silverberg GD, Mayo M, Saul T, Rubenstein E, McGuire D: Alzheimer's disease, normal-pressure hydrocephalus, and senescent changes in CSF circulatory physiology: a hypothesis. *Lancet Neurol* 2003, **2**:506–511.
  49. Zlokovic BV: Clearing amyloid through the blood–brain barrier. *J Neurochem* 2004, **89**:807–811.
  50. Privat A, Jacque C, Bourre JM, Dupouey P, Baumann N: Absence of the major dense line in myelin of the mutant mouse 'shiverer'. *Neurosci Lett* 1979, **12**:107–112.
  51. Nagaïke K, Mikoshiba K, Tsukada Y: Dysmyelination of shiverer mutant mice *in vivo* and *in vitro*. *J Neurochem* 1982, **39**:1235–1241.
  52. Gottschall PE, Yu X, Bing B: Increased production of gelatinase B (matrix metalloproteinase-9) and interleukin-6 by activated rat microglia in culture. *J Neurosci Res* 1995, **42**:335–342.
  53. Muir EM, Adcock KH, Morgenstern DA, Clayton R, von Stillfried N, Rhodes K, Ellis C, Fawcett JW, Rogers JH: Matrix metalloproteinases and their inhibitors are produced by overlapping populations of activated astrocytes. *Brain Res Mol Brain Res* 2002, **100**:103–117.
  54. Sbai O, Ould-Yahoui A, Ferhat L, Gueye Y, Bernard A, Charrat E, Mehanna A, Risso JJ, Chauvin JP, Fenouillet E, Rivera S, Khrestchatisky M: Differential vesicular distribution and trafficking of MMP-2, MMP-9, and their inhibitors in astrocytes. *Glia* 2010, **58**:344–366.
  55. Beems T, Simons KS, Van Geel WJ, De Reus HP, Vos PE, Verbeek MM: Serum- and CSF-concentrations of brain specific proteins in hydrocephalus. *Acta Neurochir (Wien)* 2003, **145**:37–43.
  56. Abdo WF, van de Warrenburg BP, Munneke M, van Geel WJ, Bloem BR, Kremer HP, Verbeek MM: CSF analysis differentiates multiple-system atrophy from idiopathic late-onset cerebellar ataxia. *Neurology* 2006, **67**:474–479.
  57. Su E, Bell MJ, Kochanek PM, Wisniewski SR, Bayir H, Clark RS, Adelson PD, Tyler-Kabara EC, Janesko-Feldman KL, Berger RP: Increased CSF concentrations of myelin basic protein after TBI in infants and children: absence of significant effect of therapeutic hypothermia. *Neurocrit Care* 2012, **17**:401–407.
  58. Lamers KJ, de Reus HP, Jongen PJ: Myelin basic protein in CSF as indicator of disease activity in multiple sclerosis. *Mult Scler* 1998, **4**:124–126.
  59. Landry CF, Ellison JA, Pribyl TM, Campagnoni C, Kampf K, Campagnoni AT: Myelin basic protein gene expression in neurons: developmental and regional changes in protein targeting within neuronal nuclei, cell bodies, and processes. *J Neurosci* 1996, **16**:2452–2462.
  60. Landry CF, Pribyl TM, Ellison JA, Givogri MI, Kampf K, Campagnoni CW, Campagnoni AT: Embryonic expression of the myelin basic protein gene: identification of a promoter region that targets transgene expression to pioneer neurons. *J Neurosci* 1998, **18**:7315–7327.
  61. Filipovic Z, Zecevic N: Interaction between microglia and oligodendrocyte cell progenitors involves Gollin proteins. *Ann N Y Acad Sci* 2005, **1048**:166–174.
  62. Harauz G, Ladizhansky V, Boggs JM: Structural polymorphism and multifunctionality of myelin basic protein. *Biochemistry* 2009, **48**:8094–8104.
  63. Chernoff GF: Shiverer: an autosomal recessive mutant mouse with myelin deficiency. *J Hered* 1981, **72**:128.
  64. Lam FC, Liu R, Lu P, Shapiro AB, Renoir JM, Sharom FJ, Reiner PB:  $\beta$ -Amyloid efflux mediated by p-glycoprotein. *J Neurochem* 2001, **76**:1121–1128.
  65. Deane R, Wu Z, Sagare A, Davis J, Du Yan S, Hamm K, Xu F, Parisi M, LaRue B, Hu HW, Spijkers P, Guo H, Song X, Lenting PJ, Van Nostrand WE, Zlokovic BV: LRP/amyloid  $\beta$ -peptide interaction mediates differential brain efflux of  $\text{A}\beta$  isoforms. *Neuron* 2004, **43**:333–344.
  66. Miners JS, Baig S, Palmer J, Palmer LE, Kehoe PG, Love S:  $\text{A}\beta$ -degrading enzymes in Alzheimer's disease. *Brain Pathol* 2008, **18**:240–252.
  67. Nagara H, Suzuki K: Chronological study of oligodendroglial alterations and myelination in quaking mice. *Neuropathol Appl Neurobiol* 1981, **7**:135–149.
  68. Best TT, Skoff RP, Bartlett WP: Astroglial plasticity in hemizygous and heterozygous jimpy mice. *Int J Dev Neurosci* 1988, **6**:39–57.
  69. Dyer CA, Kandler A, Jean-Guillaume D, Awatramani R, Lee A, Mason LM, Kamholz J: GFAP-positive and myelin marker-positive glia in normal and pathologic environments. *J Neurosci Res* 2000, **60**:412–426.
  70. Maeda A, Sobel RA: Matrix metalloproteinases in the normal human central nervous system, microglial nodules, and multiple sclerosis lesions. *J Neuropathol Exp Neurol* 1996, **55**:300–309.
  71. Liuzzi GM, Latronico T, Fasano A, Carlone G, Riccio P: Interferon-beta inhibits the expression of metalloproteinases in rat glial cell cultures: implications for multiple sclerosis pathogenesis and treatment. *Mult Scler* 2004, **10**:290–297.
  72. del Zoppo GJ, Milner R, Mabuchi T, Hung S, Wang X, Berg GJ, Koziol JA: Microglial activation and matrix protease generation during focal cerebral ischemia. *Stroke* 2007, **38**:646–651.
  73. del Zoppo GJ, Frankowski H, Gu YH, Osada T, Kanazawa M, Milner R, Wang X, Hosomi N, Mabuchi T, Koziol JA: Microglial cell activation is a source of metalloproteinase generation during hemorrhagic transformation. *J Cereb Blood Flow Metab* 2012, **32**:919–932.
  74. Roher AE, Kasunic TC, Woods AS, Cotter RJ, Ball MJ, Fridman R: Proteolysis of A  $\beta$  peptide from Alzheimer disease brain by gelatinase A. *Biochem Biophys Res Commun* 1994, **205**:1755–1761.
  75. Backstrom JR, Lim GP, Cullen MJ, Tokes ZA: Matrix metalloproteinase-9 (MMP-9) is synthesized in neurons of the human hippocampus and is capable of degrading the amyloid- $\beta$  peptide (1–40). *J Neurosci* 1996, **16**:7910–7919.
  76. Yan P, Hu X, Song H, Yin K, Bateman RJ, Cirrito JR, Xiao Q, Hsu FF, Turk JW, Xu J, Hsu CY, Holtzman DM, Lee JM: Matrix metalloproteinase-9 degrades amyloid- $\beta$  fibrils *in vitro* and compact plaques *in situ*. *J Biol Chem* 2006, **281**:24566–24574.
  77. Turk J, Cirrito JR, Yan P, Hu X, Xiao Q, Pan X, Bateman R, Song H, Hsu FF, Turk J, Xu J, Hsu CY, Mills JC, Holtzman DM, Lee JM: Matrix metalloproteinases expressed by astrocytes mediate extracellular amyloid- $\beta$  peptide catabolism. *J Neurosci* 2006, **26**:10939–10948.
  78. White AR, Du T, Laughton KM, Volitakis I, Sharples RA, Xilinas ME, Hoke DE, Holsinger RM, Evin G, Cherny RA, Hill AF, Barnham KJ, Li QX, Bush AI,

Masters CL: Degradation of the Alzheimer disease amyloid  $\beta$ -peptide by metal-dependent up-regulation of metalloprotease activity. *J Biol Chem* 2006, **281**:17670–17680.

79. Kotarba AE, Aucoin DS, Hoos MD, Smith SO, Van Nostrand WE: Fine mapping of the amyloid ss-protein binding site on myelin basic protein. *Biochemistry* 2013, **52**(15):2565–2573.

doi:10.1186/1742-2094-10-134

**Cite this article as:** Ou-Yang and Van Nostrand: The absence of myelin basic protein promotes neuroinflammation and reduces amyloid  $\beta$ -protein accumulation in Tg-5xFAD mice. *Journal of Neuroinflammation* 2013 **10**:134.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

