

CASE REPORT

Feline restrictive orbital myofibroblastic sarcoma treated with toceranib phosphate (Palladia) as adjuvant chemotherapy: A case study of one cat

Hiroyuki Komatsu¹  | Asuka Ueshima² | Yoshitaka Kobayashi^{1,2}  |
Yumiko Shimoyama³ | Naoaki Takiyama⁴ | Kenji Rimpō²

¹Animal Eye Care-Tokyo Animal Eye Clinic, Tokyo, Japan

²Saitama Animal Medical Center, Saitama, Japan

³IDEXX Laboratories, Tokyo, Japan

⁴Veterinary Eye Clinic Nagoya, Nagoya, Japan

Correspondence

Hiroyuki Komatsu, Animal Eye Care-Tokyo Animal Eye Clinic, 1-12-12, Kaminoge, Setagaya-ku, Tokyo 158-0093, Japan.

Email: v06058@gmail.com

Key Clinical Message

This is the first case report of treatment with toceranib phosphate as postsurgical adjuvant chemotherapy for advanced FROMS. This reported case highlights the need for further studies on the efficacy of toceranib phosphate as adjuvant chemotherapy for FROMS.

Abstract

Feline restrictive orbital myofibroblastic sarcoma (FROMS) is a rare aggressive tumor in cats. We explored the effectiveness of using toceranib phosphate as postsurgical adjuvant chemotherapy for advanced FROMS in a 7-year-old cat. Despite treatment, the cat died 4 months after surgery. This report highlights the need for further studies on the efficacy of toceranib phosphate as adjuvant chemotherapy for FROMS.

KEYWORDS

adjuvant chemotherapy, feline orbital tumor, feline restrictive orbital myofibroblastic sarcoma, toceranib phosphate

1 | INTRODUCTION

Most orbital diseases are caused by infection, inflammation, or neoplasia, and malignant orbital tumors are common in dogs and cats.^{1,2} In dogs, various types of tumors with different origins occur in the orbit, including mesenchymal origin, epithelial origin, and miscellaneous origin.² In cats, in contrast, epithelial tumors such as squamous cell carcinoma and round cell tumors such as lymphoma are more common orbital tumors, while mesenchymal

tumors are less common.³ In recent years, there are some reports of a rare orbital mesenchymal tumor called feline restrictive orbital myofibroblastic sarcoma (FROMS), with clinical characteristics of eyelid restriction leading to exposure keratitis without formation of an orbital mass.⁴⁻⁸ However, detailed data of this tumor including the origin and tumorigenesis have not been elucidated.

Surgical resection is often selected as the first choice in the treatment of orbital tumors in dogs and cats.^{2,9} Other treatment options include radiation therapy and

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2023 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

chemotherapy, which may be performed after surgery as adjuvant therapy.^{9,10} In dogs with orbital tumor, cases with malignant orbital tumor treated with orbitotomy followed by adjuvant radiation therapy and chemotherapy have been reported.¹ However, the efficacy of postsurgical adjuvant therapy for dogs and cats remains unclear due to a paucity of clinical reports on adjuvant therapy.¹⁰

The use of molecular targeted drugs, one type of chemotherapy, has been reported in veterinary medicine.^{11–13} Toceranib, which is one of the molecular-targeted drugs, is a tyrosine kinase receptor inhibitor and has been shown to inhibit KIT, platelet-derived growth factor receptor, and vascular endothelial growth factor receptor.^{14–16} Clinical use of toceranib in cats for the treatment of mast cell tumors,^{17,18} adenocarcinoma,^{19,20} gastrointestinal stromal tumor,²¹ pancreatic carcinoma,²² mammary carcinoma,²³ squamous cell carcinoma,^{24–26} and feline injection site-associated sarcomas (FISS)²⁷ has been reported. However, the number of these reports in cats is fewer compared with that in dogs. In addition, there are no reports of adjuvant chemotherapy using molecular-targeted drugs for FROMS. Herein, we present the first report of using toceranib as postsurgical adjuvant chemotherapy for the treatment of FROMS in a cat. The objective of this case study was to explore the potential of expanding treatment options for FROMS in cats using a molecular-targeted drug as adjuvant chemotherapy.

2 | CASE PRESENTATION

A seven-year-old neutered male American shorthair cat was referred to the Saitama Animal Medical Center with symptoms of blepharospasm in the right eye. Ophthalmic examination revealed superficial corneal ulceration with mild eyelid restriction in the right eye, but no abnormality was observed in the left eye (Figure 1). Treatment with antibiotic and hyaluronate eye drops was started, but the corneal ulceration in the right eye did not improve at follow-up 3 weeks later. In addition, entropion caused by eyelid restriction persisted. Partial right eyelid tarsorrhaphy was performed to treat the corneal ulcers. However, vision was eventually lost in the right eye due to severe exposure keratitis caused by complete eyelid restriction, and resistance to retropulsion of the globe was observed at follow-up. Orbital disease was suspected from these clinical findings. Ultrasound examination was performed, but no apparent mass was observed in the globe or in the orbit (Figure 2A).

Given the lack of improvement in clinical symptoms by medical treatment, surgical resection following computed tomography (CT) was scheduled on Day 138 after initial presentation. CT revealed a space-occupying lesion in the right orbit, with infiltration extending from the eyelid to the lips (Figure 2B). There were no osteolytic lesions in the orbital bone and no sign of abdominal or thoracic metastasis. Following the CT examination, orbital exenteration of the right eye was performed. The orbit contained



FIGURE 1 Clinical findings of the cat during the clinical course. (A) At initial presentation, blepharospasm in the right eye was observed. (B) On Day 19, corneal ulceration at dorsal site did not improve, and restriction of the eyelids was prominent. (C, D) On Day 75, exposure keratitis in the right eye deteriorated due to persisting eyelid restriction, and resistance to posterior ocular compression was observed.

multiple nodular lesions, and these were removed to the extent possible. All specimens were fixed in 10% formalin, embedded in paraffin, sectioned at 4 μ m, and subject to hematoxylin and eosin (HE), and immunohistochemical staining.

Histopathological examination of HE stained sections revealed the proliferation of atypical spindle tumor cells infiltrating the skeletal muscle in the orbital tissue with thin collagenous materials between spindle cells and moderate lymphoplasmacytic cell infiltration (Figure 3A), but there was no invasion into the globe. Neoplastic spindle cells resemble reactive fibroblasts or myofibroblasts with mild anisocytosis and anisokaryosis. These spindle cells have

a moderate amount of elongated eosinophilic cytoplasm and ovoid nuclei with finely stripped chromatin and one to two nucleoli. There were 4–5 mitotic figures in tumor cells per 10 high-power fields. Immunohistochemical staining showed that the majority of tumor cells were positive for alpha-smooth muscle actin (α SMA) staining, which is commonly used as a marker of myofibroblasts. These findings suggest that the tumor exhibited myofibroblastic differentiation. Based on these clinical and histopathological findings, the tumors were diagnosed as FROMS.

After surgery, subcutaneous infiltration of tumor cells expanded on the right side of the upper lip, leading to feeding difficulty. A gastrostomy tube was placed under general




FIGURE 2 Imaging findings. (A) On Day 75, ultrasound examination detected no obvious mass in both the globe and the orbit. (B) On Day 138, a computed tomography scan revealed a high-intensity space-occupying lesion in the orbit of right eye (*).




FIGURE 3 Histopathological findings of the tumor. (A) Hematoxylin–eosin staining shows proliferation of atypical spindle tumor cells (yellow arrow) with eosinophilic thin collagenous materials (white arrow) between spindle tumor cells and mild lymphoplasmacytic cell infiltration (red arrow). (B) Immunohistochemical staining shows moderate cytoplasmic labeling of alpha-smooth muscle Actin in majority of the spindle tumor cells. Scale bar = 200 μ m (100 \times).

anesthesia. Three weeks after surgery, toceranib phosphate (Palladia; Zoetis) 2.17 mg/kg was initiated three times per week as adjuvant chemotherapy. Toceranib administration was continued for 1 month. However, the residual tumor continued to increase in size, and adjuvant chemotherapy was deemed ineffective and was discontinued. Subsequently, bleeding from the tumor occurred, which severely deteriorated the cat's quality of life. Eventually, the cat died 4 months after surgery, 257 days after initial presentation.

3 | DISCUSSION

At first presentation, the cat showed corneal ulcers and exposure keratitis caused by eyelid restriction. Despite medical therapy, the clinical symptoms deteriorated, and CT images demonstrated an orbital mass. Orbital exenteration was performed, and histopathological examination revealed the presence of spindle tumor cells. After surgery, toceranib was administered due to enlargement of the residual mass. This is the first report of using toceranib as adjuvant chemotherapy for treating FROMS in a cat.

Feline orbital neoplasms are commonly diagnosed in clinical practice.²⁸ Isaza et al.³ reported the incidence of various types of orbital tumor in cats as follows: round cell tumors 47%, epithelial tumors 38%, mesenchymal tumors 13%, and neurologic origin tumors 1%. In their report, the most common diagnoses were lymphoma and squamous cell carcinoma. Among the mesenchymal tumors, FROMS was reported most frequently, and other malignant tumors including fibrosarcoma, osteosarcoma, unspecified sarcoma, and spindle cell tumors were also reported.³ FROMS has unique clinical characteristics marked by eyelid restriction without discrete orbital mass formation. Based on these characteristic findings, this disease was previously considered to be an inflammatory disorder, but was redefined as a neoplastic lesion by Bell et al.⁴ in 2011. FROMS is known to originate from the orbit and actively invades connective tissues of the orbit including eyelids and conjunctiva.^{4,7} Cases of the tumor invading the contralateral eye causing bilateral symptoms have also been reported.⁶ Histopathology examination of FROMS shows infiltrative proliferation of low-grade atypical spindle cells with collagen matrix in the orbit and substantia propria of conjunctival tissue.⁴ Immunohistochemistry of FROMS shows positive staining for S100, vimentin, and α SMA.⁴ From these histopathological findings, FROMS is suspected to originate from myofibroblast cells. On the other hand, fibrosarcoma is a malignant tumor originating from fibrous connective tissue and typically consists of highly atypical spindle cells with abundant mitosis. Feline fibrosarcoma usually forms a large mass with a strong infiltrate into the adjacent tissues. Other orbital mesenchymal tumors in cats, including osteosarcoma and spindle cell tumors, show

various degrees of differentiation.³ In our case, proliferation of fibroblast-like spindle cells was observed by histopathological examination, leading to a high suspicion of mesenchymal tumor. In addition, positive staining for α SMA suggested that the tumor was derived from myofibroblasts. These findings were consistent with the histopathological features of FROMS. Regarding clinical presentation, findings such as eyelid restriction and the lack of discrete orbital mass at the initial ultrasound examination were also consistent with the clinical features of FROMS. Hence, a final diagnosis of FROMS was made. However, the mitotic count in this case was relatively high compared with previous reports. These differences from previous reports may imply the existence of subtypes of FROMS.

Regarding treatment, surgical resection is often selected as the first option for feline orbital tumors. As most orbital tumors in cats are malignant, frequent local recurrence, and metastasis are a clinical issue.²⁹ Therefore, surgical resection together with other effective treatments are needed to control the progression of the disease. Additional treatment following surgery is called adjuvant therapy. Adjuvant therapy includes radiotherapy and chemotherapy. Some purposes of adjuvant radiotherapy have been proposed, including curative and palliative intent. In a previous study that evaluated adjuvant radiotherapy for FISS, a finely fractionated radiotherapy protocol significantly prolonged the survival time compared with a coarsely fractionated radiotherapy protocol.³⁰ Regarding adjuvant chemotherapy in cats, doxorubicin and lomustine have been reported for FISS, and both anticancer drugs demonstrated tumor suppression.^{31,32} In the present case, toceranib was selected as adjuvant chemotherapy. As the tumor was considered highly malignant due to the rapid enlargement, we planned therapy according to the evidence-based treatment for FISS, which is also a mesenchymal tumor. Radiation therapy was not selected due to the lack of facilities at our clinic, and frequent procedures under anesthesia could lead to further deterioration of the cat's general condition. Regarding chemotherapeutic agents, doxorubicin is known to have adverse effects including gastrointestinal and renal toxicity,³¹ and lomustine has been reported to cause severe leukopenia leading to dose reduction.³² Considering the poor general condition of the cat, use of doxorubicin or lomustine was challenging due to the possibility of severe adverse effects. Therefore, we considered using molecular-targeted drugs that are effective for sarcoma with mild adverse effects.

In recent years, molecular-targeted drugs such as imatinib, masitinib, and toceranib have been used in veterinary medicine.¹¹ Toceranib has been shown to inhibit KIT, platelet-derived growth factor receptor (PDGFR), vascular endothelial growth factor receptor, and tumor growth.^{13–16} Katayama et al.³³ reported that inhibition of PDGFR suppressed feline

fibrosarcoma in in vitro experiments. Holtermann et al.²⁷ treated FISS cases with toceranib as adjuvant therapy, but observed no significant tumor reduction. These authors suggested that toceranib may be potentially effective to some extent because all specimens of FISS were positive for PDGFR in immunohistochemistry. They also reported that gastrointestinal toxicity was the main adverse effect of toceranib, but all symptoms were mild and transient, and improved after discontinuation of toceranib. Based on the background information above, we decided to use toceranib, a molecular targeted drug with mild adverse effects and is expected to be effective to some extent against feline sarcomas. In the present case, toceranib was administered at a dose of 2.17 mg/kg three times per week for 1 month, but the residual tumor enlarged and invaded connective tissues. The response to toceranib treatment was considered poor, and there was no noticeable tumor suppression effect. Some factors may account for the lack of effectiveness. First, toceranib may be ineffective for this tumor type. As the origin of this tumor has not been identified, and molecular biological characterization remains unknown, it is unclear whether toceranib affects this tumor type. Second, the long interval between surgery and toceranib administration may be a factor. As toceranib was administered 3 weeks after surgery and the tumor had already progressed, the effect of tumor suppression may not be fully demonstrated. Third, the dosage may be low. In previous reports, toceranib was used in the range of 1.6–3.5 mg/kg.^{17,27,34,35} As the general condition of the cat was deteriorating, toceranib was used at a lower dose compared with previous reports. Therefore, the effective blood concentration of toceranib needed to suppress tumor growth might not be reached. This cat manifested restriction of the eyelids and exposed keratitis at initial presentation without obvious mass lesion in the orbit by ultrasound examination, which led to a delay in performing surgical intervention. Some orbital mesenchymal tumors in cats, such as FROMS, progress rapidly and are already advanced at the time of deciding surgical intervention, making it challenging to perform radical resection. Therefore, early surgical treatment should be considered when findings in cats suggest an orbital tumor as in the present case. In addition, adjuvant therapy should also be considered due to local recurrence and metastasis. Although toceranib did not suppress the tumor in this case, early administration or higher doses might have some effect on it.

4 | CONCLUSION

This is the first case report of treatment with toceranib as postsurgical adjuvant chemotherapy for FROMS in a cat. Although treatment was not successful to suppress the advanced tumor, further study to examine the dosage and timing of toceranib administration is warranted.

AUTHOR CONTRIBUTIONS

Hiroyuki Komatsu: Data curation; formal analysis; investigation; visualization; writing – original draft. **Asuka Ueshima:** Data curation; formal analysis; project administration. **Yoshitaka Kobayashi:** Conceptualization; supervision; writing – review and editing. **Yumiko Shimoyama:** Investigation; methodology. **Naoaki Takiyama:** Writing – review and editing. **Kenji Rimpo:** Conceptualization; project administration; writing – review and editing.

ACKNOWLEDGMENTS

All authors would like to thank the staff of Saitama Animal Medical Center for their dedicated support and to Teresa Nakatani for editorial support.

FUNDING INFORMATION

None.

CONFLICT OF INTEREST STATEMENT

All authors declare that they have no competing interests.

DATA AVAILABILITY STATEMENT


The data that support the findings of this study are available from the corresponding author, H.K., upon reasonable request.

CONSENT

All authors were conducted in accordance with the code of ethics of the Japan Veterinary Medical Association, and written informed consent was provided from the owner.

ORCID

Hiroyuki Komatsu  <https://orcid.org/0000-0002-8192-3143>

Yoshitaka Kobayashi  <https://orcid.org/0000-0002-6211-1174>

REFERENCES

1. Attali-Soussay K, Jegou JP, Clerc B. Retrobulbar tumors in dogs and cats: 25 cases. *Vet Ophthalmol.* 2001;4:19-27.
2. Meekins JM, Rankin AJ, Samuelson DA. Diseases and surgery of the canine orbit. In: Gelatt KN, ed. *Veterinary Ophthalmology*. 6th ed. Wiley Blackwell; 2021:879-922.
3. Isaza D, Robinson NA, Pizzirani S, Pumphrey SA. Evaluation of cytology and histopathology for the diagnosis of feline orbital neoplasia: 81 cases (2004–2019) and review of the literature. *Vet Ophthalmol.* 2020;23:682-689.
4. Bell CM, Schwarz T, Dubielzig RR. Diagnostic features of feline restrictive orbital myofibroblastic sarcoma. *Vet Pathol.* 2011;48:742-750.
5. Billson FM, Miller-Michau T, Mould JR, Davidson MG. Idiopathic sclerosing orbital pseudotumor in seven cats. *Vet Ophthalmol.* 2006;9:45-51.

6. Diehl KA, Pryor SG, Teixeira LBC. Orbital invasive squamous cell carcinoma with adnexal involvement clinically mimicking feline restrictive orbital myofibroblastic sarcoma: 19 cases (1990–2016). *Vet Ophthalmol.* 2018;21:281-289.
7. Thomasy SM, Cissell DD, Arzi B, et al. Restrictive orbital myofibroblastic sarcoma in a cat – cross-sectional imaging (MRI & CT) appearance, treatment, and outcome. *Vet Ophthalmol.* 2013;16(Suppl 1):123-129.
8. Miller SA, van der Woerd A, Bartick TE. Retrobulbar pseudotumor of the orbit in a cat. *J Am Vet Med Assoc.* 2000;216(356–8):345-358.
9. Gould D. Diseases of the orbit. In: Maggs DJ, Miller PE, Ofri R, Slatter D, eds. *Slatter's Fundamentals of Veterinary Ophthalmology.* 6th ed. Elsevier; 2018:101-126.
10. Withrow SJ, Vail DM, Page RL. *Withrow & MacEwen's Small Animal Clinical Oncology.* 5th ed. Elsevier/Saunders; 2013:750.
11. Londhe P, Gutwillig M, London C. Targeted therapies in veterinary oncology. *Vet Clin North Am Small Anim Pract.* 2019;49:917-931.
12. Thamm DH, Weishaar KM, Charles JB, Ehrhart EJ 3rd. Phosphorylated KIT as a predictor of outcome in canine mast cell tumours treated with toceranib phosphate or vinblastine. *Vet Comp Oncol.* 2020;18:169-175.
13. London CA, Malpas PB, Wood-Follis SL, et al. Multi-center, placebo-controlled, double-blind, randomized study of oral toceranib phosphate (SU11654), a receptor tyrosine kinase inhibitor, for the treatment of dogs with recurrent (either local or distant) mast cell tumor following surgical excision. *Clin Cancer Res.* 2009;15:3856-3865.
14. London C, Mathie T, Stingle N, et al. Preliminary evidence for biologic activity of Toceranib phosphate (Palladia(R)) in solid tumours. *Vet Comp Oncol.* 2012;10:194-205.
15. London CA, Hannah AL, Zadovskaya R, et al. Phase I dose-escalating study of SU11654, a small molecule receptor tyrosine kinase inhibitor, in dogs with spontaneous malignancies. *Clin Cancer Res.* 2003;9:2755-2768.
16. Pryer NK, Lee LB, Zadovskaya R, et al. Proof of target for SU11654: inhibition of KIT phosphorylation in canine mast cell tumors. *Clin Cancer Res.* 2003;9:5729-5734.
17. Berger EP, Johannes CM, Post GS, et al. Retrospective evaluation of toceranib phosphate (Palladia) use in cats with mast cell neoplasia. *J Feline Med Surg.* 2018;20:95-102.
18. Hasegawa Y, Shosu K, Tsuji K, et al. Intratumoral heterogeneity of c-KIT mutations in a feline splenic mast cell tumor and their functional effects on cell proliferation. *Sci Rep.* 2022;12:15791.
19. Azevedo C, Brawner W, Schleis Lindley S, Smith A. Multimodal non-surgical treatment of a feline tracheal adenocarcinoma. *JFMS Open Rep.* 2017;3:205511691668963.
20. Todd JE, Nguyen SM. Long-term survival in a cat with pancreatic adenocarcinoma treated with surgical resection and toceranib phosphate. *JFMS Open Rep.* 2020;6:205511692092491.
21. McGregor O, Moore AS, Yeomans S. Management of a feline gastric stromal cell tumour with toceranib phosphate: a case study. *Aust Vet J.* 2020;98:181-184.
22. Dedeaux AM, Langohr IM, Boudreaux BB. Long-term clinical control of feline pancreatic carcinoma with toceranib phosphate. *Can Vet J.* 2018;59:751-754.
23. Petrucci G, Henriques J, Gregorio H, et al. Metastatic feline mammary cancer: prognostic factors, outcome and comparison of different treatment modalities – a retrospective multicentre study. *J Feline Med Surg.* 2021;23:549-556.
24. Olmsted GA, Farrelly J, Post GS, Smith J. Tolerability of toceranib phosphate (Palladia) when used in conjunction with other therapies in 35 cats with feline oral squamous cell carcinoma: 2009–2013. *J Feline Med Surg.* 2017;19:568-575.
25. Wiles V, Hohenhaus A, Lamb K, Zaidi B, Camps-Palau M, Leibman N. Retrospective evaluation of toceranib phosphate (Palladia) in cats with oral squamous cell carcinoma. *J Feline Med Surg.* 2017;19:185-193.
26. Kopke MA, Gal A, Piripi SA, Poirier VJ. Squamous cell carcinoma of the anal sac in two cats. *J Small Anim Pract.* 2021;62:704-708.
27. Holtermann N, Kiupel M, Hirschberger J. The tyrosine kinase inhibitor toceranib in feline injection site sarcoma: efficacy and side effects. *Vet Comp Oncol.* 2017;15:632-640.
28. Glaze MB, Maggs DJ, Plummer CE. Feline ophthalmology. In: Gelatt KN, ed. *Veterinary Ophthalmology.* 6th ed. Wiley Blackwell; 2021:1665-1840.
29. Gilger BC, McLaughlin SA, Whitley RD, Wright JC. Orbital neoplasms in cats: 21 cases (1974–1990). *J Am Vet Med Assoc.* 1992;201:1083-1086.
30. Rossi F, Marconato L, Sabattini S, et al. Comparison of definitive-intent finely fractionated and palliative-intent coarsely fractionated radiotherapy as adjuvant treatment of feline microscopic injection-site sarcoma. *J Feline Med Surg.* 2019;21:65-72.
31. Poirier VJ, Thamm DH, Kurzman ID, et al. Liposome-encapsulated doxorubicin (Doxil) and doxorubicin in the treatment of vaccine-associated sarcoma in cats. *J Vet Intern Med.* 2002;16:726-731.
32. Saba CF, Vail DM, Thamm DH. Phase II clinical evaluation of lomustine chemotherapy for feline vaccine-associated sarcoma. *Vet Comp Oncol.* 2012;10:283-291.
33. Katayama R, Huelsmeyer MK, Marr AK, Kurzman ID, Thamm DH, Vail DM. Imatinib mesylate inhibits platelet-derived growth factor activity and increases chemosensitivity in feline vaccine-associated sarcoma. *Cancer Chemother Pharmacol.* 2004;54:25-33.
34. Harper A, Blackwood L. Toxicity and response in cats with neoplasia treated with toceranib phosphate. *J Feline Med Surg.* 2017;19:619-623.
35. Merrick CH, Pierro J, Schleis SE, et al. Retrospective evaluation of toceranib phosphate (Palladia(R)) toxicity in cats. *Vet Comp Oncol.* 2017;15:710-717.

How to cite this article: Komatsu H, Ueshima A, Kobayashi Y, Shimoyama Y, Takiyama N, Rimpo K. Feline restrictive orbital myofibroblastic sarcoma treated with toceranib phosphate (Palladia) as adjuvant chemotherapy: A case study of one cat. *Clin Case Rep.* 2023;11:e7582. doi:[10.1002/ccr3.7582](https://doi.org/10.1002/ccr3.7582)