

POSTER PRESENTATION

Open Access

The impact of radiologists' expertise on screen result decisions in CT lung cancer screening

MA Heuvelmans^{1*}, M Oudkerk¹, PA de Jong², WPTM Mali², HJM Groen³, R Vliegenthart¹

From International Cancer Imaging Society (ICIS) 14th Annual Teaching Course
Heidelberg, Germany. 9-11 October 2014

Aim

To evaluate the impact of radiological expertise on screen result decisions made in a CT lung cancer screening trial.

Methods

In the Dutch-Belgian randomized lung cancer screening trial (NELSON), the baseline CT screen result was based on the lung nodule with largest volume. According to the protocol, nodule volume <50mm³, 50-500mm³ and >500mm³ led to a negative, indeterminate and positive screen result, respectively. The protocol, however, allowed radiologists to manually adjust screen result in case of high suspicion of benign or malignant nodule nature. In this study, all participants whose baseline CT result was based on a solid nodule were included. Adjustments by radiologists at baseline were evaluated. Histology was the reference for diagnosis, or, to confirm benignity, stability on subsequent CT scans.

Results

3,268 participants (2,759 male, median age 58.0-years) were included. In 189 participants (5.8%) the initial baseline screen result was adjusted by the radiologist. Adjustment was downwards from positive or indeterminate to negative in two and 118 participants, respectively, and from positive to indeterminate in 64 participants. None of these nodules turned out malignant. In five participants (2.6%) the screen result was adjusted upwards from negative to indeterminate (N=1) or indeterminate to positive (N=4); two nodules (40%) were malignant.

Conclusion

In about one-in-twenty cases of baseline lung cancer screening, nodules were reclassified by the radiologist (97.4% downwards), leading to reduction of false-positive and false-negative screen results.

Authors' details

¹Center for Medical Imaging – North East Netherlands, Department of Radiology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands. ²Department of Radiology, University Medical Center Utrecht, Utrecht, Netherlands. ³Department of Pulmonology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands.

Published: 9 October 2014

doi:10.1186/1470-7330-14-S1-P19

Cite this article as: Heuvelmans et al.: The impact of radiologists' expertise on screen result decisions in CT lung cancer screening. *Cancer Imaging* 2014 **14**(Suppl 1):P19.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: m.a.heuvelmans@umcg.nl

¹Center for Medical Imaging – North East Netherlands, Department of Radiology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands

Full list of author information is available at the end of the article