

Oral microbial profile in oral cancer patients before and after radiation therapy in a cancer care center – A prospective study

K Anjali¹, A B Arun², T S Bastian³, R Parthiban⁴, M Selvamani³, H Adarsh⁵

Departments of ¹Microbiology and ³Oral Pathology and Microbiology, Mahe Institute of Dental Sciences, Puducherry, ²Yenepoya Research Center, Yenepoya University, Mangalore, ⁵Department of Dentistry, BGS Global Institute of Medical Sciences, Bengaluru, Karnataka, ⁴Department of Microbiology, Malabar Cancer Center, Thalassery, Kerala, India

Abstract

Background: Head and neck cancer is the sixth most common cancer reported worldwide. In many cases, the level of aggressiveness of therapy adopted in cancer patients may cause the alteration in oral microbiota; the emergence of potential pathogens may cause opportunistic infections in already immune-compromised individuals leading to increases in morbidity and mortality. Hence, this study was conducted to assess the oral microbial profile in oral cancer patients before and after radiotherapy.

Materials and Methods: A total of 145 oral swabs were collected before radiotherapy ($n = 96$), 3 months postradiotherapy ($n = 25$), 6 months postradiotherapy ($n = 12$) and controls ($n = 12$). The samples were inoculated into brain–heart infusion broth and later in different media for bacterial isolation. The isolates were subjected to phenotypic characterization by automatic identification system.

Results: Among the 96 samples studied from the preradiotherapy patient samples, *Streptococcus* species ($n = 28$) were the predominant isolate, followed by *Staphylococcus* species ($n = 16$), *Enterobacter* species ($n = 6$) and *Enterococcus* species ($n = 6$). Of the 25 samples studied 3 months after radiotherapy, *Klebsiella pneumoniae* ($n = 4$) was isolated and 12 samples studied after 6 months of radiotherapy *Candida* species ($n = 4$) and *Pediococcus* species ($n = 3$) were isolated. Among the control group ($n = 12$) screened, *Streptococcus acidominimus* ($n = 3$) is the predominant bacteria isolated.

Conclusion: High prevalence of *Streptococcus* sp. was found in patients of oral cancer before radiotherapy, while *Candida albicans* and *Klebsiella* species and *Pediococcus* species are the significant pathogens isolated in postradiotherapy cancer patients.

Keywords: Oral microbial profile, oral microbiota, oral squamous cell carcinoma, radiotherapy

Address for correspondence: Dr. A B Arun, Yenepoya Research Center, Yenepoya University, Deralakatte, Mangalore - 575 018, Karnataka, India.

E-mail: bhagwatharun@hotmail.com

Submitted: 14-Jul-2019, **Accepted:** 14-Oct-2019, **Published:** 08-May-2020

INTRODUCTION

Head-and-neck cancer is the world's sixth most common type of cancer. More than 600,000 new patients are

diagnosed and about 350,000 individuals die from this disease worldwide each year.^[1] Occurrence of oral cancer is on the rise, causing major global health problem. Incidence

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Anjali K, Arun AB, Bastian TS, Parthiban R, Selvamani M, Adarsh H. Oral microbial profile in oral cancer patients before and after radiation therapy in a cancer care center – A prospective study. J Oral Maxillofac Pathol 2020;24:117-24.

Access this article online

Quick Response Code:



Website:

www.jomfp.in

DOI:

10.4103/jomfp.JOMFP_213_19

rate of oral cancer is unevenly distributed geographically, India and France showing the highest.^[2] Increased risk of oral cancer is associated with poor oral hygiene along with the use of alcohol and tobacco.^[3] Microbial flora on the oral mucosa seems to differ between healthy and malignant sites.^[4,5] Role of specific bacterial infection causing or promoting cancer is well-known in cases of *Helicobacter pylori* with gastric cancer and other cancers of colon, gall bladder, prostate and lung.^[6-8] Rich and varied community of microbes resides on exposed surfaces (e.g., gut, skin and mouth) which outnumber the cells in human body by ten times.^[9] Changes in microbes composition and overgrowth of certain pathogen are associated with chronic periodontitis, which is a risk factor for oral cancer.^[10] Despite the availability of various successful treatment modalities such as surgery, radiation, chemotherapy or combination for oral cancer, local and systemic immunity reduced in treated patients due to swallowing dysfunction, hyposalivation and variation in quality, quantity, complexity of oral microbes leading to imbalance in oral ecosystem.^[11] Increased levels and changes in the composition of microbes in the oral cavity are reported in precancerous conditions and cancers.^[12] The predominant and abundant phyla reported in oral cancer belong to *Firmicutes*, *Proteobacteria*, *Bacteroidetes*, *Actinobacteria* and *Fusobacteria*.^[13-15] In healthy individuals, *Streptococcus* species are the frequently observed oral microbe and less frequently seen are *Prevotella*, *Veillonella*, *Neisseria* and *Haemophilus*.^[14,15] Infections result when there is disruption of balance between bacterial load and compromise in immune status, allowing certain bacteria and *Candida* species to multiply and overwhelm other resident microorganisms. Radiotherapy for oral cancer causes alteration in oral microbiota which may lead to the emergence of potential pathogen subsequently leading to other systemic problems.^[3] After radiotherapy, the patient may acquire drug-resistant opportunistic infections from altered microbes in the oral cavity which may cause systemic complications and high morbidity. As there are no sufficient microbial studies to precisely infer the presence or absence of certain microorganisms and its impact on immunocompromised patients, there is need to study the exact role played by the oral microbes and its impact on pre- and post-cancer treatment. Hence, this present study was conducted to identify microbial profile in oral cancer patients undergoing radiotherapy at tertiary care cancer center, Northern Kerala.

MATERIALS AND METHODS

This prospective study was carried out in a randomly selected cohort of total 96 oral cancer patients before radiation and 25 cases were followed up after 3 months and 12 cases after 6 months of radiation along with 12 controls at cancer

center, Thalassery, Kerala, from June 2017 to October 2018. The study was approved by the Institutional Review Board No - 1617/IRB-IEC/13/MCC/06-03-2017/12 and also ethical clearances were obtained from the ethical clearance committee before the study. All patients were informed about the study, counseled and informed consent was obtained before taking the samples. Demographic details such as age, gender and habits were collected as per the standard pro forma.

Inclusion criteria

All consenting patients suffering from oral cancer and those who underwent radiotherapy were included in this study, irrespective of age and gender. Participants having good oral health without a history or symptoms of oral cancer and precancerous lesions were taken as the control group.

Exclusion criteria

The patients suffering from diabetes mellitus, infected by human immunodeficiency virus, hepatitis and autoimmune diseases were excluded after obtaining the laboratory data from the case files.

Sample collection and processing

Oral swabs were collected from areas such as buccal mucosa, tongue, alveolar ridges, floor of the mouth and various regions in the oral cavity by scrubbing the lesions using a dry sterile cotton swab. The oral samples were immediately taken to microbiology laboratory and put in brain–heart infusion broth. The broth was incubated aerobically for overnight at 37°C. Then, it was inoculated into culture media such as MacConkey agar, blood agar and chocolate agar from the broth and incubated at 37°C. The colony characters on the grown culture plates and morphology of the microorganism by Gram staining was observed. The mixed cultures grown on the primary culture plate were subcultured into blood agar and chocolate agar separately, and individual type of colonies were subjected for rapid phenotypic identification by automated bacterial identification system (Phoenix TM – Becton Dickinson). It is fully automated system for rapid identification of bacteria and antimicrobial susceptibility testing.

RESULTS

A total of 145 oral swabs were collected from oral cancer patients. Of which, 96 were collected before radiotherapy, 25 were collected 3 months and 12 were collected 6 months of follow-up after radiotherapy along with 12 controls collected from persons with good oral hygiene and no symptoms of oral diseases. Loss of follow-up of patients was due to patient death ($n = 11$), loss of communication ($n = 25$),

clinical deterioration ($n = 11$) and patient subjected to only surgery without radiotherapy ($n = 24$).

A total of 96 cases of oral squamous cell carcinoma (OSCC) studied during this period, 72.9% (70 cases) occurred in males and 27.1% (26 cases) in females, with a male-to-female ratio of 2.7:1 [Table 1]. The mean age distribution of cases was 56.5 years (range: 32–83 years), with highest 28.1% (27 number) diagnosed in the fifth and sixth decades of life followed by 20.8% (20 number) in the fourth decade. Among 70 male patients studied, 31.4% were diagnosed in the fifth, 28.6% in the sixth decade and 8.9% in the third decade of life. Of 26 female patients studied, 26.9% were diagnosed in their sixth decade, 19.2% each in fifth and seventh decades and 7.7% each in third and eighth decades of life [Table 1].

Of 96 patients, 77 (80.2%) were with a positive history of habits which included smoke and smokeless tobacco consumption, alcohol consumption and combination of both the habits, whereas 19 patients (19.79%) were without any habits or familial history of cancer or cancer-related death. Of 70 male patients, 61 (87.14%) had a positive history of habits, whereas 9 (12.85%) were without any habit history. In case of the female patients, 16 of 26 (61.53%) were with a positive habit history and 10 (38.46%) were without any habit history.

Patients with a positive habit history 24 (31.46%) and 25 (32.46%) were diagnosed in the fifth and sixth decades of life, respectively, whereas seven patients (36.8%) without a habit history were diagnosed during the same period. A total of 28 cases (29.1%) occurred below 50 years, among them 18 (64.3%) cases were with a positive habit

history and 10 (35.7%) cases were without a habit history. The mean age of patients with and without a history of habits was 52.1 years for males and 50.8 years for females with no significant difference [Table 2].

Most common habit among the male patients was tobacco smoking and alcohol consumption, combination of both accounted for 32.78% of the total cases followed by simultaneous use of chewing tobacco (smokeless), smoking and alcohol intake that accounted for 29.5% of the total cases. While in females, the most significant habit was the tobacco (smokeless) form which accounted for about 87.5% [Table 3].

The oral swabs collected from the participants before radiation therapy ($n = 96$) showed the growth of *Klebsiella pneumoniae* (12; 12.5%) as the predominant organism in Gram-negative bacilli group followed by *Escherichia coli* (6; 6.2%), *Pseudomonas aeruginosa* (6; 6.2%), *Pasteurella species* (5; 5.2%) and others. Among Gram-positive organisms grown, *Streptococcus* species (28; 29%) was predominant followed by *Staphylococcus* species (7; 7.2%), *Staphylococcus aureus* (6; 6.2%) and others. The oral swabs collected after 3 months of radiation therapy ($n = 25$) revealed *K. pneumoniae* (4; 16%) as the predominant Gram-negative organism followed by *P. aeruginosa* (1; 4%) and *Acinetobacter baumannii* (1; 4%) while among Gram-positive organisms, *Streptococcus* species (4; 16%) was predominant followed by *Enterococcus faecalis* (4; 16%). Oral swabs ($n = 12$) collected 6 months after radiation therapy showed predominantly *Candida* species ($n = 5$; 42%), followed by *Pediococcus* species ($n = 3$; 25%). Oral swabs collected for control samples ($n = 12$) showed *Streptococcus* species ($n = 8$; 67%) as the predominant species followed by *Leuconostoc* species and *Candida* species ($n = 3$; 25%) [Tables 4-6].

Table 1: Incidence of oral squamous cell carcinoma according to age and gender

Age group (years)	Male, n (%)	Female, n (%)	Total, n (%)
30-39	6 (8.6)	2 (7.7)	8 (8.3)
40-49	15 (21.4)	5 (19.2)	20 (20.8)
50-59	22 (31.4)	5 (19.2)	27 (28.1)
60-69	20 (28.6)	7 (26.9)	27 (28.1)
70-79	7 (10)	5 (19.2)	12 (12.5)
80-89	-	2 (7.7)	2 (2.1)
Total	70 (72.9)	26 (27.1)	96 (100)

Table 2: Distribution of age and gender with reference to presence or/absences of habit

Age group (years)	Male and female combined		Male		Female	
	With habits, n (%)	Without habits, n (%)	With habits, n (%)	Without habits, n (%)	With habits, n (%)	Without habits, n (%)
30-39	5 (6.49)	3 (15.78)	5 (8.19)	1 (11.11)	-	2 (20)
40-49	13 (16.88)	7 (36.8)	11 (18.03)	4 (44.44)	2 (12.5)	3 (30)
50-59	24 (31.16)	3 (15.78)	20 (32.78)	2 (22.22)	4 (25)	1 (10)
60-69	25 (32.46)	4 (21.05)	20 (32.78)	2 (22.22)	5 (31.25)	2 (20)
70-79	8 (10.38)	2 (10.52)	5 (8.19)	-	3 (18.75)	2 (20)
80-89	2 (2.59)	-	-	-	2 (12.5)	-
Total	77 (80.2)	19 (19.79)	61 (87.14)	9 (12.85)	16 (61.53)	10 (38.46)

DISCUSSION

OSCC can be located anywhere in the oral tissues, including the lip, floor of the mouth, buccal mucosa, gingival, palate or in the tongue. The majority of squamous cell carcinoma usually arise from preexisting “potentially

Table 3: Age and sex distribution with regard to specific habits

Age (years)	TOB		Smoke	SMOK + ALCH	TOB + ALCH		TOB + ALCH + SMOK		Total	
	Male, n (%)	Female, n (%)	Male, n (%)	Male, n (%)	Male, n (%)	Female, n (%)	Male, n (%)	Female, n (%)	Male, n (%)	Female, n (%)
30-39	-	-	2 (3.27)	2 (3.27)	-	-	1 (1.6)	-	5 (8.2)	-
40-49	2 (3.27)	2 (12.5)	1 (1.6)	6 (9.8)	-	-	2 (3.27)	-	11 (18)	2 (12.5)
50-59	5 (8.2)	3 (18.75)	3 (4.9)	2 (3.27)	3 (4.9)	-	7 (11.47)	1 (6.25)	20 (32.8)	4 (25)
60-69	-	4 (25)	3 (4.9)	9 (14.75)	3 (4.9)	1 (6.25)	5 (8.2)	-	20 (32.8)	5 (31.25)
70-79	-	3 (18.75)	1 (1.6)	1 (1.6)	-	-	3 (4.9)	-	5 (8.2)	3 (18.75)
80-89	-	2 (12.5)	-	-	-	-	-	-	-	2 (12.5)
Total	7 (11.47)	14 (87.5)	10 (6.39)	20 (32.78)	6 (9.83)	1 (6.25)	18 (29.5)	1 (6.25)	61 (100)	16 (100)

TOB: Tobacco (smokeless), SMOK + ALCH: Smoke along with alcohol, TOB + ALCH: Tobacco (smokeless) and alcohol, TOB + ALCH + SMOK: Tobacco (smokeless), smoke and alcohol together

Table 4: Distribution of Gram-negative organisms in cancer patients at different intervals

Total number of organisms	Before radiation (n=96), n (%)	After radiation 3 months (n=25), n (%)	After radiation 6 months (n=12), n (%)	Control (n=12), n (%)
<i>Klebsiella pneumoniae</i>	12 (12.5)	4 (16)	1 (8.3)	-
<i>Escherichia coli</i>	6 (6.2)	-	-	-
<i>Pseudomonas aeruginosa</i>	6 (6.2)	1 (4)	-	-
<i>Pasteurella species</i>	5 (5.2)	-	-	-
<i>Enterobacter cloacae</i>	4 (4.1)	-	-	-
<i>Shigella dysenteriae</i>	3 (3.1)	-	-	-
<i>Enterobacter aerogenes</i>	2 (2)	-	-	-
<i>Acinetobacter baumannii</i>	2 (2)	1 (4)	-	-
<i>Proteus mirabilis</i>	1 (1)	-	-	-
<i>Serratia marcescens</i>	1 (1)	-	-	-
<i>Pantoea agglomerans</i>	1 (1)	-	-	-
<i>Neisseria animaloris</i>	1 (1)	-	-	-
<i>Neisseria species</i>	-	-	-	1 (8)
<i>Methylobacterium extorquens</i>	1 (1)	-	-	-
<i>Ewingella americana</i>	1 (1)	-	-	-
<i>Citrobacter species</i>	1 (1)	-	-	-
<i>Bordetella bronchiseptica</i>	1 (1)	-	-	-
<i>Chromobacterium haemolyticum</i>	1 (1)	-	-	-
<i>Yersinia pseudotuberculosis</i>	-	-	1 (8.3)	-

malignant” lesions or disorders and from normal-appearing epithelium.^[16] Although the causes for OSCC could be multifactorial, most common etiology is related to tobacco or its product in both smoke and smokeless forms. The increased consumption of tobacco has become a multidimensional problem and a threat to the health of the youth in India.^[17]

In the present study, we found that the oral cancer incidence rates are higher in males (72.9%) than females (27.1%); majority of the OSCC patients were in their fifth–sixth decades of life, and these findings are in agreement with studies done by other research groups.^[16-18] Although it is recognized that oral cancer is more prevalent among elderly age groups, the data trend in recent years indicate the increased incidence rates of oral cancer in younger age groups, and in the present study, similar trend was noted. Many recent studies have reported the prevalence rate of 4%–6% of oral cancers in the younger populations <40 years.^[3,17] In our study, 8 OSCC occurred below 40 years, among them four cases of the tongue OSCC noted corroborating the findings reported earlier by

Selvamani *et al.*,^[17] where they have recorded the prevalence of tongue cancers in the Indian population to the extent of 12% in the age group of 31–40 years, and also a review by Sarkaria and Harari identified 14 reports with three or more patients younger than 40 years of age with OSCC of the tongue.^[19]

Gender differences in the occurrence of OSCC generally show male preponderance over female, and our study also showed similar trend with a male-to-female ratio of 2.7:1. The male-to-female ratio varies widely by geographic location worldwide, 1.45:1 in Japan, 1.5:1 in Pakistan, 1.65:1 in Yemen and 10.5:1 in Taiwan patient population, and a reverse trend has been observed in the Thailand population with a male: female ratio of 1:1.56.^[20-24]

In our study, we observed that 50% of female cases showed involvement of the buccal mucosa as predominant site for OSCC in smokeless tobacco users and also the most common site observed among the female patients, similar to the study done by Singh *et al.*^[25] In this study, 47.14% cases could be attributed to the consumption of

Table 5: Distribution of Gram-positive organisms in cancer patients at different intervals

Total number of organisms Gram-positive organisms	Before radiation (n=96), n (%)	After radiation 3 months (n=25), n (%)	After radiation 6 months (n=12), n (%)	Control (n=12), n (%)
<i>Streptococcus acidominimus</i>	8 (8.3)	-	-	5 (42)
<i>Streptococcus sanguinis</i>	5 (5.2)	-	-	-
<i>Streptococcus anginosus</i>	5 (5.2)	-	-	-
<i>Streptococcus sobrinus</i>	3 (3)	-	-	3 (25)
<i>Streptococcus porcinus</i>	2 (2.08)	-	-	-
<i>Streptococcus agalactiae</i>	2 (2.08)	-	-	-
<i>Streptococcus gordonii</i>	1 (1.04)	1 (4)	-	-
<i>Streptococcus pneumoniae</i>	1 (1.04)	1 (4)	-	-
<i>Streptococcus mutans</i>	1 (1.04)	1 (4)	-	-
<i>Streptococcus oralis</i>	-	1 (4)	-	2 (16)
<i>Staphylococcus aureus</i>	6 (6.2)	1 (4)	1 (8.3)	-
<i>Staphylococcus haemolyticus</i>	1 (1.04)	-	-	-
<i>Staphylococcus lugdunensis</i>	1 (1.04)	-	-	-
<i>Staphylococcus saprophyticus</i>	1 (1.04)	-	-	-
<i>Staphylococcus epidermidis</i>	1 (1.04)	-	-	-
<i>Staphylococcus pettenkoferi</i>	1 (1.04)	1 (4)	-	-
<i>Staphylococcus capitis</i>	1 (1.04)	-	-	-
<i>Staphylococcus auricularis</i>	1 (1.04)	-	-	-
<i>Staphylococcus cohnii</i>	-	1 (4)	-	-
<i>Staphylococcus kloosi</i>	-	-	1 (8.3)	-
<i>Staphylococcus hominis</i>	-	-	1 (8.3)	-
<i>Enterococcus raffinosus</i>	1 (1.04)	-	1 (8.3)	-
<i>Enterococcus avium</i>	1 (1.04)	-	-	-
<i>Enterococcus faecalis</i>	4 (4.1)	4 (16)	-	-
<i>Aerococcus viridans</i>	1 (1.04)	-	-	-
<i>Leuconostoc mesenteroides</i>	-	1 (4)	-	2 (16)
<i>Pediococcus species</i>	-	-	3 (25)	-

Table 6: Microbial profile of oral cancer patients with complete follow-up (n=12)

Patient ID	Before RD	Follow-up	
		3 months after RD	6 months after RD
R1	<i>Staphylococcus</i> sp.	-	-
R2	<i>Pseudomonas</i> sp. <i>Neisseria</i> sp.	<i>Pasteurella</i> sp.	<i>Klebsiella</i> sp.
R3	<i>Staphylococcus</i> sp.	-	<i>Pediococcus</i> sp.
R4	<i>Streptococcus</i> sp. <i>Bacillus</i> sp.	<i>Enterococcus</i> sp.	<i>Candida</i> sp.
R5	<i>Enterococcus</i> sp.	-	<i>Bacillus</i> sp.
R6	<i>Bacillus</i> sp.	<i>Acinetobacter</i> sp.	<i>Staphylococcus</i> sp. <i>Candida</i> sp.
R7	<i>Staphylococcus</i> sp. <i>Enterococcus</i> sp.	<i>Staphylococcus</i> sp.	<i>Staphylococcus</i> sp.
R8	<i>Bacillus</i> sp. <i>Citrobacter</i> species	<i>Streptococcus</i> sp. <i>Enterococcus</i> sp.	<i>Enterococcus</i> sp.
R9	<i>Staphylococcus</i> sp.	<i>Pediococcus</i> sp.	<i>Candida</i> sp.
R10	<i>Bacillus</i> sp. <i>Pasteurella</i> sp.	<i>Leuconostoc</i> sp.	<i>Candida</i> sp.
R11	<i>Pseudomonas</i> sp. <i>Escherichia</i> sp.	-	<i>Staphylococcus</i> sp.
R12	<i>Bacillus</i> sp. <i>Pasteurella</i> sp.	<i>Streptococcus</i> sp.	<i>Candida</i> sp.

*RT: Radiation therapy, *sp: Species

alcohol and smoking, whereas Ajay *et al.* reported similar association in 28.1% of patients.^[26]

The human body is inhabited by over 100 trillion microbial cells living in association with their host.^[27] Microorganism

at certain body sites has long been believed to be involved in immune modulation, disease progression and health preservation. Some of these microorganisms have been implicated in oral diseases such as caries and periodontitis, which are among the most common bacterial infections in humans.^[14] Microorganisms and their products, including endotoxins (lipopolysaccharides), enzymes (e.g., proteases, collagenases, fibrinolysin and phospholipase) and metabolic by-products (e.g., hydrogen sulfide, ammonia and fatty acids), are toxic to host cells and may directly induce mutations or alter signaling pathways that may affect cell proliferation and/or survival of epithelial cells.^[28] A bacterial infection can play a role in initiating or promoting cancer is a well-known fact with respect to the association of *Helicobacter pylori* with gastric cancer, and other cancers of gallbladder, colon, lung and prostate have been associated with particular bacterial infections.^[12,29] Hence, the question arises whether there may be shifts in the composition of the normal oral flora and/or chronic microbial infection can support or causes oral cancer.

Although primary causes of OSCC are tobacco and alcohol, recently, the malignancy was even seen in individuals who do not have the habit of smoking and drinking, suggesting that other factors such as poor oral hygiene and chronic periodontitis may also play a role in increasing the risk of oral cancers. Numerous microorganisms, residing in the

oral region, such as bacteria, viruses and fungi, may affect the development and progression of OSCC.^[30-32] In the present study, we found *Pseudomonas*, *Neisseria*, *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacillus*, *Citrobacter*, *Pasterulla*, *Pseudomonas* and *E. coli* in oral cancer patients. Similar findings were observed by Nagy *et al.*^[33] in their study where aerobic bacterial species *Haemophilus*, *Enterobacteriaceae* and *Streptococcus* and anaerobic bacterial species such as *Veillonella*, *Fusobacterium*, *Prevotella*, *Porphyromonas*, *Actinomyces* and *Clostridium* were detected more frequently in oral cancers than in normal oral mucosa.

Binder Gallimidi *et al.*^[34] reported that chronic infection of *Porphyromonas gingivalis* and *Fusobacterium nucleatum* promotes chemically induced OSCC in mice. *Streptococcus thermophilus* and *Streptococcus mitis* hidden in deep periodontal pockets exhibit alcohol dehydrogenase activity and produce acetaldehyde, a known carcinogen derived from alcohol, suggesting one mechanism whereby the pathogenesis of oral cancer may occur.^[35] These findings concur with the present study where *Staphylococcus*, *Streptococcus* and *Enterococcus* seen in almost all patients.

Radiation therapy standard protocols play an important role in the treatment of patients with head-and-neck cancer but may result in short- and long-term side effects. Possible side effects include mucositis, osteoradionecrosis, taste loss, hyposalivation, radiation caries, periodontal disease, trismus and so on.^[36] Hyposalivation and the loss of the protective effects of saliva may predispose changes in the oral microbiome, with a shift to periodontal disease-associated flora.^[37] Kamath *et al.*^[38] studied radiotherapy effect on oropharyngeal flora in head-and-neck cancer and concluded that *Streptococcus pneumoniae* was significantly decreased at the end of irradiation, while there was an increase in *S. aureus*, *Pseudomonas*, *Bacteroides* and *Candida* species. In the present study, we found that before initiation of radiation therapy, there was increased number of *Staphylococcus*, *Streptococcus*, *Pseudomonas*, *Bacillus*, *Enterococcus*, *Neisseria*, *E. coli*, etc., but after 3 months of radiation, either decrease or no counts of *Pseudomonas* were noted; *Staphylococcus*, *Streptococcus* and *Enterococcus* could still identified in culture. Newer entities of microbes like *Acinetobacter*, *Pasteurella*, *Pediococcus* and *Leuconostoc* were observed after radiotherapy, However these micro organism were not observed prior to initiation of radiation therapy [Table 6].

A study by Sharma^[39] on salivary bacteria linked to oral cancers found that *Prevotella melaninogenica*, *Leptotrichia buccalis*, *Capnocytophaga gingivalis*, *Eubacterium saburreum* and *S. mitis* were significantly higher in patients with radiotherapy than controls. Eliasson *et al.*^[40] studied dental

plaque pH and microorganisms during hyposalivation in ten patients after 3–5 years of radiotherapy with equal controls and concluded that when compared with their controls, the irradiated group displayed increased numbers of *Lactobacilli* and *Candida* species due to reduced buffering capacity of saliva. Similar findings were noted in the study done by Srithavaj and Thaweboon^[41] and Al -Nawas and Grötz^[42] where they noted increased in the *Lactobacilli* and *Streptococci* count. In the present study, we too observed increase in *Candida* after 6 months of radiation therapy along with *Staphylococcus*, *Bacillus*, *Enterococcus*, *Pediococcus* and other microorganisms such as *Klebsiella*, *Staphylococcus kלוסי* and *Staphylococcus hominis* [Table 6]. *Streptococci* produce short-chain organic acid from carbohydrates lowering the pH of their local environment which may contribute to the acidic and hypoxic microenvironment of tumors.^[43-44] Bacterial colonization may induce alterations in microenvironment and promote chronic inflammation. The pathogenesis of approximately 15%–20% of human tumors^[45] is correlated with infection-triggered inflammations. Evidence also support the hypothesis of *E. coli* inhibiting apoptosis and thus promoting carcinogenesis. *E. coli* releases a range of virulence factors, including cytotoxic necrotizing factor type 1 which prevents apoptosis in epithelial cells by activating a cell signaling cascade and promoting the expression of antiapoptotic members of the Bcl-2 gene family.^[46]

Increase in oral *Candida* colonization in patients who received radiation for head-and-neck carcinoma was seen in the present study as reported in other studies.^[41,42] It is reported that *Candida* strains, by producing nitrosamine compounds, may directly or in concert with other carcinogens be able to activate specific proto-oncogenes, thus initiate oral cancer. It was found that *Candida* species associated with advanced precancerous lesions have relatively higher potentials for producing N-nitroso-benzylmethylamine, a compound able to induce carcinoma compared to strains associated with normal mucosa.^[47]

CONCLUSION

The present study shows that oral cancers occur commonly in the fifth and sixth decades with male preponderance and associated with smoking and drinking habits. There is a wide variation in the microbial flora of oral cancer patients' pre- and post-radiation therapy at different intervals. Gram-positive organisms are isolated more frequently than that of Gram-negative organisms. The high prevalence of *streptococcus* sp. was found in patients of oral cancer before radiotherapy and *Candida albicans* and *Klebsiella* species and *Pediococcus* species are the significant

pathogens isolated in postradiotherapy. Further studies are required to investigate the impact of alteration in microbial flora during the treatment of cancers. The continued surveillance and prompt examination, testing and careful monitoring of the imbalance in microbial ecosystem of the oral cavity and early management of complications are of prime importance in the long-term care of the patients receiving radiation treatment to reduce morbidity and give good quality of life.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010;127:2893-917.
2. Global Burden of Disease Cancer Collaboration, Fitzmaurice C, Dicker D, Pain A, Hamavid H, Moradi-Lakeh M, et al. The global burden of cancer 2013. *JAMA Oncol* 2015;1:505-27.
3. Meurman JH, Uittamo J. Oral micro-organisms in the etiology of cancer. *Acta Odontol Scand* 2008;66:321-6.
4. Napeñas JJ, Brennan MT, Bahrani-Mougeot FK, Fox PC, Lockhart PB. Relationship between mucositis and changes in oral microflora during cancer chemotherapy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007;103:48-59.
5. Gao L, Xu T, Huang G, Jiang S, Gu Y, Chen F. Oral microbiomes: More and more importance in oral cavity and whole body. *Protein Cell* 2018;9:488-500.
6. Belazi M, Velegraki A, Koussidou-Eremondi T, Andreadis D, Hini S, Arsenis G, et al. Oral *Candida* isolates in patients undergoing radiotherapy for head and neck cancer: Prevalence,azole susceptibility profiles and response to antifungal treatment. *Oral Microbiol Immunol* 2004;19:347-51.
7. Mantovani A, Allavena P, Sica A, Balkwill F. Cancer-related inflammation. *Nature* 2008;454:436-44.
8. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin* 2015;65:87-108.
9. Grice EA, Segre JA. The skin microbiome. *Nat Rev Microbiol* 2011;9:244-53.
10. Vanhoecke B, De Ryck T, Stringer A, Van de Wiele T, Keefe D. Microbiota and their role in the pathogenesis of oral mucositis. *Oral Dis* 2015;21:17-30.
11. Tolentino Ede S, Centurion BS, Ferreira LH, Souza AP, Damante JH, Rubira-Bullen IR. Oral adverse effects of head and neck radiotherapy: Literature review and suggestion of a clinical oral care guideline for irradiated patients. *J Appl Oral Sci* 2011;19:448-54.
12. Hooper SJ, Wilson MJ, Crean SJ. Exploring the link between microorganisms and oral cancer: A systematic review of the literature. *Head Neck* 2009;31:1228-39.
13. Pushalkar S, Ji X, Li Y, Estilo C, Yegnanarayana R, Singh B, et al. Comparison of oral microbiota in tumor and non-tumor tissues of patients with oral squamous cell carcinoma. *BMC Microbiol* 2012;12:144.
14. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the normal bacterial flora of the oral cavity. *J Clin Microbiol* 2005;43:5721-32.
15. Bik EM, Long CD, Armitage GC, Loomer P, Emerson J, Mongodin EF, et al. Bacterial diversity in the oral cavity of 10 healthy individuals. *ISME J* 2010;4:962-74.
16. Abdulla R, Adyanthaya S, Kini P, Mohanty V, D'Souza N, Subbannayya Y. Clinicopathological analysis of oral squamous cell carcinoma among the younger age group in coastal Karnataka, India: A retrospective study. *J Oral Maxillofac Pathol* 2018;22:180-7.
17. Selvamani M, Yamunadevi A, Basandi PS, Madhushankari GS. Prevalence of oral squamous cell carcinoma of tongue in and around Davangere, Karnataka, India: A retrospective study over 13 years. *J Pharm Bioallied Sci* 2015;7:S491-4.
18. Sharma P, Deb T, Ray JG, Singh AK, Gupta G, Das A, et al. Oral squamous cell carcinoma profile in North-Eastern regions of India from habits to histopathology: A hospital-based study. *Natl J Maxillofac Surg* 2018;9:56-60.
19. Sarkaria JN, Harari PM. Oral tongue cancer in young adults less than 40 years of age: Rationale for aggressive therapy. *Head Neck* 1994;16:107-11.
20. Ariyoshi Y, Shimahara M, Omura K, Yamamoto E, Mizuki H, Chiba H, et al. Epidemiological study of malignant tumors in the oral and maxillofacial region: Survey of member institutions of the Japanese society of oral and maxillofacial surgeons, 2002. *Int J Clin Oncol* 2008;13:220-8.
21. Bhurgri Y, Bhurgri A, Usman A, Pervez S, Kayani N, Bashir I, et al. Epidemiological review of head and neck cancers in Karachi. *Asian Pac J Cancer Prev* 2006;7:195-200.
22. Halboub ES, Al-Anazi YM, Al-Mohaya MA. Characterization of Yemeni patients treated for oral and pharyngeal cancers in Saudi Arabia. *Saudi Med J* 2011;32:1177-82.
23. Chiang CT, Hwang YH, Su CC, Tsai KY, Lian IE, Yuan TH, et al. Elucidating the underlying causes of oral cancer through spatial clustering in high-risk areas of Taiwan with a distinct gender ratio of incidence. *Geospat Health* 2010;4:230-42.
24. Kruaysawat W, Aekplakorn W, Chapman RS. Survival time and prognostic factors of oral cancer in Ubon Ratchathani cancer center. *J Med Assoc Thai* 2010;93:278-84.
25. Singh MP, Kumar V, Agarwal A, Kumar R, Bhatt ML, Misra S. Clinic-epidemiological study of oral squamous cell carcinoma: A tertiary care centre study in North India. *J Oral Biol Craniofac Res* 2016;6:31-4.
26. Ajay PR, Ashwinirani SR, Nayak A, Suragimath G, Kamala KA, Sande A, et al. Oral cancer prevalence in Western population of Maharashtra, India, for a period of 5 years. *J Oral Res Rev* 2018;10:11-4.
27. Kudo Y, Tada H, Fujiwara N, Tada Y, Tsunematsu T, Miyake Y, et al. Oral environment and cancer. *Genes Environ* 2016;38:13.
28. Pushalkar S, Mane SP, Ji X, Li Y, Evans C, Crasta OR, et al. Microbial diversity in saliva of oral squamous cell carcinoma. *FEMS Immunol Med Microbiol* 2011;61:269-77.
29. Correa P, Haenszel W, Cuello C, Tannenbaum S, Archer M. A model for gastric cancer epidemiology. *Lancet* 1975;2:58-60.
30. Llewellyn CD, Johnson NW, Warnakulasuriya KA. Risk factors for oral cancer in newly diagnosed patients aged 45 years and younger: A case-control study in Southern England. *J Oral Pathol Med* 2004;33:525-32.
31. Maier H, Zöller J, Herrmann A, Kreiss M, Heller WD. Dental status and oral hygiene in patients with head and neck cancer. *Otolaryngol Head Neck Surg* 1993;108:655-61.
32. Tezal M, Sullivan MA, Reid ME, Marshall JR, Hyland A, Loree T, et al. Chronic periodontitis and the risk of tongue cancer. *Arch Otolaryngol Head Neck Surg* 2007;133:450-4.
33. Nagy KN, Sonkodi I, Szöke I, Nagy E, Newman HN. The microflora associated with human oral carcinomas. *Oral Oncol* 1998;34:304-8.
34. Binder Gallimidi A, Fischman S, Revach B, Bulvik R, Maliutina A, Rubinstein AM, et al. Periodontal pathogens *Porphyromonas gingivalis* and *Fusobacterium nucleatum* promote tumor progression in an oral-specific chemical carcinogenesis model. *Oncotarget* 2015;6:22613-23.
35. Kurkivuori J, Salaspuro V, Kaihovaara P, Kari K, Rautemaa R, Grönroos

- L, *et al.* Acetaldehyde production from ethanol by oral streptococci. *Oral Oncol* 2007;43:181-6.
36. Jham BC, da Silva Freire AR. Oral complications of radiotherapy in the head and neck. *Braz J Otorhinolaryngol* 2006;72:704-8.
 37. Sroussi HY, Epstein JB, Bensadoun RJ, Saunders DP, Lalla RV, Migliorati CA, *et al.* Common oral complications of head and neck cancer radiation therapy: Mucositis, infections, saliva change, fibrosis, sensory dysfunctions, dental caries, periodontal disease, and osteoradionecrosis. *Cancer Med* 2017;6:2918-31.
 38. Kamath MP, Hegde MC, Sreedharan S, Salmi DK, Padmanabhan K. Radiotherapeutic effect on oropharyngeal flora in head and neck cancer. *Indian J Otolaryngol Head Neck Surg* 2002;54:111-4.
 39. Sharma DC. Salivary bacteria linked to oral cancers. *Lancet Oncol* 2005;6:547.
 40. Eliasson L, Carlén A, Almståhl A, Wikström M, Lingström P. Dental plaque pH and micro-organisms during hyposalivation. *J Dent Res* 2006;85:334-8.
 41. Srithavaj T, Thaweboon S. Determination of oral microflora in irradiated ocular deformed children. *Southeast Asian J Trop Med Public Health* 2006;37:991-5.
 42. Al-Nawas B, Grötz KA. Prospective study of the long term change of the oral flora after radiation therapy. *Support Care Cancer* 2006;14:291-6.
 43. Beighton D. The complex oral microflora of high-risk individuals and groups and its role in the caries process. *Community Dent Oral Epidemiol* 2005;33:248-55.
 44. Lunt SJ, Chaudary N, Hill RP. The tumor microenvironment and metastatic disease. *Clin Exp Metastasis* 2009;26:19-34.
 45. Allavena P, Garlanda C, Borrello MG, Sica A, Mantovani A. Pathways connecting inflammation and cancer. *Curr Opin Genet Dev* 2008;18:3-10.
 46. Lamont RJ, Jenkinson HF. Life below the gum line: Pathogenic mechanisms of *Porphyromonas gingivalis*. *Microbiol Mol Biol Rev* 1998;62:1244-63.
 47. Krogh P. The role of yeasts in oral cancer by means of endogenous nitrosation. *Acta Odontol Scand* 1990;48:85-8.