

Published online: 20 June 2018

OPEN Author Correction: Proximity of breeding and foraging areas affects foraging effort of a crepuscular, insectivorous bird

Ruben Evens¹, Natalie Beenaerts¹, Thomas Neyens², Nele Witters³, Karen Smeets¹ & Tom Artois1

Correction to: Scientific Reports https://doi.org/10.1038/s41598-018-21321-0, published online 14 February 2018

The Acknowledgements section in this Article is incomplete.

"RE is funded by a BOF-mandate (BSFFCMKDK) at Hasselt University. NW is funded as a postdoctoral researcher by FWO Flanders. Part of this research was funded by another FWO-grant (N1522015). We thank Eddy Ulenaers, Luc Lens, Dries Gorissen, Guido Winters, Koen Thijs, Lise Hendrick, Karen Vanmarcke, Fien Evens, Albert Geuens, Lyndon Kearsley, Greg Conway, Ian Henderson and Nikki Watson for their support and relevant comments on this manuscript. Research equipment was funded by the Agency for Nature and Forest (ANB; Belgium). Permissions were granted by the ANB and Royal Belgian Institute of Natural Sciences (Belgium)."

should read:

"RE is funded by a BOF-mandate (BSFFCMKDK) at Hasselt University. NW and TN are funded as postdoctoral researchers by FWO Flanders. Part of this research was funded by another FWO-grant (N1522015). We thank Eddy Ulenaers, Luc Lens, Dries Gorissen, Guido Winters, Koen Thijs, Lise Hendrick, Karen Vanmarcke, Fien Evens, Albert Geuens, Lyndon Kearsley, Greg Conway, Ian Henderson and Nikki Watson for their support and relevant comments on this manuscript. Research equipment was funded by the Agency for Nature and Forest (ANB; Belgium). Permissions were granted by the ANB and Royal Belgian Institute of Natural Sciences (Belgium)."

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2018

¹Hasselt University, Centre for Environmental Sciences, Research Group: Zoology, Biodiversity and Toxicology, Campus Diepenbeek, Agoralaan, Gebouw D, 3590, Diepenbeek, Belgium. ²Hasselt University, Centre for Statistics, Research Group: I-BIOSTAT, Campus Diepenbeek, Agoralaan, Gebouw D, 3590, Diepenbeek, Belgium. ³Hasselt University, Centre for Environmental Sciences, Research Group: Environmental Economics, Campus Diepenbeek, Agoralaan, Gebouw D, 3590, Diepenbeek, Belgium. Correspondence and requests for materials should be addressed to R.E. (email: ruben.evens@uhasselt.be)