



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

---

# Coronary artery calcium score as a prognostic factor of adverse outcomes in patients with COVID-19: a comprehensive review

Fateme Yousefimoghaddam<sup>a</sup>, Ehsan Goudarzi<sup>a</sup>,  
Alireza Ramandi<sup>b,c</sup>, and Isa Khareshi<sup>a,\*</sup>

From the <sup>a</sup> Cardiovascular Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran, <sup>b</sup> School of Medicine, Tehran University of Medical Sciences, Tehran, Iran and <sup>c</sup> Students' Scientific Research Center (SSRC), Tehran University of Medical Sciences, Tehran, Iran.

**Abstract: Background and aim:** The association of known cardiovascular risk factors and poor prognosis of coronavirus disease 2019 (COVID-19) has been recently emphasized. Coronary artery calcium (CAC) score is considered to be a risk predictor of cardiovascular events. Therefore, we have conducted a review of literature on the predictive value of CAC score predictive value in COVID-19 outcome. **Method:** A search of literature was conducted, aiming for articles published until December 2021 on PubMed and Scopus to identify potentially eligible studies. **Discussion:** A total of 18 articles were reviewed for association between higher CAC score and adverse outcomes in COVID-19. **Conclusion:** The coronary calcium score could be considered as a new radiological marker for risk assessment in COVID-19 patients and providing additional information in fields of prognosis and possible cardiovascular complications. High CAC score is associated with higher in-hospital death and adverse clinical outcomes in patients with confirmed COVID-19, which highlights the importance of calcium load

---

The authors of this manuscript have no conflicts of interest to disclose. The authors declare no relationships with any companies, whose products or services may be related to the subject matter of the article.

Curr Probl Cardiol 2022;00:101175

0146-2806/\$ – see front matter

<https://doi.org/10.1016/j.cpcardiol.2022.101175>

**testing for hospitalized COVID-19 patients and calls for attention to patients with high CAC scores. (Curr Probl Cardiol 2022;00:101175.)**

## Introduction

**S**ince the first confirmed case in Wuhan, China in December 2019, the Coronavirus disease 2019 (COVID-19) has been declared as a pandemic disease; infecting millions of people around the world.<sup>1</sup> COVID-19 infection may manifest a wide array of symptoms, including common forms of mild and nonspecific constitutional symptoms, along with more severe complications such as respiratory failure, sepsis, and acute cardiac injury. All of which may require intensive care treatments and ventilation.<sup>2-5</sup> Patients with pre-existing cardiovascular comorbidities are more presumably predisposed to COVID-19 infection complications.<sup>6-9</sup> As previous literature has noted the association of higher complication rates and mortality rates in COVID-19 patients with pre-existing hypertension, diabetes mellitus, or a past history of cardiac events.<sup>10,11</sup> Although several studies have investigated the correlation of COVID-19 infection and atherosclerosis-related diseases, there are limited number of studies that report quantitative data and measurements related to atherosclerotic features observed.<sup>12,13</sup> Coronary artery calcification (CAC) score is a reliable and quantitative predictor of cardiovascular events and mortality by utilizing various scoring systems such as Agatston score and volume score.<sup>14-16</sup> CAC scoring is a highly sensitive method with high negative predictive value which could be used as the solitary prognostic method for estimating risk of Cardiovascular disease.<sup>17</sup> However, the prognostic role of CAC score in acute systemic inflammatory diseases, such as COVID-19, has not yet been investigated. Chest CT imaging is a common work-up procedure for COVID-19 patients. This imaging modality demonstrates the extent of pulmonary dysfunction, detects COVID-19 infection in suspected patients, and may be used as screening tool for COVID-19 in patients with severe respiratory insufficiency.<sup>18,19</sup> These data may lead us to an imaging biomarker in COVID-19 patients; resulting in further insight into the disease process, reduction of unnecessary work-up procedures, and estimation of required intensive care unit beds prior to patients' complications occurrence. We herein aim to evaluate short-term prognostic value of CAC in patients with COVID-19 infection, along with discussing the probable pathophysiologic mechanisms of COVID-19 adverse effects.

## Methods and Materials

A comprehensive review of literature was performed to collect currently available data on using coronary artery calcium score in COVID-19 patients as an imaging marker for adverse outcomes. In order to identify qualified studies, a comprehensive search was conducted on PubMed and Scopus, aiming the articles published from January 2020 to December 2021. The search was carried out using the following strategy: (1) Language restriction was not applied. (2) Related articles function on PubMed was used to extend the search, followed by (3) the manual search of the bibliography to find other potentially eligible studies. The search results were screened by title and abstract, and the potentially relevant articles selected for full-text evaluation. Data extraction was conducted in a double data extraction method by two independent reviewers (F.Y, E.G), followed by a precise reevaluation of included articles by 2 other members of authoring team (A.R, I.K.)

## Review of Articles

We reviewed the association between the presence and severity of CAC as an independent risk factor and adverse outcomes in COVID-19 in 18 articles. The adverse outcomes included in this review consisted of all-cause mortality, ICU requirement, and ventilation requirement. Further details of these articles are presented in [Table](#). Among 18 articles under study, 17 articles were retrospective cohort or cross-sectional studies and 1 was letter to editor. CAC score assessment method used in the articles are 7 visual assessment (n = 7), Agatston (n = 5), volume score (n = 1), and both volume score and Agatston (n = 2). Two articles solely examined whether the calcified plaque is present or not. The association between CAC score with hypertension, dyslipidemia or smoking status was heterogeneous between the studies. *P*-value lower than 0.05 was considered statistically significant and COVID-19 was laboratory-confirmed in all patients.

### *All-Cause Mortality*

In 14 articles, higher CAC score was associated with higher all-cause mortality rate, in 1 there was no association between the two. Three articles did not state the association between all-cause mortality rate and higher CAC score.

**TABLE.** Review of literature on CAC score association with adverse outcomes in COVID-19

Article	Sample size	Average age	Female sex (%)	CACs assessment method	CACs association with		
					All-cause mortality	ICU requirements	Ventilation
Mousseaux et al, 2021 <sup>28</sup>	169	65.6 ± 15.8	30.1	V-CACs	Associated	Not Associated	Associated (both invasive and noninvasive ventilation)
Plenek et al, 202 <sup>29</sup>	245	NM	42	CAC presence or not	Associated with 60 days mortality, Not associated with In-hospital Mortality	Not Associated	Not Associated
Giannini et al, 2021 <sup>30</sup>	1093	68	32.1	Agatston and volume assessment	Associated	NM	Not associated with Noninvasive ventilation, Associated with invasive ventilation
Dillinger et al, 2020 <sup>12</sup>	209	62	28	CAC presence or not	Not Associated	NM	Associated with Noninvasive ventilation, Not associated with Invasive ventilation
Slipczuk et al, 2021 <sup>31</sup>	493	NM	50.5	V-CACs	Associated	NM	NM
Luo et al, 2021 <sup>32</sup>	2067	50 ± 15.6	46.9	V-CACs	Associated	Associated	Associated with invasive ventilation
Gupta et al, 2021 <sup>33</sup>	180	68	45.6	V-CACs	Associated	NM	Not Associated

*(continued on next page)*

TABLE. (continued)

Article	Sample size	Average age	Female sex (%)	CACs assessment method	CACs association with		
					All-cause mortality	ICU requirements	Ventilation
Fervers et al, 2021 <sup>34</sup>	89	58.1 ± 15.9	44.9	Agatston	Associated*	Associated*	Associated*
Takeshita et al, 2021 <sup>35</sup>	53	66.7	50.9	Agatston	NM	NM	NM
Cereda et al, 2021 <sup>36</sup>	1565	67.9 ± 13	33.4	volume assessment	Associated	NM	Associated with invasive ventilation
Zimmermann et al, 2020 <sup>37</sup>	109	61.8 ± 17.5	30.1	Agatston	Associated	Associated	NM
Scoccia et al, 2021 <sup>38</sup>	1625	69	32.8	Agatston	Associated	NM	NM
Fazzari et al, 2021 <sup>39</sup>	282	64 ± 13	29.8	Agatston	Associated	Not Associated	Not Associated (both noninvasive and invasive ventilation)
Luchian et al, 2021 <sup>40</sup>	280	63.2 ± 16.7	42.5	V-CACs	NM	Not Associated	NM
Colombi et al, 2020 <sup>41</sup>	248	68	29.8	V-CACs	Associated	NM	NM
Cereda et al, 2021 <sup>42</sup>	1683	67 ± 14	32.8	Agatston and volume assessment	Associated	NM	NM
Nair et al, 2021 <sup>43</sup>	67	NM	4.5	V-CACS	NM	NM	NM
Cosyns et al, 2020 <sup>44</sup>	280	NM	NM	NM	Associated	NM	NM

CACs, Coronary Artery Calcium score; NM, Not mentioned; V-CACS, visual assessment of CACs.

\*CAC score association with mortality, ICU requirements and ventilation was not evaluated separately.

## ICU

In 3 articles higher CAC score was associated with higher ICU requirement; while in 4 studies, there were no associations. 11 articles did not evaluate to the association between ICU requirement and higher CAC score.

## Ventilation

In 5 articles higher CAC score was associated with higher ventilation (invasive or noninvasive) rate. However, 4 articles have found no associations, and 9 articles did not report any data about the association between ventilation rate and higher CAC score.

Furthermore, 2 articles did not mention any associations between CAC score and mortality, ICU admission, or requirement of ventilation. However, higher CAC score had a positive correlation with deterioration of oxygenation and COVID-19 severity in these studies.

## Discussion

The Inflammatory response toward COVID-19 infection precipitates major course of the disease, including multiorgan involvement.<sup>20</sup> Early diagnosis and treatment of systemic or local inflammatory reaction is the key to eluding severe complications. Cardiovascular involvement is one of the most common complications in COVID-19 patients.<sup>21</sup>

Vessel calcification is an atherosclerotic change in the vessel walls that occurs in multiple cardiovascular diseases. CAC illustrates the lifetime aggregate burden of all for atherosclerotic risk factors.<sup>16</sup> among the risk factors of atherosclerotic calcification are older age, active smoking, diabetes mellitus, dyslipidemia and arterial hypertension;<sup>22,23</sup> All of which are also known to be associated with poor COVID-19 outcome. Since Low-dose chest CT (LDCT) is routinely obtained from COVID-19 patients in most health centers,<sup>18,19,24</sup> it can also be used to evaluate the heart and coronary vessels function to some extent. Calcified atherosclerotic plaques are detected via their higher intrinsic contrast relative to the adjacent soft tissue. Three commonly used grading systems for clinical quantification of CAC are (1) Agatston score, (2) visual assessment of coronary artery calcification, and (3) volume score. The Agatston score reflects both the volume of calcification and highest attenuation of calcified atherosclerotic plaques on a CT scan. Visual assessment of coronary artery calcification is a simple scoring system for classifying [coronary artery calcification](#) into 4 levels: none, mild, moderate or heavy.<sup>25,26</sup> The

volume score has scanning parameters similar to the Agatston score. However, it relies on estimating the true volume of calcified plaque, instead of lesion density.<sup>27</sup>

Our review has some limitations. The articles we reviewed determined different endpoints for mortality, ranging from in hospital mortality to 6 months mortality. This may cause bias in qualitative analysis of the outcomes.

## Conclusion

Our findings emphasize the potentially useful assessment of CAC during chest-CT imaging of COVID-19 patients. We noted that CAC can be used as an additional risk predictor of all-cause mortality in COVID-19 patients. The major strengths of CAC score is reporting both qualitative and quantitative data, cost-efficiency, and the ability to assess both prospectively and retrospectively in the studies. However, the association between higher CAC score and higher ICU requirement and ventilation rate is not clear yet and more studies are required.

## Authors Contribution

F.Y and E.G performed the search and data extraction and full-text review. A.R and I.K. investigated the correctness of included articles and performed second investigation on full-text of articles. E.G, F.Y contributed to table preparation. F.Y and A.R performed manuscript preparation. I.K performed final revision.

## REFERENCES

1. World Health Organization website 2021. Available from: <https://covid19.who.int/>.
2. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497–506.
3. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323:1061–9.
4. Eastin C, Eastin T. Clinical Characteristics of Coronavirus Disease 2019 in China: Guan W, Ni Z, Hu Y, et al. *N Engl J Med* 2020;58:711–2. [Online ahead of print] DOI: 10.1056/NEJMoa2002032.
5. Gupta A, Madhavan MV, Sehgal K, Nair N, Mahajan S, Sehrawat TS, et al. Extrapulmonary manifestations of COVID-19. *Nat Med* 2020;26:1017–32.
6. Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O. Potential effects of coronaviruses on the cardiovascular system: a review. *JAMA Cardiol* 2020;5:831–40.



7. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol* 2020;5:802–10.
8. Wang Y, Lu X, Li Y, Chen H, Chen T, Su N, et al. Clinical course and outcomes of 344 intensive care patients with COVID-19. *Am J Respir Crit Care Med* 2020;201:1430–4.
9. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–62.
10. Nishiga M, Wang DW, Han Y, Lewis DB, Wu JC. COVID-19 and cardiovascular disease: from basic mechanisms to clinical perspectives. *Nat Rev Cardiol* 2020;17:543–58.
11. Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. *JAMA*. 2020;323:1574–81.
12. Dillinger JG, Benmessaoud FA, Pezel T, Voicu S, Sideris G, Chergui N, et al. Coronary artery calcification and complications in patients with COVID-19. *JACC Cardiovasc Imaging* 2020;13:2468–70.
13. Nai Fovino L, Cademartiri F, Tarantini G. Subclinical coronary artery disease in COVID-19 patients. *Eur Heart J Cardiovasc Imaging* 2020;21:1055–6.
14. Xie X, Zhao Y, de Bock GH, de Jong PA, Mali WP, Oudkerk M, et al. Validation and prognosis of coronary artery calcium scoring in nontriggered thoracic computed tomography: systematic review and meta-analysis. *Circ Cardiovasc Imaging* 2013;6:514–21.
15. Azour L, Kadoch MA, Ward TJ, Eber CD, Jacobi AH. Estimation of cardiovascular risk on routine chest CT: Ordinal coronary artery calcium scoring as an accurate predictor of Agatston score ranges. *J Cardiovasc Comput Tomogr* 2017;11:8–15.
16. Greenland P, Blaha MJ, Budoff MJ, Erbel R, Watson KE. Coronary calcium score and cardiovascular risk. *J Am Coll Cardiol* 2018;72:434–47.
17. Cainzos-Achirica M, Di Carlo PA, Handy CE, Quispe R, Roura G, Pinto X, et al. Coronary artery calcium score: the “Mammogram” of the heart? *Curr Cardiol Rep* 2018;20:1–10.
18. Rubin GD, Ryerson CJ, Haramati LB, Sverzellati N, Kanne JP, Raof S, et al. The role of chest imaging in patient management during the COVID-19 pandemic: a multinational consensus statement from the Fleischner Society. *Chest*. 2020;158:106–16.
19. Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, et al. CT imaging features of 2019 novel coronavirus (2019-nCoV). *Radiology* 2020;295:202–7.
20. Thakur V, Ratho RK, Kumar P, Bhatia SK, Bora I, Mohi GK, et al. Multi-organ involvement in COVID-19: beyond pulmonary manifestations. *J Clin Med* 2021;10:446. <https://doi.org/10.3390/jcm10030446>.
21. Zaim S, Chong JH, Sankaranarayanan V, Harky A. COVID-19 and multiorgan response. *Curr Probl Cardiol* 2020;45:100618.
22. Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, et al. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020;5:811–8.

23. Tartof SY, Qian L, Hong V, Wei R, Nadjafi RF, Fischer H, et al. Obesity and mortality among patients diagnosed with COVID-19: results from an integrated health care organization. *Ann Intern Med* 2020;173:773–81.
24. Agricola E, Beneduce A, Esposito A, Ingallina G, Palumbo D, Palmisano A, et al. Heart and lung multimodality imaging in COVID-19. *JACC Cardiovasc Imaging* 2020;13:1792–808.
25. Chiles C, Duan F, Gladish GW, Ravenel JG, Baginski SG, Snyder BS, et al. Association of coronary artery calcification and mortality in the national lung screening trial: a comparison of three scoring methods. *Radiology* 2015;276:82–90.
26. Einstein AJ, Johnson LL, Bokhari S, Son J, Thompson RC, Bateman TM, et al. Agreement of visual estimation of coronary artery calcium from low-dose CT attenuation correction scans in hybrid PET/CT and SPECT/CT with standard Agatston score. *J Am Coll Cardiol* 2010;56:1914–21.
27. Blaha MJ, Mortensen MB, Kianoush S, Tota-Maharaj R, Cainzos-Achirica M. Coronary artery calcium scoring: is it time for a change in methodology? *JACC Cardiovasc Imaging* 2017;10:923–37.
28. Mousseaux E, Fayol A, Danchin N, Soulat G, Charpentier E, Livrozet M, et al. Association between coronary artery calcifications and 6-month mortality in hospitalized patients with COVID-19. *Diagn Interv Imaging* 2021;102:717–25.
29. Planek MIC, Ruge M, Du Fay de Lavallaz JM, Kyung SB, Gomez JMD, Suboc TM, et al. Cardiovascular findings on chest computed tomography associated with COVID-19 adverse clinical outcomes. *Am Heart J Plus* 2021;11:100052.
30. Giannini F, Toselli M, Palmisano A, Cereda A, Vignale D, Leone R, et al. Coronary and total thoracic calcium scores predict mortality and provides pathophysiologic insights in COVID-19 patients. *J Cardiovasc Comput Tomograp* 2021;15:421–30.
31. Slipeczuk L, Castagna F, Schonberger A, Novogrodsky E, Sekerak R, Dey D, et al. Coronary artery calcification and epicardial adipose tissue as independent predictors of mortality in COVID-19. *Int J Cardiovasc Imaging* 2021;37:3093–100.
32. Luo S, Qiu XM, Zeng XJ, Zhang DY, Wan B, Li X, et al. Coronary artery calcification and risk of mortality and adverse outcomes in patients with COVID-19: a Chinese multicenter retrospective cohort study. *Chin J Acad Radiol* 2021:1–9.
33. Gupta YS, Finkelstein M, Manna S, Toussie D, Bernheim A, Little BP, et al. Coronary artery calcification in COVID-19 patients: an imaging biomarker for adverse clinical outcomes. *Clin Imaging* 2021;77:1–8.
34. Fervers P, Kottlors J, Große Hokamp N, Bremm J, Maintz D, Tritt S, et al. Coronary artery calcification on low-dose chest CT is an early predictor of severe progression of COVID-19—a multi-center, multi-vendor study. *PLoS One* 2021;16:e0255045.
35. Takeshita Y, Terada J, Fujita R, Hirasawa Y, Kinoshita T, Isaka Y, et al. Coronary artery calcium score may be a novel predictor of COVID-19 prognosis: a retrospective study. *BMJ Open Respir Res* 2021;8:e000923. <https://doi.org/10.1136/bmjresp-2021-000923>.
36. Cereda A, Toselli M, Palmisano A, Vignale D, Khokhar A, Campo G, et al. Coronary calcium score as a predictor of outcomes in the hypertensive Covid-19 population:

- results from the Italian (S) Core-Covid-19 Registry. *Hypertens Res* 2021;45:333–43. <https://doi.org/10.1038/s41440-021-00798-9>.
37. Zimmermann GS, Fingerle AA, Müller-Leisse C, Gassert F, von Schacky CE, Ibrahim T, et al. Coronary calcium scoring assessed on native screening chest CT imaging as predictor for outcome in COVID-19: An analysis of a hospitalized German cohort. *PLoS One* 2020;15:e0244707.
  38. Scoccia A, Gallone G, Cereda A, Palmisano A, Vignale D, Leone R, et al. Impact of clinical and subclinical coronary artery disease as assessed by coronary artery calcium in COVID-19. *Atherosclerosis* 2021;328:136–43.
  39. Fazzari F, Cozzi O, Maurina M, Donghi V, Indolfi E, Curzi M, et al. In-hospital prognostic role of coronary atherosclerotic burden in COVID-19 patients. *J Cardiovasc Med (Hagerstown)* 2021;22:818–27.
  40. Luchian ML, Lochy S, Motoc A, Belsack D, Magne J, Roosens B, et al. Prognostic value of coronary artery calcium score in hospitalized COVID-19 patients. *Front Cardiovasc Med* 2021;8:684528.
  41. Colombi D, Villani GD, Maffi G, Risoli C, Bodini FC, Petrini M, et al. Qualitative and quantitative chest CT parameters as predictors of specific mortality in COVID-19 patients. *Emerg Radiol* 2020;27:701–10.
  42. Cereda A, Toselli M, Palmisano A, Vignale D, Leone R, Nicoletti V, et al. The hidden interplay between sex and COVID-19 mortality: the role of cardiovascular calcification. *GeroScience* 2021;43:2215–29.
  43. Nair AV, Kumar D, Yadav SK, Nepal P, Jacob B, Al-Heidous M. Utility of visual coronary artery calcification on non-cardiac gated thoracic CT in predicting clinical severity and outcome in COVID-19. *Clin Imaging* 2021;74:123–30.
  44. Cosyns B, Motoc A, Luchian ML, Lochy S, Belsack D. Coronary calcium score in COVID-19 hospitalized patients. *JACC: Cardiovascular Imaging* 2020;13:2698.