

Suppressive effects of quercetin on hydrogen peroxide-induced caveolin-1 phosphorylation in endothelial cells

Akari Kondo-Kawai,^{1,2,3} Tohru Sakai,² Junji Terao,^{1,4} and Rie Mukai^{1,5,*}

¹Department of Food Science, Graduate School of Biomedical Sciences and ⁵Department of Food Science, Graduate School of Technology, Industrial and Social Sciences, Tokushima University, Tokushima, Tokushima 770-8513, Japan

²Department of Public Health and Applied Nutrition, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima, Tokushima 770-8503, Japan

³Laboratory of Nutritional Science, Shikoku Junior College, Tokushima, Tokushima 771-1192, Japan

⁴Faculty of Clinical Nutrition and Dietetics, Konan Women's University, Kobe, Hyogo 658-0001, Japan

(Received 27 November, 2020; Accepted 19 December, 2020; Published online 3 April, 2021)

Caveolin-1 is a major protein of the caveolae structure in vascular endothelial cell membrane. Phosphorylation of caveolin-1 is one of the initial events leading to exacerbation of vascular permeability caused by oxidative stress. Although quercetin is known to be an anti-atherosclerosis factor that acts as a dietary antioxidant, little is known about its role in the regulation of caveolin-1 phosphorylation. In this study, we investigated the inhibitory effect of quercetin on hydrogen peroxide-induced caveolin-1 phosphorylation in human umbilical vein endothelial cells. Quercetin inhibited caveolin-1 phosphorylation in cells pretreated with quercetin for 24 h and then exposed to hydrogen peroxide. However, quercetin 3-*O*- β -glucuronide, a conjugated metabolite of quercetin, did not exert this inhibitory effect. Exposure to hydrogen peroxide increased vascular permeability and reduced mRNA expression of the intercellular adhesion protein, vascular endothelial cadherin (VE-cadherin). By contrast, pretreatment with quercetin suppressed the increase in vascular permeability and decreased VE-cadherin expression. These results indicate that deconjugated quercetin can play a role in the prevention of altered vascular permeability under oxidative stress by suppressing caveolin-1 phosphorylation. Thus, dietary quercetin may be beneficial for the maintenance of endothelial cell function.

Key Words: quercetin, caveolae, caveolin-1, hydrogen peroxide, vascular endothelium

Cardiovascular disease has provoked risk for premature death worldwide.^(1,2) It was estimated 422.7 million people was suffered from cardiovascular disease in 2015, and was estimated 17.9 million deaths worldwide in 2015. It was comprising 31% of all global deaths.^(1,3) Atherosclerosis is one of the factors that cause coronary artery disease and cerebrovascular disease. Atherosclerosis results in dysfunction of the vascular endothelium, formation of fatty streak in blood vessels, and plaque fibrosis.⁽⁴⁾ Basic diet control and exercise therapy are needed for preventing atherosclerosis.⁽⁵⁾ In addition to this, it has been suggested that several bioactive within dietary food can be is one of the solution for preventing atherosclerosis.^(6,7)

Endothelial cells play a central role in maintaining homeostasis of vascular function.^(8,9) Tight junctions and adherens junctions suppress the permeation of substances through the gap of neighboring vascular endothelial cells. However, the impairment of cell-cell adhesion in inflamed tissues enhances vascular permeability,^(10,11) and the resulting

invasion of lipoproteins and/or plasma components into the vessel walls accelerates vascular endothelial dysfunction. Vascular endothelial cells are densely populated with caveolae, which are flask-shaped invaginations measuring 50–100 nm in diameter.⁽¹²⁾ Caveolae are rich in sphingolipids and cholesterol, and are implied to serve as a platform for endocytosis and signal transduction,^(13–16) particularly in the cell membrane. Caveolin-1 (Cav-1), a major protein component of caveolae, is involved in the regulation of lipoprotein transcytosis, vascular inflammation,⁽¹⁷⁾ and progression of atherosclerosis.⁽¹⁸⁾

Phosphorylation of Cav-1 has been reported to increase the endocytosis of plasma albumin and contribute to vascular permeability by enhancing vascular endothelial cadherin (VE-cadherin) endocytosis.^(19–21) Furthermore, phosphorylation of Cav-1 in the endothelium was shown to be enhanced in response to oxidized low-density proteins (ox-LDL) and hydrogen peroxide (H₂O₂).^(22,23) It is therefore likely that the inhibition of Cav-1 phosphorylation is involved in the protective action of antioxidative food factors in the endothelium.

Flavonoids, which are composed of diphenyl propane structure, are distributed ubiquitously in the plant kingdom and have been suggested to exert a wide variety of health promoting effects.^(24–26) It has already been reported that higher flavonoid intake lowers mortality due to cardiovascular diseases.⁽²⁷⁾ Moreover, it is known that the amount of flavonoid intake is inversely correlated with the incidence rate of cardiovascular diseases.⁽²⁸⁾ Quercetin (3,3',4,5,7-pentahydroxyflavone; Q), a flavonol-type flavonoid commonly present in fruits and vegetables, is a potential candidate for use in preventing the onset of cardiovascular diseases owing to its anti-oxidative or anti-atherosclerosis functions.^(29–31)

Although Q mainly exists as a glucoside in plants,⁽³²⁾ it is hydrolyzed in epithelial cells of the small intestine during absorption,⁽³³⁾ and the resulting aglycone is further converted into various metabolites, including glucuronides, sulfates, and/or their *O*-methyl derivatives (isorhamnetin and tamarixetin) in the small intestine and liver.⁽³³⁾ Kawai *et al.*⁽³⁴⁾ and Ishisaka *et al.*⁽³⁵⁾ reported that Q 3-*O*-glucuronide (Q3GA), a major conjugated Q metabolite, is converted to its aglycone form by the action of macrophages. It is likely that Q glucuronide metabolites present

Department of Food Science, Graduate School of Biomedical Sciences, Tokushima University is the previous affiliation for Akari Kondo-Kawai, Junji Terao, and Rie Mukai. They continued this work at the present affiliation.

*To whom correspondence should be addressed.

E-mail: rmukai@tokushima-u.ac.jp

in the plasma act as anti-atherosclerotic agents in the aorta after it is converted to its aglycone form.⁽³⁶⁻³⁸⁾ In a previous study,⁽³⁹⁾ we investigated the effects of Q glucuronide metabolites and its aglycone on ox-LDL-induced expression of Cav-1 in human umbilical vein endothelial cells (HUVECs). The purpose of this study was to evaluate the protective role of Q against endothelial dysfunction, with a focus on suppression of Cav-1 phosphorylation using a H₂O₂-exposed HUVECs system.

Materials and Methods

Reagents. Q dehydrate, H₂O₂, and anti-Cav-1 (c3237) were purchased from Sigma Aldrich Co. (St. Louis, MO). Anti-glyceraldehyde-3-phosphate dehydrogenase (GAPDH, 14C10) and anti-phospho-Fyn (D49G4) were obtained from Cell Signaling Technology (Beverly, MA). Anti-phospho-Cav-1 (sc-14037) and anti-Fyn (sc-16) antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Q3GA and myricetin (M) (1127S) were obtained from Extrasynthese (Genay, France). Flavone (F) was obtained from Nacalai Tesque (16012-31) (Kyoto, Japan). Luteolin (L) was obtained from LKT Laboratories (L8377; Saint Paul, MN) and 8-prenyl Q was synthesized as previously described by Kawamura *et al.*⁽⁴⁰⁾

Cells and culture. HUVECs were purchased from Lonza Japan (Tokyo, Japan). HUVECs were cultured in endothelial cell growth medium EGM-2 (cc-3156) mixed with SingleQuots™ (cc-4176) (Lonza, Basel, Switzerland) and maintained at 37°C in a humidified atmosphere containing 5% CO₂. Cells were used at passages from #3 to #10.

Real-time reverse-transcription polymerase chain reaction (RT-PCR). HUVECs (6 × 10⁴ cells/well) were plated on 24-well plates and incubated for 24 h. Cells were treated with H₂O₂ and Q-related compounds. To estimate the effects of flavonoids, they were added to the cells before H₂O₂ treatment. Briefly, cells treated with flavonoids were washed with Hank's balanced salt solution (HBSS) and transferred to fresh medium containing H₂O₂. Total RNA was isolated from HUVECs using ISOGEN (Nippon Gene, Toyama, Japan). mRNA expression was determined by real-time reverse-transcription polymerase chain reaction (real-time PCR), as previously described.⁽⁴¹⁾ The primers used in this study are indicated in Table 1.

The relative gene expression levels in each sample compared to the control were calculated using the comparative Ct method (ΔΔCT). The gene expression levels of target genes in each sample were normalized to that of GAPDH ΔΔCT value.

Western blotting. HUVECs were seeded on 60 mm dishes (5 × 10⁵ cells/dish) and incubated for 24 h. These cells were treated with Q-related compounds. The treated cells were washed with HBSS and fresh medium containing H₂O₂ added. After incubation for 24 h, the cells were washed twice with HBSS and lysed with lysis buffer [50 mM Tris-HCl (pH 8.0), 200 mM NaCl, 20 mM EDTA, 1% sodium dodecyl sulfate polyacrylamide (SDS), 0.5% sodium deoxycholate, 0.01% Nonidet P-40, phosphatase inhibitor tablet (Phos STOP), and protease inhibitor tablet (complete EDTA-free)]. Protein concentrations were determined using the bicinchoninic acid

(BCA) protein assay (Pierce; Thermo Scientific, Waltham, MA). The protein samples were boiled with reducing buffer (Nacalai Tesque) for 5 min. The samples were separated by 15% SDS gel electrophoresis (SDS-PAGE). Proteins were transferred onto Immobilon P PVDF transfer membranes (Millipore, Bedford, MA), followed by blocking of non-specific binding with a commercial blocking buffer (Blocking One for GAPDH, Cav-1 and Fyn or Blocking One P for phospho Cav-1 and phospho Fyn; Nacalai Tesque) for 1 h. After washing with Tris-buffered saline containing 0.05% Tween 20 (TBST), the membranes were incubated with anti-Cav-1 antibody, anti-phospho-Cav-1 antibody, anti-Fyn antibody, anti-phospho-Fyn antibody, or anti-GAPDH antibody for 1 h at room temperature. After washing three times with TBST, the membranes were incubated for 1 h at room temperature with a secondary antibody. After washing with TBST, the membranes were visualized using enhanced chemiluminescence (ECL) prime detection reagents (GE Healthcare, Buckinghamshire, UK). Images were captured using Lumino Image Analyzer LAS-3000 mini (Fujifilm, Tokyo, Japan) and analyzed using ImageJ software (National Institutes of Health, Bethesda, MD). Phosphorylated proteins were normalized to the total amount of each protein.

Vascular permeability assays. Vascular permeability was assessed using CultreCoat 24 well *in vitro* vascular permeability assay kit (3475-024-K; R&D systems, Minneapolis, MN). Fresh medium was added to the upper chamber and maintained at 37°C in a humidified atmosphere containing 5% CO₂. HUVECs were seeded in the upper chamber (2 × 10⁶ cells/dish) and 500 μl of medium was added to the bottom chamber, then incubated for 72 h. Q-related compounds were then added to the cells in the upper chamber to determine their effects. After 24 h, the medium in the upper chamber was aspirated and the cells were washed with HBSS thrice. Then, the cells in the upper chamber were incubated in medium containing 2 mM H₂O₂ for 0.5 h. The medium in the upper chamber was aspirated and the cells were washed with HBSS thrice. The upper chamber was transferred to another 24 well plate and the bottom chamber was filled with 300 μl of fresh medium. Fluorescein isothiocyanate (FITC)-dextran (3475-096-02) (R&D systems) was added to the treated cells in the upper chamber and maintained for 5 min. The medium in the bottom chamber was transferred to a 96-well plate and fluorescence (485 nm excitation with 520 nm emission) was measured.

Statistical analyses. Data from at least three independent experiments are expressed as mean ± SD. Statistical analyses were performed using PASW statistics 18.0 (SPSS Inc., Chicago, IL). *P* value <0.05 were considered statistically significant.

Results

H₂O₂ induced Cav-1 phosphorylation. Exposure of HUVECs to 2 mM H₂O₂ for 30 min significantly enhanced Cav-1 phosphorylation, although exposure to the same for 15 min did not show significant enhancement of Cav-1 phosphorylation (Fig. 1A and B). Moreover, 2 mM, but not 0.5 mM H₂O₂, showed significant phosphorylation of Cav-1 (Fig. 1C and D).

Table 1. Primers used for real-time RT-PCR

Gene	Forward	Reverse
GAPDH	5'-GCACCGTCAAGGCTGAGAAC-3'	5'-TGGTGAAGACGCCAGTGGA-3'
occludin	5'-ACAGAGCAAGATCACTATGAGACA-3'	5'-TGTTGATCTGAAGTGATAGGTGGA-3'
JAM	5'-AAGACACTGGGACATACACTTGT-3'	5'-CGATGAGCTTGACCTTGACCT-3'
ZO-1	5'-TGGTGTCTCACTAATCAACTCA-3'	5'-CGCCAGCTACAAATATCCAACA-3'
VE-cadherin	5'-CAACTGGCCTGTGTTACGC-3'	5'-ATCCACTGCTGCACAGAGATGA-3'




Fig. 1. Enhancement of Cav-1 phosphorylation by H₂O₂ in HUVECs. HUVECs were treated with H₂O₂ (at 0.5 or 2 mM) for 15 or 30 min. Phosphorylated Cav-1, total Cav-1, and GAPDH were determined by Western blotting. (A, C) Typical images of Western blotting. (B, D) densities for phosphorylated Cav-1, which were normalized by total Cav-1 obtained from (A), and (C), respectively. Data are means \pm SD ($n = 3$). Different letters represent significant differences ($p < 0.05$) determined by ANOVA with the Tukey multiple comparison test.

Suppressive effect of Q and its glucuronide on phosphorylation of Cav-1. Figure 2 shows the effects of Q and Q3GA on Cav-1 phosphorylation, when HUVECs were treated with these compounds for 0.5 h (Fig. 2A and B) or 24 h (Fig. 2C and D) before the addition of 2 mM H₂O₂. Q suppressed the phosphorylation of Cav-1 when HUVECs were treated with it for 24 h before exposure to H₂O₂. Q3GA showed no suppression of Cav-1 phosphorylation in the treatment, neither at 0.5 h nor 24 h. On the contrary, Fyn phosphorylation was unaffected by both Q and Q3GA treatments (Fig. 2C and E). As shown in Fig. 2F, neither Q nor Q3GA had any effect on the phosphorylation of Cav-1 without exposure to H₂O₂.

Comparison of suppressive effects of Q analogs on H₂O₂-induced Cav-1 phosphorylation. Notably, 8PQ exerted a suppressive effect similar to that of Q, although its suppression was not significantly different. None of the other Q analogues suppressed the phosphorylation of Cav-1 (Fig. 3).

Effect of Q on H₂O₂-induced vascular permeability of HUVECs. To estimate the vascular permeability of HUVECs, they were seeded on a transwell membrane and exposed to H₂O₂ for 0.5 h after pretreatment with Q for 24 h. Then, the rate of permeation of FITC-dextran in HUVECs was measured by fluorescent assay (Fig. 4). Exposure to H₂O₂ increased vascular permeability significantly, while pretreatment with Q apparently attenuated the H₂O₂-induced increase in vascular permeability.

Effect of Q on gene expression related to intercellular junction in HUVECs. Figure 5 shows the effect of H₂O₂ exposure for different periods (0.5, 1, and 3 h) on the expression

of genes related to tight junctions and adherens junctions in HUVECs.

The expression of VE-cadherin decreased after 1 and 3 h of exposure to H₂O₂. In contrast, H₂O₂ did not affect the expression of occludin, junctional adhesion molecule (JAM), and zonula occludens-1 (ZO-1) after exposure for 0.5, 1, and 3 h. Q showed higher gene expression of VE-cadherin (Fig. 6).

Discussion

In this study, we investigated the effect of Q on H₂O₂-induced Cav-1 phosphorylation and vascular permeability in endothelial cells. Pre-treatment with Q for 24 h suppressed H₂O₂-induced Cav-1 phosphorylation and attenuated the enhancement of vascular hyper-permeability (Fig. 2). However, neither the conjugated metabolite of Q, Q3GA, nor the Q analogues L, M, and F suppressed Cav-1 phosphorylation in endothelial cells (Fig. 3). Though there have been few reports about the beneficial effect of flavonoid on phosphorylation of Cav-1, it is demonstrated Q suppressed phosphorylation of Cav-1 induced by H₂O₂ in this study. This is a novel mechanism of Q on the prevention effect of atherosclerosis.

Fyn, an Src family kinase, plays an essential role in Cav-1 phosphorylation as response to oxidative and hyperosmotic stress.⁽⁴²⁾ In this study using H₂O₂-exposed HUVECs, we confirmed that oxidative stress can induce phosphorylation of both Cav-1 and Fyn (Fig. 1 and 2). H₂O₂, a reactive oxygen species, is frequently used at high concentrations (e.g., 2 mM) to




Fig. 2. Effect of Q and Q3GA on Cav-1 phosphorylation induced by H₂O₂. HUVECs were treated with Q or Q3GA at 10 μM for 0.5 h (A) or 24 h (C) prior to the treatment with 2 mM H₂O₂ for 30 min. Phosphorylated Cav-1, total Cav-1, phosphorylated Fyn, total Fyn, and GAPDH were determined by Western blotting. (A, C) Typical images of Western blotting. (B, D, E) Densities for phosphorylated protein, which were normalized to each total protein obtained from (A), and (C), respectively. The relative density was calculated as % of the band density obtained from H₂O₂ alone. HUVECs were treated with Q or Q3GA at 1 or 10 μM for 24 h (F, G). Phosphorylated Cav-1, total Cav-1, and GAPDH were determined by Western blotting. (F) Typical images of Western blotting. (G) Densities for phosphorylated Cav-1, which were normalized by total Cav-1 obtained from (F). Data are means ± SD (n = 3). Asterisks represent significant differences (p < 0.05) between H₂O₂ alone and each treatment as determined by two-sided Student's *t* test.




Fig. 3. Effect of Q analogues on Cav-1 phosphorylation induced by H₂O₂. HUVECs were treated with Q analogues at 10 μ M for 24 h prior to treatment with 2 mM H₂O₂ for 30 min. Phosphorylated Cav-1, total Cav-1, and GAPDH were determined by Western blotting. (A) Typical images of Western blotting. (B) Densities for phosphorylated protein, which were normalized to each total protein obtained from (A). The relative density was calculated as % of the band density obtained from H₂O₂ alone. Data are means \pm SD ($n = 3$). Asterisk represents significant differences ($p < 0.05$) between H₂O₂ alone and each treatment, as determined by two-sided Student's t test.

induce oxidative stress in cultured cells, including HUVECs.⁽⁴³⁾ According to the study of Shin *et al.*⁽⁴⁴⁾ the addition of 1 μ M sodium orthovanadate to the cells inhibited cellular phosphatase activity. Therefore, the cellular system used in this study can reflect the oxidative stress in vascular endothelial cells *in vivo*.

We demonstrated that pretreatment of HUVECs with Q for 24 h exerted inhibitory effect on Cav-1 phosphorylation in these cells (Fig. 2). The inhibitory effect of Q on Cav-1 phosphorylation can be explained by both its direct antioxidant activity and indirect activity responsible for the cellular signal transduction pathway. In contrast, Q did not show any effect on Fyn phosphorylation (Fig. 2). Fyn is phosphorylated in the cytoplasm,⁽⁴⁵⁾ while Cav-1 phosphorylation occurs at the caveolae in the plasma membrane of vascular endothelial cells.⁽⁴⁶⁾ It is therefore likely that Q is capable of acting at caveolae in the plasma membrane, but not in the cytoplasm. Although few studies have demonstrated the cellular localization of Q in HUVECs, Q hardly existed in cultured cells after 24 h of treatment.^(47,48) Furthermore, direct antioxidant activity is unlikely to be responsible for cardiovascular health in humans.⁽⁴⁹⁾ Therefore, we suppose that Q may affect intracellular signaling pathways to exert its inhibitory effect. Piechota-Polanczyk *et al.*⁽⁵⁰⁾ showed that heme oxygenase-1 (HO-1), an intracellular antioxidant enzyme, is involved in the inhibition of Cav-1 phosphorylation in a mouse model. Q is known to increase HO-1 gene expression via phosphorylation of ERK, which inhibits H₂O₂-induced apoptosis.⁽⁵¹⁾ Therefore, Q may act as an inhibitor of Cav-1 phosphorylation through




Fig. 4. Suppressive effect of Q on vascular permeability. HUVECs seeded on the transwell were treated with 10 μ M Q for 24 h prior to treatment with 2 mM H₂O₂ for 30 min. Vascular permeability was estimated by the measurement of fluorescent FITC-Dextran that passes through HUVEC cell. The index was calculated based on the ratio between control and independent experiments. Data are means \pm SD ($n = 3$). Different letters represent significant differences ($p < 0.05$) determined by ANOVA with the Tukey multiple comparison test.

the induction of HO-1 expression.⁽⁵⁰⁾ Nuclear accumulation of Nrf2, a transcription factor for the gene expression of HO-1, and downregulation of the Nrf2 repressor, Keap1, reportedly participate in Q-dependent HO-1 induction.⁽⁵²⁾ Therefore, it can be hypothesized that HO-1 induction is involved in the inhibition of Cav-1 phosphorylation when HUVECs are pretreated with Q for 24 h. In HUVECs, AMPK stabilizes Prdx1 and c-Abl complexes, thus acting as a negative regulator of Cav-1 phosphorylation.⁽⁴³⁾

Cav-1 knockdown markedly reduced the localization of VE-cadherin and β -catenin at the endothelial cell binding sites.⁽⁵³⁾ Exposure to H₂O₂ causes dissociation of β -catenin from Cav-1 and VE-cadherin.⁽²³⁾ Our results revealed that H₂O₂ exacerbated vascular permeability, concomitant with both a decrease in VE-cadherin mRNA expression and Cav-1 phosphorylation. These phenomena indicate that vascular permeability is increased by the deficiency of cell-cell adhesion proteins under oxidative stress. Thus, Q appears to contribute to the maintenance of vascular function by suppressing the attenuation of VE-cadherin gene expression through inhibition of oxidative stress dependent-Cav-1 phosphorylation.

The results of the structure-activity relationship study indicated that the Q analogues tested in this study did not exert inhibitory effects on Cav-1 phosphorylation (Fig. 3). The inherent structure of Q, a flavonol structure with an o-dihydroxy group at the B-ring, may be necessary for exerting this activity. Interestingly, the present study did not show any inhibitory effects of Q3GA (Fig. 2). Q is present as conjugated glucuronide and/or sulfate metabolites in blood.⁽²⁹⁾ It is therefore implied that deconjugation and conversion into Q aglycone by β -glucuronidase at the site of vascular inflammation is required for Q glucuronide conjugates to exert their vascular function through modulation of Cav-1 phosphorylation *in vivo*.

Q mainly exists as glycosylated form such as its glucoside or rutinoside in natural sources.⁽⁵⁴⁾ Recent studies using rodents demonstrated that these glycosides exerted beneficial effect on




Fig. 5. Effect of H_2O_2 on vascular permeability related gene expression. HUVECs were treated with 2 mM H_2O_2 for 0.5, 1, or 3 h. Gene expression was normalized to that of GAPDH expression. Data were calculated as the ratio between control (without H_2O_2 treatment) and treated cells. Data are means \pm SD ($n = 3$). Different letters represent significant differences ($p < 0.05$) determined by ANOVA with the Tukey multiple comparison test.

hyperglycemia and blood stasis syndrome,^(55,56) when they were orally administered. These glycosides are hydrolyzed in the intestinal tract and resulting aglycone is transferred into the blood stream.⁽⁵⁷⁾ Therefore, Q glycosides from dietary origin seem to possess a potential to exert preventive effect on exacerbation of vascular permeability caused by oxidative stress.

In conclusion, we confirmed that H_2O_2 causes phosphorylation of Cav-1, reduces the mRNA expression of the plasma membrane-bound protein VE-cadherin, and increases vascular permeability in HUVECs. Q aglycone suppresses both H_2O_2 -dependent Cav-1 phosphorylation and increases vascular permeability. Q may protect the vascular endothelium by suppressing the phosphorylation of Cav-1 in the endothelial cells under oxidative stress. Q increased VE-cadherin expression under H_2O_2 exposure (Fig. 6C). Further study is needed for clarify whether Q protect VE-cadherin down-regulation by H_2O_2 through suppression of Cav-1 phosphorylation or not. Q glucuronide metabolite did not show such an inhibitory effect, indicating that the release of aglycone from a plasma-conjugated metabolite is necessary for dietary Q to exert vascular function.

Author Contributions

Study concept and design (JT, TS, and RM); acquisition of data and statistical analysis (AKK and RM); analysis and interpretation of data, drafting of the manuscript, obtained funding (AKK, JT, and RM); critical version of the manuscript for important intellectual content (JT and RM); obtained funding (AKK, JT, and RM); and study supervision (JT and TS).

Acknowledgments

This work was supported by the Tojuro Iijima Foundation for Food Science and Technology, and was partly supported by JSPS KAKENHI Grant Numbers 25292075 and 17K13213.

Abbreviations

BCA	bicinchoninic acid
Cav-1	caveolin-1
ECL	enhanced chemiluminescence
F	flavone
FITC	fluorescein isothiocyanate




Fig. 6. Effect of Q on vascular permeability-related gene expression in the presence of H₂O₂. HUVECs were treated with 10 μM Q for 24 h prior to treatment with 2 mM H₂O₂ for 0.5, 1, or 3 h. Gene expression was normalized to that of GAPDH expression. Data were calculated as the ratio to control (without H₂O₂ treatment). Data are means ± SD (n = 3). Asterisk represents significant differences (p < 0.05) between control (H₂O₂ alone) and Q by two-sided Student's t test.

GAPDH glyceraldehyde-3-phosphate dehydrogenase
H₂O₂ hydrogen peroxide
HUVEC human umbilical vein endothelial cells
JAM junctional adhesion molecule
L luteolin
M myricetin
ox-LDL oxidized low-density proteins
8PQ 8-prenyl quercetin
Q quercetin

Q3GA quercetin 3-O-glucuronide
RT-PCR reverse-transcription polymerase chain reaction
SDS sodium dodecyl sulfate polyacrylamide
VE-cadherin vascular endothelial cadherin
ZO-1 zonula occludens-1

Conflict of Interest

No potential conflicts of interest were disclosed.

References

- Roth GA, Johnson C, Abajobir A, *et al.* Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. *J Am Coll Cardiol* 2017; **70**: 1–25.
- Beaglehole R, Yach D. Globalisation and the prevention and control of non-communicable disease: the neglected chronic diseases of adults. *Lancet* 2003; **362**: 903–908.
- Mozaffarian D. Global scourge of cardiovascular disease: time for health care systems reform and precision population health. *J Am Coll Cardiol* 2017; **70**: 26–28.
- Madamanchi NR, Vendrov A, Runge MS. Oxidative stress and vascular disease. *Arterioscler Thromb Vasc Biol* 2005; **25**: 29–38.
- Stone NJ, Robinson JG, Lichtenstein AH, *et al.* 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014; **63** (25 Pt B): 2889–2934.
- Torres N, Guevara-Cruz M, Velázquez-Villegas LA, Tovar AR. Nutrition and Atherosclerosis. *Arch Med Res* 2015; **46**: 408–426.

- 7 Ruiz-León AM, Lapuente M, Estruch R, Casas R. Clinical advances in immunonutrition and atherosclerosis: a review. *Front Immunol* 2019; **10**: 837.
- 8 Lum H, Malik AB. Regulation of vascular endothelial barrier function. *Am J Physiol* 1994; **267** (3 Pt 1): L223–L241.
- 9 Bazzoni G, Dejana E. Endothelial cell-to-cell junctions: molecular organization and role in vascular homeostasis. *Physiol Rev* 2004; **84**: 869–901.
- 10 Mehta D, Malik AB. Signaling mechanisms regulating endothelial permeability. *Physiol Rev* 2006; **86**: 279–367.
- 11 Murata T, Lin MI, Aritake K, et al. Role of prostaglandin D2 receptor DP as a suppressor of tumor hyperpermeability and angiogenesis *in vivo*. *Proc Natl Acad Sci U S A* 2008; **105**: 20009–20014.
- 12 Lisanti MP, Scherer PE, Tang Z, Sargiacomo M. Caveolae, caveolin and caveolin-rich membrane domains: a signalling hypothesis. *Trends Cell Biol* 1994; **4**: 231–235.
- 13 Razani B, Lisanti M. The role of caveolae and the caveolins in mammalian physiology *Rice Undergraduate Science Review* 2002; **1**: 44–45.
- 14 Layne J, Majkova Z, Smart EJ, Toborek M, Hennig B. Caveolae: a regulatory platform for nutritional modulation of inflammatory diseases. *J Nutr Biochem* 2011; **22**: 807–811.
- 15 Schnitzer JE, Oh P, McIntosh DP. Role of GTP hydrolysis in fission of caveolae directly from plasma membranes. *Science* 1996; **274**: 239–242.
- 16 Liu P, Rudick M, Anderson RG. Multiple functions of caveolin-1. *J Biol Chem* 2002; **277**: 41295–41298.
- 17 Pavlides S, Gutierrez-Pajares JL, Iturrieta J, Lisanti MP, Frank PG. Endothelial caveolin-1 plays a major role in the development of atherosclerosis. *Cell Tissue Res* 2014; **356**: 147–157.
- 18 Fernández-Hernando C, Yu J, Dávalos A, Prendergast J, Sessa WC. Endothelial-specific overexpression of caveolin-1 accelerates atherosclerosis in apolipoprotein E-deficient mice. *Am J Pathol* 2010; **177**: 998–1003.
- 19 Hu G, Vogel SM, Schwartz DE, Malik AB, Minshall RD. Intercellular adhesion molecule-1-dependent neutrophil adhesion to endothelial cells induces caveolae-mediated pulmonary vascular hyperpermeability. *Circ Res* 2008; **102**: e120–e131.
- 20 Hu G, Minshall RD. Regulation of transendothelial permeability by Src Kinase. *Microvasc Res* 2009; **77**: 21–25.
- 21 Zhang Y, Zhang L, Li Y, Sun S, Tan H. Different contributions of clathrin- and caveolae-mediated endocytosis of vascular endothelial cadherin to lipopolysaccharide-induced vascular hyperpermeability. *PLoS One* 2014; **9**: e106328.
- 22 Lin F, Pei L, Zhang Q, et al. Ox-LDL induces endothelial cell apoptosis and macrophage migration by regulating caveolin-1 phosphorylation. *J Cell Physiol* 2018; **233**: 6683–6692.
- 23 Sun Y, Hu G, Zhang X, Minshall RD. Phosphorylation of caveolin-1 regulates oxidant-induced pulmonary vascular permeability via paracellular and transcellular pathways. *Circ Res* 2009; **105**: 676–685.
- 24 Míean KH, Mohamed S. Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem* 2001; **49**: 3106–3112.
- 25 Aherne SA, O'Brien NM. Dietary flavonols: chemistry, food content, and metabolism. *Nutrition* 2002; **18**: 75–81.
- 26 Landberg R, Naidoo N, van Dam RM. Diet and endothelial function: from individual components to dietary patterns. *Curr Opin Lipidol* 2012; **23**: 147–155.
- 27 Hertog MG, Hollman PC, Katan MB, Kromhout D. Intake of potentially anticarcinogenic flavonoids and their determinants in adults in The Netherlands. *Nutr Cancer* 1993; **20**: 21–29.
- 28 Hertog MG, Kromhout D, Aravanis C, et al. Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. *Arch Intern Med* 1995; **155**: 381–386.
- 29 Terao J. Factors modulating bioavailability of quercetin-related flavonoids and the consequences of their vascular function. *Biochem Pharmacol* 2017; **139**: 15–23.
- 30 Salvamani S, Gunasekaran B, Shaharuddin NA, Ahmad SA, Shukor MY. Antiatherosclerotic effects of plant flavonoids. *Biomed Res Int* 2014; **2014**: 480258.
- 31 Deng Q, Li XX, Fang Y, Chen X, Xue J. Therapeutic potential of quercetin as an antiatherosclerotic agent in atherosclerotic cardiovascular disease: a review. *Evid Based Complement Alternat Med* 2020; **2020**: 5926381.
- 32 Starke H, Herrmann K. Flavonols and flavones of vegetables. VI. On the changes of the flavonols of onions (authors transl). *Z Lebensm Unters Forsch* 1976; **161**: 137–142. (in Deutsch)
- 33 Day AJ, Cañada FJ, Díaz JC, et al. Dietary flavonoid and isoflavone glycosides are hydrolysed by the lactase site of lactase phlorizin hydrolase. *FEBS Lett* 2000; **468**: 166–170.
- 34 Kawai Y, Nishikawa T, Shiba Y, et al. Macrophage as a target of quercetin glucuronides in human atherosclerotic arteries: implication in the anti-atherosclerotic mechanism of dietary flavonoids. *J Biol Chem* 2008; **283**: 9424–9434.
- 35 Ishisaka A, Kawabata K, Miki S, et al. Mitochondrial dysfunction leads to deconjugation of quercetin glucuronides in inflammatory macrophages. *PLoS One* 2013; **8**: e80843.
- 36 Menendez C, Dueñas M, Galindo P, et al. Vascular deconjugation of quercetin glucuronide: the flavonoid paradox revealed? *Mol Nutr Food Res* 2011; **55**: 1780–1790.
- 37 Galindo P, Rodríguez-Gómez I, González-Manzano S, et al. Glucuronidated quercetin lowers blood pressure in spontaneously hypertensive rats via deconjugation. *PLoS One* 2012; **7**: e32673.
- 38 Perez A, Gonzalez-Manzano S, Jimenez R, et al. The flavonoid quercetin induces acute vasodilator effects in healthy volunteers: correlation with beta-glucuronidase activity. *Pharmacol Res* 2014; **89**: 11–18.
- 39 Kamada C, Mukai R, Kondo A, Sato S, Terao J. Effect of quercetin and its metabolite on caveolin-1 expression induced by oxidized LDL and lysophosphatidylcholine in endothelial cells. *J Clin Biochem Nutr* 2016; **58**: 193–201.
- 40 Kawamura T, Hayashi M, Mukai R, Terao J, Nemoto H. An efficient method for C8-prenylation of flavonols and flavanones. *Synthesis* 2012; **44**: 1308–1314.
- 41 Mukai R, Nakao R, Yamamoto H, Nikawa T, Takeda E, Terao J. Quercetin prevents unloading-derived disused muscle atrophy by attenuating the induction of ubiquitin ligases in tail-suspension mice. *J Nat Prod* 2010; **73**: 1708–1710.
- 42 Sanguinetti AR, Cao H, Corley Mastick C. Fyn is required for oxidative- and hyperosmotic-stress-induced tyrosine phosphorylation of caveolin-1. *Biochem J* 2003; **376** (Pt 1): 159–168.
- 43 Takeuchi K, Morizane Y, Kamami-Levy C, et al. AMP-dependent kinase inhibits oxidative stress-induced caveolin-1 phosphorylation and endocytosis by suppressing the dissociation between c-Abl and Prdx1 proteins in endothelial cells. *J Biol Chem* 2013; **288**: 20581–20591.
- 44 Shin J, Jo H, Park H. Caveolin-1 is transiently dephosphorylated by shear stress-activated protein tyrosine phosphatase mu. *Biochem Biophys Res Commun* 2006; **339**: 737–741.
- 45 Resh MD. Fyn, a Src family tyrosine kinase. *Int J Biochem Cell Biol* 1998; **30**: 1159–1162.
- 46 Lee H, Volonte D, Galbiati F, et al. Constitutive and growth factor-regulated phosphorylation of caveolin-1 occurs at the same site (Tyr-14) *in vivo*: identification of a c-Src/Cav-1/Grb7 signaling cassette. *Mol Endocrinol* 2000; **14**: 1750–1775.
- 47 Mukai R, Fujikura Y, Murota K, et al. Prenylation enhances quercetin uptake and reduces efflux in Caco-2 cells and enhances tissue accumulation in mice fed long-term. *J Nutr* 2013; **143**: 1558–1564.
- 48 Mukai R, Kawabata K, Otsuka S, et al. Effect of quercetin and its glucuronide metabolite upon 6-hydroxydopamine-induced oxidative damage in Neuro-2a cells. *Free Radic Res* 2012; **46**: 1019–1028.
- 49 Hollman PCH, Cassidy A, Comte B, et al. The biological relevance of direct antioxidant effects of polyphenols for cardiovascular health in humans is not established. *J Nutr* 2011; **141**: 989S–1009S.
- 50 Piechota-Polanczyk A, Zielińska M, Piekliński D, Fichna J. The influence of lipoic acid on caveolin-1-regulated antioxidant enzymes in the mouse model of acute ulcerative colitis. *Biomed Pharmacother* 2016; **84**: 470–475.
- 51 Chow JM, Shen SC, Huan SK, Lin HY, Chen YC. Quercetin, but not rutin and quercitrin, prevention of H₂O₂-induced apoptosis via anti-oxidant activity and heme oxygenase 1 gene expression in macrophages. *Biochem Pharmacol* 2005; **69**: 1839–1851.
- 52 Luo M, Tian R, Yang Z, Peng YY, Lu N. Quercetin suppressed NADPH oxidase-derived oxidative stress via heme oxygenase-1 induction in macrophages. *Arch Biochem Biophys* 2019; **671**: 69–76.
- 53 Song L, Ge S, Pachter JS. Caveolin-1 regulates expression of junction-associated proteins in brain microvascular endothelial cells. *Blood* 2007; **109**: 1515–1523.

- 54 Tsushida T, Suzuki M. Isolation of flavonoid-glycosides in onion and identification by chemical synthesis of the glycosides flavonoid in fruits and vegetables Part I. *Nippon Shokuhin Kagaku Kogaku Kaishi* 1995; **42**: 100–108. (in Japanese)
- 55 Oku H, Maeda M, Kitagawa F, Ishiguro K. Effect of polyphenols from *syringa vulgaris* on blood stasis syndrome. *J Clin Biochem Nutr* 2020; **67**: 84–88.
- 56 Jiang H, Horiuchi Y, Hironao KY, Kitakaze T, Yamashita Y, Ashida H. Prevention effect of quercetin and its glycosides on obesity and hyperglycemia through activating AMPK α in high-fat diet-fed ICR mice. *J Clin Biochem Nutr* 2020; **67**: 74–83.
- 57 Murota K, Terao J. Antioxidative flavonoid quercetin: implication of its intestinal absorption and metabolism. *Arch Biochem Biophys* 2003; **417**: 12–17.



This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
