



# Genome Sequences and Methylation Patterns of *Natrinema versiforme* BOL5-4 and *Natrinema pallidum* BOL6-1, Two Extremely Halophilic Archaea from a Bolivian Salt Mine

Priya DasSarma,<sup>a</sup> Brian P. Anton,<sup>b</sup> Satyajit L. DasSarma,<sup>a</sup> Fabiana L. Martinez,<sup>a,c</sup> Daniel Guzman,<sup>d</sup> Richard J. Roberts,<sup>b</sup> Shiladitya DasSarma<sup>a</sup>

<sup>a</sup>Institute of Marine and Environmental Technology, Department of Microbiology and Immunology, University of Maryland, Baltimore, Maryland, USA

<sup>b</sup>New England Biolabs, Ipswich, Massachusetts, USA

<sup>c</sup>Instituto de Investigaciones para la Industria Química, Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad Nacional de Salta, Salta, Argentina

<sup>d</sup>Centro de Biotecnología, Faculty of Sciences and Technology, Universidad Mayor de San Simón, Cochabamba, Bolivia

**ABSTRACT** Two extremely halophilic archaea, namely, *Natrinema versiforme* BOL5-4 and *Natrinema pallidum* BOL6-1, were isolated from a Bolivian salt mine and their genomes sequenced using single-molecule real-time sequencing. The GC-rich genomes of BOL5-4 and BOL6-1 were 4.6 and 3.8 Mbp, respectively, with large chromosomes and multiple megaplasms. Genome annotation was incorporated into HaloWeb and methylation patterns incorporated into REBASE.

Halophilic microbes capable of surviving extreme conditions are of interest for biotechnology and astrobiology (1, 2). Two extremely halophilic archaea, members of the *Natrinema* genus, were isolated from pink salt obtained from a salt mine in the Department of Tarija, O'Connor Province, Bolivia. Together with the novel *Halorubrum* sp. strain BOL3-1 from Salar de Uyuni, our collection of halophilic archaea from Bolivia has expanded from one to three species (3). These strains also provide the basis for a wider comparative genomic analysis of haloarchaea from the subsurface to high elevation.

Pink salt was sampled at a remote salt mine (21°24'19.73"S, 64°07'51.52"W) at 1,230-meter elevation, where temperatures range from –10°C to 37°C. The salt samples were dissolved in CM<sup>+</sup> medium, and growth was stimulated with shaking at 220 rpm at 37°C under illumination (4). The enrichment cultures were plated on CM<sup>+</sup> agar plates and two isolates, namely, *Natrinema versiforme* BOL5-4, a nearly unpigmented strain, and *Natrinema pallidum* BOL6-1, a pigmented strain, were purified by 3 rounds of streaking.

Nucleic acids were extracted using a standard method (5), and sequencing was performed using the Sequel platform (Pacific Biosciences, Menlo Park, CA). SMRTbell libraries were prepared from unsharded genomic DNA (2 μg BOL5-4 and 0.9 μg BOL6-1), and each library was sequenced on 1 single-molecule real-time (SMRT) cell with a Sequel binding kit 3.0, with 10-h collection and 2-h preextension times (6). Sequencing subreads were filtered and assembled *de novo* using Hierarchical Genome Assembly Process (HGAP) version 4, with default parameters. There were 553,417 filtered subreads (mean length, 4,931 bp; coverage, 780×) for BOL5-4 and 551,878 filtered subreads (mean length, 4,717 bp; coverage, 520×) for BOL6-1.

The *N. versiforme* BOL5-4 and *N. pallidum* BOL6-1 genome sequences were annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) build 3190 (7), which was incorporated into HaloWeb (version r1559404112; <https://halo.umbc.edu/>) and analyzed further using EMBOSS version 6.6.0.0 (<http://www.bioinformatics.nl/cgi-bin/emboss/>) (8). Default parameters were used for all software. The BOL5-4

**Citation** DasSarma P, Anton BP, DasSarma SL, Martinez FL, Guzman D, Roberts RJ, DasSarma S. 2019. Genome sequences and methylation patterns of *Natrinema versiforme* BOL5-4 and *Natrinema pallidum* BOL6-1, two extremely halophilic archaea from a Bolivian salt mine. *Microbiol Resour Announc* 8:e00810-19. <https://doi.org/10.1128/MRA.00810-19>.

**Editor** Frank J. Stewart, Georgia Institute of Technology

**Copyright** © 2019 DasSarma et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Shiladitya DasSarma, [sdassarma@som.umaryland.edu](mailto:sdassarma@som.umaryland.edu).

**Received** 5 July 2019

**Accepted** 26 July 2019

**Published** 15 August 2019

**TABLE 1** Motifs containing methylated bases <sup>m6</sup>A and <sup>m4</sup>C

Motif <sup>a</sup>	<i>N. versiforme</i> BOL5-4		<i>N. pallidum</i> BOL6-1	
	% modified	Gene <sup>b</sup>	% modified	Gene <sup>b</sup>
GACGAAC	100	FEJ81_15005		
CATTC	100	FEJ81_07280	99.9	FGF80_04050
CCWGG	99.3	FEJ81_16560		
GAACA <b>YC</b>	100	FEJ81_15230		
CTAG	97.0	FEJ81_09745	98.7	FGF80_01935
TCCTCGG	96.0	FEJ81_19855		
GCAAT	71.4	FEJ81_20490		
GTAYTCG			98.8	FGF80_00870
CAGYAAC			100	FGF80_10950

<sup>a</sup> Locations of methylated bases are in bold for the top strand and underlined for the bottom strand.

<sup>b</sup> Putative assignments of MTases responsible for the modification in the first column based on sequence comparison with known enzymes of that specificity in REBASE.

genome was 4,674,473 bp (G+C content, 63.4%) and included a 3,747,116-bp circular chromosome (G+C content, 64.7%) and 4 plasmids, namely, pNVE500 (492,102-bp linear contig; G+C content, 56.3%), pNVE414 (413,865-bp circle; G+C content, 60.0%), pNVE19 (18,925-bp circle; G+C content, 63.5%), and pNVE2 (2,465-bp circle; G+C content, 69.2%). The BOL6-1 genome was 3,778,093 bp (G+C content, 64.3%) and included a 3,503,953-bp circular chromosome (G+C content, 64.6%) and 2 plasmids, namely, pNPA200 (203,201-bp circle) and pNPA70 (70,939-bp circle), both with a G+C content of 60.6%.

The BOL5-4 genome contained 4,589 genes, including 3 rRNA operons and 64 tRNA genes, whereas the BOL6-1 genome contained 3,785 genes, including 3 rRNA operons and 50 tRNA genes. Both proteomes were highly acidic (9), with calculated mean pI values of 4.6 to 4.7, and all but 5 of nearly 800 core haloarchaeal orthologous groups were encoded in the genomes (10, 11). Both contained expanded gene families, e.g., Orc/Cdc6, TATA-binding, and transcription factor B (TFB) genes (12), as well as a gene cluster for gas vesicle nanoparticles (13) and polyhydroxyalkanoate synthesis genes (14). Both genomes encode many transposases, namely, a total of 100 in BOL5-4 and 80 in BOL6-1 (15).

Methylated DNA motifs and the methyltransferases (MTases) predicted to be responsible for some were deposited in REBASE (Table 1) (16). Both genomes contained the methylated motifs <sup>m4</sup>CTAG and C<sup>6m</sup>ATTC, which are common to halophilic archaea.

**Data availability.** The *N. versiforme* BOL5-4 genome sequence has been deposited in GenBank with the accession numbers [CP040329](#), [CP040330](#), [CP040331](#), [CP040332](#), and [CP040333](#). Raw data are available in the NCBI Sequence Read Archive with the accession number [SRX5888851](#). The *N. pallidum* BOL6-1 genome sequence has been deposited in GenBank with the accession numbers [CP040637](#), [CP040638](#), and [CP040639](#). Raw data are available in the NCBI Sequence Read Archive with the accession number [SRX6057204](#).

## ACKNOWLEDGMENTS

The DasSarma laboratory was supported by NASA grant NNH18ZDA001N. B.P.A. and R.J.R. work for New England BioLabs, a company that sells research reagents, including restriction enzymes and DNA methyltransferases, to the scientific community. F.L.M. was supported by the Fulbright Fellowship Program and D.G. by the Swedish International Development Cooperation Agency (ASDI).

## REFERENCES

- DasSarma S, DasSarma P. 2017. Halophiles, encyclopedia of life science. In eLS. John Wiley & Sons Ltd., Chichester, United Kingdom. <https://doi.org/10.1002/9780470015902.a0000394.pub4>.
- DasSarma S, Schwieterman EW. 2018. Early evolution of purple retinal pigments on Earth and implications for exoplanet biosignatures. *Int J Astrobiol* 1–10. <https://doi.org/10.1017/S1473550418000423>.
- DasSarma P, Anton BP, DasSarma S, Laye VJ, Guzman D, Roberts RJ, DasSarma S. 2019. Genome sequence and methylation patterns of *Halorubrum* sp. strain BOL3-1, the first haloarchaeon isolated and cultured from Salar de Uyuni, Bolivia. *Microbiol Resour Announc* 8:e00386-19. <https://doi.org/10.1128/MRA.00386-19>.
- Berquist BR, Müller JA, DasSarma S. 2006. Chapter 27. Genetic sys-

- tems for halophilic archaea, p 649–680. *In* Oren A, Rainey F (ed), *Methods in microbiology*, vol. 35. Elsevier Academic Press, San Diego, CA.
5. Ng WL, Yang CF, Halladay JT, Arora P, DasSarma S. 1995. Protocol 25. Isolation of genomic and plasmid DNAs from *Halobacterium halobium*, p 179–184. *In* DasSarma S, Fleischmann EM (ed), *Archaea, a laboratory manual: halophiles*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
  6. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
  7. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
  8. DasSarma SL, Capes MD, DasSarma P, DasSarma S. 2010. HaloWeb: the haloarchaeal genomes database. *Saline Systems* 6:12. <https://doi.org/10.1186/1746-1448-6-12>.
  9. DasSarma S, DasSarma P. 2015. Halophiles and their enzymes: negativity put to good use. *Curr Opin Microbiol* 25:120–126. <https://doi.org/10.1016/j.mib.2015.05.009>.
  10. Capes MD, DasSarma P, DasSarma S. 2012. The core and unique proteins of haloarchaea. *BMC Genomics* 13:39. <https://doi.org/10.1186/1471-2164-13-39>.
  11. Kennedy SP, Ng WV, Salzberg SL, Hood L, DasSarma S. 2001. Understanding the adaptation of *Halobacterium* species NRC-1 to its extreme environment through computational analysis of its genome sequence. *Genome Res* 11:1641–1650. <https://doi.org/10.1101/gr.190201>.
  12. Capes MD, Coker JA, Gessler R, Grinblat-Huse V, DasSarma SL, Jacob CG, Kim J-M, DasSarma P, DasSarma S. 2011. The information transfer system of halophilic archaea. *Plasmid* 65:77–101. <https://doi.org/10.1016/j.plasmid.2010.11.005>.
  13. DasSarma S, DasSarma P. 2015. Gas vesicle nanoparticles for antigen display. *Vaccines (Basel)* 3:686–702. <https://doi.org/10.3390/vaccines3030686>.
  14. Mahansaria R, Dhara A, Saha A, Haldar S, Mukherjee J. 2018. Production enhancement and characterization of the polyhydroxyalkanoate produced by *Natrinema ajinwuensis* (as synonym) ≡ *Natrinema altunense* strain RM-G10. *Int J Biol Macromol* 107:1480–1490. <https://doi.org/10.1016/j.ijbiomac.2017.10.009>.
  15. DasSarma S. 2004. Genome sequence of an extremely halophilic archaeon, p 383–399. *In* Fraser CM, Read T, Nelson KE (ed), *Microbial genomes*. Humana Press, Inc., Totowa, NJ.
  16. Roberts RJ, Vincze T, Posfai J, Macelis D. 2015. REBASE—a database for DNA restriction and modification: enzymes, genes and genomes. *Nucleic Acids Res* 43:D298–D299. <https://doi.org/10.1093/nar/gku1046>.