



# Complete Genome Sequences of Three of the Earliest Community-Associated Methicillin-Resistant *Staphylococcus aureus* Strains Isolated in Remote Western Australia

Nicola M. Karakatsanis,<sup>a,b</sup> Elena Colombi,<sup>a,b</sup>  Shakeel Mowlaboccus,<sup>c,d</sup> Julie C. Pearson,<sup>d</sup> Geoffrey W. Coombs,<sup>c,d</sup>  Joshua P. Ramsay<sup>a,b</sup>

<sup>a</sup>Curtin Health Innovation Research Institute, Curtin University, Bentley, Western Australia, Australia

<sup>b</sup>Curtin Medical School, Curtin University, Bentley, Western Australia, Australia

<sup>c</sup>College of Science, Health, Engineering, and Education, Murdoch University, Murdoch, Western Australia, Australia

<sup>d</sup>Department of Microbiology, PathWest Laboratory Medicine WA, Fiona Stanley Hospital, Murdoch, Western Australia, Australia

**ABSTRACT** Initially reported in Western Australia in the 1980s, community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA) has become a major cause of *S. aureus* infections globally. We report the complete genome sequences of three of the earliest CA-MRSA strains isolated from remote Australian Indigenous communities in the Kimberley region of Western Australia.

Methicillin-resistant *Staphylococcus aureus* (MRSA) is a leading cause of health care-associated infections worldwide and has emerged as a major cause of infections in the community (1). Community-associated MRSA (CA-MRSA) is genetically distinct and was first detected in Western Australia in 1984, in remote Indigenous communities (2, 3). Here, we present the complete genome sequences of three of the earliest isolated CA-MRSA strains, i.e., WBG7583 (sequence type 8 [ST8]), WBG8381 (ST5), and WBG8366 (ST78) (4, 5).

Previously isolated *S. aureus* strains (4) were rehydrated from lyophilized stocks in tryptic soy broth (TSB) and grown overnight on blood agar at 37°C. Strains were grown overnight in 5 ml TSB at 37°C, and cells were harvested by centrifugation. Cells were resuspended in 200  $\mu$ l Tris-EDTA (TE) buffer containing 1 mg/ml RNase (Sigma-Aldrich, USA) and 100  $\mu$ g/ml lysostaphin (Sigma-Aldrich) and were incubated at 37°C for 10 min. DNA for Oxford Nanopore Technologies (ONT) MinION sequencing was extracted using the FavorPrep blood/cultured cell genomic DNA extraction minikit (product number FABGK-001-2; Favorgen Biotech Corp., Taiwan). For Illumina NextSeq sequencing, DNA from a separate culture was extracted using the DNeasy blood and tissue kit (product number 69506; Qiagen, Germany). No DNA shearing or size selection was carried out before sequencing.

Genomes were sequenced using the MinION Mk1B system (ONT, UK). The ONT SQK-RAD004 kit was used to create DNA sequencing libraries, which were loaded onto an ONT FLO-MIN106 flow cell and sequenced using ONT MinKNOW v20.10.3 and MinKNOW Core v4.1.2. Base calling was performed with Guppy v4.4.1+1c81d62 (ONT), using the dna\_r9.4.1\_450bps\_hac.cfg model. NanoFilt (6) was used to filter reads for size (>4 kb) and average quality (scores of >Q10), and genomes were assembled using Flye v2.8.3-b1695 (7). Unfiltered MinION reads were mapped to each genome assembly with Minimap2 v2.17-r941 (8) and used to polish the assembly six times with Racon v1.4.15 ([github.com/lcb-science/racon](https://github.com/lcb-science/racon)). Each genome was also sequenced with the Illumina NextSeq 500 platform using the Nextera XT DNA library preparation kit (150-bp paired-end chemistry) and the NextSeq 500/550 kit v2.5 (300-cycle format; Illumina, USA). Neson clip ([github.com/Victorian-Bioinformatics-Consortium/neson](https://github.com/Victorian-Bioinformatics-Consortium/neson)) was used to remove adaptor sequences and to quality filter the reads. Illumina reads were mapped to the genome assembly with Minimap2 v2.17-r941 (8), and mapped reads were used to polish the assembly five times using Pilon v1.23 (9).

**Citation** Karakatsanis NM, Colombi E, Mowlaboccus S, Pearson JC, Coombs GW, Ramsay JP. 2021. Complete genome sequences of three of the earliest community-associated methicillin-resistant *Staphylococcus aureus* strains isolated in remote Western Australia. *Microbiol Resour Announc* 10:e00797-21. <https://doi.org/10.1128/MRA.00797-21>.

**Editor** Steven R. Gill, University of Rochester School of Medicine and Dentistry

**Copyright** © 2021 Karakatsanis et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Geoffrey W. Coombs, [g.coombs@murdoch.edu.au](mailto:g.coombs@murdoch.edu.au), or Joshua P. Ramsay, [josh.ramsay@curtin.edu.au](mailto:josh.ramsay@curtin.edu.au).

**Received** 5 August 2021

**Accepted** 19 August 2021

**Published** 16 September 2021

**TABLE 1** Genome and mapped-read statistics for WBG7583, WBG8381, and WBG8366

Strain and replicon type	Data for MinION sequencing					Data for Illumina sequencing					
	Genome size (bp)	GC content (%)	GenBank accession no.	Total no. of reads (mapped)	Read $N_{50}$ (bases)	Mean read depth (x)	SRA accession no.	Total no. of reads (mapped)	Mean read length (bp)	Mean read depth (x)	SRA accession no.
WBG7583 (ST8)				1,139,489	2,373		SRX11246052	3,279,211	147		SRX11246051
Chromosome	2,790,388	33	CP070989.1			593				100	
pWBG753	30,029	30	CP070990.1			2,061				232	
WBG8381 (ST5)				1,900,172	2,493		SRX11247744	2,400,046	147		SRX11247745
Chromosome	2,820,507	33	CP071046.1			1,005				62	
Circular prophage WBG8381	42,493	35	CP071049.1			2,226				151	
pWBG749	38,087	30	CP071047.1			2,312				193	
pWBG8381	2,539	31	CP071048.1			31,299				409	
WBG8366 (ST78)				189,095	3,057		SRX11246056	1,880,412	148		SRX11246055
Chromosome	2,768,386	33	CP070983.1			114				62	
pWBG763	20,730	28	CP070984.1			516				213	
pWBG764	2,393	28	CP070985.1			11,503				330	

Circlator v1.5.5 (10) was used to set the start position of each genome, and Qualimap v2.2.2-dev (11) was used to generate genome statistics (Table 1). Genome annotation was performed using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v5.1 (12). Default parameters were used for all software unless otherwise specified.

**Data availability.** GenBank accession numbers for the genome assemblies are provided in Table 1. The MinION and Illumina sequencing reads have been deposited under BioProject accession numbers [PRJNA703734](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA703734) and [PRJNA703736](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA703736) and Sequence Read Archive (SRA) accession numbers [SRX11246052](https://www.ncbi.nlm.nih.gov/sra/SRX11246052), [SRX11246051](https://www.ncbi.nlm.nih.gov/sra/SRX11246051), [SRX11247745](https://www.ncbi.nlm.nih.gov/sra/SRX11247745), [SRX11247744](https://www.ncbi.nlm.nih.gov/sra/SRX11247744), [SRX11246056](https://www.ncbi.nlm.nih.gov/sra/SRX11246056), and [SRX11246055](https://www.ncbi.nlm.nih.gov/sra/SRX11246055), as indicated in Table 1.

## ACKNOWLEDGMENT

N.M.K. thanks the Faculty of Health Science, Curtin University, for the Faculty of Health Science Summer Scholarship award.

## REFERENCES

1. International Working Group on the Classification of Staphylococcal Cassette Chromosome Elements. 2009. Classification of staphylococcal cassette chromosome *mec* (SCC*mec*): guidelines for reporting novel SCC*mec* elements. *Antimicrob Agents Chemother* 53:4961–4967. <https://doi.org/10.1128/AAC.00579-09>.
2. Riley TV, Rouse IL. 1995. Methicillin-resistant *Staphylococcus aureus* in Western Australia, 1983–1992. *J Hosp Infect* 29:177–188. [https://doi.org/10.1016/0195-6701\(95\)90327-5](https://doi.org/10.1016/0195-6701(95)90327-5).
3. Turnidge JD, Bell JM. 2000. Methicillin-resistant *Staphylococcal aureus* evolution in Australia over 35 years. *Microb Drug Resist* 6:223–229. <https://doi.org/10.1089/mdr.2000.6.223>.
4. O'Brien FG, Coombs GW, Pearman JW, Gracey M, Moss F, Christiansen KJ, Grubb WB. 2009. Population dynamics of methicillin-susceptible and -resistant *Staphylococcus aureus* in remote communities. *J Antimicrob Chemother* 64:684–693. <https://doi.org/10.1093/jac/dkp285>.
5. Udo EE, Pearman JW, Grubb WB. 1993. Genetic analysis of community isolates of methicillin-resistant *Staphylococcus aureus* in Western Australia. *J Hosp Infect* 25:97–108. [https://doi.org/10.1016/0195-6701\(93\)90100-e](https://doi.org/10.1016/0195-6701(93)90100-e).
6. De Coster W, D'Hert S, Schultz DT, Cruts M, Broeckhoven CV. 2018. NanoPack: visualising and processing long-read sequencing data. *Bioinformatics* 34:2666–2669. <https://doi.org/10.1093/bioinformatics/bty149>.
7. Kolmogorov M, Yuan J, Lin Y, Pevzner P. 2019. Assembly of long error-prone reads using repeat graphs. *Nat Biotechnol* 37:540–546. <https://doi.org/10.1038/s41587-019-0072-8>.
8. Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100. <https://doi.org/10.1093/bioinformatics/bty191>.
9. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
10. Hunt M, De Silva N, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
11. Okonechnikov K, Conesa A, Garcia-Alcalde F. 2016. Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* 32:292–294. <https://doi.org/10.1093/bioinformatics/btv566>.
12. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.