

Preplanned Studies

Mortality and Causes of Death in Primary and Secondary School Students — China, 2018

Wei Wang¹; Yunning Liu¹; Jiangmei Liu¹; Peng Yin¹; Jinlei Qi¹; Jinling You¹; Lijun Wang¹; Maigeng Zhou^{1,*}

Summary

What is already known about this topic?

There were approximately 1.23 million deaths reported among individuals aged 5–19 years worldwide in 2017. Limited attention has been paid to current mortality among primary and secondary school students aged 6–18 years in China.

What is added by this report?

In 2018, an estimated 28,519 deaths occurred among primary and secondary school students aged 6–18 years in China with an age-standardized mortality rate (ASMR) of 17.66 per 100,000. Substantial disparities existed among sexes, age groups, areas, and regions for different causes of death. Drowning, road traffic injuries, and leukemia were three leading causes of death.

What are the implications for public health practice?

Government support, legislation enforcement, multisectoral cooperation, and stakeholder engagement should be initiated to reduce premature deaths among primary and secondary school students in China, especially those due to injuries.

Premature death among primary and secondary school students aged 6–18 years is an important public health issue in China. In this study, we used data from China's Cause-of-Death Reporting System (CDRS) to estimate students' mortality. In 2018, it was estimated that there were 28,519 student deaths between the ages of 6–18 years, with age-standardized mortality rate (ASMR) of 17.66 per 100,000. Substantial disparities existed among different sexes, age groups, areas, and regions. Drowning, road traffic injuries, and leukemia were three leading causes of death. Multisectoral efforts should be initiated to reduce premature death among students, especially those due to injuries.

Deaths in older children (5–9 years) and younger adolescents (10–14 years) have received little attention. The neglect of these groups are presumably associated with the fact that this is the age range during which

human mortality risks reach their minimum but are not insubstantial (1). It was reported that there were approximately 1.23 million deaths among individuals aged 5–19 years worldwide in 2017 (2). Primary and secondary school students aged 6–18 years mostly consist of older children and adolescents. Estimating their mortality is of great necessity to identify health priorities for reducing premature death.

This report used deaths in 2018 as collected by CDRS. Detailed descriptions of CDRS have been reported elsewhere (3). After primary quality control, 512 out of 605 were identified as eligible disease surveillance points, with 1,822 thousand reported deaths, covering 272 million population. Individuals aged 6–18 years were extracted and stratified by sex, age group, occupation, area, and region. National data of total population and current students in primary and secondary schools were obtained from National Bureau of Statistics (4). International Classification of Disease, 10th revision (ICD-10) was used to re-classify cause-of-death.

Crude mortality rate was calculated by using deaths and the respective populations. In consideration of CDRS under-reporting, mortality rate was adjusted for under-reporting through following formula: mortality rate=crude mortality rate/(1-under-reporting rate) (5). The Sixth National Population Census in 2010 was used for age-standardized mortality rate (ASMR) estimation (4). Estimated death count nationwide was generated through scaled-up aggregation data in each stratum acquired. Years of life lost (YLLs) is a measure of premature death calculated as the sum of each death multiplied by the standard life expectancy at each age (2). Software R (version 3.6.2, The R foundation for Statistical Computing) was applied for statistical analysis.

In 2018, a total of 45,385 deaths among individuals aged 6–18 years nationwide were estimated with 28,519 students and 16,867 non-students. The ASMR for those aged 6–18 years was 23.55 per 100,000 with students estimated at 17.66 and non-students at 54.13 per 100,000. Non-students showed higher ASMR

compared with its counterpart in all age groups, especially for those aged 6–12 years (136.25 *vs.* 14.12 per 100,000). Deaths occurring in those aged 6–18 years were estimated to occupy 0.47% of total population and calculated YLLs constituted 2.19% of total YLLs in all age. (Table 1)

Male students showed higher ASMR than female students (21.50 *vs.* 13.30 per 100,000). Rural areas showed a higher value than urban areas (19.72 *vs.* 14.28 per 100,000). Western regions showed the highest ASMR (22.07 per 100,000), followed by central regions (18.23 per 100,000) and eastern regions (14.07 per 100,000). Injury (52.87%) and non-communicable diseases (NCDs) (38.71%) contributed over 90% deaths to overall mortality in students. Drowning (2.93 per 100,000, 16.59%), road traffic injuries (2.74 per 100,000, 15.51%), and leukemia (1.37 per 100,000, 7.75%) were the three leading causes of death. (Table 2)

The ASMR of students varied as age changed. The ASMR of drowning reached its peak in primary school students with downward trends in junior high school students. Following a similar trend as age increased, suicide mortality came reached its peak in secondary school students aged 15 years and older. For other conditions, a comparatively stable trend was observed in adjacent ages such as road traffic injuries, falls, poisonings, and most NCDs. Female students showed more frequent mortality fluctuations in comparison with male students, especially for malignant tumor after 13 years. (Figure 1)

DISCUSSION

This study presents an up-to-date overview of mortality among primary and secondary school students aged 6–18 years in China in 2018. Differences in ASMR was noticed among students and non-students, especially for those aged 6–12 years for primary school. The health gap was large in relation to dropout, which was presumed to be induced by socio-economic structural transformation, household unemployment and poverty, as well as stereotype conventions. For disadvantaged groups who encountered obstacles for education and suffered from inferior health conditions such as migrant, left-behind, disabled, and impoverished children, providing economic support, promoting universal health coverage, and popularizing compulsory education could be effectively reduce premature death in non-students (6). The results concerning the age-specific

percentage of deaths and YLLs were consistent with the Global Burden of Disease 2017 (GBD2017) (2). Given the five times higher age-specific YLLs percentage compared to deaths, the value of health losses occurring in the youth were inferred to be considerably higher than that of the elderly, which reminds of the urgent need to prevent premature death in order to minimize social labor force damage.

Obvious mortality disparities in population distribution and geographical patterns could be found among primary and secondary school students aged 6–18 years. Sex differences could be primarily driven by variations in exposure prevalence to cause-specific risk factors, particularly for injury (6). It was reported that intrinsic characteristics with aggressive and adventurous spirits in boys made them inclined to actively encounter hazards, and in the absence of cognition and self-control, they tended to experience more injuries than female students (7). Suicide was a typical condition that increased considerably with age, which brings attention to issues of mental health in senior high school students (6). A higher ASMR in remote areas was consistent with previous studies (8). Reasons could be partially attributed to economic involvement, healthcare coverage, infrastructure assurance, effective guardian supervision, and education popularization in developed areas and regions. For example, students in developed areas were inclined to receive multi-source health protection and security training, including courses taught by professionals from health, transportation, and fire-fighting departments (8).

Injury was the leading cause of death among students. Premature death as a result of unintentional injuries were discussed at length in most studies, with the majority deemed preventable (6). A “5E comprehensive strategy”, which covers education, environmental modification, engineering, enforcement, and evaluation, was introduced for injury control and prevention worldwide to effectively reduce injury incidence and mortality (6). NCDs also accounted for a large proportion of student mortality, most of which might be explained by biological factors, inferior living environments, and unhealthy lifestyle; for instance, leukemia was regarded as being related to indoor environment pollution caused by household decorations (7). Although premature death caused by communicable diseases seemed to not be so substantial, guardians should still take precautions as existing risk factors continued to rise and potential new risk factors emerge. Planned immunizations would be an effective

TABLE 1. Age-standardized mortality rate, estimated deaths in students and non-students aged 6–18 years in China, 2018.

Age (Years)	Education level [†]	Age-standardized mortality rate (per 100,000)				Total population (10,000)				Estimated deaths				Age-specific deaths/All-age deaths (%)		Age-specific YLLs/All-age YLLs (%)		
		Total		Non-students		Total		Non-students		Total		Students		Non-students		Total	Male	Female
		Students	Non-students	Students	Non-students	Students	Non-students	Students	Non-students	Students	Non-students	Students	Non-students					
6–12	Primary school	19.61	14.12	136.25	10,568.86	10,093.70	475.16	20,730	14,256	6,474	0.21	1.02						
13–15	Junior high school	27.03	22.46	55.51	4,386.04	3,780.05	605.99	11,855	8,491	3,364	0.12	0.54						
16–18	Senior high school	28.88	25.78	32.35	4,314.39	2,279.49	2,034.91	12,458	5,876	6,583	0.14	0.63						
	Total	23.55	17.66	54.13	19,269.30	16,153.24	3,116.06	45,385	28,519	16,867	0.47	2.19						

Abbreviation: YLLs=Years of life lost.

[†] Education level for school students aged 6–18 years only.

TABLE 2. The age-standardized mortality rate (per 100,000) of primary and secondary school students aged 6–18 years in China, 2018.

Item	Sex						Urban			Rural			Eastern			Central			Western					
	Total		Male		Female		Total		Male		Female		Total		Male		Female		Total		Male		Female	
	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female
All-cause	17.66	21.50	13.30	14.28	16.91	11.34	19.72	24.27	14.51	14.07	16.76	11.00	18.23	22.62	13.16	22.07	27.07	16.59						
Injury	9.33	12.14	6.15	7.29	9.30	5.04	10.58	13.86	6.83	6.60	8.61	4.30	9.72	12.85	6.11	12.75	16.43	8.71						
Drowning	2.93	4.34	1.33	1.92	2.93	0.78	3.55	5.20	1.67	1.86	2.76	0.82	3.20	4.56	1.63	4.16	6.37	1.73						
Road traffic injuries	2.74	3.46	1.93	2.19	2.60	1.74	3.07	3.98	2.04	2.03	2.48	1.52	2.93	3.80	1.94	3.53	4.49	2.47						
Suicide	1.10	1.31	0.86	1.11	1.24	0.96	1.09	1.35	0.79	0.86	1.23	0.43	1.05	1.21	0.86	1.49	1.54	1.44						
Falls	0.91	1.10	0.69	0.83	1.07	0.56	0.95	1.12	0.76	0.62	0.73	0.50	1.08	1.43	0.67	1.13	1.28	0.96						
Poisoning	0.50	0.53	0.47	0.30	0.29	0.30	0.62	0.68	0.56	0.37	0.36	0.37	0.43	0.48	0.37	0.76	0.82	0.70						
Other injuries	1.15	1.40	0.88	0.95	1.18	0.69	1.28	1.53	0.99	0.87	1.05	0.67	1.03	1.36	0.65	1.68	1.94	1.40						
Non-communicable diseases	6.83	7.79	5.75	5.72	6.28	5.09	7.52	8.70	6.16	6.21	6.91	5.42	7.01	8.12	5.72	7.53	8.70	6.24						
Leukemia	1.37	1.61	1.11	1.17	1.31	1.00	1.50	1.78	1.17	1.39	1.60	1.14	1.47	1.73	1.16	1.25	1.48	1.01						
Cardiovascular diseases	1.05	1.24	0.83	0.94	1.00	0.88	1.11	1.38	0.81	1.01	1.24	0.74	1.03	1.10	0.95	1.13	1.38	0.85						
Congenital abnormality	0.82	0.90	0.74	0.81	0.77	0.86	0.83	0.98	0.66	0.57	0.63	0.50	1.03	1.19	0.84	0.97	0.98	0.96						
Nervous systems diseases	0.76	0.87	0.63	0.48	0.59	0.36	0.93	1.04	0.79	0.62	0.73	0.50	0.77	0.93	0.58	0.94	1.02	0.85						
Respiratory diseases	0.50	0.50	0.50	0.33	0.36	0.29	0.61	0.59	0.64	0.39	0.43	0.35	0.42	0.35	0.49	0.74	0.76	0.72						
Immune system diseases	0.26	0.25	0.27	0.19	0.15	0.24	0.30	0.30	0.29	0.25	0.15	0.35	0.28	0.30	0.26	0.25	0.32	0.18						
Digestive system diseases	0.17	0.15	0.19	0.10	0.09	0.10	0.22	0.19	0.25	0.13	0.08	0.19	0.10	0.15	0.04	0.30	0.26	0.35						
Other malignant tumor	1.90	2.27	1.48	1.71	2.01	1.37	2.02	2.43	1.55	1.86	2.04	1.66	1.92	2.36	1.40	1.94	2.52	1.31						
Communicable diseases	0.31	0.32	0.30	0.20	0.21	0.19	0.38	0.39	0.37	0.22	0.20	0.24	0.29	0.35	0.22	0.47	0.48	0.46						
Infectious & parasitic diseases	0.31	0.32	0.30	0.20	0.21	0.19	0.38	0.39	0.37	0.22	0.20	0.24	0.29	0.35	0.22	0.47	0.48	0.46						
Others	1.18	1.24	1.11	1.07	1.12	1.02	1.24	1.32	1.16	1.04	1.05	1.04	1.22	1.30	1.12	1.33	1.46	1.18						

approaches in preventing many communicable diseases (9).

During past years, several programs for protecting

and promoting children’s health have been launched and yielded profound benefits. Related clauses were embodied in legislations and regulations, like “The




FIGURE 1. Sex and age differences in mortality in primary and secondary school students aged 6–18 years in China by cause-of-death, 2018.

Law of Education”, “The Law of Protection of Juveniles”, “The Law of Road Traffic Safety”, “China Children’s Development Outlines”. With multi-sectional cooperation at its burgeoning phase, measures such as health literature advocacy, regular lifestyle development, and mental health establishment were provided for students in some regions (6). Nevertheless, enforcement was a demanding bottleneck. Due to the complexity of multi-sector enabler management, such as health, education, security, transportation, and urban construction, no specific department or agency was assigned to centralize and coordinate the efforts of societal partners to work together (10). Therefore, the government should maintain a leading role to present strategies that aim to eliminate modifiable risk factors for early intervention, such as providing financial and technical support, implementing enforcement for legislation and promoting community engagement, especially to narrow the gaps between developed and remote areas and considerable differences between subgroups at different ages. These initiatives could, in return, inform individual-level change, decrease the demand on tertiary services, and operate effectively to reduce premature death among students (9).

The findings of this study were subject to some limitations. Ascertainment bias in cause-of-death diagnosis remained the greatest concern, which required correction for redistribution algorithms for implausible diagnostic codes (2).

Understanding mortality in primary and secondary school students remained as an essential concern in effectively reducing premature death. This study helped to identify heterogeneity in mortality patterns and is of great value for tailoring priorities to promote students’ health in China.

Corresponding author: Maigeng Zhou, zhoumaigeng@ncncd.chinacdc.cn.

¹ National Center for Chronic and Noncommunicable Disease Control

and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China.

Submitted: March 04, 2020; Accepted: March 16, 2020

REFERENCES

1. Fadel SA, Boschi-Pinto C, Yu SC, Reynales-Shigematsu LM, Menon GR, Newcombe L, et al. Trends in cause-specific mortality among children aged 5–14 years from 2005 to 2016 in India, China, Brazil, and Mexico: an analysis of nationally representative mortality studies. *Lancet* 2019;393(10176):1119 – 27. [http://dx.doi.org/10.1016/S0140-6736\(19\)30220-X](http://dx.doi.org/10.1016/S0140-6736(19)30220-X).
2. GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018;392(10159):1736 – 88. [http://dx.doi.org/10.1016/S0140-6736\(18\)32203-7](http://dx.doi.org/10.1016/S0140-6736(18)32203-7).
3. Liu SW, Wu XL, Lopez AD, Wang LJ, Cai Y, Page A, et al. An integrated national mortality surveillance system for death registration and mortality surveillance, China. *Bull World Health Organ* 2016; 94(1):46 – 57. <http://dx.doi.org/10.2471/BLT.15.153148>.
4. National Bureau of Statistics of China. National data of total population and number of students enrollment by level and type of school. <http://data.stats.gov.cn/>. [2020-02-29]. (In Chinese).
5. Guo K, Yin P, Wang LJ, Ji YB, Li QF, Bishai D, et al. Propensity score weighting for addressing under-reporting in mortality surveillance: a proof-of-concept study using the nationally representative mortality data in China. *Popul Health Metr* 2015;13:16. <http://dx.doi.org/10.1186/s12963-015-0051-3>.
6. Duan LL, Ye PP, Haagsma J, Jin Y, Wang Y, Er YL, et al. The burden of injury in China, 1990–2017: findings from the Global Burden of Disease Study 2017. *Lancet Public Health* 2019;4(9):e449 – 61. [http://dx.doi.org/10.1016/S2468-2667\(19\)30125-2](http://dx.doi.org/10.1016/S2468-2667(19)30125-2).
7. Xie YF, Luo YN, Luo D, Ma X, Liu Q, Wu YP. Analysis of cause-of-death and years of life lost among children and adolescents in Cixi, 2007–2016. *Chin J Prev Control Chron Dis* 2018;26(6):477 – 9. <http://dx.doi.org/10.16386/j.cjpcd.issn.1004-6194.2018.06.022>. (In Chinese).
8. Xu XD. Analysis of temporal trend of primary and middle school students in urban/rural areas in Dalian, 2004–2013. *Chin J Public Health Manag* 2014;30(6):925 – 7. <http://dx.doi.org/10.19568/j.cnki.23-1318.2014.06.061>. (In Chinese).
9. Kuruvilla S, Bustreo F, Kuo T, Mishra C, Taylor K, Fogstad H, et al. The Global strategy for women’s, children’s and adolescents’ health (2016–2030): a roadmap based on evidence and country experience. *Bull World Health Organ* 2016;94(5): 398-400. <http://dx.doi.org/10.2471/BLT.16.170431>.
10. Ning PS, Schwebel DC, Hu GQ. Healthy China 2030: a missed opportunity for injury control. *Inj Prev* 2017;23(6):363. <http://dx.doi.org/10.1136/injuryprev-2017-042314>.