

Minireview

Yeast Polo-like kinase substrates are nailed with the right tools

Vincent Archambault and David M Glover

Address: Department of Genetics, University of Cambridge, Downing Street, Cambridge CB2 3EH, UK.

Correspondence: Vincent Archambault. Email: va228@cam.ac.uk

Published: 30 January 2008

Genome Biology 2008, **9**:203 (doi:10.1186/gb-2008-9-1-203)

The electronic version of this article is the complete one and can be found online at <http://genomebiology.com/2008/9/1/203>

© 2008 BioMed Central Ltd

Abstract

A platform has been built combining chemical genetics and bioinformatics to screen the proteome for physiological substrates of the Polo-like kinase in budding yeast. A novel role for this kinase in regulating the mitotic spindle is revealed.

Protein kinases are important regulators of almost every cellular function in eukaryotes. They catalyze the transfer of a phosphate group onto serine, threonine or tyrosine amino acid residues, resulting in a change of activity of the substrate protein. The identification of the physiological substrates of a kinase constitutes an important but intimidating challenge for many biologists. A recent paper by Jennifer Snead and colleagues in *Chemistry and Biology* [1] reports the development of a multidisciplinary approach to the identification of protein kinase substrates that has led to the discovery of a novel mitotic role for the Polo-like kinase in the budding yeast *Saccharomyces cerevisiae* (where it is called Cdc5), implicating novel molecular substrates.

Progression and regulation of the cell-division cycle relies strongly on kinases, including the cyclin-dependent kinases (Cdks), Aurora kinases and Polo-like kinases [2]. Each of these families has been conserved in eukaryotes, from yeasts to humans. Originally discovered in *Drosophila* [3], Polo-like kinases are known to promote and coordinate several events of mitosis and cytokinesis in a multitude of models [4,5]. Among their most crucial functions, Polo-like kinases promote centrosome maturation and separation (in animal cells), sister-chromatid separation in prophase, activation of the equally important Cdks, establishment of a bipolar spindle and cytokinesis. To facilitate their targeting to the appropriate subcellular locations, Polo-like kinases possess a carboxy-terminal Polo-box domain (PBD), which enables these enzymes to dock to proteins that have been pre-phosphorylated (primed) at a specific motif by Polo-like

kinases themselves or by other kinases. Humans have four Polo-like kinases (Plk1-4). Plk1-3 are most closely related to *Drosophila* Polo and these have partially diverging functions, with Plk1 fulfilling most of the functions in cell division. Plk4 has a separate function in centriole duplication in humans and flies. The yeasts have single Plks: Cdc5 in budding yeast and Plo1 in fission yeast. Plk1 is often deregulated in many cancers, which has stimulated the development of several chemical inhibitors with therapeutic potential [6-8]. Nevertheless, surprisingly few phosphorylation substrates of the Polo kinases are known, and it is suspected that many more remain to be identified before we have a full picture of how these kinases impact on cell division at the molecular level.

Existing approaches for identifying kinase substrates

Searching the protein sequence complement for particular sequences can often aid the identification of kinase substrates. Kinases differ in substrate specificity, preferring different sequence motifs around the residue to be phosphorylated. For some of the better-characterized kinases, their substrate-motif preference is known, making it possible to predict whether a given protein is likely to be phosphorylated. The primary sequence of a potential sequence does not, however, allow prediction with high confidence of whether a particular segment of the protein will be accessible to phosphorylation by a kinase (for example, it may be buried in the protein core or sterically hindered by an interacting protein). Moreover, kinase and substrates need to meet in space and time in the cell for the

reaction to occur. For these reasons, genomic analysis alone has a very limited capacity to identify physiological substrates of protein kinases.

Various strategies have been used to attempt the systematic identification of kinase substrates experimentally. *In vitro* screens exploiting the change in electrophoretic mobility of proteins after phosphorylation have been carried out, starting with pools of radiolabeled proteins transcribed and translated *in vitro*, with some success [9,10]. Substrates identified in such *in vitro* reactions must, however, then be validated *in vivo*. Another approach is to purify the kinase by affinity-based methods and identify co-purifying proteins using mass spectrometry [11]. Proteins co-purifying with a kinase are often physiological phosphorylation substrates, but this strategy is most effective for kinase-substrate complexes of relatively high binding affinity and abundance. A technique that allows the identification of the physiological substrates of a kinase *in vivo* and with no bias for affinity or abundance is needed.

In a previous study from the laboratories of Kevan Shokat and David Morgan, Ubersax *et al.* [12] reported a screen using chemical biology to identify substrates of Cdk1, a master cell-cycle regulator. The authors designed a modified Cdk1 (Cdk1-as, analog sensitive) with an enlarged ATP-binding site, capable of accommodating and using a bulkier radiolabeled ATP analog in phosphorylation reactions. The ATP analog was designed to be unable to bind any cellular, unmodified kinases. Because the ATP analog was cell-impermeable, however, Ubersax *et al.* [12] could not assay for Cdk1 substrates *in vivo* and instead carried out the reactions in cell extracts (as close to *in vivo* as possible). The reactions combined recombinant cyclin B-Cdk1-as and multiple cell extracts from yeast strains overexpressing single epitope-tagged candidate substrates that were then purified and assayed for their level of radioisotope incorporation. The study tested a selected group of candidate proteins chosen for their known involvement in the cell cycle and for the presence of Cdk1 phosphorylation motifs, as well as a random group. The result was the identification of some 200 potential substrates, and constitutes a milestone in cell-cycle research. However, their strategy fell short of directly identifying *in vivo* substrates.

A new strategy that goes *in vivo*

Now, they have done it. The same groups now report a strategy [1] that allows the systematic identification of *in vivo* kinase substrates. Again, the trick is to combine elements of chemical biology and substrate prediction using bioinformatics. As in the previous work, Snead *et al.* [1] began by generating a modified but functional Cdc5 (Polo) kinase with an enlarged binding site (L158G-substituted) capable of accepting a cell-permeable inhibitor that has a low affinity for the unmodified, wild-type Cdc5 (Figure 1a) and is not

predicted to be accommodated in the binding sites of any other kinases of the genome. The resulting kinase, Cdc5-as1, is still named 'as' for 'analog-sensitive' [13]. This general approach has been used previously to selectively inhibit several kinases [14]. However, the particularities of the Cdc5 ATP-binding site make it resistant to the usual pyrazolopyrimidine (PP1) analog inhibitors and forced the use of a pyrrolopyrimidine inhibitor containing a chloromethylketone (referred to as CMK).

With the new kinase-inhibitor pair in hand, Snead *et al.* [1] constructed a strain of yeast in which the endogenous *CDC5* gene was replaced with *cdc5-as1* at the natural locus, enabling the *in vivo* inhibition of Cdc5-as1, the only source of Cdc5 in the cell. This was achieved without any significant risk of simultaneously inhibiting other cellular kinases, which is usually a problem when working with inhibitors developed to target natural kinases.

Treatment of *cdc5-as1* cells with CMK led to inhibition of proliferation (Figure 1b). Cytological examination revealed that cells arrested at or after anaphase. This on its own was not surprising, as Cdc5 is known to be required for exit from mitosis and cytokinesis [15]. But on close examination, an increased percentage of cells had an elongated spindle that was completely enclosed in the mother cell, instead of normally spanning both the mother cell and the bud (although this defect was eventually corrected). Moreover, the early, short spindle was often misaligned and microtubules were often misoriented or sometimes detached from the spindle pole bodies (SPBs). Therefore, Cdc5 is required for normal positioning and function of the mitotic spindle. Inhibition of Cdc5 also led to premature sister-chromatid separation, as expected. Altogether, these experiments revealed both known and new roles for Cdc5 in the cell cycle. This novel role for Cdc5 at the spindle level was not identified in previous studies using temperature-sensitive (*ts*) alleles of *CDC5*; this may be because such alleles often retain some activity even at the restrictive temperature or because the increase in temperature required to inactivate a *ts* allele can alter cell-cycle progression.

Snead *et al.* [1] then set out to screen for substrates of Cdc5 *in silico* and *in vivo*. Using a sequence-profile-scanning algorithm [16,17], they searched all yeast predicted proteins (over 6,000 proteins) for Plk phosphorylation motifs and for PBD-binding motifs (defined using published information available for human and yeast Plks). As an output, each protein was assigned a 'Cdc5 substrate likelihood score'. From among the highest-scoring proteins, functional criteria (for example, known involvement in mitosis or cell cycle) were used to choose a list of 192 candidates to be tested (Figure 1c).

The experimental part of the screen made use of a library of strains in which genes are fused at their natural loci with a tag that can be used for tandem affinity purification (TAP),




Figure 1

Strategy used by Snead *et al.* [1] to identify substrates of yeast Polo-like kinase (Cdc5). **(a)** A modified Cdc5 (Cdc5-as1) is generated to be selectively inhibited by a small-molecule inhibitor (CMK) that does not affect wild-type Cdc5. K, kinase domain; PBD, Polo-box domain. **(b)** Treatment of *cdc5Δ cdc5-as1* cells with CMK reveals the cellular phenotypes associated with Cdc5 inhibition. **(c)** A sequence-scanning algorithm identifies proteins containing potential Plk phosphorylation motifs (blue), potential PBD-binding motifs (red) and satisfying functional criteria in database annotations (yellow). **(d)** Individual strains expressing candidate substrates (P1, P2, P3, and so on) from their normal loci in fusion with the TAP tag are screened for electrophoretic mobility shifts of the fusion protein between a G1 arrest (Cdc5 inactive) and an M-phase arrest (Cdc5 and other kinases active). **(e)** For fusion proteins showing a shift between G1 and M, strains combining the TAP-tagged gene with the *cdc5Δ cdc5-as1* allele are generated. Those are then screened for a CMK-dependent (Cdc5-dependent) mobility shift in M-phase-arrested cells. Hits from that final step can be considered to be physiological Cdc5 substrates. The cellular phenotypes observed following Cdc5 inhibition in (b) may be hypothesized to result from a failure of phosphorylation of one or more of the Cdc5 substrates identified.

providing easy detection in western blotting. Tagged proteins were then assayed for their phosphorylation state as detected by a shift in electrophoretic mobility. A first screen retained only those proteins from the candidate list that changed in mobility (shifted) between G1 (Cdc5 inactive) and mitosis (Cdc5 and other kinases active) (Figure 1d). Hits from that stage (74) were finally assayed for a shift that was dependent on treatment with CMK in mitotically arrested *cdc5-as1* cells (Figure 1e). Only five proteins passed this final selection step; two of them are known Cdc5 substrates, while three are novel. One of the novel substrates is *Spc72*, a SPB component with known roles in microtubule nucleation and spindle positioning [18-21]. Since it was known to interact with Cdc5 [22,23], *Spc72* was an obvious candidate for a molecular target of Cdc5 in the regulation of the mitotic

spindle. Moreover, Cdc5 is known to localize to the SPB. Snead *et al.* [1] showed that Cdc5 requires its PBD to interact with *Spc72* and can phosphorylate *Spc72 in vitro*. Finally, they also assayed the binding of recombinant Cdc5 to a set of 90 TAP-tagged proteins annotated (or reported) as SPB-localized in databases. Several of these proteins associated with Cdc5, and *Spc72* was among the most efficient interactors, further suggesting a direct interaction.

A significant new advance

Further experiments will be required to clarify how Cdc5 regulates *Spc72* and how this affects SPB and spindle function. It is likely that disruption of the Cdc5-*Spc72* pathway is at least partly responsible for the cellular phenotypes obtained

when Cdc5 is inhibited *in vivo*. In animal cells, Plk1 is known to promote centrosome maturation and microtubule nucleation activity by various mechanisms [4,24]. Despite marked differences between budding yeasts and animal cells in how the spindle is assembled and regulated, it is becoming clear in the light of the paper by Snead *et al.* that the yeast Polo-like kinase Cdc5 is a major regulator of the mitotic spindle.

Because many phosphoproteins do not produce a marked shift in electrophoretic mobility when phosphorylated at a given site, several Cdc5 substrates are likely to have been missed in this screen. This may be even more problematic for Cdc5 substrates that are also phosphorylated by other mitotic kinases. Nonetheless, the experimental approach presented by Snead *et al.* provides a powerful means of systematically identifying physiological substrates of a kinase in its natural environment - without overexpressing the kinase or the candidate substrates to be screened. This platform should be readily amenable to similar screens for substrates of any kinase of choice in *S. cerevisiae* and therefore constitutes a powerful handle on many signaling pathways in cell biology. The technique is not yet easily transferable to other organisms where exact gene replacement and genomic tagging at the natural loci are, at present, impracticable or challenging, including *Drosophila*, *Caenorhabditis elegans* or mammalian cells. Nonetheless, a recent paper reported the use of a Plk1-as allele (and a specific inhibitor) to analyze Plk1 functions in cytokinesis in human cells where endogenous Plk1 has been knocked out by homologous recombination [25]. The viral expression of Plk1-as in these experiments may not, however, follow the normal levels or cell-cycle profile of endogenous Plk1 expression. In mammalian cells a screen for substrates such as the one presented by Snead *et al.* would almost certainly require the tagged candidates to be artificially expressed. Again, the little budding yeast is first to stick its neck out and one has to hope that technological advances will soon allow other organisms to follow its path.

Acknowledgements

VA is supported by a Long-Term Fellowship from the Human Frontier Science Program.

References

- Snead JL, Sullivan M, Lowery DM, Cohen MS, Zhang C, Randle DH, Taunton J, Yaffe MB, Morgan DO, Shokat KM: **A coupled chemical-genetic and bioinformatic approach to Polo-like kinase pathway exploration.** *Chem Biol* 2007, **14**:1261-1272.
- Morgan DO: *The Cell Cycle: Principles of Control*. London: New Science Press; 2007.
- Sunkel CE, Glover DM: **polo, a mitotic mutant of Drosophila displaying abnormal spindle poles.** *J Cell Sci* 1988, **89**:25-38.
- van de Weerd BC, Medema RH: **Polo-like kinases: a team in control of the division.** *Cell Cycle* 2006, **5**:853-864.
- Barr FA, Silje HH, Nigg EA: **Polo-like kinases and the orchestration of cell division.** *Nat Rev Mol Cell Biol* 2004, **5**:429-440.
- Gumireddy K, Reddy MV, Cosenza SC, Boominathan R, Baker SJ, Papathi N, Jiang J, Holland J, Reddy EP: **ON01910, a non-ATP-competitive small molecule inhibitor of Plk1, is a potent anticancer agent.** *Cancer Cell* 2005, **7**:275-286.
- McInnes C, Mazumdar A, Mezna M, Meades C, Midgley C, Scaerou F, Carpenter L, Mackenzie M, Taylor P, Walkinshaw M, Fischer PM, Glover D: **Inhibitors of Polo-like kinase reveal roles in spindle-pole maintenance.** *Nat Chem Biol* 2006, **2**:608-617.
- Lenart P, Petronczki M, Steegmaier M, Di Fiore B, Lipp JJ, Hoffmann M, Rettig WJ, Kraut N, Peters JM: **The small-molecule inhibitor BI 2536 reveals novel insights into mitotic roles of polo-like kinase I.** *Curr Biol* 2007, **17**:304-315.
- Gao ZH, Metherall J, Virshup DM: **Identification of casein kinase I substrates by in vitro expression cloning screening.** *Biochem Biophys Res Commun* 2000, **268**:562-566.
- Lee LA, Lee E, Anderson MA, Vardy L, Tahinci E, Ali SM, Kashevsky H, Benasutti M, Kirschner MW, Orr-Weaver TL: **Drosophila genome-scale screen for PAN GU kinase substrates identifies Mat89Bb as a cell cycle regulator.** *Dev Cell* 2005, **8**:435-442.
- Archambault V, Chang EJ, Drapkin BJ, Cross FR, Chait BT, Rout MP: **Targeted proteomic study of the cyclin-Cdk module.** *Mol Cell* 2004, **14**:699-711.
- Ubersax JA, Woodbury EL, Quang PN, Paraz M, Blethrow JD, Shah K, Shokat KM, Morgan DO: **Targets of the cyclin-dependent kinase Cdk1.** *Nature* 2003, **425**:859-864.
- Zhang C, Kenski DM, Paulson JL, Boushtien A, Sessa G, Cross JV, Templeton DJ, Shokat KM: **A second-site suppressor strategy for chemical genetic analysis of diverse protein kinases.** *Nat Methods* 2005, **2**:435-441.
- Bishop AC, Ubersax JA, Petsch DT, Matheos DP, Gray NS, Blethrow J, Shimizu E, Tsien JZ, Schultz PG, Rose MD, Wood JL, Morgan DO, Shokat KM: **A chemical switch for inhibitor-sensitive alleles of any protein kinase.** *Nature* 2000, **407**:395-401.
- Lee KS, Park JE, Asano S, Park CJ: **Yeast polo-like kinases: functionally conserved multitask mitotic regulators.** *Oncogene* 2005, **24**:217-229.
- Obenaus JC, Cantley LC, Yaffe MB: **Scansite 2.0: Proteome-wide prediction of cell signaling interactions using short sequence motifs.** *Nucleic Acids Res* 2003, **31**:3635-3641.
- Yaffe MB, Leparc GG, Lai J, Obata T, Volinia S, Cantley LC: **A motif-based profile scanning approach for genome-wide prediction of signaling pathways.** *Nat Biotechnol* 2001, **19**:348-353.
- Hoepfner D, Schaerer F, Brachat A, Wach A, Philippsen P: **Reorientation of mispositioned spindles in short astral microtubule mutant spc72Delta is dependent on spindle pole body outer plaque and Kar3 motor protein.** *Mol Biol Cell* 2002, **13**:1366-1380.
- Knop M, Schiebel E: **Receptors determine the cellular localization of a gamma-tubulin complex and thereby the site of microtubule formation.** *EMBO J* 1998, **17**:3952-3967.
- Soues S, Adams IR: **SPC72: a spindle pole component required for spindle orientation in the yeast Saccharomyces cerevisiae.** *J Cell Sci* 1998, **111**:2809-2818.
- Chen XP, Yin H, Huffaker TC: **The yeast spindle pole body component Spc72p interacts with Stu2p and is required for proper microtubule assembly.** *J Cell Biol* 1998, **141**:1169-1179.
- Ho Y, Gruhler A, Heilbut A, Bader GD, Moore L, Adams SL, Millar A, Taylor P, Bennett K, Boutillier K, Yang L, Wolting C, Donaldson I, Schandorff S, Shewnarane J, Vo M, Taggart J, Goudreaux M, Muskat B, Alfarano C, Dewar D, Lin Z, Michalickova K, Willems AR, Sassi H, Nielsen PA, Rasmussen KJ, Andersen JR, Johansen LE, Hansen LH, et al.: **Systematic identification of protein complexes in Saccharomyces cerevisiae by mass spectrometry.** *Nature* 2002, **415**:180-183.
- Park JE, Park CJ, Sakchaisri K, Karpova T, Asano S, McNally J, Sunwoo Y, Leem SH, Lee KS: **Novel functional dissection of the localization-specific roles of budding yeast polo kinase Cdc5p.** *Mol Cell Biol* 2004, **24**:9873-9886.
- Glover DM: **Polo kinase and progression through M phase in Drosophila: a perspective from the spindle poles.** *Oncogene* 2005, **24**:230-237.
- Burkard ME, Randall CL, Laroche S, Zhang C, Shokat KM, Fisher RP, Jallepalli PV: **Chemical genetics reveals the requirement for Polo-like kinase I activity in positioning RhoA and triggering cytokinesis in human cells.** *Proc Natl Acad Sci USA* 2007, **104**:4383-4388.